



HAL
open science

Volume-constrained minimizers for the prescribed curvature problem in periodic media

Michael Goldman, Matteo Novaga

► **To cite this version:**

Michael Goldman, Matteo Novaga. Volume-constrained minimizers for the prescribed curvature problem in periodic media. 2012. hal-00580074v3

HAL Id: hal-00580074

<https://hal.science/hal-00580074v3>

Preprint submitted on 27 Nov 2012 (v3), last revised 28 Nov 2012 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Volume-constrained minimizers for the prescribed curvature problem in periodic media

M. Goldman ^{*} M. Novaga [†]

Abstract

We establish existence of compact minimizers of the prescribed mean curvature problem with volume constraint in periodic media. As a consequence, we construct compact approximate solutions to the prescribed mean curvature equation. We also show convergence after rescaling of the volume-constrained minimizers towards a suitable Wulff Shape, when the volume tends to infinity.

1 Introduction

In recent years, a lot of attention has been drawn towards the problem of constructing surfaces with prescribed mean curvature. More precisely, given an assigned function $g : \mathbb{R}^d \rightarrow \mathbb{R}$, the problem is finding a hypersurface having mean curvature κ satisfying

$$\kappa = g. \tag{1}$$

To our knowledge, this problem was first posed by S.T. Yau in [30], under the additional constraint of the hypersurface being diffeomorphic to a sphere, and a solution was provided in [27, 16] when the function g satisfies suitable decay conditions at infinity, namely that it decays faster than the mean curvature of concentric spheres. Another approach was presented in [5, 15], by means of conformal parametrizations and a clever use of the mountain pass lemma. A serious limitation of this method is the impossibility to extend it to dimension higher than three, due to the lack of a good equivalent of a conformal parametrization.

Motivated by some homogenization problems in front propagation [22], in this paper we look for solutions to (1) without any topological constraint but with a periodic function

^{*}CMAP, CNRS UMR 7641, Ecole Polytechnique, 91128 Palaiseau, France, email: goldman@cmap.polytechnique.fr

[†]Dipartimento di Matematica, Università di Padova, via Trieste 63, 35121 Padova, Italy, email: novaga@math.unipd.it

g , so that in particular, it does not decay to zero at infinity. A natural idea is to look for critical points of the prescribed curvature functional

$$F(E) = P(E) - \int_E g \, dx,$$

as it is well-known that such critical points solve (1), whenever they are smooth [14]. Observe that, in general, it is not possible to construct solutions of (1) by a direct minimization of the functional F , because such minimizers may not exist or be empty.

The first result in this setting was obtained by Caffarelli and de la Llave in [7] (see also [9]) where the authors construct planelike solutions of (1) under the assumption that g is small and has zero average, by minimizing F among sets with boundary contained in a given strip, and then show that the constraint does not affect the curvature of the solution.

Here we are interested instead in compact solutions of (1). This problem seems difficult in this generality and only some preliminary results, in the two-dimensional case, are presently available [17]. However, the following perturbative result has been proved in [22]: given a periodic function g with zero average and small L^∞ -norm and ε arbitrarily small, there exists a compact solution of

$$\kappa = g_\varepsilon$$

where $\|g_\varepsilon - g\|_{L^1} \leq \varepsilon$. Since the L^1 -norm does not seem very well suited for this problem, a natural question raised in [22] was whether the same result holds when the L^1 -norm is replaced by the L^∞ -norm.

In this paper we answer this question. More precisely, we prove the following result (see Theorem 4.4): let g be a periodic Hölder continuous function with zero average on the unit cell $Q = [0, 1]^d$ and such that

$$\int_E g \, dx \leq (1 - \Lambda)P(E, Q) \quad \forall E \subset Q \quad (2)$$

for some $\Lambda > 0$, where $P(E, Q)$ is the relative perimeter of E in Q . Then for every $\varepsilon > 0$ there exist $0 < \varepsilon' < \varepsilon$ and a compact solution of

$$\kappa = g + \varepsilon'. \quad (3)$$

We observe that (2) is the same assumption made in [9] in order to prove existence of planelike minimizers. This condition is for instance verified if $\|g\|_{L^d(Q)}$ is smaller than the isoperimetric constant of Q , and allows g to take large negative values.

We construct approximate solutions of (3) as volume constrained minimizers of F for big volumes. This motivates the study of the isovolumetric function $f : [0, +\infty) \rightarrow \mathbb{R}$ defined as

$$f(v) = \min_{|E|=v} F(E). \quad (4)$$

As a by-product of our analysis, we are able to characterize the asymptotic shape of minimizers as the volume tends to infinity, showing that they converge after appropriate rescaling to the Wulff Shape (i.e. the solution of the isoperimetric problem) relative to an anisotropy ϕ_g depending on g . We mention that, in the small volume regime, the contribution of g becomes irrelevant and the minimizers converge to standard spheres (see [13] and references therein).

The plan of the paper is the following: in Section 2 we show existence of compact minimizers of (4). In Section 3 we prove that the function f is locally Lipschitz continuous and link its derivative to the curvature of the minimizers. We also provide an example of a function f which is not differentiable everywhere. Let us notice that in these first two parts no assumption is made on the average of g or on its size. In Section 4 we use the isovolumetric function to find solutions of (3). Eventually, in Section 4.1 we investigate the behavior of the constrained minimizers of (4) as the volume goes to infinity.

Notation and general assumptions. We shall assume that g is a $\mathcal{C}^{0,\alpha}$ periodic function, with periodicity cell $Q = [0, 1]^d$. We shall also suppose that the dimension of the ambient space is smaller or equal to 7, so that quasi-minimizers of the perimeter have boundary of class $\mathcal{C}^{2,\alpha}$ [14]. We believe that this restriction is not relevant for the results of this work, but we were not able to remove it. For a set of finite perimeter we denote by $P(E)$ its perimeter and by ∂^*E its reduced boundary (see [14] for precise definitions). Given an open set Ω , we denote by $P(E, \Omega)$ the relative perimeter of E in Ω . We take as a convention that the mean curvature (which we define as the sum of all principal curvatures) of a convex set is positive. If ν is the outward normal to a set with smooth boundary, this amounts to say that the mean curvature κ is equal to $\operatorname{div}(\nu)$.

Acknowledgements. The authors would like to thank Antonin Chambolle for very helpful comments and suggestions. They thank Luca Mugnai for pointing out a mistake in a previous version of (8). The first author wish to thank Gilles Thouroude for several discussions on this problem.

2 Existence of minimizers

In this section we prove existence of compact volume-constrained minimizers of F , by showing that for every volume v , the problem is equivalent to the unconstrained problem

$$\min_{E \subset \mathbb{R}^d} F_\mu(E) = \min_{E \subset \mathbb{R}^d} P(E) - \int_E g \, dx + \mu ||E| - v|, \quad (5)$$

for $\mu > 0$ large enough. We start by studying (5), showing existence of smooth compact minimizers. We then show that there exists μ_0 such that, for $\mu \geq \mu_0$, every compact

minimizer of F_μ has volume v . In particular, this will provide existence of minimizers of (4), since $f(v) \leq \min_E F_\mu(E)$ for every $\mu \geq 0$.

Denoting by Q_R the cube $[-R/2, R/2]^d$ of sidelength R , we consider the spatially constrained problem

$$\min_{E \subset Q_R} F_\mu(E). \quad (6)$$

Having restrained our problem to a bounded domain, we gain compactness of minimizing sequences and thus existence of minimizers for (6) by the direct method [14]. We want to show that these minimizers do not depend on R for R big enough. In order to do so, we need density estimates as [7].

Proposition 2.1. *There exist two constants $C(d)$ and γ depending only on the dimension d such that, if we set $r_0(\mu) = \frac{C(d)}{\mu + \|g\|_\infty}$, then for every minimizer E of (6) and every $x \in \mathbb{R}^d$,*

- $|E \cap B_r(x)| \geq \gamma r^d$ for every $r \leq r_0$ if $|B_r(x) \cap E| > 0$ for any $r > 0$,
- $|B_r(x) \setminus E| \geq \gamma r^d$ for every $r \leq r_0$ if $|B_r(x) \setminus E| > 0$ for any $r > 0$.

Proof. Let $x \in \partial^* E$ then by minimality of E we have

$$P(E) - \int_E g \, dx + \mu|E| - v \leq P(E \setminus B_r(x)) - \int_{E \setminus B_r(x)} g \, dx + \mu|E \setminus B_r(x)| - v,$$

hence

$$\begin{aligned} P(E) &\leq \int_{E \cap B_r} g \, dx + P(E \setminus B_r) + \mu|E| - |E \setminus B_r| \\ &= \int_{E \cap B_r} g \, dx + P(E \setminus B_r) + \mu|E \cap B_r| \\ &\leq |E \cap B_r|(\|g\|_\infty + \mu) + P(E \setminus B_r). \end{aligned}$$

On the other hand we have

$$P(E) = \mathcal{H}^{d-1}(\partial^* E \cap B_r) + \mathcal{H}^{d-1}(\partial^* E \cap B_r^c)$$

and

$$P(E \setminus B_r) = \mathcal{H}^{d-1}(E \cap \partial B_r) + \mathcal{H}^{d-1}(\partial^* E \cap B_r^c).$$

From these inequalities we get

$$\mathcal{H}^{d-1}(\partial^* E \cap B_r) \leq \mathcal{H}^{d-1}(E \cap \partial B_r) + (\|g\|_\infty + \mu)|E \cap B_r|.$$

Letting $U(r) = |E \cap B_r|$ and using the isoperimetric inequality [14], we have

$$\begin{aligned} c(d)U(r)^{\frac{d-1}{d}} &\leq P(E \cap B_r) \\ &= \mathcal{H}^{d-1}(\partial^* E \cap B_r) + \mathcal{H}^{d-1}(\partial B_r \cap E) \\ &\leq 2\mathcal{H}^{d-1}(\partial B_r \cap E) + (\|g\|_\infty + \mu)U(r). \end{aligned}$$

Recalling that $\mathcal{H}^{d-1}(\partial B_r \cap E) = U'(r)$ for a.e. $r > 0$, we find

$$c(d)U(r)^{\frac{d-1}{d}} \leq 2U'(r) + (\|g\|_\infty + \mu)U(r). \quad (7)$$

The idea is that, when U is small, the term $U^{\frac{d-1}{d}}$ dominates the term which is linear in U so that we can get rid of it. Letting ω_d be the volume of the unit ball and $r \leq \omega_d^{-\frac{1}{d}} \left(\frac{c(d)}{2(\mu + \|g\|_\infty)} \right)$, we then have

$$U(r) \leq |B_r| = \omega_d r^d \leq \left(\frac{c(d)}{2(\mu + \|g\|_\infty)} \right)^d.$$

Raising each side of the inequality to the power $-\frac{1}{d}$ and multiplying by U we get

$$U(r)^{\frac{d-1}{d}} \geq \frac{2(\mu + \|g\|_\infty)}{c(d)}U$$

and from this

$$\frac{c(d)}{2}U(r)^{\frac{d-1}{d}} - (\mu + \|g\|_\infty)U \geq 0$$

thus finally

$$c(d)U(r)^{\frac{d-1}{d}} - (\mu + \|g\|_\infty)U \geq \frac{c(d)}{2}U(r)^{\frac{d-1}{d}}.$$

Putting this back in (7) and letting $C(d) = c(d)\omega_d^{-\frac{1}{d}}/2$ we have

$$\frac{c(d)}{4}U(r)^{\frac{d-1}{d}} \leq U'(r) \quad \forall r \leq \frac{C(d)}{(\mu + \|g\|_\infty)}.$$

If we set $V(r) = U^{\frac{1}{d}}(r)$ we have

$$V'(r) = \frac{1}{d}U'(r)U^{\frac{1-d}{d}}(r) \geq \frac{c(d)}{4d}.$$

Integrating we get

$$V(r) \geq \frac{c(d)}{4d}r \quad \text{hence} \quad U(r) \geq \left(\frac{c(d)}{4d} \right)^d r^d.$$

The second inequality is obtained by repeating the argument with $E \cup B_r(x)$ instead of $E \setminus B_r(x)$. \square

We now estimate the error made by relaxing the constraint on the volume.

Lemma 2.2. *For every set of finite perimeter E and every $\mu > \|g\|_\infty$ we have*

$$||E| - v| \leq \frac{F_\mu(E) + v\|g\|_\infty}{\mu - \|g\|_\infty}.$$

Proof. If $|E| > v$ we have

$$F_\mu(E) = P(E) - \int_E g + \mu(|E| - v)$$

thus

$$\mu(|E| - v) \leq F_\mu(E) + \|g\|_\infty |E|$$

and from this we find

$$(\mu - \|g\|_\infty)(|E| - v) \leq F_\mu(E) + v\|g\|_\infty.$$

Dividing by $\mu - \|g\|_\infty$ we get

$$||E| - v| \leq \frac{F_\mu(E) + v\|g\|_\infty}{\mu - \|g\|_\infty}.$$

If $|E| \leq v$ we similarly get

$$(\mu + \|g\|_\infty)(|E| - v) \leq F_\mu(E) + v\|g\|_\infty$$

hence

$$||E| - v| \leq \frac{F_\mu(E) + v\|g\|_\infty}{\mu + \|g\|_\infty} \leq \frac{F_\mu(E) + v\|g\|_\infty}{\mu - \|g\|_\infty}.$$

□

We now prove that the minimizers do not depend on R , for R big enough. Here the periodicity of g is crucial.

Proposition 2.3. *For every $\mu > \|g\|_\infty$, there exists $R_0(\mu)$ such that for every $R \geq R_0$, there exists a minimizer E_R of (6) verifying $\text{diam}(E_R) \leq R_0$. Equivalently we have*

$$\min_{E \subset Q_R} F_\mu(E) = \min_{E \subset Q_{R_0}} F_\mu(E)$$

for all $R \geq R_0$.

Proof. Let E_R be a minimizer of (6). Let Q be the unit square and

$$N = \#\{z \in \mathbb{Z}^d / |\{z + Q\} \cap E_R| \neq 0\}.$$

We want to bound N from above by a constant independent of R .

Let $r_0 = \frac{C(d)}{\mu + \|g\|_\infty}$ as in Proposition 2.1. For all $x \in E_R$ we have

$$|E_R \cap B_r(x)| \geq \gamma r^d \quad \forall r \leq r_0.$$

Letting $r_1 = \min(r_0, \frac{1}{2})$, for all $x \in \mathbb{R}^d$ we have

$$\#\{z \in \mathbb{Z}^d / \{z + Q\} \cap B_{r_1}(x) \neq \emptyset\} \leq 2^d.$$

Therefore, we can find at least $N/2^d$ points x_i in E_R such that $B_{r_1}(x_i) \cap B_{r_1}(x_j) = \emptyset$ for every $i \neq j$ and such that $x_i \in Q + z_i$ with $|\{z_i + Q\} \cap E_R| \neq 0$ for some $z_i \in \mathbb{Z}$.

We thus have

$$|E_R| \geq \sum_i |B_{r_1}(x_i) \cap E_R| \geq \frac{N}{2^d} \gamma r_1^d.$$

This gives us

$$N \leq \frac{2^d |E_R|}{\gamma r_1^d}.$$

Letting B^v be a ball of volume v , by Lemma 2.2 and $F_\mu(E_R) \leq F_\mu(B^v)$, we have

$$\begin{aligned} ||E_R| - v| &\leq \frac{F_\mu(B^v) + v\|g\|_\infty}{\mu - \|g\|_\infty} \\ &\leq \frac{c(d)v^{\frac{d-1}{d}} + 2v\|g\|_\infty}{\mu - \|g\|_\infty}. \end{aligned}$$

This shows that

$$|E_R| \leq v + \frac{c(d)v^{\frac{d-1}{d}} + 2v\|g\|_\infty}{\mu - \|g\|_\infty}$$

so that N is bounded by a constant independent of R .

We now prove that $\text{diam}(E_R) \leq C(d)N$. Indeed let $x \in E_R$ and let $P_0 = [0, 1] \times [-R/2, R/2]^{d-1}$ be a slice of Q_R orthogonal to the direction e_1 . For $i \in \mathbb{Z}$ we also set $P_i = P_0 + ie_1$. Our aim is showing that E_R is contained in a box of size N in the direction e_1 . Up to translation we can suppose that $E_R \cap P_i = \emptyset$ for all $i < 0$. We want to show that we can choose $E_R \subset \bigcup_{0 \leq i \leq N} P_i$.

Let $I \leq R$ be the least integer such that $E_R \subset \bigcup_{0 \leq i \leq I} P_i$ and suppose $I \geq N$. Because of the definition of N , there is at most N slices P_i such that $P_i \cap E_R \neq \emptyset$. Hence there exists




Figure 1: the construction in the proof of Proposition 2.3.

i between 0 and N such that $P_i \cap E_R = \emptyset$. Let $E_i^+ = \bigcup_{j>i} E_R \cap P_j$ and $E_i^- = \bigcup_{j<i} E_R \cap P_j$ then if we set $\tilde{E}_R = E_i^- \cup \{E_i^+ - e_1\}$ we have $F_\mu(\tilde{E}_R) = F_\mu(E_R)$ and $\tilde{E}_R \subset \bigcup_{0 \leq i \leq I-1} P_i$ giving the claim by iterating the procedure (see Figure 1).

The same argument applies to any orthonormal direction e_k , hence $E_R \subset Q_{2N}$. \square

We now prove existence of minimizers for F_μ .

Proposition 2.4. *For $\mu > \|g\|_\infty$, there exists a bounded minimizer of F_μ . Moreover such minimizer has boundary of class $C^{2,\alpha}$, where α is the Hölder exponent of the function g .*

Proof. By Proposition 2.3 there exists R_0 such that $E_R \subset B_{R_0}$ for every $R > 0$. Suppose

now that there exists E with $F_\mu(E) < F_\mu(E_{R_0})$. Then there exists $\varepsilon > 0$ such that

$$F_\mu(E) + \varepsilon \leq F_\mu(E_{R_0}).$$

Let us show that there exists $R > R_0$ such that

$$F_\mu(E \cap B_R) + \frac{\varepsilon}{2} \leq F_\mu(E_{R_0}).$$

We start by noticing that $|E \cap B_R|$ tends to $|E|$ and that $\int_{E \cap B_R} g \, dx$ tends to $\int_E g \, dx$ when $R \rightarrow +\infty$. On the other hand,

$$P(E \cap B_R) = \mathcal{H}^{d-1}(E \cap \partial B_R) + \mathcal{H}^{d-1}(\partial^* E \cap B_R)$$

and we have

$$\lim_{R \rightarrow +\infty} \mathcal{H}^{d-1}(\partial^* E \cap B_R) = P(E)$$

and

$$\lim_{R \rightarrow +\infty} \int_0^R \mathcal{H}^{d-1}(E \cap \partial B_s) \, ds = \lim_{R \rightarrow +\infty} |E \cap B_R| = |E|.$$

The last equality shows that $\mathcal{H}^{d-1}(E \cap \partial B_R)$ is integrable so that, for every $R > 0$, there exists $R' > R$ such that $\mathcal{H}^{d-1}(E \cap \partial B_{R'})$ is arbitrarily small. This implies that we can find a R large enough so that

$$F_\mu(E \cap B_R) + \frac{\varepsilon}{2} \leq F_\mu(E_{R_0}).$$

The minimality of E_{R_0} yields to a contradiction.

We now focus on the regularity. Let E be a minimizer of F_μ then for every G ,

$$P(E) - \int_E g \, dx + \mu||E| - v| \leq P(G) - \int_G g \, dx + \mu||G| - v|.$$

Hence

$$\begin{aligned} P(E) &\leq P(G) + \|g\|_\infty |E \Delta G| + \mu||E| - |G|| \\ &\leq P(G) + (\|g\|_\infty + \mu) |E \Delta G|. \end{aligned}$$

E is thus a quasi-minimizer of the perimeter so that, by classical regularity theory [14] (see also [20]), we get that ∂E is of class $\mathcal{C}^{2,\alpha}$. \square

Before stating the equivalence between the constrained and unconstrained problems, we prove an inequality which in the case of convex sets, directly follows from the Alexandrov-Fenchel inequality (see Schneider [23]).

Lemma 2.5. *Let E be a compact set with \mathcal{C}^2 boundary and assume that $\kappa > 0$ on ∂E , where κ denotes the mean curvature of ∂E . Then*

$$(d-1)P(E) \geq |E| \min_{\partial E} \kappa. \quad (8)$$

Proof. Let $\Lambda = \min_{\partial E} \kappa$ then no point of E is at distance of ∂E greater than $\frac{d-1}{\Lambda}$. Indeed, if $x \in E$, considering the ball $B(x, R)$ centered in x and of radius R , with R the smallest radius such that $\partial E \cap B(x, R) \neq \emptyset$ then $R \leq \frac{d-1}{\Lambda}$ since the points of $\partial E \cap B(x, R)$ have curvature less than $\frac{d-1}{R}$. Let now $b(x) = \text{dist}(x, \mathbb{R}^d \setminus E)$ be the distance function to the complementary of E . By the Coarea Formula [3], we have

$$|E| = \int_0^{\frac{d-1}{\Lambda}} P(\{b > t\}) dt$$

from which we deduce (8) provided that

$$P(\{b > t\}) \leq P(\{b > 0\}) = P(E)$$

for a.e. $t > 0$. We now prove this inequality.

As b is locally semi-concave in E (see [18]), that is $D^2b \leq C \text{Id}$ in the sense of measures, the singular part of D^2b is a negative measure. Moreover, letting $Sing$ be the set where b is not differentiable and letting $S = \overline{Sing}$, we have that $Sing$ corresponds to the set of points having more than one projection on ∂E , b is \mathcal{C}^2 out of S , and S is of zero Lebesgue measure [11] (and even $(d-1)$ -rectifiable if ∂E is \mathcal{C}^3 [18]). The hypothesis that ∂E is \mathcal{C}^2 is sharp since there exists sets with $\mathcal{C}^{1,1}$ boundary such that the cut locus is of positive Lebesgue measure [18]. The set S is sometime called the *cut locus* of ∂E . We refer to [2, 18] for a proof of these properties of the distance function b .

If $x \in \{b = t\}$ is a point out of S , by the smoothness of b and by classical formulas there holds [2]

$$-\Delta b(x) = \kappa_{\{b=t\}}(x) = \sum_{i=1}^{d-1} \frac{\kappa_i(\pi(x))}{1 - b(x)\kappa_i(\pi(x))}$$

where $\pi(x)$ is the (unique) projection of x on ∂E and where κ_i are the principal curvatures of ∂E . By the convexity of the function $\kappa \rightarrow \kappa/(1 - b\kappa)$, and recalling that the mean curvature of ∂E is positive, we get that $\Delta b(x) \leq 0$ on $E \setminus S$. Finally, since the singular part of the measure Δb (which is concentrated on S) is non positive, we find that $\Delta b \leq 0$ in the sense of measures.

By the Coarea Formula, for a.e. $t > 0$ we have $\mathcal{H}^{d-1}(\partial\{b > t\} \cap S) = 0$, so that for such t 's

$$P(\{b > t\}) - P(E) = \int_{\{b=t\}} \nabla b \cdot \nu + \int_{\partial E} \nabla b \cdot \nu = \int_{\{0 < b < t\}} \Delta b \leq 0, \quad (9)$$

where ν denotes the exterior unit normal to the set $\{0 < b < t\}$, so that $\nu = -\nabla b$ on ∂E and $\nu = \nabla b$ on $\{b = t\} \setminus S$.

As the vector field ∇b is bounded and its divergence Δb is a Radon measure, the integration by part formula in (9) is justified by a result of Chen, Torres and Ziemer [10, Th. 21.1 (g)]. Notice also that, since ∇b is continuous on $\{b = t\} \setminus S$, the (weak) normal trace of ∇b on $\{b = t\}$ coincides with $\nabla b \cdot \nu$ on $\{b = t\} \setminus S$ [10, Th. 27.1]. \square

Remark 2.6. Under the hypothesis $\Lambda := \min_{\partial E} \kappa > 0$, one could also replace (8) by

$$P(E)R_{\max} \geq |E|$$

where $R_{\max} \leq \frac{d-1}{\Lambda}$ is the radius of the largest ball contained in E .

Remark 2.7. Notice that the inequality

$$\frac{d-1}{d} P(E)^2 \geq |E| \int_{\partial E} \kappa \quad (10)$$

which is one of the Alexandrov-Fenchel inequalities (and which implies (8) does not hold for a general smooth compact set. Indeed, for $d = 2$ we can consider a disjoint union of N balls of radius r_i , so that the left hand-side is of order $(\sum_i r_i)^2$ and the right hand-side is of order $N (\sum_i r_i^2)$. Hence, if we let $r_i = 1/i^2$, we get that the left hand-side remains bounded while the right hand-side blows-up when the number of balls N increases, thus violating (10).

We are finally in position to prove existence of minimizers of problem (4).

Theorem 2.8. *Let $d \leq 7$, then for all $v > 0$ there exists a compact minimizer E_v of (4) with ∂E_v of class $C^{2,\alpha}$. Moreover, E_v is also a minimizer of F_μ for all*

$$\mu \geq C_1(d)\|g\|_\infty + C_2(d)v^{-\frac{1}{d}} \quad (11)$$

where $C_1(d)$ and $C_2(d)$ are two positive constants depending only on d .

Proof. Letting E_μ be a bounded and smooth minimizer of F_μ , given by Proposition 2.4, We will show that $|E_\mu| = v$, for μ large enough. Let μ be larger than $\|g\|_\infty$ and suppose by contradiction $|E_\mu| \neq v$. Then, if $|E_\mu| > v$, the Euler-Lagrange equation for F_μ writes

$$\kappa_{E_\mu} = g - \mu$$

where κ_{E_μ} is the mean curvature of E_μ . But this is impossible since $\mu > \|g\|_\infty$, which would lead to $\kappa_{E_\mu} < 0$, contradicting the compactness of E_μ .

Thus for $\mu > \|g\|_\infty$, we have $|E_\mu| < v$ and

$$\kappa_{E_\mu} = g + \mu.$$

Using inequality (8) with $E = E_\mu$, and the fact that $|E_\mu| \geq v/2$ by Lemma 2.2, we get

$$\begin{aligned} F_\mu(E_\mu) &\geq \frac{1}{d-1}(\mu - \|g\|_\infty)|E_\mu| - \|g\|_\infty|E_\mu| \\ &\geq \frac{1}{d-1}(\mu - \|g\|_\infty)\frac{v}{2} - \|g\|_\infty v. \end{aligned}$$

On the other hand, $F_\mu(E_\mu) \leq F_\mu(B^v)$, where B^v is a ball of volume v , so that

$$C(d)v^{\frac{d-1}{d}} + \|g\|_\infty v \geq F_\mu(B^v) \geq \frac{1}{d-1}(\mu - \|g\|_\infty)\frac{v}{2} - \|g\|_\infty v$$

and we finally obtain

$$\mu \leq C_1(d)\|g\|_\infty + C_2(d)v^{-\frac{1}{d}}.$$

□

Remark 2.9. The minimizer E_v satisfies the Euler-Lagrange equation

$$\kappa_E = g + \lambda_v \quad \text{with } |\lambda_v| \leq \mu,$$

where μ verifies (11). In particular, λ_v and thus also $\|\kappa_E\|_\infty$ are uniformly bounded in v , for $v \in [\varepsilon, +\infty)$.

The regularity of ∂E_v also follows from the works of Rigot [24] and Xia [29] on quasi-minimizers of the perimeter with a volume constraint.

3 Properties of the isovolumetric function

We show here some of the properties of the isovolumetric f defined by (4).

Proposition 3.1. *The function f is sub-additive and locally Lipschitz continuous. Let v be a point of differentiability of f and E_v be a minimizer of (4) then $f'(v) = \lambda_v$ where λ_v is the Lagrange multiplier associated to E_v , that is, $\kappa_{E_v} = g + \lambda_v$. As a consequence, λ_v is unique for almost every $v > 0$, in the sense that it does not depend on the specific minimizer E_v .*

Proof. Let E_v and $E_{v'}$ be compact minimizers associated to v and v' . Up to a translation we can suppose that $F(E_v \cup E_{v'}) = F(E_v) + F(E_{v'})$, so that

$$f(v + v') \leq F(E_v \cup E_{v'}) = F(E_v) + F(E_{v'}) = f(v) + f(v')$$

and f is sub-additive.

By Theorem 2.8, for every $\alpha > 0$ there exists μ_α such that, for every $v \geq \alpha$, the constrained problem (4) and the relaxed one (5) are equivalent for $\mu \geq \mu_\alpha$. Let $v, v' \in [\alpha, +\infty)$, then

$$f(v) = F(E_v) \leq P(E_{v'}) - \int_{E_{v'}} g \, dx + \mu_\alpha |v - v'| = f(v') + \mu_\alpha |v - v'|$$

thus $|f(v) - f(v')| \leq \mu_\alpha |v - v'|$ and f is Lipschitz continuous on $[\alpha, +\infty)$.

We now compute the derivative of f . For $v, \varepsilon > 0$ we have

$$f(v + \varepsilon) - f(v) \leq F((1 + \varepsilon/v)^{\frac{1}{d}} E_v) - F(E_v).$$

Let $\delta_\varepsilon = (1 + \varepsilon/v)^{\frac{1}{d}} - 1$; then $(1 + \varepsilon/v)^{\frac{1}{d}} E_v = E_v + \delta_\varepsilon E_v$. Recalling that $\kappa_{E_v} = g + \lambda_v$ we get

$$\begin{aligned} P((1 + \delta_\varepsilon)E_v) &= P(E_v) + \delta_\varepsilon \int_{\partial E_v} \kappa_{E_v} x \cdot \nu \, d\mathcal{H}^{d-1} + o(\delta_\varepsilon) \\ &= P(E_v) + \delta_\varepsilon \int_{\partial E_v} g(x) x \cdot \nu \, d\mathcal{H}^{d-1} + \delta_\varepsilon \int_{\partial E_v} \lambda_v x \cdot \nu \, d\mathcal{H}^{d-1} + o(\delta_\varepsilon) \\ &= P(E_v) + \delta_\varepsilon \int_{\partial E_v} g(x) x \cdot \nu \, d\mathcal{H}^{d-1} + \delta_\varepsilon \lambda_v d|E_v| + o(\delta_\varepsilon) \end{aligned}$$

and

$$\int_{(1+\delta_\varepsilon)E_v} g = \int_{E_v} g \, dx + \delta_\varepsilon \int_{\partial E_v} g(x) x \cdot \nu \, d\mathcal{H}^{d-1} + o(\delta_\varepsilon).$$

From this we obtain

$$F((1 + \varepsilon/v)^{\frac{1}{d}} E_v) - F(E_v) = \delta_\varepsilon v d \lambda_v + o(\delta_\varepsilon).$$

As $\delta_\varepsilon = \varepsilon/(vd) + o(\varepsilon)$, we find

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0^+} \frac{f(v + \varepsilon) - f(v)}{\varepsilon} &\leq \lambda_v \\ \liminf_{\varepsilon \rightarrow 0^-} \frac{f(v + \varepsilon) - f(v)}{\varepsilon} &\geq \lambda_v. \end{aligned}$$

In particular, if f is differentiable in v we have

$$f'(v) = \lambda_v.$$

□

In fact, the isovolumetric function f is slightly more regular.

Proposition 3.2. *Let λ_v^{\max} and λ_v^{\min} be respectively the bigger and the smaller Lagrange multipliers associated with v then f has left and right derivatives in v and*

$$\lim_{h \rightarrow 0^+} \frac{f(v+h) - f(v)}{h} = \lambda_v^{\min} \leq \lambda_v^{\max} = \lim_{h \rightarrow 0^-} \frac{f(v+h) - f(v)}{h}. \quad (12)$$

The proof is based on the following lemma:

Lemma 3.3. *Let v_n be a sequence converging to v . Then there exist sets E_n with $|E_n| = v_n$ and*

$$f(v_n) = F(E_n),$$

and a set E with $|E| = v$ and

$$f(v) = F(E),$$

such that, up to extraction, E_n tends to E in the L^1 -topology, ∂E_n tends to ∂E in the Hausdorff sense, and λ_n tends to λ , where λ_n (resp. λ) is the Lagrange multiplier corresponding to E_n (resp. to E).

Proof. By Theorem 2.8, we can find minimizers E_n of (4), with $|E_n| = v_n$. Moreover, by Proposition 2.3 we can assume that $E_n \subset B_R$ with R independent of n . Since $P(E_n)$ is uniformly bounded from above, it then follows that there exists a (not relabelled) subsequence of E_n converging in the L^1 -topology to a set $E \subset B_R$ with volume $v = \lim_n v_n$. Moreover, by the lower-semi-continuity of the perimeter and the continuity of f , the set E verifies

$$f(v) = F(E).$$

Let us now prove that the convergence also occurs in the sense of Hausdorff.

Let $\varepsilon > 0$ be fixed and let $x \in E \cap \{y / d(y, \partial E) > \varepsilon\}$. If x is not in E_n then by Proposition 2.1 we have

$$|E_n \Delta E| \geq |B_\varepsilon(x) \setminus E_n| \geq \gamma \varepsilon^d.$$

This is impossible if n is big enough because $|E_n \Delta E|$ tends to zero. Similarly, we can show that for n big enough, all the points of $E^c \cap \{y / d(y, \partial E) > \varepsilon\}$ are outside E_n . This shows that $\partial E_n \subset \{y / d(y, \partial E) \leq \varepsilon\}$. Inverting the rôles of E_n and E , the same argument proves that $\partial E \subset \{y / d(y, \partial E_n) \leq \varepsilon\}$ giving the Hausdorff convergence of ∂E_n to ∂E . Now if λ_n is the Lagrange multiplier associated with E_n , it is uniformly bounded and we can extract a converging subsequence which converges to some $\lambda \in \mathbb{R}$.

To conclude the proof we must show that $\kappa_E = g + \lambda$. As proved for instance in [25], for every $x \in \partial E$ there exists $r > 0$ such that for n large enough the set $B_r(x) \cap \partial E_n$ is the graph of a function φ_n , and the set $B_r(x) \cap \partial E$ is the graph of a function φ , in a suitable coordinate system. We then have that φ_n tends uniformly to φ , as $n \rightarrow +\infty$, and

$$-\operatorname{div} \left(\frac{\nabla \varphi_n}{\sqrt{1 + |\nabla \varphi_n|^2}} \right) = g(x, \varphi_n(x)) + \lambda_n \quad (13)$$

for all n big enough. By elliptic regularity [8], we can pass to the limit in (13) and obtain that ϕ solves

$$-\operatorname{div} \left(\frac{\nabla \phi}{\sqrt{1 + |\nabla \phi|^2}} \right) = \kappa_E = g(x, \phi(x)) + \lambda.$$

□

Proof of Proposition 3.2. Let $v > 0$ and let

$$\lambda = \liminf_{\varepsilon \rightarrow 0^+} f'(v + \varepsilon) \quad (14)$$

Notice that, for every $\varepsilon > 0$, there exists a $v_\varepsilon \in]v, v + \varepsilon[$ such that

$$f'(v_\varepsilon) \leq \frac{f(v + \varepsilon) - f(v)}{\varepsilon}. \quad (15)$$

From (15) we get

$$\lambda \leq \liminf_{\varepsilon \rightarrow 0^+} \frac{f(v + \varepsilon) - f(v)}{\varepsilon}.$$

Let ε_n be a sequence realizing the infimum in (14) and let $E_n \subset B_R$ be a set of volume $v_n = v + \varepsilon_n$ such that

$$f(v_n) = F(E_n).$$

By Lemma 3.3 the sets E_n converge, up to a subsequence in the L^1 -topology, to a limit set E , with $|E| = v$ and $\kappa_E = g + \lambda$, where $\lambda = \lim_n \lambda_n$. Reasoning as in Proposition 3.1, we see that

$$\liminf_{\varepsilon \rightarrow 0^+} \frac{f(v + \varepsilon) - f(v)}{\varepsilon} \geq \lambda \geq \limsup_{\varepsilon \rightarrow 0^+} \frac{f(v + \varepsilon) - f(v)}{\varepsilon}$$

hence f admits a right derivative which is equal to λ_v^{min} . Analogously one can show that f has a left derivative equal to λ_v^{max} . □

Remark 3.4. Notice that (12) implies that f is differentiable at any local minimum so that, if equation (1) has no solution, either f is increasing on $[0, +\infty)$, or there exists $\bar{v} > 0$ such that f is increasing on $[0, \bar{v}]$, decreasing on $[\bar{v}, +\infty)$, and is not differentiable at \bar{v} .

We now give an example of a isovolumetric function f which has a point of nondifferentiability. It is not clear to which extent this is a generic phenomenon.

Example. Consider a periodic function g which is equal to 0 everywhere in the unit cell Q , except in the neighborhood of two points a and b . Around these points, g is taken to be equal to radial parabolas centered at the point, one parabola high and thin, and the other small and large (see Figure 2).

It is shown in [13] that, when the volume v is sufficiently small, the minimizer E_v is connected. Since the bound on v depends only on $\|g\|_\infty$, which can be fixed as small as we want, we can suppose that the minimizers E_v are connected and are located near a or b . By the isoperimetric inequality [14] we then get that E_v is a disk with volume v centered at a or b , and will be denoted by $D_v(a)$, $D_v(b)$, respectively.

Therefore, for small volumes the global minimizer is $D_v(a)$ and, once the equality

$$\int_{D_v(a)} g = \int_{D_v(b)} g$$

is attained, it switches to the disk $D_v(b)$. When this transition occurs, there is a jump singularity of the derivative f' .




Figure 2: example of a function f with a point of nondifferentiability.

4 Existence of surfaces with prescribed mean curvature

In this section we shall assume that g has zero average and satisfies

$$\int_E g \leq (1 - \Lambda)P(E, Q) \quad \forall E \subset Q \quad (16)$$

for some $\Lambda > 0$. Notice that (16) is always satisfied if $\|g\|_{L^d(Q)}$ is small enough, and is precisely the assumption needed in [9] (see also [7]) to prove existence of planelike minimizers of F . Notice also that, if g satisfies (16), then the inequality in (16) holds for all sets $E \subset \mathbb{R}^d$ of finite perimeter. In particular, this implies the following estimate on the function f :

$$c v^{\frac{d-1}{d}} \leq f(v) \leq C v^{\frac{d-1}{d}} \quad \text{for some } 0 < c < C. \quad (17)$$

In the sequel we will need a representation result for the functional F , due to Bourgain and Brezis [6].

Theorem 4.1. *Let g be a function verifying (16) then there exists a periodic and continuous function σ with $\max \sigma(x) < 1$ satisfying $\operatorname{div} \sigma = g$. The energy F can thus be written as an anisotropic perimeter:*

$$F(E) = \int_{\partial^* E} (1 + \sigma(x) \cdot \nu).$$

Theorem 4.1 implies that

$$\Lambda P(E) \leq F(E) \leq 2P(E) \quad (18)$$

for all sets E of finite perimeter.

The next Lemma gives an upper bound on the number of “large” connected components of a volume-constrained minimizer.

Lemma 4.2. *Let g be a periodic $C^{0,\alpha}$ function with zero average and satisfying (16). Let E_v be a compact minimizer of (4), and let E_i be the connected components of E_v . We can order the sets E_i in such a way that $|E_i|$ is decreasing in i . Given $\delta > 0$ let*

$$N_\delta = \left[1 + \left(\frac{C}{c} \right)^d \frac{1}{\delta^d} \right].$$

Then

$$\sum_{i=N_\delta}^{\infty} |E_i| \leq \delta v. \quad (19)$$

Proof. Let $x_i = \frac{|E_i|}{v} \in [0, 1]$. Recalling (17), we have

$$cv^{\frac{d-1}{d}} \sum_{i=1}^{\infty} x_i^{\frac{d-1}{d}} \leq \sum_{i=1}^{\infty} f(|E_i|) = f(v) \leq Cv^{\frac{d-1}{d}},$$

hence

$$\sum_{i=1}^{\infty} x_i^{\frac{d-1}{d}} \leq \frac{C}{c} \quad \text{and} \quad \sum_{i=1}^{\infty} x_i = 1.$$

Let now M be the smallest integer such that

$$\sum_{i=M+1}^{\infty} x_i < \delta,$$

we want to prove that $M < N_\delta$. Indeed, we have

$$\delta \leq \sum_{n=M}^{\infty} x_n = \sum_{n=M}^{\infty} x_n^{\frac{1}{d}} x_n^{\frac{d-1}{d}} \leq x_M^{\frac{1}{d}} \sum_{n=M}^{\infty} x_n^{\frac{d-1}{d}} \leq \frac{C}{c} x_M^{\frac{1}{d}}.$$

We then obtain

$$x_M \geq \left(\frac{c}{C}\right)^d \delta^d.$$

Hence, as

$$1 \geq \sum_{i=1}^M x_i \geq \sum_{i=1}^M x_M = Mx_M,$$

by the decreasing property of x_i , we get

$$1 \geq Mx_M \geq M \left(\frac{c}{C}\right)^d \delta^d,$$

which gives

$$M \leq \left(\frac{C}{c}\right)^d \frac{1}{\delta^d} < N_\delta.$$

□

4.1 Compact solutions with big volume.

From (17), Proposition 3.2 and Remark 3.4, we immediately obtain the following result.

Proposition 4.3. *Let g be a periodic $C^{0,\alpha}$ function of zero average satisfying (16). Assume that $f'(v) \leq 0$ for some $v > 0$. Then there exists $w > 0$ such that $f'(w) = 0$, therefore problem (1) admits a compact solution.*

Theorem 4.4. *Let g be a periodic $C^{0,\alpha}$ function with zero average and satisfying (16). There exist $v_n \rightarrow +\infty$ and compact minimizers E_n of (4) such that $|E_n| = v_n$ and E_n solves*

$$\kappa = g + \lambda_n$$

with $\lambda_n \geq 0$ and $\lambda_n \rightarrow 0$ as $n \rightarrow +\infty$.

Proof. Two situations can occur:

Case 1. There exists a sequence $\tilde{v}_n \rightarrow +\infty$ such that $f'(\tilde{v}_n) \leq 0$. Recalling (17) we have $f(v) \geq cv^{\frac{d-1}{d}}$, which implies that we can find $v_n \geq \tilde{v}_n$ such that f has a local minimum in v_n , hence $\lambda_{v_n} = f'(v_n) = 0$.

Case 2. There exists $v_0 > 0$ such that $f'(v) > 0$ for every $v \geq v_0$. By (17) we have $f(v) \leq Cv^{\frac{d-1}{d}}$, and

$$f(v) = f(v_0) + \int_{v_0}^v f'(s) ds.$$

It follows that there exists a sequence $v_n \rightarrow +\infty$ such that

$$\lim_{n \rightarrow +\infty} f'(v_n) = 0.$$

□

Corollary 4.5. *Let g be a periodic $C^{0,\alpha}$ function with zero average and satisfying (16). Then for every $\varepsilon > 0$ there exists $\varepsilon' \in [0, \varepsilon]$ such that there exists a compact solution of*

$$\kappa = g + \varepsilon'.$$

Notice that for a general function g we cannot let $\varepsilon' = 0$ in Corollary 4.5. Indeed, as shown in [4], there are no compact solutions to (1) for periodic functions g , of zero average, which are translation invariant in some direction and of sufficiently small lipschitz norm.

We expect that condition (16) is not necessary for the thesis of Corollary 4.5 to hold, as suggested by the following result:

Theorem 4.6. *Let g be a periodic $C^{0,\alpha}$ function with zero average and such that $g|_{\partial Q} = 0$. Then for every $\varepsilon > 0$ there exists a compact solution of*

$$\kappa = g + \varepsilon.$$

Proof. Fix $\varepsilon > 0$. For $N \in \mathbb{N}$ we let E_N be a minimizer of the problem

$$\min_{E \subset Q_N} P(E) - \int_E (g(x) + \varepsilon) dx.$$

Since $g|_{\partial Q} = 0$, by strong maximum principle, E_N is contained in the interior of Q_N and either $E_N = \emptyset$ or ∂E_N is a $C^{2,\alpha}$ solution of $\kappa = g + \varepsilon$.

However, from the inequality

$$P(E_N) - \int_{E_N} (g(x) + \varepsilon) dx \leq P(Q_N) - \varepsilon N^d = N^{d-1} (2^d - \varepsilon N) < 0$$

which holds for all $N > 2^d/\varepsilon$, it follows $E_N \neq \emptyset$. \square

4.2 Asymptotic behavior of minimizers.

For $\varepsilon > 0$ and $E \subset \mathbb{R}^d$ of finite perimeter, we let

$$F_\varepsilon(E) = \varepsilon^{(d-1)} F(\varepsilon^{-1}E) = P(E) - \frac{1}{\varepsilon} \int_E g\left(\frac{x}{\varepsilon}\right) dx.$$

Notice that, given a minimizer E_ν of (4), the set εE_ν is a volume-constrained minimizer of F_ε . We recall from [9, Theorem 2] the following result.

Theorem 4.7. *Let g be a periodic $C^{0,\alpha}$ function with zero average and satisfying (16). Then there exists a convex positively one-homogeneous function $\phi_g : \mathbb{R}^d \rightarrow [0, +\infty)$, with $\phi_g(x) > 0$ for all $x \neq 0$, such that the functionals F_ε Γ -converge, with respect to the L^1 -convergence of the characteristic functions, to the anisotropic functional*

$$F_0(E) = \int_{\partial^* E} \phi_g(\nu) d\mathcal{H}^{d-1} \quad E \subset \mathbb{R}^d \text{ of finite perimeter.}$$

We remark that, with a minor modification of the proof, the result of Theorem 4.7 also holds if we restrict the functionals F_ε and F_0 to set of prescribed volume. In particular, by a general property of Γ -converging sequences [12], we have the following consequence of Theorem 4.7.

Corollary 4.8. *Let \tilde{E}_ε be minimizers of F_ε with volume constraint $|\tilde{E}_\varepsilon| = v$, then*

$$\limsup_{\varepsilon \rightarrow 0} F_\varepsilon(\tilde{E}_\varepsilon) \leq \min_{|\tilde{E}|=v} F_0(\tilde{E}). \quad (20)$$

Moreover, if $|\tilde{E}_\varepsilon \Delta \tilde{E}| \rightarrow 0$ for some $\tilde{E} \subset \mathbb{R}^d$, as $\varepsilon \rightarrow 0$, then $|\tilde{E}| = v$ and \tilde{E} is a volume-constrained minimizer of F_0 . More generally, if $\tilde{E}_\varepsilon \rightarrow \tilde{E}$ in the L^1_{loc} topology, then \tilde{E} is a minimizer of F_0 with volume constraint $|\tilde{E}| \leq v$.

Given the function ϕ_g as above, we let

$$W_g = \left\{ x \in \mathbb{R}^d : \max_{\phi_g(y) \leq 1} x \cdot y \leq 1 \right\}$$

be the Wulff Shape corresponding to ϕ_g . It is well-known that W_g is the unique minimizer of F_0 with volume constraint, up to homothety and translation [28, 26].

By Theorem 4.7 we can characterize the asymptotic shape of the constrained minimizers as the volume tend to infinity.

Theorem 4.9. *Let $d \leq 7$. For $v > 0$ we let E_v be volume-constrained minimizers of (4), whose existence is guaranteed by Theorem 2.8. Then, there exist points $z_v \in \mathbb{R}^d$ such that letting*

$$\tilde{E}_v = \left(\frac{|W_g|}{v} \right)^{\frac{1}{d}} E_v + z_v$$

it holds

$$\lim_{v \rightarrow +\infty} \left| \tilde{E}_v \Delta W_g \right| = 0. \quad (21)$$

Proof. Notice first that \tilde{E}_v is a minimizer of $F_{\left(\frac{|W_g|}{v}\right)^{\frac{1}{d}}}$, with volume constraint $|\tilde{E}_v| = |W_g|$. Moreover, by (17) the perimeter of \tilde{E}_v is uniformly bounded in v .

Case 1. Let us consider the case $d = 2$. Assume first that \tilde{E}_v is connected. Then we have

$$\text{diam}(\tilde{E}_v) \leq P(\tilde{E}_v)/\pi,$$

hence the sets \tilde{E}_v are all contained, up to a translation, in a fixed ball centered in the origin. By the compactness theorem for sets of finite perimeter [14], there exist a bounded set \tilde{E}_∞ of finite perimeter and a sequence $v_k \rightarrow \infty$ such that $|\tilde{E}_\infty| = |W_g|$ and

$$\lim_{k \rightarrow +\infty} \left| \tilde{E}_{v_k} \Delta \tilde{E}_\infty \right| = 0.$$

Since by Theorem 4.7 the set \tilde{E}_∞ is also a volume-constrained minimizer of F_0 , by uniqueness of the minimizer it follows that \tilde{E}_∞ is equal to W_g up to a translation.

We now consider the general case when the sets \tilde{E}_v are not necessarily connected. In particular we can write $\tilde{E}_v = \cup_{i \geq 1} \tilde{E}_v^i$, with $|\tilde{E}_v^i|$ a decreasing sequence and $\sum_{i \geq 1} |\tilde{E}_v^i| = 1$. Reasoning as before, there exists a sequence $v_k \rightarrow +\infty$ such that for all $i \in \mathbb{N}$ the sets $\tilde{E}_{v_k}^i$ converge to $\rho_i W_g$, up to a translation, where $\rho_i \in [0, 1]$ is a decreasing sequence. Moreover, by Lemma 4.2, for all $\delta > 0$ there exists $N_\delta \in \mathbb{N}$ such that $\sum_{i=N_\delta}^\infty |\tilde{E}_v^i| \leq \delta |W_g|$ for all $\delta > 0$, which implies in the limit

$$\sum_{i=1}^{\infty} \rho_i^2 = 1. \quad (22)$$

We claim that $\rho_1 = 1$ and $\rho_i = 0$ for all $i > 1$. Indeed, from (20) we have

$$F_0(W_g) \geq \limsup_{k \rightarrow +\infty} F_{\left(\frac{|W_g|}{v_k}\right)^{\frac{1}{2}}}(\tilde{E}_{v_k}) \geq \sum_{i=1}^{+\infty} F_0(\rho_i W_g) = F_0(W_g) \sum_{i=1}^{+\infty} \rho_i.$$

Recalling (22), this implies

$$\sum_{i=1}^{+\infty} \rho_i = \sum_{i=1}^{+\infty} \rho_i^2 = 1$$

which proves the claim.

Case 2. We now turn to the general case. Let $v_k \rightarrow +\infty$ and let $\varepsilon_k = (|W_g|/v_k)^{\frac{1}{d}}$. For all k , let $\{Q_{i,k}\}_{i \in \mathbb{N}}$ be a partition of \mathbb{R}^d into disjoint cubes of equal volume larger than $2|W_g|$, such that the sets $\tilde{E}_{v_k} \cap Q_{i,k}$ are of decreasing measure, and let $x_{i,k} = |\tilde{E}_{v_k} \cap Q_{i,k}|/|W_g|$. By the isoperimetric inequality [14], there exist $0 < c < C$ such that

$$\begin{aligned} c \sum_i x_{i,k}^{\frac{d-1}{d}} &= c \sum_i \min \left(\frac{|\tilde{E}_{v_k} \cap Q_{i,k}|}{|W_g|}, \frac{|Q_{i,k} \setminus \tilde{E}_{v_k}|}{|W_g|} \right)^{\frac{d-1}{d}} \\ &\leq \sum_i P(\tilde{E}_{v_k}, Q_{i,k}) \\ &\leq \sum_i \frac{1}{\Lambda} \int_{\partial \tilde{E}_{v_k} \cap Q_{i,k}} \left(1 + \sigma \left(\frac{x}{\varepsilon_k} \right) \cdot \nu \right) d\mathcal{H}^{d-1} \\ &\leq \frac{1}{\Lambda} F_{\varepsilon_k}(\tilde{E}_{v_k}) \leq C \end{aligned}$$

hence

$$\sum_{i=1}^{+\infty} x_{i,k} = 1 \quad \text{and} \quad \sum_{i=1}^{+\infty} x_{i,k}^{\frac{d-1}{d}} \leq \frac{C}{c}.$$

Reasoning as in Lemma 4.2 we obtain that for all $\delta > 0$ there exists $N_\delta \in \mathbb{N}$ such that

$$\sum_{i=N_\delta}^{\infty} x_{i,k} \leq \delta. \quad (23)$$

Up to extracting a subsequence, we can suppose that $x_{i,k} \rightarrow \alpha_i^d \in [0, 1]$ as $k \rightarrow +\infty$ for every $i \in \mathbb{N}$, so that by (23) we have

$$\sum_i \alpha_i^d = 1. \quad (24)$$

Let $z_{i,k} \in Q_{i,k}$. Up to extracting a further subsequence, we can suppose that $d(z_{i,k}, z_{j,k}) \rightarrow c_{ij} \in [0, +\infty]$, and

$$\left(\tilde{E}_{v_k} - z_{i,k} \right) \rightarrow E_i \quad \text{in the } L_{\text{loc}}^1\text{-convergence}$$

for every $i \in \mathbb{N}$ (see Figure 3). By Corollary 4.8 we thus have

$$E_i = \rho_i W_g \quad \rho_i \in [0, 1].$$

We say that $i \sim j$ if $c_{ij} < +\infty$ and we denote by $[i]$ the equivalence class of i . Notice that E_i equals E_j up to a translation, if $i \sim j$. We want to prove that

$$\sum_{[i]} \rho_i^d \geq 1, \quad (25)$$

where the sum is taken over all equivalence classes. For all $R > 0$ let $Q_R = [-R/2, R/2]^d$ be the cube of sidelength R . Then for every $i \in \mathbb{N}$,

$$|E_i| \geq |E_i \cap Q_R| = \lim_{k \rightarrow +\infty} \left| \left(\tilde{E}_{v_k} - z_{i,k} \right) \cap Q_R \right|.$$

If j is such that $j \sim i$ and $c_{ij} \leq \frac{R}{2}$, possibly increasing R we have $Q_{j,k} - z_{i,k} \subset Q_R$ for all $k \in \mathbb{N}$, so that

$$\lim_{k \rightarrow +\infty} \left| \left(\tilde{E}_{v_k} - z_{i,k} \right) \cap Q_R \right| \geq \lim_{k \rightarrow +\infty} \sum_{c_{ij} \leq \frac{R}{2}} |\tilde{E}_{v_k} \cap Q_{j,k}| = \sum_{c_{ij} \leq \frac{R}{2}} \alpha_j^d |W_g|.$$

Letting $R \rightarrow +\infty$ we then have

$$|E_i| \geq \sum_{i \sim j} \alpha_j^d |W_g|$$

hence, recalling (24),

$$\sum_{[i]} |E_i| \geq |W_g|,$$

thus proving (25).

Let us now show that

$$\sum_{[i]} \rho_i^{d-1} = 1. \quad (26)$$

Up to passing to a subsequence, from now on we shall assume that $c_{ij} = +\infty$ for all $i \neq j$. Let $I \in \mathbb{N}$ be fixed. Then for every $R > 0$ there exists $K \in \mathbb{N}$ such that for every $k \geq K$ and i, j less than I , we have

$$d(z_{i,k}, z_{j,k}) > R.$$

For $k \geq K$ we thus have

$$\begin{aligned} F_{\varepsilon_k}(\tilde{E}_{v_k}) &\geq \sum_{i=1}^I \int_{\partial \tilde{E}_{v_k} \cap (B_R + z_{i,k})} \left(1 + \sigma \left(\frac{x}{\varepsilon_k} \right) \cdot \nu \right) d\mathcal{H}^{d-1} \\ &= \sum_{i=1}^I \int_{\partial(\tilde{E}_{v_k} - z_{i,k}) \cap B_R} \left(1 + \sigma \left(\frac{x}{\varepsilon_k} \right) \cdot \nu \right) d\mathcal{H}^{d-1} \\ &= \sum_{i=1}^I F_{\varepsilon_k}(\tilde{E}_{v_k} - z_{i,k}, B_R) \end{aligned}$$




Figure 3: the construction in the proof of Theorem 4.9.

where

$$F_\varepsilon(E, B_R) = \int_{\partial E \cap B_R} \left(1 + \sigma \left(\frac{x}{\varepsilon_k} \right) \cdot \nu \right) d\mathcal{H}^{d-1}.$$

From this, (20) and the Γ -convergence of $F_\varepsilon(\cdot, B_R)$ to $F_0(\cdot, B_R)$, we get

$$F_0(W_g) \geq \limsup_{\varepsilon_k \rightarrow 0} F_{\varepsilon_k}(\tilde{E}_{v_k}) \geq \sum_{i=1}^I \liminf_{\varepsilon_k \rightarrow 0} F_{\varepsilon_k}(\tilde{E}_{v_k} - z_{i,k}, B_R) \geq \sum_{i=1}^I F_0(E_i, B_R).$$

For $R > \text{diam}(W_g)$ we have $F_0(E_i, B_R) = F_0(E_i)$ because $E_i = \rho_i W_g$ and therefore

$$F_0(W_g) \geq \sum_{i=1}^I F_0(E_i) = \sum_{i=1}^I \rho_i^{d-1} F_0(W_g).$$

Letting $I \rightarrow +\infty$ we get (26).

Recalling (25), from (26) we then obtain

$$\sum_i \rho_i^{d-1} = \sum_i \rho_i^d = 1.$$

As before, this implies $\rho_1 = 1$ and $\rho_i = 0$ for all $i > 1$, thus giving

$$\lim_{k \rightarrow +\infty} \left| \left(\tilde{E}_{v_k} - z_{1,k} \right) \Delta W_g \right| = 0.$$

By the uniqueness of the limit this shows that the whole sequence \tilde{E}_v tends to W_g as $v \rightarrow +\infty$, up to suitable translations. \square

Remark 4.10. Let us point out that, if uniform density estimates for \tilde{E}_v were available, we would get Hausdorff convergence instead of L^1 convergence in (21), showing in particular that the sets \tilde{E}_v are connected for v large enough (see [21]). We believe that such estimates are true even if we were not able to prove them.

Remark 4.11. The asymptotic behavior of minimizers of (4), in the small volume regime, have been considered in [13], where the authors prove a result similar to Theorem 4.9, with the Wulff Shape W_g replaced by the Euclidean ball, showing in particular that the volume term becomes irrelevant for small volumes.

Remark 4.12. Notice that the results of this paper can be extended with minor modifications of the proofs to anisotropic perimeters of the form

$$P_\phi(E) = \int_{\partial^* E} \phi(\nu) d\mathcal{H}^{d-1}$$

where $\phi : \mathbb{R}^d \rightarrow [0, +\infty)$ is a smooth and uniformly convex norm on \mathbb{R}^d , with $d \leq 3$ [1].

References

- [1] F.J. ALMGREN, R. SCHOEN AND L. SIMON, *Regularity and singularity estimates on hypersurfaces minimizing elliptic variational integrals*, Acta Math., vol. 139, pp. 217-265, 1977.
- [2] L. AMBROSIO AND N. DANCER, *Calculus of variations and partial differential equations, Topics on geometrical evolution problems and degree theory*, papers from the Summer School held in Pisa, September 1996. Edited by G. Buttazzo, A. Marino and M. K. V. Murthy. Springer-Verlag, Berlin, 2000.
- [3] L. AMBROSIO, N. FUSCO AND D. PALLARA, *Functions of bounded variation and free discontinuity problems*, Oxford Science Publications, 2000.
- [4] G. BARLES, A. CESARONI AND M. NOVAGA, *Homogenization of fronts in highly heterogeneous media*, SIAM J. on Math. Anal., vol. 43 n. 1, 212-227, 2011.

- [5] F. BETHUEL, P. CALDIROLI AND M. GUIDA, *Parametric Surfaces with Prescribed Mean Curvature*, Rend. Sem. Mat. Univ. Torino, vol. 60 n. 4, 175-231, 2002.
- [6] J. BOURGAIN AND H. BREZIS, *On the Equation $\operatorname{div} Y = f$ and Application to Control of Phases*, Jour. of Amer. Math. Soc., vol. 16 n. 2, 393-426, 2002.
- [7] L. CAFFARELLI AND R. DE LA LLAVE, *Planelike Minimizers in Periodic Media*, Comm. Pure Appl. Math., vol. 54, 1403-1441, 2001.
- [8] L. CAFFARELLI AND X. CABRÉ, *Fully Nonlinear Elliptic Equations*, AMS, 1991.
- [9] A. CHAMBOLLE AND G. THOUROUDE, *Homogenization of interfacial energies and construction of plane-like minimizers in periodic media through a cell problem*, Netw. Heterog. Media, vol. 4 n. 1, 127-152, 2009.
- [10] G.-Q. CHEN, W. TORRES AND W.P. ZIEMER, *Gauss-Green theorem for weakly differentiable vector fields, sets of finite perimeter, and balance laws*, Comm. Pure Appl. Math. 62, no. 2, 242-304, 2009.
- [11] G. CRASTA AND A. MALUSA, *The distance function from the boundary in a Minkowski space*, Trans. Amer. Math. Soc., 359, 2007.
- [12] G. DAL MASO, *An introduction to Γ -convergence*, Progress in Nonlinear Differential Equations and their Applications, vol. 8, Birkhäuser, 1993.
- [13] A. FIGALLI AND F. MAGGI, *On The Equilibrium Shapes Of Liquid Drops And Crystals*, to appear on Arch. Rat. Mech. Anal.
- [14] E. GIUSTI, *Minimal Surfaces and functions of Bounded Variation*, Monographs in Mathematics, vol. 80, Birkhäuser, 1984.
- [15] M. GUIDA AND S. ROLANDO, *Symmetric κ -loops*, Diff. Int. Equations, vol. 23, 861-898, 2010.
- [16] X. HUANG, *Closed Surface with Prescribed Mean Curvature in \mathbb{R}^3* , Science in China, vol. 34 n. 10, 1991.
- [17] S. KIRSCH, *PhD Thesis*, Université Pierre et Marie Curie, 2007.
- [18] C. MANTEGAZZA AND A.C. MENNUCCI, *Hamilton-Jacobi equations and distance functions on Riemannian manifolds* Appl. Math. Optim. 47, no. 1, 1-25, 2003.
- [19] F. MORGAN, *Geometric Measure Theory. A Beginner's Guide*, Fourth Edition, Elsevier Academic Press, 2009.

- [20] F. MORGAN, *Regularity of isoperimetric hypersurfaces in Riemannian manifolds*, Trans. AMS, vol. 355 n. 12, pp. 5041-5052, 2003.
- [21] F. MORGAN AND A. ROS, *Stable constant-mean-curvature hypersurfaces are area minimizing in small L^1 neighborhoods*, Interfaces Free Bound., vol. 12 n. 2, pp. 151-155, 2010.
- [22] M. NOVAGA AND E. VALDINOCI, *Bump solutions for the mesoscopic Allen-Cahn equation in periodic media*, Calc. Var. PDE, vol. 40 n. 1-2, pp. 37-49, 2011.
- [23] R. SCHNEIDER, *Convex Bodies: The Brunn-Minkowski Theory*, Encyclopedia of Mathematics and its Applications, Cambridge university Press, 1993.
- [24] S. RIGOT, *Ensembles Quasi-Minimaux avec Contrainte de Volume et Rectifiabilité Uniforme*, Mémoires de la Société Mathématiques de France, vol. 82, 2000.
- [25] I. TAMANINI, *Boundaries of Caccioppoli sets with Hölder continuous normal vector*, J. Reine Angew. Math., vol. 334, 27-39, 1982.
- [26] J. TAYLOR, *Crystalline variational problems*, Bull. Amer. Math. Soc., vol. 84 n. 4, 568-588, 1978.
- [27] A.E. TREIBERGS AND S.W. WEI, *Embedded hyperspheres with prescribed mean curvature*, J. Differential Geom., vol. 18 n. 3, 513-521, 1983.
- [28] G. WULFF, *Zur Frage der Geschwindigkeit des Wachstums und der Auflösung der Kristallflächen*, Z. Kristallogr., vol. 34, 449-530, 1901.
- [29] Q. XIA, *Regularity of minimizers of quasi perimeters with a volume constraint*, Interfaces and Free Boundaries, vol. 7 n. 3, 2005.
- [30] S.T. YAU, *Problem section. Seminar on Differential Geometry*. Ann. of Math. Stud., vol. 102, 669-706, Princeton Univ. Press, Princeton, N.J., 1982.