HAL open science

Central limit theorem by moments

René Blacher

To cite this version:

René Blacher. Central limit theorem by moments. Statistics and Probability Letters, 2009, 77 (17), pp.1647. 10.1016/j.spl.2007.04.003 . hal-00580098

HAL Id: hal-00580098

https://hal.science/hal-00580098v1

Submitted on 26 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Author's Accepted Manuscript



www.elsevier.com/locate/stapro

Central limit theorem by moments

René Blacher

PII: S0167-7152(07)00146-0
DOI: doi:10.1016/j.spl.2007.04.003
Reference: STAPRO 4649

To appear in: *Statistics & Probability Letters*

Received date: 23 March 2004
Revised date: 22 February 2006
Accepted date: 10 April 2007

Cite this article as: René Blacher, Central limit theorem by moments, *Statistics & Probability Letters* (2007), doi:10.1016/j.spl.2007.04.003

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Central Limit Theorem by Moments

René BLACHER

January 21, 2007

René BLACHER
Laboratoire LMC, BP 53 38.041 Grenoble Cedex 9 FRANCE

Summary : In a previous central limit theorem by moments, it has been proved that the moments converge to those of the normal distribution if the moments of sums are asymptotically independent (cf Theoreme de la limite centrale par les moments, Blacher R. Cpte rendus Acad. Sci Paris. t-311- I, 465-468). . In this paper we generalize this result by adding a negligible sequence to these sums. So, we can prove that the moments of some functionals of strong mixing sequences converge.

KeyWords : Central Limit Theorem, moments, strongly mixing sequence, higher order correlation coefficients.

Accepted manuscript

1 Introduction

Let $\{X_n\}$ be a sequence of random variables defined on a probability space (Ω, A, P) such that, for all $n \in N$, $E\{X_n\} = 0$ and $0 < E\{|X_n|^k\} < \infty$ for all $k \in N$.

On decompose $X_1 + X_2 + ... + X_n$ en $X_1 + X_2 + ... + X_{u(n)}$, $X_{u(n)+1} + X_{u(n)+2} + ... + X_{u(n)+t(n)}$ and $X_{u(n)+t(n)+1} + X_{u(n)+t(n)+2} + ... + X_{u(n)+t(n)+u(n)}$, where $u(n) + t(n) + u(n) = n$.

Notations 1.1 *: We denote by $\kappa(n) \in N$, an increasing sequence such that $\kappa(1) = 0$, $\kappa(n) \leq n$ and $\kappa(n)/n \to 0$ as $n \to \infty$. We define the sequences u(n) and t(n) by : u(1)=1, $u(n) = max\{m \in N^* | 2m + \kappa(m) \leq n\}$ and t(1)=0, t(n) = n-2u(n) if $n \geq 2$. Moreover, we simplify u(n) and t(n) as u and t.*

Let $\sigma(u)^2$ be the variance of $X_1+X_2+...+X_u$. One sets $S_u = \frac{X_1+X_2+.....+X_u}{\sigma(u)}$, $\xi_u = \frac{X_{u+1}+X_{u+2}+.....+X_{u+t}}{\sigma(u)}$ and $S'_u = \frac{X_{u+t+1}+X_{u+t+2}+.....+X_{u+t+u}}{\sigma(u)}$. If t(n)=0, we set $\xi_u = 0$. Moreover, one tolerates the notation $S_n = \frac{X_1+X_2+.....+X_n}{\sigma(n)}$.

We denote also by $\sigma^o(t)^2$ the variance of $X_{u+1} + X_{u+2} + ... + X_{u+t}$.

ACCEPTED MANUSCRIPT

In [?] , one has proved that $u \to \infty$, $n/u \to 2$ and $t/u \to 0$ as $n \to \infty$.

Moreover, if $\kappa(n) \to \infty$, $t \to \infty$ as $n \to \infty$. Finally, ξ_u is negligible if $\sigma^o(t)/\sigma(u) \to 0 : E\{(\xi_u)^2\} \to 0$.

Notations 1.2 *: We define conditions H_{mS} and H_{mI} by the following way.*

H_{mS} : $\forall p \in \mathbf{N}$, $E\big\{(S_u + v_u)^p\big\} - E\big\{(S_u + v_u)^p\big\} \to 0$ as $n \to \infty$.

H_{mI} : $\forall (p,q) \in \mathbf{N^2}$, $E\big\{(S_u + v_u)^p (S'_u + v'_u)^q\big\} - E\big\{(S_u + v_u)^p\big\} E\big\{(S'_u + v'_u)^q\big\} \to 0$ as $n \to \infty$,

where $\{v_u\}$ and $\{v'_u\}$ are two sequences of random variables such that $E\{(v_u)^{2k}\} + E\{(v'_u)^{2k}\} \to 0$ as $n \to \infty$ for all $k \in N$.

Now, we want to study the MCLT (Moment Central Limit theorem) for $\{X_n\}$. Generally, one assumes that $\{X_n\}$ is a martingale or is strong mixing. Then, one proves that under some assumptions, moments converge or are bounded : cf Bernstein, [?] , Brown [?] , Eissein-Janson [?], Hernndorf [?], Birkel [?], Krugov [?], Mairoboda [?], Yokohama ([?],[?]), Cox-Kim [?], Ibragimov-Lifshits [?], Soulier [?], Rozovsky [?].

So one can deduce that H_{mI} holds. For example, if $\{X_n\}$ is φ-mixing, H_{mI} holds if $t(n) \to \infty$ as $n \to \infty$ enough slowly (cf [?],[?]). If $\{X_n\}$ is strong mixing, H_{mI} holds if $E\{|S_n|^p\}$ is bounded for all $p \in N$. Of course, if $\{X_n\}$ is strictly stationary, H_{mS} holds.

Now, the martingale condition is a strong condition. It is the same for strong mixing conditions : even they does not hold for some AR(1) processes. Then, some authors have introduce weaker hypotheses : Versik Ornstein ([?], [?]), Withers [?], Cogburn [?] Rosenblatt [?], Pinskers [?]. But they did not studied the moments's convergence. Now another look is possible : one can define asymptotical independence conditions by using moments. In this way, one can use higher order correlation coefficients (cf Lancaster [?], Blacher [?]). Then, in [?] we have turned the convergence of moments into an equivalent condition on these coefficients. These results have led to the introduction of the condition H_{mI} : that is, rather than to assume that $\{X_n\}$ is strong mixing, we choose H_{mI} as asymptotical independence assumptions. Then, under this assumption, $S_n \to^M N(0,1)$, that is, for all $p \in N$, $E\{(S_n)^p\} \to \mu_p$ as $n \to \infty$, the moment of order p of N(0,1).

Théorème 1 *: Assume that $E\{|\xi_u|^k\} \to 0$ as $n \to \infty$ for all $k \in N$. Assume that H_{mS} and H_{mI} hold. Then, $S_n \to^M N(0,1)$.*

In [?], we have proved this theorem when $v_u = v'_u = 0$. In section III, we prove theorem **??** for any sequences v_u and v'_u . In II-a, we explain why we need this generalization. In II-b, we apply these results to functionals of IID sequences. In II-C we prove the MCLT for some functionals of strong mixing sequences or more general sequences.

Accepted manuscript

ACCEPTED MANUSCRIPT

2 Moments Central Limit Theorems for functionals of some strictly stationary sequences

2.1 Discussion

Notations 2.1 *: Let Ψ_t be a strictly stationary process. Assume that $X_t = \sum_{i=0}^{\infty} k_{i+1}(\Psi_{t+i})$, where the k_i's are measurable functions such that $E\{k_i(\Psi_1)\} = 0$ and $0 < E\{k_i(\Psi_1)^{2p}\} < \infty$ for all $i \in N$ and all $p \in N$.*

We set $M_n = \frac{1}{\sigma(n)} \sum_{r=1}^{n} \sum_{s=1}^{r} k_s(\Psi_r)$ and $r_n = \frac{1}{\sigma(n)} \sum_{r=1}^{\infty} \sum_{s=r+1}^{r+n} k_s(\Psi_{n+r})$. Then, one can write $S_n = M_n + r_n$ with $E\{M_n\} = E\{r_n\} = 0$.

Of course, $S_{u(n)} = M_{u(n)} + r_{u(n)}$. Moreover, one can write $S'_{u(n)} = M'_{u(n)} + r'_{u(n)}$ with $M'_u = \frac{1}{\sigma(u)} \sum_{r=1}^{u} \sum_{s=1}^{r} k_s(\Psi_{r+u+t})$ and $r'_u = \frac{1}{\sigma(u)} \sum_{r=1}^{\infty} \sum_{s=r+1}^{r+u} k_s(\Psi_{u+r+u+t})$.

Now one wants to know if the MCLT hold. For example let $\{\Theta_t\}$ be an IID sequence of random variables. Assume that $\Psi_t = \Theta_t$ and that, for all $p \in N$, $E\{|r_n|^p\} \to 0$ as $n \to \infty$.

In order to prove the MCLT, one can try to apply the results of [21], i.e. theorem **??** with $v_u = v'_u = 0$: we write

$$E\{(S_u)^p (S'_u)^q\} = E\{(M_u)^p\} E\{(M'_u)^q\} + \sum_{s \neq q, r \neq p} A_{r,s} E\big\{(M_u)^r (r_u)^{p-r} (M'_u)^s (r'_u)^{q-s}\big\}.$$

By Holder Inequality, $E\big\{(M_u)^r (r_u)^{p-r} (M'_u)^s (r'_u)^{q-s}\big\}$ converges to 0 if $E\{(M_u)^r\}$ is bounded. Then, in order to use the theorem **??** with $v_u = v'_u = 0$, we have to prove that $E\{(M_u)^r\}$ is bounded. It is also difficult as well to prove that $E\{(S_n)^r\}$ is bounded. But, if one uses theorem **??** with $v_u = -r_u$ and $v'_u = -r'_u$, H_{mI} holds obviously : $E\big\{(S_u+v_u)^p (S'_u+v'_u)^q\big\} = E\big\{(S_u+v_u)^p\big\} E\big\{(S'_u+v'_u)^q\big\}$. For this reason we generalize theorem **??** to the case where v_u and v'_u are not equal to 0.

2.2 II-b) Functional of some IID sequences.

Proposition 2.1 *: Assume that $\Psi_t = \Theta_t$ and that $\left|\sum_{s=0}^{\infty} \sum_{i=s}^{\infty} k_{i+1}(\Psi_1)\right| < C < \infty$. Then, $S_n \to^M N(0,1)$.*

Proof : First, we need the following lemma.

Lemma 2.2 *: Assume that $E\big\{k_i(\Psi_r) k_j(\Psi_s)\big\} = 0$ for all $(i,j,r,s) \in N^4$, $r \neq s$. Assume $\sum_{s=0}^{\infty} \left|\sum_{i=s}^{\infty} k_{i+1}(\Psi_1)\right| < C < \infty$ and that, in $L^2(\Omega)$, $\sum_{i=1}^{r} k_i(\Psi_1) \to K$ as $r \to \infty$ where $\int K^2.dP > 0$. Then, $\sigma(n) \to \infty$ and $sup|r_n| \to 0$.*

Proof : There exists N_0 such that $E\Big\{\Big(\sum_{r=1}^{n} \sum_{s=1}^{r} k_s(\Psi_r)\Big)^2\Big\} = \sum_{r=1}^{n} E\Big\{\Big(\sum_{s=1}^{r} k_s(\Psi_r)\Big)^2\Big\} \geq (n/2) \int K^2.dP$ if $n \geq N_0$. Moreover, $\sigma(n)|r_n| \leq 2C$. ★

Accepted manuscript

ACCEPTED MANUSCRIPT

We choose $\kappa(n) = 0$. Because $\sigma(n) \to \infty$, $E\{|\xi_u|^p\} \to 0$ as $n \to \infty$ for all $p \in N$. Moreover, $E\{(M_u)^p\} = E\{(M'_u)^p\}$ and $E\{(M_u)^p(M'_u)^q\} = E\{(M_u)^p\}E\{(M'_u)^q\}$. Then we apply theorem **??** with $v_u = -r_u$ and $v'_u = -r'_u$. ★

Example 2.3 *: Assume that Θ_1 has the standard normal distribution and that $k_i(\Psi_1) = i^{-3}sin(2\pi\Theta_1)$. Then, $S_n \to^M N(0,1)$.*

One can also prove the asymptotic normality in some cases where r_n is not bounded .

Example 2.4 : **AR(1)Process**. *Assume $X_t = \sum_{i=0}^{\infty} a_i\Theta_{t+i}$ where $0 < a < 1$. Then, $M_n = \frac{1}{\sigma(n)}\sum_{j=1}^{n}\frac{1-a^j}{1-a}\Theta_j$ and $r_n = \frac{1}{\sigma(n)}\sum_{j=n+1}^{\infty}\frac{a^{j-n}(1-a^n)}{1-a}\Theta_j$. Therefore, for all $r \in N$, $\kappa u_r(r_u) = \frac{\kappa u_r(\Theta_1)}{\sigma(u)^r}\sum_{j=u+1}^{\infty}\left(\frac{a^{j-u}(1-a^u)}{1-a}\right)^r \to 0$, where κu_r is the r-th cumulant. Therefore, $E\{|r_u|^p\} \to 0$ for all $p \in N^*$.*

*Then, the condition H_{mI} is satisfied for any sequence $\{u(n)\}$ defined in notation **??**. Then, $S_n \to^M N(0,1)$.*

Recall, there exists AR(1) processes which are not strongly mixing : e.g. a=1/2, $P\{\Theta_t = 1/2\} = P\{\Theta_t = -1/2\} = 1/2$(cf [**?**] p 360-362). Then, these processes satisfy H_{mI} but are not strongly mixing. Moreover, $S_n \to^M N(0,1)$.

2.3 Functional of some strong mixing sequences

Strong mixing sequences can often be approached by functional of IID sequences. For example, let $\tau_t = \sum_{i=0}^{\infty} b^{i+1}h_i(\overline{\Theta_{t+i}})$, where $|h_i(\overline{\Theta_1})| \leq 1$, $|b| \leq 1/2$ and where $\overline{\Theta_t}$ is an IID sequence independent of Θ_t.

For example, if $h_i(\overline{\Theta_1}) = a^i\Theta_1$, τ_t can be strong mixing or not. Then, one can prove MCLT even for functionals of some non strong mixing sequence.

Example 2.5 *: Assume that $k_i(\Psi_t) = i^{-5/2}g_i(\Theta_t).sin(e(i)\tau_t)$ where $E\{g_i(\Theta_1)\} = 0$ and $|e(i)| \leq 2\pi$ for all $i \in N$. Assume that $\sum_{s=0}^{\infty}\left|\sum_{i=s}^{\infty}k_{i+1}(\psi_1)\right| < C < \infty$.*

*Then, by lemma **??**, $\sigma(n) \to \infty$ and $sup|r_n| \to 0$ as $n \to \infty$. Moreover, $|\tau_t| \leq 1$ and $\tau_t = \tau_{t,u-t} + b^{u-t+1}\rho$ where $|\rho| \leq 1$ and $\tau_{t,u-t} = \sum_{i=0}^{u-t} b^{i+1}h_i(\overline{\Theta_{t+i}})$. Therefore, $sin(e(s)\tau_r) = sin(e(s)\tau_{r,u-r}) + 2\pi b^{u-r+1}\rho^*$ where $|\rho^*| \leq 1$.*

*Then, $M_u = \sigma(u)^{-1}\sum_{r=1}^{u}\sum_{s=1}^{r} s^{-5/2}g_s(\Theta_r)sin(e(s)\tau_{r,u-r}) + r^*_u$, where $Sup|r^*_u| \to 0$ as $n \to \infty$. Then, one applies theorem **??** with $v_u = -r_u - r^*_u$ and a similar decomposition for M'_u : $E\{(S_u + v_u)^p(S'_u + v'_u)^q\} = E\{(S_u + v_u)^p\}E\{(S'_u + v'_u)^q\}$. One deduces that $S_n \to N(0,1)$.*

By using theorem **??**, the MCLT can be proved for many other functionals, in particular , if $\{\Psi_t\}$ is a q-dependent sequence.

Accepted manuscript

ACCEPTED MANUSCRIPT

3 Proof of theorem ??

At first, for example, by Schwarz inequality, $E\{S_u\xi_u\}^2 \leq E\{S_u^2\}E\{\xi_u^2\} \leq E\{\xi_u^2\} = o(1)$ where "o" is the classical "o" : $o(1) \to 0$ as $n \to \infty$.

We deduce that $\sigma(n)^2 = \sigma(u)^2 E\{(S_u + v_u + S'_u + v'_u + \xi_u - v_u - v'_u)^2\} = \sigma(u)^2[2E\{(S_u)^2\} + o(1)]$. Then, for all $h \in N$, $\sigma(u)^h/\sigma(n)^h = 2^{-h/2} + o(1)$.

For all $p \in N$, we set $M_n^p = E\{(S_n)^p\}$.Clearly, $M_n^0 = 1$,$M_n^1 = 0$ and $M_n^2 = 1$.

Let $h \in N$, $2 \leq h \leq 2k$. Assume that, for all $p \in N$, $p \leq 2h-2$, $M_n^p \to \mu_p$, the moment or order p of N(0,1) .

By Holder inequality, $|E\{(S_u)^r(v_u)^{p-r}\}| \leq E\{|S_u|^p\}^{r/p}E\{|v_u|^p\}^{(p-r)/p}$ if $r < p$. Therefore, if $r < p \leq 2h-2$, $\left|E\{(S_u)^r(v_u)^{p-r}\}\right| = o(1)$. If $r < 2h$, $\left|E\{(S_u)^r(v_u)^{2h-r}\}\right| = M_s^{2h}o(1)$, where $M_s^{2h}(n) = sup\{M_{u(n)}^{2h}, 1\}$.

Then, by using H_{mI} and H_{mS} ,

$$E\{(S_u + v_u + S'_u + v'_u)^{2h}\}$$

$$= E\{(S_u+v_u)^{2h}\}+E\{(S'_u+v'_u)^{2h}\}+\sum_{m=1}^{2h-1}\frac{2h!}{m!(2h-m)!}E\{(S_u+v_u)^{2h-m}\}E\{(S'_u+v'_u)^m\}+o(1)$$

$$= 2E\{(S_u+v_u)^{2h}\} + \sum_{m=2}^{2h-2}\frac{2h!}{m!(2h-m)!}E\{(S_u+v_u)^{2h-m}\}E\{(S_u+v_u)^m\} + o(1)$$

$$= 2E\{S_u^{2h}\}+2\sum_{r=0}^{2h-1}\frac{2h!}{r!(2h-r)!}E\{S_u^r\}E\{v_u^{2h-r}\}+\sum_{m=2}^{2h-2}\frac{2h!}{m!(2h-m)!}E\{S_u^{2h-m}\}E\{S_u^m\}+o(1)$$

$$= 2M_u^{2h} + M_s^{2h}o(1) + \sum_{m=2}^{2h-2}\frac{2h!}{m!(2h-m)!}\mu_{2h-m}\mu_m + o(1)$$

$$\leq 2M_u^{2h} + M_s^{2h}|o(1)| + 2(2^{h-1}-1)\mu_{2h} + |o(1)|.$$

Moreover, if $m \geq 1$,

$$|E\{(S_u + v_u + S'_u + v'_u)^{2h-m}(\xi_u - v_u - v'_u)^m\}|$$

$$\leq |E\{(S_u + v_u + S'_u + v'_u)^{2h}\}|^{(2h-m)/(2h)}|E\{(\xi_u - v_u - v'_u)^{2h}\}|^{m/(2h)}$$

$$\leq |o(1)|\left|2M_s^{2h} + M_s^{2h}|o(1)| + 2(2^{h-1}-1)\mu_{2h} + |o(1)|\right|^{(2h-m)/(2h)}$$

$$\leq |o(1)|M_s^{2h} + |o(1)|.$$

Accepted manuscript

ACCEPTED MANUSCRIPT

Therefore,

$$M_n^{2h} = \frac{\sigma(u)^{2h}}{\sigma(n)^{2h}} E\{(S_u + v_u + S'_u + v'_u + \xi_u - v_u - v'_u)^{2h}\}$$

$$= (2^{-h} + o(1)) \sum_{m=0}^{2h} \frac{2h!}{m!(2h-m)!} E\{(S_u + v_u + S'_u + v'_u)^{2h-m} (\xi_u - v_u - v'_u)^m\}$$

$$\leq 2^{1-h} \big| M_s^{2h} + M_s^{2h} |o(1)| + (2^{h-1} - 1)\mu_{2h} + |o(1)| \big| + |o(1)| M_s^{2h} + |o(1)|$$

$$\leq 2^{1-h} \big| M_s^{2h} + (2^{h-1} - 1)\mu_{2h} \big| [1 + \epsilon(n)],$$

where $\epsilon(n) \to 0$ as $n \to \infty$.

The same equality has been obtained in the proof of [**?**] when $v_u = v'_u = 0$. We deduce by the same way that M_n^{2h} is bounded. Therefore, $M_n^{2h} = 2^{1-h}\big[M_u^{2h} + (2^{h-1} - 1)\mu_{2h}\big] + o(1)$. We deduce the convergence of M_n^{2h-1} and M_n^{2h} to μ_{2h-1} and μ_{2h}. $\bigstar$

References

[1] IBRAGIMOV I.A. LINNIK Yu V.(1971) Independent and stationary sequences of random variables. Wolters-Noordhoff, Groningen.

[2] BRADLEY R.C. (1984) On a very weak Bernouilli condition. Stochastics, 13, 61-81.

[3] DEHLING H. DENKER M. PHILLIPPS W. (1984) Versik Processes and very weak Bernouilli processes with summable rates are in,dependent. Proc. Amer. Math Soc. 91, 618-624.

[4] WITHERS C. S. (1981) Central limit theorems for dependent random variables. I. Z. Wahrsch. verw. Gebiete, 54, 509-534.

[5] COGBURN R. (1960) Asymptotic properties of stationary sequences. Univ. Calif. Publ. Statist. 3, 99-146.

[6] ROSENBLATT M. (1972) Uniform ergodicity and strong mixing. Z. Wahrsch. Werw. Gebiete. 24, 79-84.

[7] PINSKER M.S. (1964) Information and information stability of random variables and processes. Holden Day, San Francisco.

[8] BERNSTEIN (1939) Quelques remarques sur le thorme limite Liapounoff. Dokl. Akad, Nauk SSSR, 24, 3-8.

[9] BROWN (1970) Characteristics functions, moments and the Central Limit Theorem. Ann. Math. Statist. 41 658-664.

[10] ESSEEN C.G. JANSON S. (1985) On moments conditions for normed sums of independent random variables and martingales differences. Stochastics Processes and their Applications, 19, 173-182.

[11] HERRNDORF N. (1984) A functional Central Limit Theorem for r-mixing sequences. Journal of multivariate analysis. 15, 141-146.

Accepted manuscript

ACCEPTED MANUSCRIPT

[12] BIRKEL T. (1988) Moment bounds for associated sequences. Ann. Prob. 16-3, 1184-1193.

[13] KRUGOV V.M. (1988) The convergence of moments of random sums. Theory of Probability and its applications. 33-2, 339-342.

[14] MAIROBODA R. E. (1989) The Central limit Theorem for empirical moment generating functions. Theory of Probability and its applications, vol 34-2, 332-335.

[15] YOKOHAMA R. (1980) Moment bounds for stationary mixing sequences. Z Wahrscheinlichkeistheorie, ver. Gebiete, 52, 45-57.

[16] YOKOHAMA R. (1983) The convergence of moments in the Central limit Theorem for stationary f-mixing processes. Analysis mathematica, 9, 79-84.

[17] COX D. KIM T.Y. (1995) Moment bounds for mixing random variables useful in nonparametric function estimation. Stochastics Processes and their Applications. 56, 151-158.

[18] LANCASTER H. O. (1960) Orthogonal models for contingency tables. Developments in statistics. Academic Press, New York.

[19] BLACHER R. (1993) Higher Order Correlation Coefficients. Statistics 25, 1-15.

[20] BLACHER R. (1995) Central limit theorem by polynomial dependence coefficients. Journal of computational and applied mathematics 57, 45-56.

[21] BLACHER R. (1990) Theoreme de la limite centrale par les moments. Compte rendus de l'Acadmie des Sciences de Paris. t-311 srie I, p 465-468.

[22] IBRAGIMOV I. LIFSHITS M. (1998) On the convergence of generalized moments in almost sure central limit theorem. Stat and Prob Letters 343-351.

[23] SOULIER P. (2001) Moment bounds and the central limit theorem for functions of Gaussian vectors. Stat and Prob Letters 193-203.

[24] ROZOVSKY L.V. . (2002) An estimate of the remainder in the central limit theorem for a sum of independent random variables with infinite moments of a higher order. Theor Proba and its appl, 174-183.

Accepted manuscript