



HAL
open science

Synthesis and Characterization of 3,5-Diamino-1,2,4-triazolium Dinitramide

Jörg Stierstorfer, Thomas Klapoetke, Franz Martin, Norbert Mayr

► **To cite this version:**

Jörg Stierstorfer, Thomas Klapoetke, Franz Martin, Norbert Mayr. Synthesis and Characterization of 3,5-Diamino-1,2,4-triazolium Dinitramide. *Journal of Inorganic and General Chemistry / Zeitschrift für anorganische und allgemeine Chemie*, 2010, 636 (15), pp.2555. 10.1002/zaac.201000298 . hal-00580684

HAL Id: hal-00580684

<https://hal.science/hal-00580684>

Submitted on 29 Mar 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Synthesis and Characterization of 3,5-Diamino-1,2,4-
triazolium Dinitramide**

| | |
|-------------------------------|---|
| Journal: | <i>Zeitschrift für Anorganische und Allgemeine Chemie</i> |
| Manuscript ID: | zaac.201000298.R1 |
| Wiley - Manuscript type: | Article |
| Date Submitted by the Author: | 23-Aug-2010 |
| Complete List of Authors: | Stierstorfer, Jörg; LMU Munich, Chemistry Klapoetke, Thomas; LMU Munich, Chemistry Martin, Franz; LMU Munich, Chemistry Mayr, Norbert; LMU Munich, Chemistry |
| Keywords: | Energetic Materials, Azoles, Dinitramide, Sensitivities, Detonation parameters |
| | |

SCHOLARONE™
Manuscripts

ARTICLE

DOI: 10.1002/zaac.200((will be filled in by the editorial staff))

Synthesis and Characterization of 3,5-Diamino-1,2,4-triazolium Dinitramide

Thomas M. Klapötke,^{*[a]} Franz A. Martin,^[a] Norbert T. Mayr^[a] and Jörg Stierstorfer^[a]**Keywords:** Energetic Materials, Azoles, Dinitramide, Sensitivities, Detonation parameters.

3,5-Diamino-1,2,4-triazole (**1**, guanozol) was protonated with diluted hydrochloric acid, nitric acid, as well as perchloric acid forming 3,5-diamino-1,2,4-triazolium chloride hemihydrate (**2**), 3,5-diamino-1,2,4-triazolium nitrate (**3**) and 3,5-diamino-1,2,4-triazolium perchlorate (**4**), respectively. In a second step **4** was reacted with potassium dinitramide forming 3,5-diamino-1,2,4-triazolium dinitramide (**5**) and low soluble potassium perchlorate. **2–5** were characterized by low temperature single X-ray diffraction, IR and Raman as well as multinuclear NMR spectroscopy, mass spectrometry and differential scanning calorimetry.

The heat of formation of **1–5** were calculated by the CBS-4M method to be 81.1 (**1**), 124.7 (**2**), -76.1 (**3**), -25.2 (**4**) and 138.7 (**5**) kJ mol⁻¹. With these values as well as the X-ray densities several detonation parameters were calculated using both computer codes EXPLO5.03 and EXPLO5.04. In addition, the sensitivities of **1–5** were determined by the BAM drophammer and friction tester as well as a small scale electrical discharge device.

* Corresponding Author

Fax: +49 89 2180 77492

E-Mail: tmk@cup.uni-muenchen.de

[a] Ludwig-Maximilian University Munich, Energetic Materials Research, Department of Chemistry, Butenandtstr. 5-13, D-81377, Germany

dinitramide,^[13] 1,5-diaminotetrazolium dinitramide^[14] 1,4-dimethyl-5-aminotetrazolium dinitramide,^[15] 1,5-diamino-4-methyltetrazolium dinitramide^[16] are described as energetic materials in literature.^[17] In this work we present the synthesis and characterization of the new energetic compound 3,5-diamino-1,2,4-triazolium dinitramide (**5**) which is based on the commercially available guanozol (**1**).


Introduction

The development of azole based new energetic materials^[1] for civil and military applications is of great interest in many research programs worldwide due to their unique properties: shock waves producing pressure up to 500,000 times that of Earth's atmosphere, detonation waves travelling at 10 kilometers per second, temperatures soaring to 5500 Kelvin, and power approaching 20 billion watts per square centimetre. Advances in energetic materials, which include high explosives, propellants and pyrotechnics have been gained on nitrogen- and oxygen rich materials.^[2] Especially energetic ionic salts of tri- and tetrazoles have been utilized in energetic roles owing to their higher heats of formation, density and oxygen balance compared to those of their carbocyclic analogues. Azolium cations paired with nitrate, perchlorate, dinitramide, or picrate anions form highly energetic salts with mostly good thermal stabilities. With the exception of perchlorate they and their decomposition products are often environmentally benign. Probably most suitable as high explosives, gas generators or components in propellants or propellant charges^[3-6] are azolium salts containing the dinitramide anion, N(NO₂)₂⁻ (DN).^[7] They often show excellent oxygen balances by combining both the fuel (tetrazole heterocycle) and the oxidizer (dinitramide). Many N-rich dinitramides like guanidinium dinitramide,^[8] aminoguanidinium dinitromide,^[9] bisguanidinium dinitramide^[10], guanylurea dinitramide (FOX-12)^[11] and triaminoguanidinium dinitramide^[4,12] were synthesized and characterized as energetic materials. Also tetrazolium derivatives, e.g. 5-aminotetrazolium

Results and Discussion

Synthesis

3,5-Diamino-1,2,4-triazole (**1**) was protonated with diluted hydrochloric and nitric acid, forming the compounds 3,5-diamino-1,2,4-triazolium chloride hemihydrate (**2**) and 3,5-diamino-1,2,4-triazolium nitrate (**3**). **2** was then recrystallized from water and **3** from an ethanolic solution (water/ethanol: 1:1). The synthesis of 3,5-diamino-1,2,4-triazolium dinitramide (**5**) was performed according to Scheme 1 by the reaction of 3,5-diamino-1,2,4-triazolium perchlorate (**4**) with potassium dinitramide (KDN). Compound **4** was synthesized by protonation of **1** with an equimolar amount of 1N perchloric acid. After removing the solvent, single crystals of **4** were obtained by recrystallization from hot ethanol. Compound **4** is readily soluble in water, slightly soluble in MeOH, DMSO as well as DMF and slightly soluble in cold EtOH, acetone, diethyl ether or THF. In the second step, aqueous solutions of **4** and potassium dinitramide were combined forming **5** under precipitation of potassium perchlorate. The suspension was evaporated to dryness and extracted with ethanol yielding clean **5** after evaporation of the solvent. The solubility of **5** is comparable to that of **4**. Single crystals were obtained from a wet ethanolic solution.



Scheme 1 Protocol of the syntheses of 3,5-diamino-1,2,4-triazolium nitrate (2), chloride (3), perchlorate (4) and dinitramide (5).

NMR spectroscopy

The formation of the 3,5-diamino-1,2,4-1*H*-triazolium ions can easily be monitored and observed due to the differences in the NMR spectra. The neutral compound **1** shows two unique peaks for each of the amine groups at chemical shifts of 5.56 ppm and 4.68 ppm and one broad peak at a chemical shift of 10.68 ppm for the nitrogen bonded hydrogen atom (N1, heterocyclic ring). This can be explained with the slow exchange of the N-H hydrogen atom between nitrogen atoms N1 and N2 in DMSO-*d*₆ which generates two different chemical surroundings for each amine group. The same is observed for the carbon atoms. In the ¹³C-¹H NMR spectra we observed two peaks at chemical shifts of 161.8 ppm and 156.4 ppm, respectively.

In contradiction to the neutral compound the triazolium ions always show a different pattern in the NMR spectra. When protonated, only one signal is observed for the two amine groups, shifted well to lower field, at a chemical shift of 7.07 ppm (2), 6.97 ppm (3), 6.94 ppm (4) and 6.94 ppm (5), respectively. The two hydrogen atoms located on the heterocyclic ring show only one broad singlet signal at chemical shifts of 12.03 ppm (2), 12.08 ppm (3), 11.93 ppm (4) and 12.13 ppm (5). The same is observed in the ¹³C{¹H} NMR spectra where we observe only one signal for the two equivalent carbon atoms of the heterocyclic ring at a chemical shift of 151.4 ppm (2), 151.8 ppm (3), 151.8 ppm (4) and 152.0 ppm (5), shifted to higher fields compared with **1**.

In addition unique peaks for the nitrate and dinitramide anions are observed in the ¹⁴N NMR spectra of compounds **3** and **5** at chemical shifts of -4 ppm for compound **3** and -10 ppm for compound **5**, respectively.

Structures

The single crystal X-ray diffraction data of **1–5** were collected using an Oxford Xcalibur3 diffractometer with a

Spellman generator (voltage 50 kV, current 40 mA) and a KappaCCD detector. The data collection was undertaken using the CRYSLIS CCD software [18] and the data reduction was performed with the CRYSLIS RED software. [19] The structures were solved with SIR-92 [20] and refined with SHELXL-97 [21] implemented in the program package WinGX [22] and finally checked using PLATON. [23] Further information regarding the crystal-structure determination have been deposited with the Cambridge Crystallographic Data Centre [24] as supplementary publication Nos. 784960 (1), 784963 (2), 784962 (3), 784959 (4) and 784961 (5).

The structure of guanzol (**1**) was redetermined, since the solution published by G. L. Starova *et al.* [25] (i) has been measured at room temperature and (ii) was published in the non standard space group *P*2₁/*b*. The asymmetric unit is shown in Figure 1. The bond distances, angles and torsion angles agree with the previously measured structure.




Figure 1 Molecular moiety of **1**. Thermal ellipsoids represent the 50% probability level. Selected bond lengths (Å): N5–C2 1.375(2), N3–C1 1.334(3), N3–C2 1.354(2), N2–C2 1.324(6), N4–C1 1.350(3), N2–N1 1.395(3), N1–C1 1.336(4); selected bond angles (°): N3–C1–N1 110.27(11), C2–N2–N1 101.28(11), N3–C1–N4 125.44(12), C1–N1–N2 109.67(12), N1–C1–N4 124.18(14), N2–C2–N3 115.76(13), N2–C2–N5 122.62(12), C1–N3–C2 103.02(11), N3–C2–N5 121.44(13).

Protonation of **1** using diluted hydrochloric acid yields 3,5-diamino-1,2,4-triazolium chloride as its semihydrate, which crystallizes in the monoclinic space group *P*2₁/*c*. The asymmetric unit, four times in a unit cell, includes two moieties of cations and anions resulting in a density of 1.584 g cm⁻³. The chlorides participate in a strong hydrogen bond network interacting with all of the existing hydrogen atoms. The structure of the cation is influenced only very marginal by protonation at the nitrogen atom N3. With this, the angle C1–N3–C2 is slightly elongated to 106.6° (C3–N8–C4 106.4°; 1: C1–N3–C2 103.02(11)°).

3,5-Diamino-1,2,4-triazolium chloride hemihydrate (**2**) crystallizes in the monoclinic space group *P*2₁/*n* with eight molecular moieties in the unit cell. The asymmetric unit (Figure 2) consists of two anion/cation pairs and one crystal water. The density of 1.584 g cm⁻³ is in the range of other azole chlorides, e.g. 5-azido-3-amino-1,2,4-triazolium hydrochloride mono-hydrate (1.572 g cm⁻³). [26] The two independent chloride ligands are coordinated by six (C11) as well as four (C12) hydrogen atoms forming no regular polyhedron.

Table 1 Crystallographic data and parameters.

| | 1 | 2 | 3 | 4 | 5 |
|--|--|--|---|---|---|
| Formula | C ₂ H ₅ N ₅ | C ₄ H ₁₄ N ₁₀ Cl ₂ O | C ₂ H ₆ N ₆ O ₃ | C ₂ H ₆ N ₅ ClO ₄ | C ₂ H ₆ N ₈ O ₄ |
| FW [g mol ⁻¹] | 99.11 | 289.15 | 162.13 | 199.57 | 206.15 |
| Crystal system | monoclinic | monoclinic | monoclinic | triclinic | monoclinic |
| Space Group | <i>P</i> 2 ₁ / <i>c</i> (No. 14) | <i>P</i> 2 ₁ / <i>n</i> (No. 14) | <i>P</i> 2 ₁ / <i>c</i> (No. 14) | <i>P</i> -1 (No. 2) | <i>P</i> 2 ₁ / <i>n</i> (No. 14) |
| Color / Habit | colorless rods | colorless plates | colorless rods | colorless prismn | colorless rods |
| Size [mm] | 0.13 x 0.14 x 0.16 | 0.05 x 0.17 x 0.18 | 0.13 x 0.15 x 0.30 | 0.14 x 0.17 x 0.18 | 0.12 x 0.14 x 0.20 |
| <i>a</i> [Å] | 10.6366(6) | 6.0570(5) | 9.5615(7) | 5.3035(5) | 13.0772(7) |
| <i>b</i> [Å] | 4.3042(2) | 24.662(2) | 9.1082(6) | 7.6267(7) | 17.4883(5) |
| <i>c</i> [Å] | 10.8114(6) | 8.1341(6) | 7.4390(6) | 9.2813(8) | 14.1722(5) |
| α [°] | 90 | 90 | 90 | 74.399(8) | 90 |
| β [°] | 118.784(7) | 93.516(7) | 96.362(8) | 84.825(7) | 107.678(4) |
| γ [°] | 90 | 90 | 90 | 83.683(8) | 90 |
| <i>V</i> [Å ³] | 433.81(5) | 1212.78(16) | 643.86(8) | 358.67(6) | 3088.1(2) |
| <i>Z</i> | 4 | 4 | 4 | 2 | 16 |
| ρ_{calc} [g cm ⁻³] | 1.518 | 1.584 | 1.673 | 1.848 | 1.774 |
| μ [mm ⁻¹] | 0.114 | 0.541 | 0.150 | 0.520 | 0.162 |
| <i>F</i> (000) | 208 | 600 | 336 | 204 | 1696 |
| $\lambda_{\text{MoK}\alpha}$ [Å] | 0.71073 | 0.71073 | 0.71073 | 0.71073 | 0.71073 |
| <i>T</i> [K] | 200 | 200 | 200 | 200 | 200 |
| Theta Min-Max [°] | 3.8, 26.0 | 4.2, 26.0 | 4.3, 26.0 | 3.9, 26.0 | 3.7, 26.0 |
| Dataset | -12:13; -5:5; -13:8 | -7:4; -30:26; -9:10 | -11:11; -11:7; 9:9 | -6:6; -9:9; -11:11 | -16:16; -21:21; - 17:17 |
| Reflections collected | 2134 | 4625 | 3272 | 3539 | 30659 |
| Independent reflections | 852 | 2375 | 1264 | 1419 | 6040 |
| <i>R</i> _{int} | 0.019 | 0.038 | 0.033 | 0.023 | 0.026 |
| Observed reflections | 677 | 1518 | 1264 | 1011 | 4065 |
| No. parameters | 84 | 106 | 124 | 133 | 602 |
| <i>R</i> ₁ (obs) | 0.0315 | 0.0350 | 0.0276 | 0.0327 | 0.0319 |
| w <i>R</i> ₂ (all data) | 0.0862 | 0.0623 | 0.0639 | 0.0885 | 0.0984 |
| <i>S</i> | 1.04 | 0.82 | 0.92 | 1.00 | 1.03 |
| Resd. Dens. [e Å ⁻³] | -0.22, 0.14 | -0.29, 0.31 | -0.16, 0.15 | -0.28, 0.27 | -0.31, 0.40 |
| Device type | Oxford Xcalibur3 CCD | Oxford Xcalibur3 CCD | Oxford Xcalibur3 CCD | Oxford Xcalibur3 CCD | Oxford Xcalibur3 CCD |
| Solution | SIR-92 | SIR-92 | SIR-92 | SIR-92 | SIR-92 |
| Refinement | SHELXL-97 | SHELXL-97 | SHELXL-97 | SHELXL-97 | SHELXL-97 |
| Absorption correction | multi-scan | multi-scan | multi-scan | multi-scan | multi-scan |
| CCDC | 784960 | 784963 | 784962 | 784959 | 784961 |




Figure 2 Molecular moiety of **2**. Thermal ellipsoids represent the 50% probability level. Selected bond lengths (Å): N1–C1 1.3173, N1–N2 1.3966, N2–C2 1.3074, N3–C1 1.3545, N3–C2 1.3725, N4–C1 1.3229, N5–C2 1.3532, N6–C3 1.3187, N6–N7 1.3948, N7–C4 1.3131, N8–C3 1.3445, N8–C4 1.3781, N9–C3 1.3326, N10–C4 1.3351.

3,5-Diamino-1,2,4-triazolium nitrate (**3**) crystallizes in the monoclinic space group $P2_1/c$ with four molecular moieties in the unit cell. The density of 1.672 g cm^{-3} is in the range of other azole nitrate derivatives, e.g. 5-aminotetrazolium nitrate.^[27]




Figure 3 Molecular moiety of **3**. Thermal ellipsoids represent the 50% probability level. Selected bond lengths (Å): N1–C2 1.317(2), N1–N2 1.401(6), N5–C1 1.338(5), N2–C1 1.305(2), N3–C2 1.348(4), N3–C1 1.375(4), O1–N6 1.257(5), O2–N6 1.240(2), N4–C2 1.318(5), O3–N6 1.257(3); selected bond angles (°): C2–N1–N2 111.75(11), N2–C1–N5 125.99(13), C1–N2–N1 103.49(11), N2–C1–N3 111.23(12), N4–C2–N3 124.93(12), O2–N6–O1 120.92(11), O2–N6–O3 120.13(11), O1–N6–O3 118.95(10).

In the structure of **3** layers are formed, which are formed by a strong hydrogen-bond network. All of the oxygen atoms of the nitrate anions participate in strong hydrogen bonds listed in Table 2.

Table 2 Hydrogen bonds based on the nitrate anions in the structure of **3**.

| Atoms D,H,A | Dist. D,H [Å] | Dist. H,A [Å] | Dist. D,A [Å] | Angle D,H,A [°] |
|------------------------------|------------------|------------------|---------------|--------------------|
| N3–H3– O1 ⁱ | 0.891(15) | 1.945(16) | 2.8362(15) | 178.1(12) |
| N4– H4A–O3 ⁱ | 0.869(18) | 2.123(18) | 2.9549(17) | 159.9(14) |
| N5–H5B– O1 ⁱⁱⁱ | 0.843(18) | 2.094(19) | 2.9350(18) | 175.9(16) |
| N4–H4B– O3 ^{iv} | 0.867(17) | 2.057(17) | 2.9038(17) | 165.3(15) |
| N4–H4B– O2 ^{iv} | 0.867(17) | 2.643(16) | 3.0789(16) | 112.3(13) |
| N1–H1– O2 ^v | 0.874(17) | 2.048(17) | 2.9190(16) | 175.3(15) |

(i) $x, 0.5-y, 0.5+z$; (ii) $-x, -0.5+y, 0.5-z$; (iii) $-x, 1-y, -z$; (iv) $1-x, 1-y, 1-z$; (v) $x, 1.5-y, 0.5+z$.

By analyzing the graph sets^[28] with the software RPLUTO^[29] several chain (**C1,1(6)**, **C1,1(4)** and **C2,2(5)**) and ring motifs (e.g. **R2,2(8)**) can be found within the layers, which are illustrated in Figure 4.




Figure 4 Hydrogen bonding within the layers in the structure of **3**. Symmetry codes: (i) $-x, 1-y, -z$; (ii) $x, 1.5-y, -0.5+z$; (iii) $1-x, 1-y, 1-z$; (iv) $x, 0.5-y, -0.5+z$.

3,5-Diamino-1,2,4-triazolium perchlorate (**4**) crystallizes in the triclinic space group $P-1$ with two molecular moieties in the unit cell. The density of 1.848 g cm^{-3} is in the range of other azole perchlorate derivatives, e.g. 5-aminotetrazolium perchlorate.^[30] The molecular structure (Figure 5) of the cation is in accordance to that in **2** and **3**. The perchlorate anions show a regular tetrahedral structure (all O–Cl–O angles between 109° and 111°).




Figure 5 Molecular moiety of **5**. Thermal ellipsoids represent the 50% probability level. Selected bond lengths (Å): Cl–O4 1.431(6), N5–C2 1.354(8), Cl–O2 1.431(6), Cl–O3 1.440(5), Cl–O1 1.440(6), N2–C1 1.318(6), N3–C2 1.296(5), N3–N2 1.399(8), N4–C1 1.329(6), N1–C1 1.341(7), N1–C2 1.380(6); selected bond angles (°): C1–N2–N3 111.81(18), C2–N3–N2 103.62(20), C1–N1–C2 107.48(19), N2–C1–N4 127.29(20), N2–C1–N1 105.98(20), N3–C2–N5 126.72(23), N3–C2–N1 111.10(21).

Again a layer-like structure is formed in which three of the oxygen atoms of the perchlorate anions participate in hydrogen bonds. The fourth oxygen atom is directed alternating to both the upper and lower layers. Analyzing the graph sets yields several chain and ring motives, some of them marked in Figure 6.




Figure 6 Hydrogen bonding within the layers in the structure of **4**. Symmetry codes: (i) $-1-x, -y, 1-z$; (ii) $-1+x, -1+y, z$; (iii) $-2-x, 1-y, -z$.

3,5-Diamino-1,2,4-triazolium dinitramide (**5**) crystallizes in the monoclinic space group $P2_1/n$ with 16 (!) molecular moieties in the unit cell. For better clearness only one of the four independent molecular units is shown in Figure 7. The density observed is 1.774 g cm^{-3} , which is comparable to

that of 1,5-diaminotetrazolium dinitramide (1.771 g cm^{-3}) described in literature.^[14]




Figure 7 Molecular moiety of **5**. The ellipsoids represent the 50% probability level. Selected bond lengths (Å): O2–N8 1.240(4), O1–N8 1.251(4), N7–N8 1.354(4), N7–N6 1.368(4), O3–N6 1.238(4), O4–N6 1.218(4), N1–C1 1.317(4), N2–C2 1.308(5), N2–N1 1.403(4), N4–C1 1.313(5), C1–N3 1.358(5), N3–C2 1.367(5), C2–N5 1.343(5).

Compound **5** crystallizes in accordance to **3** and **4** forming a layer-like structure. Except for the outer nitrogen atoms in the dinitramide anions all atoms participate in hydrogen bonds, shown in Figure 8. Due to that fact, several remarkable also bifurcated hydrogen-bonds are formed.




Figure 8 Hydrogen bonding of one dinitramide anion in the structure of **5**.

Heat of Formation

Due to the highly energetic character of **3** and **5**, bomb calorimetric measurements could only performed with small amounts, consequently doubtful combustion energies were obtained. Therefore an extensive computational study was accomplished for **1–5**, which is presented in the following.

All calculations were carried out using the Gaussian G03W (revision B.03) program package.^[31] The enthalpies (H) and free energies (G) were calculated using the complete basis set (CBS) method of Petersson and coworkers in order to obtain very accurate energies. The CBS models use the known asymptotic convergence of pair natural orbital expressions to extrapolate from calculations using a finite basis set to the estimated complete basis set limit. CBS-4 begins with a HF/3-21G(d) geometry optimization; the zero point energy is computed at the same level. It then uses a large basis set SCF calculation as a base energy, and a MP2/6-31+G calculation with a CBS extrapolation to correct the energy through second order. A MP4(SDQ)/6-31+(d,p) calculation is used to approximate higher order contributions. In this study we applied the modified CBS-4M method (M referring to the use of Minimal Population localization) which is a re-parametrized version of the original CBS-4 method and also includes some additional empirical corrections.^[32] The enthalpies of the gas-phase species M were computed according to the atomization energy method (eq. 1) (Tables 3–5).^[33]

$$\Delta_f H^\circ_{(g, M, 298)} = H_{(Molecule, 298)} - \sum H^\circ_{(Atoms, 298)} + \sum \Delta_f H^\circ_{(Atoms, 298)} \quad (1)$$

Table 3. CBS-4M results

| | point group | el. state | $-H^{298}$ / a.u. | NIMAG |
|------------------------------------|-------------|------------|-------------------|-------|
| DATr (1) | C_s | | 352.484409 | 0 |
| HDATr⁺ | C_s | 1A_1 | 352.839733 | 0 |
| Cl⁻ | K_h | | 459.809901 | 0 |
| NO₃⁻ | D_{3h} | 1A | 280.080446 | 0 |
| ClO₄⁻ | T_d | | 760.171182 | 0 |
| DN⁻ | C_2 | | 464.499549 | 0 |
| H | | $^2A_{1g}$ | 0.500991 | 0 |
| C | | | 37.786156 | 0 |
| N | | $^4A_{1g}$ | 54.522462 | 0 |
| O | | | 74.991202 | 0 |
| Cl | | | 459.674576 | 0 |

Table 4 Literature values for atomic $\Delta_f H^\circ_{f, 298}$ / kcal mol⁻¹

| | | NIST ^[34] |
|----|--|----------------------|
| H | | 52.1 |
| C | | 171.3 |
| N | | 113.0 |
| O | | 59.6 |
| Cl | | 29.0 |

Table 5 Enthalpies of the gas-phase species M.

| M | M | $\Delta_f H^\circ_{(g, M)}$ / kcal mol ⁻¹ |
|-----------------------------------|---|--|
| DATr (1) | C ₂ H ₅ N ₅ | 41.8 |
| HDATr⁺ | C ₂ H ₆ N ₅ ⁺ | 185.3 |
| Cl⁻ | Cl ⁻ | -55.9 |
| NO₃⁻ | NO ₃ ⁻ | -74.9 |

| | | |
|------------------------------------|--|-------|
| ClO₄⁻ | ClO ₄ ⁻ | -66.1 |
| DN⁻ | N ₃ O ₄ ⁻ | -29.6 |

The solid state energy of formation (Table 7) of **DATr** was calculated by subtracting the gas-phase enthalpy with the heat of sublimation (22.5 kcal mol⁻¹) obtained by the TROUTMAN rule ($\Delta H_{\text{sub}} = 188 \cdot T_m$) ($T_m = 204$ °C).^[35] In the case of **ADNQ**, the lattice energy (U_L) and lattice enthalpy (ΔH_L) were calculated from the corresponding molecular volumes (Table 9) according to the equations provided by Jenkins *et al.*^[36] With the calculated lattice enthalpy (Table 6) the gas-phase enthalpy of formation (Table 5) was converted into the solid state (standard conditions) enthalpy of formation. These molar standard enthalpies of formation (ΔH_m) were used to calculate the molar solid state energies of formation (ΔU_m) according to equation 2 (Table 7).

$$\Delta U_m = \Delta H_m - \Delta n RT \quad (2)$$

(Δn being the change of moles of gaseous components)

Table 6 Lattice energies and lattice enthalpies.

| | V_{M_3} / nm ³ | U_L / mol ⁻¹ / kJ | ΔH_L / mol ⁻¹ / kJ | ΔH_L / mol ⁻¹ / kcal |
|----------|-----------------------------|--------------------------------|---------------------------------------|---|
| 2 | 0.151 | 543.8 | 545.8 | 130.3 |
| 3 | 0.161 | 535.0 | 538.5 | 128.6 |
| 4 | 0.179 | 520.1 | 523.6 | 125.0 |
| 5 | 0.193 | 509.8 | 513.2 | 122.6 |

Table 7 Solid state energies of formation ($\Delta_f U^\circ$)

| | $\Delta_f H^\circ(s)$ / kcal mol ⁻¹ | $\Delta_f H^\circ(s)$ / kJ mol ⁻¹ | Δn | $\Delta_f U^\circ(s)$ / kJ mol ⁻¹ | M / g mol ⁻¹ | $\Delta_f U^\circ(s)$ / kJ kg ⁻¹ |
|----------|--|--|------------|--|-------------------------|---|
| 1 | 19.4 | 81.1 | 5 | 93.4 | 99.12 | 942.8 |
| 2 | -29.8 | -124.7 | 6.5 | -108.6 | 144.57 | -751.0 |
| 3 | -18.2 | -76.1 | 7.5 | -57.6 | 162.14 | -354.9 |
| 4 | -6.0 | -25.2 | 8 | -5.4 | 199.58 | -26.8 |
| 5 | 33.1 | 138.7 | 9 | 161.0 | 206.16 | 781.1 |

Detonation parameters

The calculation of the detonation parameters was performed with the program package EXPLO5 (version 5.03 and 5.04).^[37] The program is based on the chemical equilibrium, steady-state model of detonation. It uses the Becker-Kistiakowsky-Wilson's equation of state (BKW EOS) for gaseous detonation products and Cowan-Fickett's equation of state for solid carbon. The calculation of the equilibrium composition of the detonation products is done by applying modified White, Johnson and Dantzig's free energy minimization technique. The program is designed to enable the calculation of detonation parameters at the CJ point. The BKW equation in the following form was used with the BKWN set of parameters (α , β , κ , θ) as stated below the equations and X_i being the mol fraction of i -th gaseous product, k_i is the molar covolume of the i -th gaseous product^[38]:

$$pV/RT = 1 + xe^{\beta x} \quad x = (\kappa \sum X_i k_i) / [V(T + \theta)]^\alpha$$

$$\alpha = 0.5, \beta = 0.176, \kappa = 14.71, \theta = 6620.$$

The detonation parameters calculated with the EXPLO5 versions V5.03 and V5.04 using the experimentally

determined densities (X-ray) are summarized in Table 8. It is not possible to calculate detonation parameters of compounds containing chlorine; therefore, the corresponding cells have been abandoned.

Table 8 Physico-chemical properties of **1–5** in comparison with trinitrotoluene (**TNT**), nitropenta (**PETN**) and hexogen (**RDX**).

| | 1 | 2 | 3 | 4 | 5 | TNT* | PETN* | RDX* |
|--|--|--|---|---|---|---|--|---|
| Formula | C ₂ H ₅ N ₅ | C ₂ H ₇ N ₅ ClO ₁ <i>12</i> | C ₂ H ₆ N ₆ O ₃ | C ₂ H ₆ ClN ₅ O ₄ | C ₂ H ₆ N ₈ O ₄ | C ₇ H ₅ N ₃ O ₆ | C ₅ H ₈ N ₄ O ₁₂ | C ₃ H ₆ N ₆ O ₇ |
| Molecular Mass [g mol ⁻¹] | 99.12 | 144.57 | 162.14 | 199.58 | 206.12 | 227.13 | 316.14 | 222.12 |
| Impact sensitivity [J] ^a | > 40 | > 40 | 40 | 25 | > 3 | 15 | 3 | 7.5 |
| Friction sensitivity [N] ^b | > 360 | > 360 | 288 | 240 | 192 | 353 | 60 | 120 |
| ESD–test [J] | > 5 | > 10 | 0.3 | 0.8 | 0.3 | n.d. | n.d. | 0.1 - 0.2 |
| <i>N</i> [%] ^c | 70.67 | 48.44 | 51.84 | 35.10 | 54.36 | 18.50 | 17.72 | 37.8 |
| Ω [%] ^d | -104.9 | -71.91 | -39.48 | -20.04 | -23.28 | -73.96 | -10.1 | -21.6 |
| <i>T</i> _{dec.} [°C] ^e | 204 (mp) | 282 | 276 | 252 | 164 | >160 | 202 | 210 |
| ρ [g cm ⁻³] ^f | 1.520 | 1.584 | 1.672 | 1.848 | 1.774 | 1.654 | 1.778 | 1.800 |
| $\Delta_f H_m^\circ$ [kJ mol ⁻¹] ^g | 81.1 | -124.7 | -76.1 | -25.2 | 138.7 | -59.1 | -539.0 | 70 |
| $\Delta_f U^\circ$ [kJ kg ⁻¹] ^h | 942.8 | -751.0 | -354.9 | -33.1 | 781.1 | -184.9 | -1611.7 | 417 |
| EXPLO5 values: | | | | | | | | |
| V5.03 (V5.04) | | | | | | | | |
| $-\Delta_f U^\circ$ [kJ kg ⁻¹] ⁱ | 1853 (1674) | --- | 4052 (4060) | --- | 5101 (5157) | 5112 (5227) | 5979 (6190) | 6038 (6125) |
| <i>T</i> _E [K] ^j | 1152 (1561) | --- | 3104 (2996) | --- | 3810 (3705) | 3756 (3657) | 4423 (4306) | 4368 (4236) |
| <i>p</i> _{C-J} [kbar] ^k | 113 (134) | --- | 233 (246) | --- | 302 (321) | 205 (216) | 321 (320) | 341 (349) |
| <i>V</i> _{Det.} [m s ⁻¹] ^l | 6058 (6516) | --- | 7789 (7927) | --- | 8624 (8681) | 7178 (7253) | 8665 (8320) | 8906 (8748) |
| Gas vol. [L kg ⁻¹] ^m | 847 (758) | --- | 833 (826) | --- | 833 (810) | 617 (574) | 765 (688) | 793 (739) |

[a] BAM drophammer, grain size (75–150 μm); [b] BAM friction tester, grain size (75–150 μm); [c] Nitrogen content; [d] Oxygen balance [39]; [e] Temperature of decomposition by DSC ($\beta = 5^\circ\text{C}$); [f] X-ray structure; [g] Molar enthalpy of formation; [h] Energy of formation; [i] Energy of Explosion; [j] Explosion temperature; [k] Detonation pressure; [l] Detonation velocity; [m] Assuming only gaseous products; * values based on Ref. [40] and the EXPLO5 database; n.d.: not determined.

Especially the detonation parameters of compound **5** show promising values, higher than those of trinitrotoluene (**TNT**) and in the range of those of pentaerythryl tetranitrate (**PETN**). Also compound **3** exceeds the values of TNT in addition to its great thermal stability of 276 °C and low sensitivities (Table 8). The most important criteria of high explosives are the detonation velocity ($V_{\text{Det.}} = \mathbf{3}: 7927, \mathbf{5}: 8681, \text{TNT}: 7253, \text{PETN}: 8320, \text{RDX}: 8748 \text{ m s}^{-1}$), the detonation pressure ($p_{\text{det.}} = \mathbf{3}: 246, \mathbf{5}: 321, \text{TNT}: 216, \text{PETN}: 320, \text{RDX}: 349 \text{ kbar}$) and the energy of explosion ($\Delta_f U^\circ = \mathbf{3}: -4060, \mathbf{5}: -5157, \text{TNT}: 5227, \text{PETN}: 6190, \text{RDX}: 6125 \text{ kJ kg}^{-1}$).

For application of new energetic compounds important values for safety, handling and processing are the sensitivity data. All values were determined according to BAM standard methods described in the NATO STANAG 4487, 4489 and 4490 specifications for energetic materials.^[41–47]

Unfortunately, well performing **5** is very sensitive towards impact (> 3 J) which is slightly below the commercial available PETN (3 J) and RDX (7.5 J). However, it is not as sensitive in terms of friction (**5**: > 192 N, RDX: > 120 N) and electrostatic discharge (**5**: > 0.3 J, RDX: > 0.1–0.2 J). The latter values are most important for handling and processing on an industrial scale and therefore compound **5** could be considered for application.

The most promising compound regarding to the sensitivity data is compound **3** which shows rather low sensitivity to friction (288 N), to impact (40 J) and to electrostatic discharge (0.3 J). According to the UN Recommendations on the Transport of Dangerous Goods compound **3** is regarded insensitive.^[48,49]

Thermostability of energetic compounds is considered important especially in processing and storing the material. Because of the diverse use of energetic materials e.g. under extreme climatic conditions like in deserts, for oil drilling or military ammunition high temperature stability is desired. The introduced compounds **3** and **5** display good values for melting (**3**: 159 °C, **5**: 155 °C) and decomposition (**3**: 276 °C, **5**: 164 °C). Compound **5** melts at 155 °C and decomposes subsequently at 164 °C, whereas compound **3** shows a melting point at 159 °C followed by a wide liquid range of 110 °C and decomposes finally at 276 °C. The high difference between melting and decomposition point leaves compound **3** very suitable as melt cast explosive. Comparing compounds **3** and **5** to common explosives, they exceed both TNT (m.p. 80 °C, $T_{dec.} > 160$ °C) and compound **3** passes even RDX (m.p./ $T_{dec.}$ 210 °C) and PETN (m.p. 140 °C, $T_{dec.}$ 200 °C), regarding the thermal stability against decomposition.

Conclusions

From this combined experimental and theoretical study the following conclusions have been drawn:

- 3,5-Diamino-1,2,4-triazole (**1**) can be protonated using diluted mineralic acids such as hydrochloric, nitric and perchloric acid. The hydrodinitramide salt (**5**) of **1** can be synthesized by metathesis reaction of the perchlorate salt (**4**) with potassium dinitramide. The obtained compounds 3,5-diamino-1,2,4-triazolium chloride hemihydrate (**2**), 3,5-diamino-1,2,4-triazolium nitrate (**3**) and 3,5-diamino-1,2,4-triazolium perchlorate (**4**), and 3,5-diamino-1,2,4-triazolium dinitramide (**5**) are air-stable, not hygroscopic colorless solids.

- All salts were recrystallized from ethanol/water mixtures yielding single crystals, which were analyzed by X-ray diffraction. The salts crystallizes in common space groups (**2**, **5**: $P2_1/n$, **3**: $P2_1/c$, **4**: $P-1$). All structures are dominated by a strong hydrogen-bond network.

- The energetic properties of **1–5** were determined and compared with trinitrotoluene (TNT), pentaerythryl tetranitrate (PETN) and hexogen (RDX). The sensitivities towards impact, friction and electrostatic discharge were measured. The sensitivities escalate from **1** to **5**. The dinitramide salt **5** is very sensitive towards impact (3 J) and moderately towards friction (192 N).

- The detonation parameters of **1**, **3**, and **5** were calculated with the computer codes EXPLO5.03 and EXPLO5.04, respectively. The detonation velocities and pressures of **3** and **5** are greater than those of TNT and in the case of **5** in the range of PETN.

Experimental Part

Caution: Although all 3,5-diamino-1,2,4-triazolium salts reported in this publication are rather stable against friction, impact and electric discharge, proper safety precautions should be taken when handling dinitramide salts. The derivatives are energetic materials and tend to explode under certain conditions, especially under physical stress. Laboratories and personnel should be properly grounded, and safety equipment such as Kevlar gloves, leather coats, face shields and ear plugs are recommended.

General. All chemical reagents, except 3,5-diamino-1,2,4-*1H*-triazole and ammonium dinitramide, and solvents were obtained from Sigma-Aldrich Inc. or Acros Organics (analytical grade) and were used as supplied. 3,5-diamino-1,2,4-*1H*-triazole was obtained from ABCR and ammonium dinitramide was supplied by EURENCO Bofors AB. Potassium dinitramide was prepared from ammonium dinitramide following known literature procedures[Error! Bookmark not defined.]. ^1H , $^{13}\text{C}\{^1\text{H}\}$, and ^{14}N NMR spectra were recorded on a JEOL Eclipse 400 instrument in DMSO- d_6 at or near 25 °C. The chemical shifts are given relative to tetramethylsilane (^1H , ^{13}C) or nitromethane (^{14}N) as external standards and coupling constants are given in Hertz (Hz). Infrared (IR) spectra were recorded on a Perkin-Elmer Spectrum BX FT-IR instrument equipped with an ATR unit at 25 °C. Transmittance values are qualitatively described as “very strong” (vs), “strong” (s), “medium” (m) and “weak” (w). Raman spectra were recorded on a Bruker RAM II spectrometer equipped with a Nd:YAG laser (1064 nm) and a reflection angle of 180°. The intensities are reported as percentages of the most intense peak and are given in parentheses. Elemental analyses were performed with a Netsch Simultaneous Thermal Analyzer STA 429. Melting points were determined by differential scanning calorimetry (Setaram DSC141 instrument, calibrated with standard pure indium and zinc). Measurements were performed at a heating rate of 5 °C min⁻¹ in closed aluminum sample pans with a 1 μm hole in the top for gas release under a nitrogen flow of 20 mL/min with an empty identical aluminum sample pan as a reference. For initial safety testing, the impact and friction sensitivities as well as the electrostatic sensitivity were determined.^[48] The impact sensitivity tests were carried out according to STANAG 4489^[41] modified according to instruction^[44] using a BAM^[46] drophammer. The friction sensitivity tests were carried out according to STANAG 4487^[42] modified according to instruction^[45] using the BAM friction tester. The electrostatic sensitivity tests were carried according to STANAG 4490^[43] out using an electric spark testing device ESD 2010EN (OZM Research) operating with the “Winspark 1.15 software package”.^[48]

3,5-Diamino-1,2,4-triazole (**1**)

As obtained from ABCR.

DSC (5 °C min⁻¹): 205 °C (mp., onset); ^1H NMR: δ (ppm) = 10.68 (s, 1H, NH), 5.56 (s, 2H, NH₂), 4.68 (s, 2H, NH₂); $^{13}\text{C}\{^1\text{H}\}$ NMR (D₆-DMSO): δ (ppm) = 161.8, 156.4; m/z (DEI) = 99.09 ([M]⁺); **IS**: > 40 J; **FS**: > 360 N; **ESD**: > 1.5 J.

3,5-Diamino-1,2,4-triazolium chloride hemihydrate (**2**)

0.99 g (10 mmol) of 3,5-diamino-1,2,4-*1H*-triazole was added with stirring to 10 mL (10 mmol) of 1 M hydrochloric acid solution. The

1
2 resulting solution was heated slightly to 50 °C and was kept at this
3 temperature for 10 minutes. After cooling to room temperature the
4 solvent was evaporated completely and the white residue was
5 recrystallized from water, yielding 1.34 g (99%) of pure **2** as white
6 crystalline needles.

7 **DSC** (5 °C min⁻¹): 105 °C (mp., onset), 145 °C (-H₂O, onset),
8 282 °C (dec., onset); **IR** (ATR): 3328 (s), 3224 (s), 3157 (vs), 2899
9 (m), 2651 (m), 1690 (m), 1657 (s), 1651 (ms), 1646 (m), 1635 (m),
10 1352 (w), 1297 (w), 1160 (w), 1059 (w), 1011 (w), 794 (w),
11 711(w); **Raman** (1064 nm, 25 °C): 3322 (17), 3229 (24), 3181 (22),
12 1688 (75), 1657 (31), 1593 (15), 1568 (18), 1540 (12), 1455 (36),
13 1333 (26), 1305 (17), 1166 (28), 1071 (100), 1051 (65), 797 (50),
14 724 (16), 664 (74), 500 (33), 355 (42); **¹H NMR**: δ (ppm) = 12.03
15 (s, br, 2H, NH), 7.07 (s, 4H, NH₂); **¹³C{¹H} NMR**: δ (ppm) =
16 151.4; **MS**: FAB⁺ = 100.12 ([M]⁺), FAB⁻ = 34.95 ([Cl]⁻); **EA**:
17 calcd.: C 16.62, H 4.88, N 48.45; found: C 16.74, H 4.58, N 48.21;
18 **IS**: > 40 J; **FS**: 360 N; **ESD**: > 1.5 J.

20 3,5-Diamino-1,2,4-triazolium nitrate (**3**)

21 0.99 g (10 mmol) of 3,5-diamino-1,2,4-*H*-triazole was added with
22 stirring to 5 mL (10 mmol) of 2 M nitric acid solution. The
23 resulting solution was heated slightly to 50 °C and was kept at this
24 temperature for 10 minutes. After cooling to room temperature the
25 solvent was reduced to about one third of its original volume and
26 the slightly yellow solution was left standing for crystallization. **3**
27 separated from the solution over the course of one week, left
28 standing on air, yielding 1.46 g (91%). Crystals of **3** suitable for X-
29 ray diffraction measurements were obtained by recrystallization
30 from a water/ethanol mixture (1:1).

31 **DSC** (5 °C min⁻¹): 159 °C (mp., onset), 276 °C (dec., onset); **IR**
32 (ATR): 3792 (vw), 3423 (m), 3326 (m), 3282 (m), 3166 (m), 2936
33 (m), 2792 (m), 2688 (m), 1680 (s), 1662 (vs), 1612 (m), 1537 (w),
34 1425 (m), 1321 (s), 1297 (s), 1040 (w), 1010 (w), 812 (w), 794 (w),
35 716 (w), 657 (w); **Raman** (1064 nm, 25 °C): 3274 (7), 3175 (14),
36 1705 (20), 1689 (37), 1660 (22), 1601 (20), 1551 (7), 1468 (18),
37 1418 (11), 1354 (8), 1172 (16), 1076 (77), 1053 (100), 1042 (95),
38 1016 (51), 796 (32), 718 (26), 665 (65), 499 (15), 347 (26); **¹H**
39 **NMR**: δ (ppm) = 12.08 (s, br, 2H, NH), 6.97 (s, 4H, NH₂); **¹³C{¹H}**
40 **NMR**: δ (ppm) = 151.8; **¹⁴N NMR**: δ (ppm) = -4 (NO₃⁻); **MS**:
41 FAB⁺ = 100.09 ([M]⁺), FAB⁻ = 62.01 ([NO₃]⁻); **EA**: calcd.: C 14.82,
42 H 3.73, N 51.84; found: C 15.07, H 3.57, N 51.64; **IS**: 40 J; **FS**:
43 288 N; **ESD**: 0.3 J.

46 3,5-Diamino-1,2,4-triazolium perchlorate (**4**)

47 0.99 g (10 mmol) of 3,5-diamino-1,2,4-*H*-triazole was added with
48 stirring to 10 mL (10 mmol) of 1 M perchloric acid solution. After
49 complete dissolution of **1**, the solvent was evaporated to dryness
50 and the remaining white residue was recrystallized from hot ethanol,
51 yielding 1.95 g (98%) of pure **4**. Crystals suitable for X-
52 ray diffraction measurements were also obtained from hot ethanol.
53 **DSC** (5 °C min⁻¹): 185 °C (mp., onset), 252 °C, 294 °C, 335 °C
54 (dec., onset); **IR** (ATR): 3561 (w), 3468 (m), 3455 (m), 3408 (s),
55 3366 (s), 3276 (m), 3216 (s), 3178 (s), 1672 (s), 1601 (m), 1541
56 (m), 1456 (m), 1344 (w), 1079 (s), 1050 (s), 1011 (m), 932 (w),
57 667 (w), 621 (w); **Raman** (1064 nm, 25 °C): 3460 (2), 3414 (4),
58 3369 (5), 3217 (6), 3176 (5), 1698 (18), 1664 (13), 1600 (14), 1458
59 (18), 1342 (7), 1170 (11), 1063 (26), 1016 (10), 935 (100), 797 (20),
60 667 (30), 630 (23), 491 (14), 470 (17), 453 (24), 343 (16); **¹H**
NMR: δ (ppm) = 11.93 (s, br, 2H, NH), 6.94 (s, 4H, NH₂); **¹³C{¹H}**
NMR: δ (ppm) = 151.8; **MS**: FAB⁺ = 100.10 ([M]⁺), FAB⁻ = 99.00

([ClO₄]⁻); **EA**: calcd.: C 12.04, H 3.03, N 35.10; found: C 12.00, H
3.13, N 35.34; **IS**: 25 J; **FS**: 240 N; **ESD**: 0.8 J.

3,5-Diamino-1,2,4-triazolium dinitramide (**5**)

Aqueous solutions of 0.997 g (5 mmol) of **4** in 3 mL of water and
0.725 g (5 mmol) potassium dinitramide in 3 mL of water were
combined under stirring, resulting in the formation of a yellow
precipitate. The suspension was then stirred for an additional 15
minutes and the solvent was evaporated completely afterwards.
The remaining yellow (slightly wet) residue was extracted with 40
mL of ethanol. The ethanolic solution was concentrated to one
fourth of its original volume and left for crystallization, yielding
0.86g (83%) of clean **5**. Crystals suitable for X-ray diffraction
measurements were obtained from a wet ethanolic solution.

DSC (5 °C min⁻¹): 155 °C (mp., onset), 164 °C (dec., onset); **IR**
(ATR): 3788 (vw), 3460 (w), 3425 (w), 3397 (m), 3319 (w), 3269
(m), 3180 (m), 3114 (m), 1707 (m), 1691 (m), 1652 (s), 1512 (s),
1469 (m), 1429 (m), 1342 (m), 1173 (vs), 1150 (s), 1054 (w), 1034
(m), 1004 (s), 960 (m), 815 (w), 799 (w), 761 (m), 723 (w), 688
(m), 673 (m); **Raman** (1064 nm, 25 °C): 3408 (3), 3346 (7), 3323
(6), 3284 (6), 3224 (5), 1699 (23), 1659 (19), 1607 (8), 1589 (9),
1544 (10), 1527 (9), 1470 (22), 1435 (13), 1326 (100), 1285 (20),
1213 (7), 1182 (17), 1151 (21), 1050 (51), 963 (27), 819 (53), 802
(22), 767 (7), 670 (30), 499 (35), 486 (18), 452 (10), 352 (25), 306
(23); **¹H NMR**: δ (ppm) = 12.13 (s, br, 2H, NH), 6.94 (s, 4H,
NH₂); **¹³C{¹H} NMR**: δ (ppm) = 152.0; **¹⁴N NMR**: δ (ppm) = -10
(N₃O₄⁻); **MS**: FAB⁺ = 100.10 ([M]⁺), FAB⁻ = 106.00 ([N₃O₄]⁻);
EA: calcd.: C 11.65, H 2.93, N 54.36; found: C 11.46, H 2.80, N
53.93; **IS**: > 3 J; **FS**: 192 N; **ESD**: 0.3 J.

Acknowledgments


Financial support of this work by the Ludwig-Maximilian
University of Munich (LMU), the U.S. Army Research
Laboratory (ARL), the Armament Research, Development
and Engineering Centre (ARDEC), the Strategic
Environmental Research and Development Program
(SERDP) and the Office of Naval Research (ONR Global,
title: "Synthesis and Characterization of New High Energy
Dense Oxidizers (HEDO) - NICOP Effort ") under contract
nos. W911NF-09-2-0018 (ARL), W911NF-09-1-0120
(ARDEC), W011NF-09-1-0056 (ARDEC) and 10
WPSEED01-002 / WP-1765 (SERDP) is gratefully
acknowledged. The authors acknowledge collaborations
with Dr. Mila Krupka (OZM Research, Czech Republic) in
the development of new testing and evaluation methods for
energetic materials and with Dr. Muhamed Sucasca
(Brodarski Institute, Croatia) in the development of new
computational codes to predict the detonation and
propulsion parameters of novel explosives. We are indebted
to and thank Drs. Betsy M. Rice and Brad Forch (ARL,
Aberdeen, Proving Ground, MD) and Mr. Gary Chen
(ARDEC, Picatinny Arsenal, NJ) for many helpful and
inspired discussions and support of our work.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- [1] a) T. M. Klapötke, in *Moderne Anorganische Chemie*, E. Riedel (Hrsg.), 3. Aufl., Walter de Gruyter, Berlin, New York, **2007**, 99-104. b) R. P. Singh, R. D. Verma, D. T. Meshri, J. M. Shreeve, *Ang. Chem. Int. Ed.* **2006**, *45*(22), 3584. c) T. M. Klapötke, in *High Energy Density Materials*, T. M. Klapötke (Hrsg.), Springer, Berlin, Heidelberg, **2007**, 85-122. d) R. D. Chapman, in *High Energy Density Materials*, T. M. Klapötke (Hrsg.), Springer, Berlin, Heidelberg, **2007**, 123-152.
- [2] (a) Y.-H. Joo, J. M. Shreeve, *Angew. Chem. Int. Ed.* **2009**, *48*, 564-567; (b) J. M. Veauthier, D. E. Chavez, B. C. Tappan, D. A. Parrish, *J. Energ. Mater.* **2010**, *28*, 229-249; (c) T. M. Klapötke, M. Göbel, *Adv. Funct. Mater.* **2009**, *19*, 347-365.
- [3] Y. N. Matyushin, T. S. Kon'kova, A. B. Vorob'ev, Y. A. Lebedev, *International Annual Conference of ICT 2005*, 36th, 92/1-92/9.
- [4] (a) T. M. Klapötke, J. Stierstorfer, *Dalton Transactions*, **2008**, *4*, 643-653. (b) T. M. Klapötke, J. Stierstorfer, *New Trends in Research of Energetic Materials, Proceedings of the Seminar, 11th, Pardubice, Czech Republic*, **2008**, *2*, 810-831.
- [5] N. Wingborg, N. V. Latypov, *Propellants, Explos., Pyrotech.* **2003**, *28*(6), 314-318.
- [6] H. R. Blomquist, U.S. **1999**, 9 pp. US 6004410 A 19991221 CAN 132:24485 AN 1999:808506
- [7] (a) K. O. Christe, W. W. Wilson, M. A. Petrie, H. H. Michels, J. C. Bottaro, R. Gilardi, *Inorg. Chem.* **1996**, *35*, 5068-5071. (b) V. A. Shlyapochnikov, M. A. Tafiopolsky, I. V. Tokmakov, E. S. Baskir, O. V. Anikin, Yu.A. Štrelenko, O. A. Luk'yanov, V. A. Tartakovskiy, *J. Molec. Struct.* **2001**, *559*, 147-166. (c) J. C. Bottaro, P. E. Penwell, R. J. Schmitt, *Synth. Commun.* **1991**, *21*, 945; (d) O. A. Luk'yanov, V. P. Gorelik, V. A. Tartakovskii, *Russ. Chem. Bull.* **1994**, *43*, 89.
- [8] R. Gilardi, R. J. Butcher *J. Chem. Cryst.* **2002**, *32*(11), 477-484.
- [9] M. E. Sitzmann, R. Gilardi, R. J. Butcher, W. M. Koppes, A. G. Stern, J. S. Trasher, N. J. Trivedi, Z.-Y. Yang, *Inorg. Chem.* **2000**, *39*, 843-850.
- [10] N. B. Bolotina, M. J. Hardie, A. A. Pinkerton, *J. Appl. Crystallogr.* **2003**, *36*(6), 1334-1341.
- [11] (a) H. Östmark, U. Bemm, H. Bergman, A. Langlet, *Thermochim. Acta* **2002**, *384*, 253-259; (b) P. B. Kempa, M. Herrmann, I. Fuhr, H. Östmark, *International Annual Conference of ICT, 40th*, **2009**, 40/1-40/11.
- [12] T. M. Klapötke, J. Stierstorfer, *Phys. Chem. Chem. Phys.* **2008**, *10*, 4340-4346.
- [13] (a) V. P. Sinditskii, A. I. Levshenkov, V. Y. Egorshv, V. V. Serushkin, *Proceedings of the Seminar, 7th, Pardubice, Czech Republic*, **2004**, *2*, 649-658. (b) C. J. Hinshaw, R. B. Wardle, T. K. Highsmith, U.S. patent **1998**, US 5741998, A 19980421.
- [14] T. M. Klapötke, J. Stierstorfer, *Eur. J. Inorg. Chem.* **2008**, *26*, 4055-4062.
- [15] K. Karaghiosoff, T. M. Klapötke, P. Mayer, C. M. Sabate, A. Penger, J. M. Welch, *Inorg. Chem.* **2008**, *47*, 1007-1019.
- [16] (a) T. M. Klapötke, P. Mayer, A. Schulz, J. J. Weigand, *J. Am. Chem. Soc.* **2005**, *127*(7), 2032-2033. (b) G. Fischer, G. Holl, T. M. Klapötke, P. Mayer, J. J. Weigand, *New Trends in Research of Energetic Materials, Proceedings of the Seminar, 8th, Pardubice, Czech Republic*, 2005, *1*, 190-199.
- [17] J. J. Weigand, *Dissertation*, Ludwig Maximilian University Munich, **2005**.
- [18] CrysAlis CCD, Oxford Diffraction Ltd., Version 1.171.27p5 beta (release 01-04-2005 CrysAlis171.NET).
- [19] CrysAlis RED, Oxford Diffraction Ltd., Version 1.171.27p5 beta (release 01-04-2005 CrysAlis171.NET).
- [20] A. Altomare, G. Cascarano, C. Giacovazzo, A. Guagliardi, SIR-92, A program for crystal structure solution, *J. Appl. Cryst.* **1993**, *26*, 343.
- [21] G. M. Sheldrick, **1997**, *SHELXL-97*, Program for the Refinement of Crystal Structures. University of Göttingen, Germany.
- [22] L. J. Farrugia, WinGX suite for small molecule single-crystal crystallography, *J. Appl. Cryst.* **1999**, *32*, 837-838.
- [23] A. L. Spek, **1999**, Platon, A Multipurpose Crystallographic Tool, Utrecht University, Utrecht, The Netherlands.
- [24] Crystallographic data for the structure(s) have been deposited with the Cambridge Crystallographic Data Centre. Copies of the data can be obtained free of charge on application to The Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (Fax: int.code (1223)336-033; e-mail for inquiry: fileserv@ccdc.cam.ac.uk; e-mail for deposition: deposit@ccdc.cam.ac.uk).
- [25] G. L. Starova, O. V. Frank-Kamenetskaya, E. F. Shibanova, V. A. Lopyrev, M. G. Voronkov, V. V. Makarskii, *Chem. Heterocyclic Comp.* **1979**, *15*(10), 1149-1150.
- [26] M. Bichay, J. W. Fronabarger, R. Gilardi, R. J. Butcher, W. B. Sanborn, M. E. Sitzmann, M. D. Williams, *Tetrahedron Lett.* **2006**, *47*, 6663-6666.
- [27] M. v. Denffer, T. M. Klapötke, G. Kramer, G. Spies, J. M. Welch, G. Heeb, *Propellants, Explos. Pyrotech.* **2005**, *30*(3), 191-195.
- [28] J. Bernstein, R. E. Davis, L. Shimoni, N.-L. Chang, *Angew. Chem. Int. Ed.* **1995**, *34*, 1555-1573.
- [29] RPluto - Graphical Display of Molecular and Crystal Structures. RPluto, The Cambridge Crystallographic Data Centre. **2004**.
- [30] T. M. Klapötke, C. M. Sabaté, J. Stierstorfer, *Z. Anorg. Allg. Chem.* **2008**, *634*, 1867-1874.
- [31] M. J. Frisch et al., Gaussian 03, Revision B04, Gaussian Inc., Wallingford, CT, **2004**.
- [32] (a) J. W. Ochterski, G. A. Petersson, and J. A. Montgomery Jr., *J. Chem. Phys.* **1996**, *104*, 2598; (b) J. A. Montgomery Jr., M. J. Frisch, J. W. Ochterski, G. A. Petersson, *J. Chem. Phys.* **2000**, *112*, 6532.
- [33] (a) L. A. Curtiss, K. Raghavachari, P. C. Redfern, J. A. Pople, *J. Chem. Phys.* **1997**, *106*(3), 1063; (b) E. F. C. Byrd, B. M. Rice, *J. Phys. Chem. A* **2006**, *110*(3), 1005-1013; (c) B. M. Rice, S. V. Pai, J. Hare, *Comb. Flame* **1999**, *118*(3), 445-458.
- [34] <http://webbook.nist.gov/chemistry/>
- [35] (a) M. S. Westwell, M. S. Searle, D. J. Wales, D. H. Williams, *J. Am. Chem. Soc.* **1995**, *117*, 5013-5015; (b) F. Trouton, *Philos. Mag.* **1884**, *18*, 54-57.
- [36] (a) H. D. B. Jenkins, H. K. Roobottom, J. Passmore, L. Glasser, *Inorg. Chem.* **1999**, *38*(16), 3609-3620. (b) H. D. B. Jenkins, D. Tudela, L. Glasser, *Inorg. Chem.* **2002**, *41*(9), 2364-2367.
- [37] a) M. Sućeska, EXPLO5.3 program, Zagreb, Croatia, **2009**; b) M. Sućeska, EXPLO5.4 program, Zagreb, Croatia, **2010**.
- [38] (a) M. Sućeska, *Materials Science Forum*, **2004**, 465-466, 325-330. (b) M. Sućeska, *Propellants, Explos., Pyrotech.* **1991**, *16*, 197-202. (c) M. Sućeska, *Propellants, Explos., Pyrotech.* **1999**, *24*, 280-285. (d) M. L. Hobbs, M. R. Baer, *Proc. of the 10th Symp. (International) on Detonation*, ONR 33395-12, Boston, MA, July 12-16, 1993, p. 409.
- [39] Calculation of oxygen balance: Ω (%) = $(wO - 2xC - 1/2yH - 2zS)/1600/M$. (w: number of oxygen atoms, x: number of carbon atoms, y: number of hydrogen atoms, z: number of sulfur atoms, M: molecular weight).
- [40] J. Köhler, R. Meyer, in „Explosivstoffe“ 9th edn., Wiley-VCH, Weinheim, **1998**.
- [41] NATO standardization agreement (STANAG) on explosives, impact sensitivity tests, no. 4489, 1st ed., Sept. 17, **1999**.
- [42] NATO standardization agreement (STANAG) on explosive, friction sensitivity tests, no. 4487, 1st ed., Aug. 22, **2002**.

- 1
2 [43] NATO standardization agreement (STANAG) on explosive,
3 electrostatic discharge sensitivity tests, no. 4490, 1st ed., Feb.
4 19, **2001**.
5 [44] WIWEB-Standardarbeitsanweisung 4-5.1.02, Ermittlung der
6 Explosionsgefährlichkeit, hier der Schlagempfindlichkeit mit
7 dem Fallhammer, Nov. 8, **2002**.
8 [45] WIWEB-Standardarbeitsanweisung 4-5.1.03, Ermittlung der
9 Explosionsgefährlichkeit oder der Reibeempfindlichkeit mit
10 dem Reibeapparat, Nov. 8, **2002**.
11 [46] <http://www.bam.de>
12 [47] Impact: Insensitive ≥ 40 J, less sensitive ≥ 35 J, sensitive ≥ 4
13 J, very sensitive ≤ 3 J; friction: Insensitive > 360 N, less
14 sensitive = 360 N, sensitive < 360 N a. > 80 N, very sensitive
15 ≤ 80 N, extreme sensitive ≤ 10 N; According to the UN
16 Recommendations on the Transport of Dangerous Goods (+)
17 indicates: not safe for transport.
18 [48] (a) [http://www.ozm.cz/testing-instruments/small-](http://www.ozm.cz/testing-instruments/small-scaleelectrostatic-discharge-tester.htm)
19 [scaleelectrostatic-discharge-tester.htm](http://www.ozm.cz/testing-instruments/small-scaleelectrostatic-discharge-tester.htm); (b) V. Pelikán, OZM
20 research, Czech Republic, private communication.
21 [49] (a) REICHEL & PARTNER GmbH, [http://www.reichelt-](http://www.reichelt-partner.de)
22 [partner.de](http://www.reichelt-partner.de) (b) Test methods according to the UN
23 Recommendations on the Transport of Dangerous Goods,
24 Manual of Test and Criteria, fourth revised edition, United
25 Nations Publication, New York and Geneva, **2003**, ISBN 92-
1-139087-7, Sales No. E.03.VIII.2; 13.4.2 Test 3(a) (ii) BAM
26 Fallhammer.

26 Received: ((will be filled in by the editorial staff))
27 Published online: ((will be filled in by the editorial staff))
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Entry for the Table of Contents



Thomas M. Klapötke,* Franz A. Martin, Norbert T. Mayr and
Jörg Stierstorfer
..... Page No. – Page No.

Synthesis and Characterization of 3,5-Diamino-1,2,4-triazolium
Dinitramide