

Empowerment: Qualitative Underpinning of a New Clinical Genetics-Specific Patient Reported Outcome

Marion Mcallister, Graham Dunn, Chris Todd

▶ To cite this version:

Marion Mcallister, Graham Dunn, Chris Todd. Empowerment: Qualitative Underpinning of a New Clinical Genetics-Specific Patient Reported Outcome. European Journal of Human Genetics, 2010, 10.1038/ejhg.2010.160 . hal-00583531

HAL Id: hal-00583531 https://hal.science/hal-00583531

Submitted on 6 Apr 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Title: Empowerment: Qualitative Underpinning of a New Clinical Genetics-Specific
2	Patient Reported Outcome.
3	
4	Short running title: Empowerment: A New Patient Reported Outcome
5	
6	Marion McAllister PhD (corresponding author), Medical Genetics Research Group,
7	School of Clinical and Laboratory Sciences, The University of Manchester, The
8	Nowgen Centre for Genetics in Healthcare, 29 Grafton Street, Manchester M13 9WU,
9	United Kingdom. Telephone number: +44(0)161 276 8979, fax number: +44(0)161 276
10	4058; e-mail address: marion.mcallister@manchester.ac.uk
11	
12	Graham Dunn PhD, School of Community-Based Medicine, The University of
13	Manchester
14	
15	Chris Todd PhD, School of Nursing, Midwifery & Social Work, The University of
16	Manchester
17	
18	This research was funded by the Medical Research Council.

1 ABSTRACT

2 Recent qualitative research developed a new construct labelled Empowerment 3 describing a new patient outcome from using clinical genetics services that included 4 four dimensions: Knowledge and Understanding, Decision-Making, Instrumentality and 5 Future-Orientation. The aim in this study was to explore the validity, relevance and 6 importance of the Empowerment construct for use as a Patient Reported Outcome for 7 clinical genetics services, and to refine the construct if necessary. Qualitative research 8 (interviews and focus groups) was conducted in the UK with 12 patients, 15 9 representatives from patient support groups, 10 genetics clinicians and 4 service 10 commissioners. Participants were asked to (1) describe what they think are the patient 11 benefits from using clinical genetics services and (2) critique the Empowerment 12 construct as a Patient Reported Outcome. Interviews and focus groups were transcribed 13 in full and analysed using grounded theory. Findings confirmed the relevance and 14 importance of the Empowerment construct, and identified Emotional Regulation as a 15 further dimension of Empowerment. Data analysis also resulted in refinement of the 16 construct, by renaming the other four dimensions to be Cognitive Control, Decisional 17 Control, Behavioural Control and Hope. Empowerment has potential to be a useful 18 Patient Reported Outcome to evaluate interventions in clinical genetics, and for use in 19 clinical practice to generate data for continuous quality improvement. A study is 20 underway to operationalise Empowerment by developing a psychometrically sound 21 Patient Reported Outcome Measure that will take the form of a short questionnaire. 22 23 **KEYWORDS:** Patient Reported Outcomes, empowerment, qualitative research, 24 emotional regulation, grounded theory

25

1 INTRODUCTION

2 Evaluating interventions in clinical genetics is difficult because there is lack of clarity about which are the best outcome domains to measure¹⁻³, although efforts are underway 3 to define and achieve consensus on outcomes^{3,4}. Traditional measures of health status 4 e.g. EQ-5D or SF36 may not be appropriate because many genetic conditions can be 5 neither treated nor cured⁵. Often, the interventions offered in clinical genetics relate to 6 7 provision of information about a diagnosis, or a genetic test result, or empiric risk 8 information. One approach has been to measure patient knowledge, or information 9 recall, but this approach has been criticized because of the substantial assumptions that 10 specific items of knowledge are either valued by patients, or contribute to effective 11 decision-making. In effect, these studies are tests of memory and understanding, not 12 evaluations of patient benefit. Another approach has been to use generic measures of 13 psychological constructs, e.g. anxiety, but these approaches have not been shown to 14 discriminate effectively between different models of service delivery in clinical trials⁶. 15 A third, more recent approach is to measure effectiveness of decisions using measures 16 of informed decision-making or decisional conflict. These approaches are limited to 17 evaluating effectiveness of a single decision, and may not be relevant for evaluating 18 clinical genetics services, where outcomes relate to the capacity to make many decisions⁷. Measures of satisfaction have also been used to evaluate clinical genetics⁸⁻¹⁰. 19 20 Although an important component in assessing healthcare quality, patient satisfaction is 21 not sufficient to capture all the important patient benefits, and furthermore, is influenced by expectations¹¹. 22

Our recent systematic review of validated outcome measures used in clinical genetics
identified that 67 different outcome measures have been used to evaluate clinical

1	genetics. 46 of those measures were used in just one study each ⁸ , highlighting the
2	confusion about appropriate measures. Outcome domains captured by the measures
3	ranged from health status through knowledge to quality of life. The review concluded
4	that for clinical genetics services (a) there was no single validated outcome measure
5	available to capture all potential patient benefits (b) some important aspects of patient
6	benefit are not captured by available measures and (c) there is scope for new measures.
7	Our previous qualitative research ^{6,7,12,13} suggests that patient benefits from clinical
8	genetics services could be summarised using a new construct labelled Empowerment ⁷ .
9	Empowerment was defined as a set of beliefs that enable a person from a family
10	affected by a genetic condition, to feel that they have some control over and hope for
11	the future. Empowerment included four dimensions: the beliefs that one (1) can make
12	important life decisions in an informed way (Decision-Making), (2) has sufficient
13	information about the condition, including risks to oneself and one's relatives, and any
14	treatment, prevention and support available (Knowledge and Understanding), (3) can
15	make effective use of the health and social care systems for the benefit of the whole
16	family (Instrumentality) and (4) can look to the future with hope for a fulfilling family
17	life, for oneself, one's family, and/or one's future descendents (Future Orientation) ⁷ .
18	When compared with outcome measures used in evaluations of clinical genetics,
19	Empowerment was found to be conceptually most similar to the construct captured by
20	the Perceived Personal Control (PPC) measure ¹⁴ , which includes three dimensions:
21	Cognitive Control, Decisional Control and Behavioural Control. However, the PPC
22	questionnaire does not capture all the important patient benefits summarised in the
23	Empowerment construct e.g. benefits to other relatives and to future generations.

1	There is scope to develop a new Patient Reported Outcome Measure (PROM) to capture
2	the Empowerment construct. PROMs are short validated questionnaires completed by
3	patients to evaluate health interventions. A PROM of Empowerment could be useful in
4	evaluating new interventions in clinical genetics, thus contributing to establishing a
5	robust evidence base for practice. However, PROMs development is difficult and time-
6	consuming ¹⁵⁻¹⁷ , and justifiable only if there is evidence that the new measure will
7	capture a Patient Reported Outcome (PRO) that is novel, useful, relevant and important.
8	The aims in the present study were to (1) explore the validity, relevance and importance
9	of the Empowerment construct and refine the construct as necessary and (2) inform the
10	decision whether or not to develop a clinical genetics-specific PROM of Empowerment.
11	METHODS
12	To meet the research objectives, further qualitative data were collected. Qualitative
13	methods are suitable for exploratory research in areas that are not well understood, and
14	where the relevant variables have not been fully identified ¹⁸ . Since one purpose of this
15	research was to explore whether the Empowerment model includes all relevant facets of
16	patient benefit from using clinical genetics services, a qualitative approach was
17	appropriate. Patients were recruited through Genetic Medicine, St Mary's Hospital,
18	Manchester, UK. The term 'patient' is used here to refer to all users of a clinical
19	genetics service, including unaffected parents and unaffected at risk families members.
20	Patients were asked to participate in a focus group, but also had the option to be
21	interviewed. Genetics clinicians were recruited through professional networks, and were
22	only asked to participate in a focus group. Previous research indicated that focus groups
23	work well with genetics clinicians, whereas a combined focus group / interview
24	approach works best with patients of clinical genetics services ^{4 5 11 12} . Commissioners

(who purchase clinical genetics services on behalf of primary care) and patient
 representatives (members of patient support groups) were recruited through UK
 professional networks, to participate in an interview only because of geographical
 spread. Focus groups were conducted in Nowgen, Manchester. Interviews were
 conducted in person or by telephone, according to participant preference. Ethical
 approval was obtained from the NHS National Research Ethics Service.

7 Participants were asked to describe their views about the benefits and harms of using 8 clinical genetics services, and then given a page describing the Empowerment construct, with its four dimensions, and the dimension definitions⁷. Participants were asked to read 9 10 and critique this, in particular to identify any benefits / harms of using a clinical 11 genetics service that were missing. Focus groups and interviews were transcribed in full, and analysed using tools of grounded theory methodology^{19,20} including: (1) open 12 13 coding: labels (codes) significant pieces of text as important (2) constant comparison: 14 each new instance of a code is compared with every other instance of that code, and the 15 concept definition revised accordingly and (3) questioning the data in relation to the 16 research question. Glaser and Strauss, the developers of grounded theory methodology, 17 recommend that any emerging grounded theory (model) should be "checked out" for usefulness with "experts" familiar with the substantive area under investigation^{19,21}. In 18 19 this study, the emerging construct (model) of Empowerment is checked out with 20 patients, patient representatives, clinicians, and commissioners of clinical genetics 21 services to clarify whether the construct provides a plausible and useful summary of 22 patient benefit. Feedback of this kind can indicate whether or not the construct (model) has "grab"²¹ i.e. whether it provides a novel and useful framework. 23

24 **RESULTS**

1	138 people were approached and asked to participate. 54 (39.2%) agreed to participate,
2	and 18 (13%) declined. Of the 54 who agreed to participate, 13 (9.4%) were excluded
3	because they could not be accommodated in a focus group because of scheduling
4	difficulties, and did not wish to be interviewed instead. The final sample was composed
5	of 41 participants (29.8%) (Tables 1&2). Due to ethical constraints, those approached
6	were not asked to give reasons for their decision, so reasons for the low response rate
7	amongst patients are not known. Three focus groups were conducted, two with genetics
8	clinicians ($n = 5$ in both) and one with patients ($n = 4$). 27 interviews were conducted, 4
9	with service commissioners, 8 with patients, and 15 with patient representatives.
10	Participants' views about the 'Empowerment' construct
11	The qualitative study generated data confirming the importance of the Empowerment
12	construct, and the previously identified four dimensions of Empowerment (Knowledge
13	and Understanding, Decision-Making, Instrumentality and Future-Orientation). All 41
14	participants, except for one patient representative, endorsed use of the word
15	"Empowerment" as a good overarching term to summarise patient benefits, and clarified
16	further what Empowerment means in the context of clinical genetics:
17	"I think that's the best word you could think of, yes it is, having empowerment -
18	having that knowledge is the biggest benefit that you can gain. And yes I would
19	recommend a genetic service to anybody, well I do when they have a new baby and
20	they're all sort of all bewildered [] 'empowerment,' yes that is [] a very good word
21	to describe it." (patient, Congenital adrenal hyperplasia).
22	Results suggest that genetic diagnosis may engender a state of disempowerment, which
23	throws families into a new state, perhaps best described as social alienation, which can
24	be addressed in genetic counseling, as illustrated by this participant:

1	"I just feel though that in a way the system of the diagnosis is what has disempowered
2	parents in the first place, so it actually about re-establishing confidence rather than
3	just giving it them for the first time." (patient representative, D/deaf charity)
4	A good outcome from using a clinical genetics service could then be seen to involve
5	learning about their new situation, and regaining some confidence and control over their
6	lives, so that they "know what they are doing", which is experienced as empowering,
7	expressed as follows:
8	"the information empowers you to make decisions, and provided us with the
9	opportunity to have a family, it gave us the confidence to try for another baby, which
10	we would not have done otherwise." (patient, previously had a late termination at 23
11	weeks gestation of a pregnancy affected by Edwards syndrome)
12	The views expressed by patients, and patient representatives, were endorsed by the
13	service commissioner participants, who confirmed that they want to commission clinical
14	genetics services that can offer choice to patients:
15	" the patients [] have the choice to consider their own lifestyles, their own social
16	decisions, the future of their own children and that chance of things if they wish to
17	[] carry on with the pregnancy if they've already conceived, or to terminate or to not
18	conceive in future, to adopt, to go through PGD (preimplantation genetic diagnosis)
19	whatever, their decision, it is the parents decision." (service commissioner)
20	One participant, although she validated the importance of the four dimensions of
21	Empowerment, was unsure about the appropriateness of the word "Empowerment":
22	"I have slight reservations about the word but certainly in terms of those - what you've
23	laid out there as the what underpins that I think all are extremely valid, you know, I
24	think that would [] definitely [be] a useful way of measuring (outcomes) []

1	'empowerment' suggests that you are able to be proactive and that I don't think is
2	always going to be the way or would be a potential outcome of a clinical genetic
3	service, I don't think that's feasible. (patient representative, charity supporting couples
4	who have terminated pregnancies for fetal abnormalities)
5	Because all 41 participants, except for this one, endorsed use of the word
6	"Empowerment" to summarise patient benefits from using clinical genetics services,
7	and some did so strongly, it was decided to retain the term "Empowerment" as an
8	overarching term to describe the model.
9	Refinement of the Empowerment construct
10	All participants endorsed the relevance and importance of the four dimensions of
11	Empowerment when shown the definitions, but the analysis resulted in renaming three
12	of these dimensions to further clarify the Empowerment construct. Data analysis
13	confirmed that Knowledge and Understanding, as conceptualised in Empowerment, is
14	more akin to "Cognitive Control", a dimension of PPC, than to information recall.
15	Knowledge about the condition is valued for its own sake because it provides an
16	explanation for previously unexplained, unexpected and distressing events (recurrence
17	of disease in the family, loss of a child, disease symptoms), and this in itself is a benefit.
18	Simply having a diagnosis provides reassurance and enables families to prepare for the
19	future.
20	"[] The benefits to me of having the diagnosis confirmed that what I was
21	experiencing that it actually had a name, and despite the fact that there was very little
22	research done in to the condition twelve years ago it enabled us to do some research
23	ourselves and to keep up with the latest what was going on. So we could put a focus
24	on it, you know " (patient representative, affected by Porphyria)

1	It appeared that labeling symptoms with a scientific name had the quality of a "latin
2	incantation" which reduced uncertainty/anxiety, and validated the illness experience.
3	Similarly, the data confirmed that Decision-Making, as conceptualised in
4	Empowerment, is more like "Decisional Control", a dimension of PPC, than to
5	effectiveness of any single decision. Finally, Instrumentality, as conceptualised in
6	Empowerment, is more similar to "Behavioural Control", a dimension of PPC, than to
7	any specific action taken. Understanding the implications of the genetic condition to
8	oneself, one's relatives, and future generations, and knowing about available medical
9	interventions and support enables decisions to be made, and actions to be taken:
10	"it's the information that enables you to make the decisions, whether to go ahead
11	and have another baby, or whether to have more children, and what sort of tests to
12	have." (patient, baby affected by chromosome abnormality)
13	For these reasons, the PPC terminology of Cognitive, Decisional and Behavioural
14	Control was adopted for the Empowerment dimensions previously known as
15	Knowledge and Understanding, Decision-making and Instrumentality respectively.
16	Future-Orientation was also endorsed by all participants in this study as relevant and
17	important to good patient outcomes. Understanding the genetic condition, and knowing
18	about services and interventions available to help manage the effects of the condition
19	enables families to orient themselves towards the future with more confidence:
20	" it comes back to this control isn't it that you are able to map out your future life
21	you know, it's not that much of an unknown [] (and) feeling comfortable about
22	one's children's future" (patient representative, child affected by an unbalanced
23	chromosome translocation)

1	One participant suggested that the Future-Orientation dimension be renamed "Hope"
2	because she felt that providing Hope was an important part of any diagnosis of a life-
3	limiting condition. The word Hope came up a few times in interviews as an important
4	benefit from using clinical genetics, and the decision was made to change the name of
5	this dimension from Future-Orientation to Hope, partly because the term Future-
6	Orientation required some explanation to participants, whereas the word Hope is more
7	widely understood.
8	Further development of the Empowerment construct
9	The qualitative analysis resulted in identification of one further dimension of
10	Empowerment: Emotional Regulation. Genetics clinicians in one focus group, and a
11	number of patient representatives, some of whom were affected by genetic conditions,
12	and some of whom were parents of affected people, commented in interviews that the
13	model omitted any mention of emotions, which they felt were important. No other
14	omissions were identified. Patient representatives, all of whom had a genetic condition
15	in their own family, and were also able to provide an overview of the variability in
16	experiences of their patient group members, described how a genetic diagnosis throws
17	the family into emotional turmoil, and highlighted the need for families to be given
18	emotional support:
19	" for families to have emotional support around a point of diagnosis [] I would
20	say that everybody is depressed around the time of using the service". (patient
21	representative, charity supporting families affected by Battens disease)
22	They felt strongly that clinical genetics services should be instrumental in addressing
23	these difficult emotions, and in helping families to reach a new equilibrium:

1	" when you're first given a diagnosis it's like your whole life is spinning out of
2	control you don't know where you're going, you don't know what's going to happen to
3	you or your family, what the future holds, it's incredibly frightening having been
4	through it myself [] a very important outcome (is) that the emotions are dealt with."
5	(patient group representative, child has unbalanced chromosome translocation)
6	Genetic counsellors commented that a large part of their work focuses on offering
7	emotional support to patients, and so they appeared to recognise the importance to
8	families of addressing and dealing with difficult emotions in genetic counselling.
9	No other new issues were identified in the final 5 interviews, and so theoretical
10	saturation was assumed, and recruitment stopped.
11	The new Empowerment construct
12	In summary, the qualitative analysis endorsed the Empowerment construct, and refined
13	the construct by (a) adopting the PPC terminology of Cognitive, Decisional and
14	Behavioural Control for the dimensions previously labeled Knowledge and
15	Understanding, Decision-Making and Instrumentality respectively, (2) adopting the
16	term "Hope" for the dimension previously labeled "Future-Orientation", and (3)
17	extending the model to include a fifth dimension: Emotional Regulation.
18	DISCUSSION
19	This research has validated and further developed and refined the Empowerment
20	construct, to summarise the patient benefits from using clinical genetics services ⁷ . A
21	sample of patients and patient representatives, clinicians and commissioners of clinical
22	genetics services in the UK endorsed use of the term "Empowerment" to summarise the
23	patient benefits, and identified one further dimension of Empowerment, Emotional
24	Regulation, that was missing from the four-dimensional model. Furthermore, the data

1	have enabled the meaning of the previously identified four dimensions of Knowledge
2	and Understanding, Decision-Making, Instrumentality and Future-Orientation to be
3	refined. The revised construct is composed of 5 dimensions: Cognitive Control,
4	Decisional Control, Behavioural Control, Hope and Emotional Regulation, and has
5	clarified the relationship between Empowerment, and the concept of Perceived Personal
6	Control (PPC) ^{7,12,20} . Findings support the importance of PPC as a PRO, and identified
7	that Empowerment includes the three PPC dimensions (Cognitive, Decisional and
8	Behavioural Control), and two further dimensions: Hope and Emotional Regulation.
9	In contrast to other research, for example exploring autism in France ²² , neither this nor
10	our previous research ^{6,7,12,13} identified any differences between the views of healthy
11	parents of a child affected by a condition, patients affected by a condition, and those at
12	risk for developing a condition, or between adults at different stages of development.
13	There is scope to explore this further in future research.
14	The Empowerment PRO, if operationalised as a PROM, could be used for evaluation of
15	whether some of the criteria in the recently developed EuroGentest "Instrument for
16	quality self-assessment in provision of genetic counselling"
17	(http://www.eurogentest.org/professionals/assessment_tool/assessment_tool/form.xhtm)
18	have been met. For example, the EuroGentest instrument includes the criteria (1)
19	"Counsellees are provided with decision-making support", which could provide
20	counsellees with improved decisional control and (2) "Counsellees understand what
21	they are being told" which could provide counsellees with improved cognitive control.
22	A PROM of Empowerment would enable these criteria to be measured by patient self-
23	report, rather than by service self-assessment.

1	The constituent dimensions of Empowerment identified in the present study also
2	overlap with some of the outcomes identified in a consensus study recently reported by
3	the US Western States Genetics Services Collaborative (WSGSC) ⁴ , and the present
4	study provides further support for the relevance and importance of these initiatives.
5	For example, the WSGSC outcome "Individuals and families feel supported in
6	managing grief, stress, and emotional challenges of living with a genetic condition"
7	could be interpreted as Emotional Regulation. The WSGSC outcome "Individuals and
8	families are able to make informed health and life decisions based upon diagnosis"
9	could be interpreted as Decisional Control. However, the present study goes further than
10	identifying patient outcomes by integrating 5 different patient outcomes in the
11	overarching patient-centred construct of Empowerment. The present study provides
12	sufficient evidence to suggest that Empowerment is a novel PRO for clinical genetics
13	services.
13 14	services. Identification of Emotional Regulation as an additional dimension of Empowerment,
14	Identification of Emotional Regulation as an additional dimension of Empowerment,
14 15	Identification of Emotional Regulation as an additional dimension of Empowerment, and as an important outcome from using clinical genetics services, is not surprising.
14 15 16	Identification of Emotional Regulation as an additional dimension of Empowerment, and as an important outcome from using clinical genetics services, is not surprising. Good patient outcomes from using healthcare services could be conceptualised as
14 15 16 17	Identification of Emotional Regulation as an additional dimension of Empowerment, and as an important outcome from using clinical genetics services, is not surprising. Good patient outcomes from using healthcare services could be conceptualised as alleviation of those disease effects experienced as problematic by patients and their
14 15 16 17 18	Identification of Emotional Regulation as an additional dimension of Empowerment, and as an important outcome from using clinical genetics services, is not surprising. Good patient outcomes from using healthcare services could be conceptualised as alleviation of those disease effects experienced as problematic by patients and their families. From this perspective, the finding that Emotional Regulation is an important
14 15 16 17 18 19	Identification of Emotional Regulation as an additional dimension of Empowerment, and as an important outcome from using clinical genetics services, is not surprising. Good patient outcomes from using healthcare services could be conceptualised as alleviation of those disease effects experienced as problematic by patients and their families. From this perspective, the finding that Emotional Regulation is an important outcome in clinical genetics is consistent with research evidence that the emotional
14 15 16 17 18 19 20	Identification of Emotional Regulation as an additional dimension of Empowerment, and as an important outcome from using clinical genetics services, is not surprising. Good patient outcomes from using healthcare services could be conceptualised as alleviation of those disease effects experienced as problematic by patients and their families. From this perspective, the finding that Emotional Regulation is an important outcome in clinical genetics is consistent with research evidence that the emotional effects of genetic diseases are problematic for families ¹² . These emotional effects may
14 15 16 17 18 19 20 21	Identification of Emotional Regulation as an additional dimension of Empowerment, and as an important outcome from using clinical genetics services, is not surprising. Good patient outcomes from using healthcare services could be conceptualised as alleviation of those disease effects experienced as problematic by patients and their families. From this perspective, the finding that Emotional Regulation is an important outcome in clinical genetics is consistent with research evidence that the emotional effects of genetic diseases are problematic for families ¹² . These emotional effects may include feelings of powerlessness and a threatened sense of security and identity,

2that patients come to clinical genetics services for information only. In contrast, the3psycho-educational approach assumes that patients are seeking a therapeutic4relationship in which they are supported in using genetic information in a personally5meaningful way that minimizes psychological distress and increases personal control.6Hope has not previously been identified as a valued outcome from clinical genetics7services, and its inclusion in the model is novel. On a theoretical level, Emotional8Regulation is likely to interact with Hope, and with Perceived Personal Control, and it's9three dimensions of Cognitive, Decisional and Behavioural Control. A genetic10diagnosis, and a scientific explanation for the previously unexplained death of her child,11information about risks to future pregnancies, and options for prenatal diagnosis could12help a bereaved mother to (1) understand why her child died (Cognitive Control) (2)13make decisions about whether to have further children (Decisional Control) (3) use14prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel15that it is possible to have healthy children and a rewarding family life in the future16(Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to17cause her child's condition (Emotional Regulation). Clearly, it may not be possible to18bring about positive change in all 5 dimensions for every patient who attends a clinical19genetics service. However it is plausible to hypothesise that the 5 dimensions of	1	educational model emphasises 'educating' the patient, and is based on the assumption
 relationship in which they are supported in using genetic information in a personally meaningful way that minimizes psychological distress and increases personal control. Hope has not previously been identified as a valued outcome from clinical genetics services, and its inclusion in the model is novel. On a theoretical level, Emotional Regulation is likely to interact with Hope, and with Perceived Personal Control, and it's three dimensions of Cognitive, Decisional and Behavioural Control. A genetic diagnosis, and a scientific explanation for the previously unexplained death of her child, information about risks to future pregnancies, and options for prenatal diagnosis could help a bereaved mother to (1) understand why her child died (Cognitive Control) (2) make decisions about whether to have further children (Decisional Control) (3) use prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	2	that patients come to clinical genetics services for information only. In contrast, the
 meaningful way that minimizes psychological distress and increases personal control. Hope has not previously been identified as a valued outcome from clinical genetics services, and its inclusion in the model is novel. On a theoretical level, Emotional Regulation is likely to interact with Hope, and with Perceived Personal Control, and it's three dimensions of Cognitive, Decisional and Behavioural Control. A genetic diagnosis, and a scientific explanation for the previously unexplained death of her child, information about risks to future pregnancies, and options for prenatal diagnosis could help a bereaved mother to (1) understand why her child died (Cognitive Control) (2) make decisions about whether to have further children (Decisional Control) (3) use prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	3	psycho-educational approach assumes that patients are seeking a therapeutic
 Hope has not previously been identified as a valued outcome from clinical genetics services, and its inclusion in the model is novel. On a theoretical level, Emotional Regulation is likely to interact with Hope, and with Perceived Personal Control, and it's three dimensions of Cognitive, Decisional and Behavioural Control. A genetic diagnosis, and a scientific explanation for the previously unexplained death of her child, information about risks to future pregnancies, and options for prenatal diagnosis could help a bereaved mother to (1) understand why her child died (Cognitive Control) (2) make decisions about whether to have further children (Decisional Control) (3) use prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	4	relationship in which they are supported in using genetic information in a personally
 services, and its inclusion in the model is novel. On a theoretical level, Emotional Regulation is likely to interact with Hope, and with Perceived Personal Control, and it's three dimensions of Cognitive, Decisional and Behavioural Control. A genetic diagnosis, and a scientific explanation for the previously unexplained death of her child, information about risks to future pregnancies, and options for prenatal diagnosis could help a bereaved mother to (1) understand why her child died (Cognitive Control) (2) make decisions about whether to have further children (Decisional Control) (3) use prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	5	meaningful way that minimizes psychological distress and increases personal control.
 Regulation is likely to interact with Hope, and with Perceived Personal Control, and it's three dimensions of Cognitive, Decisional and Behavioural Control. A genetic diagnosis, and a scientific explanation for the previously unexplained death of her child, information about risks to future pregnancies, and options for prenatal diagnosis could help a bereaved mother to (1) understand why her child died (Cognitive Control) (2) make decisions about whether to have further children (Decisional Control) (3) use prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	6	Hope has not previously been identified as a valued outcome from clinical genetics
three dimensions of Cognitive, Decisional and Behavioural Control. A genetic diagnosis, and a scientific explanation for the previously unexplained death of her child, information about risks to future pregnancies, and options for prenatal diagnosis could help a bereaved mother to (1) understand why her child died (Cognitive Control) (2) make decisions about whether to have further children (Decisional Control) (3) use prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct.	7	services, and its inclusion in the model is novel. On a theoretical level, Emotional
 diagnosis, and a scientific explanation for the previously unexplained death of her child, information about risks to future pregnancies, and options for prenatal diagnosis could help a bereaved mother to (1) understand why her child died (Cognitive Control) (2) make decisions about whether to have further children (Decisional Control) (3) use prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	8	Regulation is likely to interact with Hope, and with Perceived Personal Control, and it's
information about risks to future pregnancies, and options for prenatal diagnosis could help a bereaved mother to (1) understand why her child died (Cognitive Control) (2) make decisions about whether to have further children (Decisional Control) (3) use prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct.	9	three dimensions of Cognitive, Decisional and Behavioural Control. A genetic
 help a bereaved mother to (1) understand why her child died (Cognitive Control) (2) make decisions about whether to have further children (Decisional Control) (3) use prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	10	diagnosis, and a scientific explanation for the previously unexplained death of her child,
make decisions about whether to have further children (Decisional Control) (3) use prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by	11	information about risks to future pregnancies, and options for prenatal diagnosis could
prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by	12	help a bereaved mother to (1) understand why her child died (Cognitive Control) (2)
 that it is possible to have healthy children and a rewarding family life in the future (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	13	make decisions about whether to have further children (Decisional Control) (3) use
 (Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	14	prenatal testing in future pregnancies to avoid recurrence (Behavioural Control) (4) feel
 cause her child's condition (Emotional Regulation). Clearly, it may not be possible to bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	15	that it is possible to have healthy children and a rewarding family life in the future
 bring about positive change in all 5 dimensions for every patient who attends a clinical genetics service. However it is plausible to hypothesise that the 5 dimensions of Empowerment are facets of a single overarching construct. The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	16	(Hope) and (5) alleviate feelings of guilt that she had done something in pregnancy to
 19 genetics service. However it is plausible to hypothesise that the 5 dimensions of 20 Empowerment are facets of a single overarching construct. 21 The main weakness of this research is that it is limited to the perspectives of families 22 living in the UK, who speak English. Empowerment is likely to be influenced by 	17	cause her child's condition (Emotional Regulation). Clearly, it may not be possible to
 20 Empowerment are facets of a single overarching construct. 21 The main weakness of this research is that it is limited to the perspectives of families 22 living in the UK, who speak English. Empowerment is likely to be influenced by 	18	bring about positive change in all 5 dimensions for every patient who attends a clinical
 The main weakness of this research is that it is limited to the perspectives of families living in the UK, who speak English. Empowerment is likely to be influenced by 	19	genetics service. However it is plausible to hypothesise that the 5 dimensions of
22 living in the UK, who speak English. Empowerment is likely to be influenced by	20	Empowerment are facets of a single overarching construct.
	21	The main weakness of this research is that it is limited to the perspectives of families
culture ⁷ , and so Empowerment would benefit from validation in other cultures and	22	living in the UK, who speak English. Empowerment is likely to be influenced by
	23	culture ⁷ , and so Empowerment would benefit from validation in other cultures and
24 languages. A further weakness is that patient participation rates were low. However,	24	languages. A further weakness is that patient participation rates were low. However,

1	because this work builds on extensive previous qualitative research ⁴⁵¹¹¹² , and because			
2	care was taken to reach data saturation, it is unlikely that significant issues were missed.			
3	The main strength of this research is that the new PRO identified – Empowerment - has			
4	been developed through consultation with (1) people struggling with the challenges of			
5	having a genetic condition in their family and (2) clinicians with experience of helping			
6	these families. They are the "experts" consulted in this research to identify the patient			
7	benefits from using clinical genetics services. The traditional approach to PROMs			
8	development is to use theoretical frameworks to identify PROs, and this approach was			
9	used in identifying PPC as a patient outcome from genetic counselling ^{14,26} . This			
10	0 approach is useful, but as the present study shows, consulting patients and clinicians can			
11	enable new constructs to be developed, and existing constructs to be extended, to ensure			
12	that PROs include all benefits valued by patients.			
13	A measure of PPC was developed and evaluated in Hebrew ¹⁴ . When the paper reporting			
14	development of the PPC questionnaire was published in a US journal, the PPC was			
15	translated into English ¹⁴ . Whilst the English translation of the PPC has been used with			
16	English speaking samples ²⁷ , it has not yet been validated for use with a UK population.			
17	A study is underway to operationalise Empowerment by developing a new genetics-			
18	specific PROM of Empowerment, using the PPC measure and the qualitative data			
19	collected in this and previous studies ^{4 5 11 12} . Both PROMs and patient empowerment			
20	have been identified as key aspects of current UK health policy ²⁸ . In addition, there was			
21	a recent call for better measures to capture aspects of Health Related Quality of Life			
22	(HRQoL) in clinical trials in medical genetics ²⁹ . These developments strengthen the			
23	argument in support of investing the time and resources needed to develop a new			
24	genetics-specific PROM to capture Empowerment. This PROM would then be available			

- 1 to capture HRQoL outcomes in clinical trials, and in clinical practice to generate data to
- 2 support continuous quality improvement³⁰. As part of PROM development¹⁶, factor
- 3 analysis will investigate the dimensional structure of Empowerment to confirm whether
- 4 or not Empowerment is a single multidimensional construct.

5 ACKNOWLEDGEMENTS

- 6 We are grateful to the patients and clinicians who participated. This research was
- 7 funded by the Medical Research Council. Ethical approval was provided by Central
- 8 Manchester NHS Research Ethics Committee (ref.07/H1008/226). The views expressed
- 9 in this paper are those of the authors, and not those of the funding body.

10 **CONFLICT OF INTEREST:**

11 The authors declare no conflict of interest.

12 **REFERENCES**

13 14 15	1.	Wang C, Gonzalez R, Merajver SD: Assessment of genetic testing and related counseling services: current research and future directions. <i>Soc Sci Med</i> 2004; 58: 1427-1442.
16		
17	2.	Williams MS: Quality in clinical genetics. Am J Med Genet Part C: Seminars in
18		Medical Genetics 2009; 151C: 175-178.
19		
20	3.	Chou AF, Norris AI, Williamson L, Garcia K, Baysinger J, Mulvihill MJ:
21		Quality Assurance in Medical and Public Health Genetics Services: A
22		Systematic Review. AM J Med Genet Part C (Seminars in Medical Genetics)
23		2009; 151C: 215-234.
24		
25	4.	Silvey K, Stock J, Hasegawa LE, Mann Au S: Outcomes of Genetics Services:
26		Creating an Inclusive Definition and Outcomes Menu for Public health and
27		Clinical Genetics Services. AM J Med Genet Part C (Seminars in Medical
28		Genetics) 2009; 151C: 207-213.
29		
30	5.	Jarrett J, Mugford M: Genetic health technology and economic evaluation. A
31		critical review. Health Econ Health Policy 2006; 5: 27-35.
32		
33	6.	McAllister M, Payne K, MacLeod R, Nicholls S, Donnai D, Davies L: What
34		process attributes of clinical genetics services could maximise patient benefits?
35		<i>Eur J Hum Genet</i> 2008; 16: 1467-1476.
36		

1 2 3	7.	McAllister M, Payne K, Nicholls S, MacLeod R, Donnai D, Davies L: Patient Empowerment in Clinical Genetics Services. <i>J Hlth Psychol</i> 2008; 13 : 895-905.
4 5 6 7	8.	Payne K, Nicholls S, McAllister M, MacLeod R, Donnai D, Davies L: Outcome measurement in clinical genetics services: a systematic review of validated measures. <i>Value in Health</i> 2008; 11 : 497-508.
8 9 10 11	9.	Shiloh S, Avdor O, Goodman RM: Satisfaction with genetic counseling: dimensions and measurement. <i>American Journal of Medical Genetics</i> 1990; 37: 522-529.
12 13 14 15 16	10.	Zellerino B, Milligan SA, Brooks R, Freedenberg DL, Collingridge DS, Williams MS: Development, Testing and Validation of a Patient Satisfaction Questionnaire for Use in the Clinical Genetics Setting. <i>Am J Med Genet Part C</i> (<i>Seminars in Medical Genetics</i>) 2009; 151C: 191-198.
17 18 19	11.	Coulter A: <i>Tha autonomous patient: Ending paternalism in medical care</i> , London: The Nuffield Trust, 2002.
20 21 22 23	12.	McAllister M, Davies L, Payne K, Nicholls S, Donnai D, MacLeod R: The emotional effects of genetic diseases: implications for clinical genetics. <i>American Journal of Medical Genetics A</i> 2007; 143 : 2651-2661.
24 25 26 27 28	13.	McAllister M, Payne K, Nicholls S, MacLeod R, Donnai D, Davies LM: Improving service evaluation in clinical genetics: identifying effects of genetic diseases on individuals and families. <i>Journal of Genetic Counseling</i> 2007; 16 : 71-83.
29 30 31 32	14.	Berkenstadt M, Shiloh S, Barkai G, Bat-Miriam-Katznelson M, Goldman B: Percieved Personal Control (PPC) : A new concept in measuring outcome of genetic counseling. <i>Am J Med Genet</i> 1999; 82: 53-59.
33 34 35	15.	Clark LA, Watson DW: Constructing validity: Basic Issues in Objective Scale Development. <i>Psychological Assessment</i> 1995; 7: 309-319.
36 37 38	16.	Streiner DL, Norman GR: <i>Health measurement scales: A practical guide to their development and use.</i> , Oxford.: Oxford University Press., 1995.
39 40 41 42	17.	McAllister M: Outcome Measures in Clinical Genetics Services; in: Kristoffersson U, Schmidtke J, Cassiman J-J (eds): <i>Quality Issues in Clinical Genetics Services</i> Springer, 2010.
43 44 45	18.	Miles MB, Huberman AM: Qualitative data analysis: An expanded sourcebook., London.: Sage., 1994.
46 47 48	19.	Glaser BG, Strauss AL: <i>The discovery of grounded theory.</i> , London: Weidenfield & Nicholson., 1967.

1 2 3	20.	20. McAllister M: Grounded Theory in Genetic Counseling Research. <i>Journal of Genetic Counseling</i> 2001; 10: 18.			
4 5	21.	Glaser B: Theoretical Sensitivity, Mill Valley CA: Sociology Press, 1978.			
6 7 8	22.	Chamak B: Autism and social movements: French parents associations and international autistic individuals' organisations. <i>Soc Hlth Illness</i> 2008; 30: 76-96.			
9 10 11 12	23.	Aujoulat I, Marcolongo R, Bonadiman L, Deccache A: Reconsidering patient empowerment in chronic illness: a critique of models of self-efficacy and bodily control. <i>Soc Sci Med</i> 2008; 66 : 1228-1239.			
12 13 14 15	24.	Biesecker BB, Peters K: Process studies in genetic counseling: peering into the black box. <i>Am J Med Genet</i> 2001; 106: 191-198.			
16 17 18	25.	Harper PS: General Aspects of Genetic Counseling; in: Harper PS (ed): <i>Practical Genetic Counseling</i> : Oxford: Butterworth Heinemann, 1998.			
19 20 21 22 23	26.	Shiloh S, Berkenstadt M, Meiran N, Bat-Miriam-Katznelson M, Goldman B: Mediating effects of Percieved Personal Control in coping with a health threat: The case of genetic counselling. <i>Journal of Applied Social Psychology</i> 1997; 27: 1146-1174.			
24 25 26 27	27.	Davey A, Rostant K, Harrop K, Goldblatt J, O'Leary P: Evaluating genetic counselling: Client expectations, psychological adjustment and satisfaction with service. <i>Journal of Genetic Counseling</i> 2005; 14: 197-206.			
28 29 30 31 32	28.	Darzi A. High Quality Care for All: NHS Next Stage Review final report. Available from: <u>http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_085825</u> [accessed 1 July 2009], 2008.			
33 34 35 36	29.	Stevenson DA, Carey JC: Health-Related Quality of Life Measures in Genetic Disorders: An Outcome Variable for Consideration in Clinical Trials. <i>AM J Med Genet Part C (Seminars in Medical Genetics)</i> 2009; 151C: 255-260.			
 37 38 39 40 41 42 	30.	Gray JR, Berwick DM: Peering into the Chasm: Improving the Quality of Clinical Genetics Services. <i>Am J Med Genet Part C: Sem Med Genet</i> 2009; 151C: 173-174.			

Table 1: Participation rates and sample composition					
	Patients	Patient representatives	Genetics clinicians	Commissioners	TOTAL
	(choice of focus group or interview)	(interviews only)	(focus groups only)	(interviews only)	(interviews &
	n (%)	n (%)	n (%)	n (%)	focus groups)
		<u> </u>			n (%)
Approached	79 (100)	20 (100)	32 (100)	7 (100)	138 (100)
No response	47 (59.5)	5 (25)	12 (37.5)	2 (28.6)	66 (47.8)
Declined	17 (21.5)	0 (0)	0 (0)	1 (14.3)	18 (13)
Agreed to	15 (19)	15 (75)	20 (62.5)	4 (57.1)	54 (39.2)
participate					
Subsequently	3 (3.8)	0	10 (31.25)	0	13 (9.4)
unable to					1
participate ^I					l
Participated	12 (15.2)	15 (75)	10 (31.25)	4 (57.1)	41 (29.8)
Genetic	• Hereditary cardiomyopathy	Chromosome abnormalities			
conditions	• Ehlers Danlos syndrome	Hereditary Deafness			
represented	Congenital adrenal hyperplasia	• Sickle cell disease			
	• BRCA2	• Alpha-1 antitrypsin deficiency			
	• Zellweger syndrome	Cystic fibrosis			
	Chromosome abnormalities	Laurence Moon Bardet Biedl			
	Retinitis pigmentosa	syndrome			
	• Marfan syndrome	Muscular dystrophy			
	 Neurofibromatosis Type 1 	• Porphyria			
	• Fragile X syndrome	• Pregnancy termination for fetal			
	 Undiagnosed rare conditions 	abnormalities			
	ondugnosed rate conditions	Charcot Marie Tooth disease			
		Battens disease			
		 Ectodermal dysplasia 			
		Ectodermar ay sphasta		<u> </u>	·ـــــا

Table 1: Participation rates and sample composition

^IThis group is made up of individuals who wanted to participate but could not be accommodated because of the logistical difficulties associated with trying to find a suitable time to meet

Table 2. Sample characteristics. Fatients and patient representatives			
Genetic status	Patients	Patient representatives	TOTAL
	(choice of focus group or	(interviews only)	(interviews &
	interview)	n = 15	focus groups)
	n = 12		n = 27
*Patient advocate	0	3	3
Diagnosed with a genetic	4	7	11
condition			
At risk for developing a genetic	1	0	1
condition			
Parent/grandparent of an affected	7	5	12
child			

Table 2: Sample characteristics: Patients and patient representatives

*Not from a family affected by a genetic condition