



Long-term follow-up of patients with chronic myeloid leukemia having received autologous stem cell transplantation

Björn Hackanson, Cornelius F. Waller

► To cite this version:

Björn Hackanson, Cornelius F. Waller. Long-term follow-up of patients with chronic myeloid leukemia having received autologous stem cell transplantation. *Annals of Hematology*, 2010, 90 (4), pp.395-399. <10.1007/s00277-010-1094-y>. <hal-00583540>

HAL Id: hal-00583540

<https://hal.science/hal-00583540v1>

Submitted on 6 Apr 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Long-term follow-up of patients with chronic myeloid leukemia having received autologous stem cell transplantation

Björn Hackanson ¹ and Cornelius F. Waller ¹

¹Department of Internal Medicine, Division of Hematology and Oncology, Albert- Ludwigs- University Freiburg, 79106 Freiburg, Germany;

Corresponding Author: Cornelius F. Waller, MD
Department of Internal Medicine
Division of Hematology and Oncology
Albert-Ludwigs-University, Freiburg,
Hugstetter Str. 55
79106 Freiburg, Germany
Phone: (49) 761-270-3422
Fax: (49) 761-270-3418
E-mail: cornelius.waller@uniklinik-freiburg.de

Abstract

Prior to introduction of imatinib mesylate, the median survival of chronic myeloid leukemia (CML) patients was approximately 60 months and the standard treatment with interferon-alpha (IFN- α) resulted in major cytogenetic responses of 20-25 %. As an alternative treatment approach besides allogeneic transplantation intensive chemotherapy followed by autologous hematopoietic stem cell transplantation (autoHSCT) was investigated at that time with the rationale of debulking disease burden and mobilisation and transplantation of Philadelphia chromosome-negative (Ph-) stem cells.

In the era of tyrosine kinase inhibitors (TKIs) as state-of-the-art therapy for CML, the concept of autoHSCT has attracted only little interest and long-term follow-up and outcome data after autoHSCT in CML patients is scarce. In this long-term analysis, we report on 21 CML patients, mobilized in early chronic phase (ECP) and transplanted with largely Ph- grafts,

who received IFN α as maintenance therapy. Imatinib mesylate was administered upon cytogenetic relapse or disease progression while on IFN- α . The 10-year survival was 61% and 11 patients (52%) were alive at a median follow-up of 12.5 years (range 0.3 - 13.8) with 8 patients in complete hematologic remission (CHR) and 3 of 8 in major molecular remission (MMR). While all patients in MMR and 2 of 5 patients in CHR received imatinib, it is noteworthy that three patients remaining in CHR only received IFN- α maintenance after autoHSCT.

With the limitations of a small patient population, this is the longest follow-up analysis demonstrating that autoHSCT in CML is very efficient to debulk the disease, restore Ph-negative hematopoiesis and is able to induce major and sustained molecular responses in the majority of patients with substantial long-term survival rates.

Keywords: CML; autologous transplantation; long-term follow up

Introduction

Chronic myeloid leukemia (CML) is characterized by the presence of the Philadelphia chromosome, a reciprocal translocation that results in the BCR-ABL fusion gene (1-3). Deciphering of the BCR-ABL protein as a tyrosine kinase (TK) and the development of several substances, which are able to block BCR-ABL's increased tyrosine kinase activity, have revolutionized the treatment of CML (4, 5). Currently, Imatinib mesylate is considered standard of care for first-line treatment of chronic phase CML (6, 7).

However, in the 1990's, before the introduction of TKIs, treatment intensification of CML patients in early chronic phase (ECP) with high-dose chemotherapy (e.g. busulfan, melphalan and cyclophosphamide combined with total body irradiation in some studies) followed by autoHSCT and IFN α maintenance was a promising and intensively investigated treatment regimen (8-14). The strong rationale behind this treatment approach was the concept of debulking the disease burden, mobilization of Ph- hematopoietic stem cells and consecutive transplantation of a largely Ph- autograft.

While some observations pointed to increased remission rates and prolonged survival when autografting was performed in ECP, a meta-analysis of six randomized studies was unable to show a benefit for autoHSCT compared to IFN α -based therapy with regard to event free survival and overall survival (15). Whether there still exists a role for autoHSCT in the current management of CML, e.g. to reverse imatinib resistance or to eliminate clones harbouring BCR-ABL kinase domain mutations remains to be determined (15, 16).

Interestingly, only few long-term data on autografted patients are available (15, 18). Therefore, we report our long-term follow-up analysis on 21 CML patients who were autografted in ECP between 1995 and 1999.

Patients and Methods

Patient selection, mobilization of CD 34⁺ stem cells and high-dose chemotherapy with autologous transplantation have been described in detail previously (10, 11). Briefly, after priming with rhG-CSF (5 µg/kg s.c.; Filgrastim, Amgen, Germany), mobilization chemotherapy consisted of idarubicin (8 mg/m², day 1-3), cytosine arabinoside (400 mg/m², day 1-3) and etoposide (150 mg/m², day 1-4). On day 7 after chemotherapy, administration of rhG-CSF (10 µg/kg s.c. daily) in all patients and rhIL-3 (5 µg/kg s.c, Novartis, Germany) in 12 patients was started until stem cell harvest. After apheresis, positive selection for CD 34⁺ cells was performed using avidin-immunoaffinity columns (CellPro, Washington, USA) or Isolex magnetic cell separation (Baxter, Unterschleissheim, Germany). High-dose chemotherapy contained busulphan (4 mg/kg day -7 to -4) and cyclophosphamide (60 mg/kg day -1 to -2). After autologous transplantation of a minimum of 1.5 x 10⁶ CD 34⁺ cells, intravenous rhG-CSF treatment was started on day +3 until neutrophil recovery (> 1.5 x 10⁹/l). 20 patients were transplanted in first chronic phase and one patient in accelerated phase with a median time from initial diagnosis of 12 months (**Table 1**). Following platelet regeneration, patients were treated with IFN-α (initially 3 x 3x10⁶ units/week; dose reduction to 3 x 1.5x10⁶ units/week or less if side effects occurred) until complete cytogenetic/molecular remission, onset of intolerance or progressive disease. If imatinib was started, the initial dose was 400 mg daily. One patient received Nilotinib 2 x 400 mg daily and Dasatinib 2 x 70 mg daily after imatinib failure and one patient received Nilotinib 2 x 400 mg daily upon imatinib failure. Patient's characteristics are provided in **Table 1**.

Overall survival and median follow-up were estimated using the Kaplan-Meier method.

Results

Of 21 CML patients transplanted with an autograft between 1995 and 1999 at our institution, 11 patients were alive at follow-up analysis in 2009. Three patients (one receiving imatinib, one nilotinib and one dasatinib) were in complete hematologic remission (CHR) but without complete regeneration. Five patients (two on imatinib, three on IFN α) were in CHR and three patients were in major molecular remission (MMR) (**Table 2**). It is noteworthy that three patients remaining in CHR only received IFN α maintenance after autoHSCT, while all patients in MMR and 2/5 patients in CHR received imatinib.

Five patients had died after a median of 36 months (range 20-150) due to progressive disease with only one patient having received imatinib (**Table 2**). Three patients had received allogeneic stem cell transplantation (alloHSCT) in complete hematologic remission, chronic or accelerated phase, three, four and nine years after autoHSCT, respectively. Unfortunately, all three died due to infectious complications or graft-versus-host disease following alloHSCT. One patient died two months after autoHSCT in CP due to toxic alveolitis and lung bleeding and one patient died seven years after autoHSCT of undetermined cause.

In our patient cohort, the 10-year survival rate after autologous transplantation was 61% (13/21) and overall survival after a median follow-up of 12.5 years was 52 % (11/21) (**Figure 1**).

Discussion

Due to the impact of TKI in the first-line therapy of CP-CML only few data are available on long-term outcome after autoHSCT in CML patients. However, as some patients have primary or develop secondary resistance to imatinib and other TKIs and are not eligible for alloHSCT, autoHSCT might still be a treatment option in a defined subset of CML patients.

Simonsson et al. showed a median survival of 7.5 years after autoHSCT in the pre-imatinib era (16). However, in a meta-analysis of this and other trials from several groups, a benefit of autoHSCT in the initial treatment of CML as compared to interferon- α (alone or in combination) could not be demonstrated (15). Furthermore, it has to be mentioned that all trials involved in that analysis closed early because of the introduction of imatinib. Of those trials, the German CML III A trial is still ongoing for analysis.

However, intensive chemotherapy for HSC-mobilization followed by autoHSCT of Ph-negative grafts has been shown to be effective in debulking leukemia burden and to induce good cytogenetic and molecular responses maintained by low-dose IFN- α in some patients as demonstrated by our data and others' (10-14). Recently, some very promising studies have shown that mobilization of Ph-negative peripheral blood progenitor cells by G-CSF alone while patients are being treated with imatinib is feasible (19, 20). However, data on successful transplantation of such cells has not been investigated thoroughly, yet (21).

In our study all patients had received mobilization chemotherapy plus filgrastim in order to collect CD34⁺ stem cells. While in this study CML patients were treated with TKIs several years after autoHSCT, in the era of imatinib and 2nd and 3rd generation TKIs a potential treatment concept would rather consist of initial treatment with TKIs and a stem cell harvest in CCR and/or MMR. In case of TKI-resistance later on in the course of CML, autoHSCT could be one possible treatment option for patients who are ineligible for alloHSCT.

Compared to the median survival in CP-CML of 60 months prior to introduction of imatinib our data on a small cohort of selected patients show substantial overall survival after

treatment with intensive chemotherapy and autologous stem cell transplantation, followed by TKI therapy in the majority of patients. However, the significance of maintenance therapy with IFN- α and that of subsequent therapy with imatinib or another TKI is difficult to interpret (22). The initial debulking by chemotherapy might be of primary importance in this setting. As the precise role of transplantation of autologous Ph-negative PBPC in the management of CML patients resistant to current first-line therapy with TKIs is still unknown further evaluation in suitably designed clinical trials is warranted.

Literature

1. Koeffler HP, Golde DW. Chronic myelogenous leukemia--new concepts. *N Engl J Med* 1981; **304**: 1201-1209.
2. Daley GQ, Van Etten RA, Baltimore D. Induction of chronic myelogenous leukemia in mice by the P210bcr/abl gene of the Philadelphia chromosome. *Science* 1990 16;**247**:824-830.
3. Sawyers CL. Chronic myeloid leukemia. *N Engl J Med* 1999; **340**: 1330-1340.
4. Druker BJ, Tamura S, Buchdunger E, Ohno S, Segal GM, Fanning S et al. Effects of a selective inhibitor of the Abl tyrosine kinase on the growth of Bcr-Abl positive cells. *Nat Med* 1996; 2: 561-566.
5. Baccarani M, Cortes J, Pane F, Niederwieser D, Saglio G, Apperley J et al. Chronic myeloid leukemia: an update of concepts and management recommendations of European LeukemiaNet. *J Clin Oncol* 2009; **27**:6041-6051.
6. Peggs K, Mackinnon S. Imatinib mesylate--the new gold standard for treatment of chronic myeloid leukemia. *N Engl J Med* 2003; 348: 1048-1050.
7. Hochhaus A, O'Brien SG, Guilhot F, Druker BJ, Branford S, Foroni L et al. Six-year follow-up of patients receiving imatinib for the first-line treatment of chronic myeloid leukemia. *Leukemia* 2009; **23**: 1054-1061.
8. McGlave PB, De Fabritiis P, Deisseroth A, Goldman J, Barnett M, Reiffers J et al. Autologous transplants for chronic myelogenous leukaemia: results from eight transplant groups. *Lancet* 1994; 343:1486-1488.
9. Bhatia R, Verfaillie CM, Miller JS, McGlave PB. Autologous Transplantation Therapy for Chronic Myelogenous Leukemia. *Blood*. 1997; 89:2623-2634.
10. Waller CF, Heinzinger M, Rosenstiel A, Lange W. Mobilization and transplantation of Philadelphia chromosome-negative peripheral blood progenitor cells in patients with CML. *Br J Haematol* 1998; 103: 227-234.
11. Heinzinger M, Waller CF, Rosenstiel A, Scheid S, Burger KJ, Lange W. Quality of IL-3 and G-CSF-mobilized peripheral blood stem cells in patients with early chronic phase CML. *Leukemia* 1998; **12**: 333-339.
12. Bergamaschi G, Podestà M, Frassoni F, Rosti V, Carella AM, Saglio G et al. Restoration of normal polyclonal haemopoiesis in patients with chronic myeloid leukaemia autografted with Ph-negative peripheral stem cells. *Br J Haematol* 1994;**87**:867-870.
13. Carella AM, Cunningham I, Lerma E, Dejana A, Benvenuto F, Podestà M et al. Mobilization and transplantation of Philadelphia-negative peripheral-blood progenitor cells early in chronic myelogenous leukemia. *J Clin Oncol* 1997;**15**:1575-1582.

14. Carella AM, Beltrami G, Corsetti MT. Autografting in chronic myeloid leukemia. *Semin Hematol* 2003; **40**: 72-78.
15. CML Autograft Trials Collaboration. Autologous stem cell transplantation in chronic myeloid leukaemia: a meta-analysis of six randomized trials. *Cancer Treat Rev* 2007; **33**: 39-47.
16. Olavarria E. Autologous stem cell transplantation in chronic myeloid leukemia. *Semin Hematol* 2007; **44**: 252-258.
17. Perseghin P, Gambacorti-Passerini C, Tornaghi L, Dassi M, Pioltelli P, Parma M et al. Peripheral blood progenitor cell collection in chronic myeloid leukemia patients with complete cytogenetic response after treatment with imatinib mesylate. *Transfusion* 2005; **45**: 1214-1220.
18. Simonsson B, Oberg G, Bjoreman M, Bjorkholm M, Carneskog J, Karlsson K et al. Intensive treatment and stem cell transplantation in chronic myelogenous leukemia: long-term follow-up. *Acta Haematol* 2005; **113**: 155-62.
19. Kreuzer KA, Klühs C, Baskaynak G, Movassaghi K, Dörken B, le Coutre P. Filgrastim-induced stem cell mobilization in chronic myeloid leukaemia patients during imatinib therapy: safety, feasibility and evidence for an efficient in vivo purging. *Br J Haematol* 2004; **124**: 195-9.
20. Hui CH, Goh KY, White D, Branford S, Grigg A, Seymour JF et al. Successful peripheral blood stem cell mobilisation with filgrastim in patients with chronic myeloid leukaemia achieving complete cytogenetic response with imatinib, without increasing disease burden as measured by quantitative real-time PCR. *Leukemia* 2003; **17**: 821-8.
21. le Coutre P, Kreuzer KA, Massenkeil G, Baskaynak G, Zschieschang P, Genvresse I et al. Autologous peripheral blood stem cell transplantation of stem cells harvested in imatinib-induced complete cytogenetic remission: an example of in vivo purging in CML. *Leukemia* 2003; **17**: 2525-6.
22. Fischer T, Reifenrath C, Hess GR, Corsetti MT, Kreil S, Beck J et al. Safety and efficacy of STI-571 (imatinib mesylate) in patients with bcr/abl-positive chronic myelogenous leukemia (CML) after autologous peripheral blood stem cell transplantation (PBSCT). *Leukemia* 2002; **16**: 1220-1228.

Tables

Table 1. Patient characteristics

Patients characteristics	
Male/female	16/5
Age at transplantation (years)	46 (25-58)
Remission status prior to transplantation (19 patients evaluable)	18 CP, 1 AP
Sokal Score prior to transplantation (data of 11 patients evaluable)	0,7 (0,52-1,74)
Time from initial diagnosis until transplantation (months)	12 (5-64)
Time from transplantation until first imatinib treatment (11 patients; years)	7 (4-13)
Follow-up after transplant (years)	12.5 (0.3-13.8)

Values are given as median and range
CP= chronic phase, AP= accelerated phase

Table 2. Time course of 21 CML patients with remission status (where available) from transplantation to last follow-up or death.

	1995	96	97	98	99	2000	01	02	03	04	05	06	07	08	09
1	Tx	▲●			▲			▲	▲	▲	▲		▲	▲	▲ a
2		Tx	BC	BC ж											
3	Tx	▲■	▲■		CP	CP	CP	▲□			▲	▲	▲□♦	▲□	▲□ a
4	Tx	CP			CP	CP	CP	AC♦	▲	▲al.Tx	ж				
5		Tx	▲●	AC	CP	al.Tx	ж								
6		Tx	▲■			▲		▲	●	▲■	▲	▲	BC♦	BC	
7		Tx	AC		2.Tx	BC ж								ж	
8		Tx	▲●	▲○		▲○	▲■		▲■		▲	▲		▲	▲ a
9		Tx			CP	♦	CP	▲■	▲□	▲		▲■	▲	▲	▲□ a
10		Tx		CP	AC al.Tx	ж									
11		Tx	▲	▲■	▲■	▲■	▲■	▲■	▲■						▲ a
12		Tx	▲	▲●	▲	▲●	▲	▲	▲	▲●	▲	▲	▲	▲○	▲♦ a
13		Tx	CP	CP	CP	CP	CP	CP♦			CP●	CP●			▲x a
14		Tx	CP	CP	CP	CP	CP♦	CP	CP	CP	CP	CP	CP♦♦	CP	▲x a
15		Tx	▲	▲	▲	CP	CP	CP	CP	▲♦	▲	▲		CP	▲x♦ a
16		Tx	CP	CP	CP	CP	CP	▲	▲♦	▲	▲	▲	▲	▲□	▲ a
17			Tx	ж											
18				Tx	CP	BC ж									
19				Tx	CP	CP	CP	CP	CP	CP	CP	CP	CP	CP	BC ж
20					Tx	CP	▲	CP	CP	CP♦	▲	ж			
21					Tx	▲	▲	▲	▲	▲	▲	▲	▲	▲□♦	▲□ a

Tx = transplantation; ж = death; a = alive
 ▲ = CHR (complete hematologic response)
 ▲x = CHR without complete regeneration
 ○ = MCR (minimal cytogenetic remission)
 ● = PCR (partial cytogenetic remission)
 ■ = CCR (complete cytogenetic remission)

□ = MMR (major molecular remission)
 ■ = CMR (complete molecular remission)
 CP = chronic phase
 AC = acceleration
 BC = blast crisis
 al.Tx = allogeneic stem cell transplantation
 ♦ = Start with Imatinib
 ♦ = Start with Nilotinib
 ♦ = Start with Dasatinib

Figure legends

Figure 1. Overall survival of CML patients from time of transplantation using the Kaplan-Meier method.

Figure 1.

