



**HAL**  
open science

# Eigenvectors of Wigner matrices: universality of global fluctuations

Florent Benaych-Georges

► **To cite this version:**

Florent Benaych-Georges. Eigenvectors of Wigner matrices: universality of global fluctuations. 2011. hal-00583889v1

**HAL Id: hal-00583889**

**<https://hal.science/hal-00583889v1>**

Preprint submitted on 7 Apr 2011 (v1), last revised 28 Sep 2012 (v6)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# EIGENVECTORS OF WIGNER MATRICES: UNIVERSALITY OF GLOBAL FLUCTUATIONS

FLORENT BENAYCH-GEORGES

ABSTRACT. We prove that for  $[u_{i,j}]_{i,j=1}^n$  the eigenvectors matrix of a Wigner matrix, under some moments conditions, the bivariate random process

$$\left( \sum_{\substack{1 \leq i \leq ns, \\ 1 \leq j \leq nt}} (|u_{i,j}|^2 - 1/n) \right)_{(s,t) \in [0,1]^2}$$

converges in distribution to a bivariate Brownian bridge.

## 1. INTRODUCTION

It is well known that the matrix  $U_n = [u_{i,j}]_{i,j=1}^n$  whose columns are the eigenvectors of a GOE or GUE matrix  $X_n$  can be chosen to be distributed according to the Haar measure on the orthogonal or unitary group. As a consequence, much can be said about the  $u_{i,j}$ 's: their joint moments can be computed via the so-called *Weingarten calculus* developed in [7, 8], any finite (or not too large) set of  $u_{i,j}$ 's can be approximated, as  $n \rightarrow \infty$ , by independent Gaussian variables (see [14, 6]) and the global asymptotic fluctuations of the  $|u_{i,j}|$ 's are governed by a theorem of Donati-Martin and Rouault, who proved in [9] that as  $n \rightarrow \infty$ , the bivariate càdlàg process

$$\left( B_{s,t}^n := \sqrt{\frac{2}{\beta}} \sum_{\substack{1 \leq i \leq ns, \\ 1 \leq j \leq nt}} (|u_{i,j}|^2 - 1/n) \right)_{(s,t) \in [0,1]^2}$$

(where  $\beta = 1$  in the real case and  $\beta = 2$  in the complex case) converges in distribution, for the Skorokhod topology, to the *bivariate Brownian bridge*, i.e. the centered continuous Gaussian process  $(B_{s,t})_{(s,t) \in [0,1]^2}$  with covariance

$$(1) \quad \mathbb{E}[B_{s,t} B_{s',t'}] = (\min\{s, s'\} - ss')(\min\{t, t'\} - tt').$$

A natural question is the following:

What can be said beyond the Gaussian case, when the entries of the Wigner matrix  $X_n$  are not Gaussian anymore ?

---

*Date:* April 7, 2011.

*2000 Mathematics Subject Classification.* 15A52, 60F05.

*Key words and phrases.* Random matrices, Haar measure, eigenvectors, Wigner matrices, bivariate Brownian motion, bivariate Brownian bridge.

This work was partially supported by the *Agence Nationale de la Recherche* grant ANR-08-BLAN-0311-03.

For a general Wigner matrix<sup>1</sup>, the exact distribution of the matrix  $U_n$  cannot be computed and few works had been devoted to this subject until quite recently. However, in the last three years, the eigenvectors of Wigner matrices have been the object of a growing interest, due in part to some relations with the universality conjecture for the eigenvalues. In several papers (see, among others, [10, 11, 12]), a *delocalization property* was shown for the eigenvectors of random matrices. More recently, Knowles and Yin in [16] and Tao and Vu in [19] proved that if the first four moments of the atom distributions of  $X_n$  coincide with the ones of a GOE or GUE matrix, then under some tail assumptions on these distributions, the  $u_{i,j}$ 's can be approximated by independent Gaussian variables as long as we only consider a finite (or not too large) set of  $u_{i,j}$ 's.

In this paper, we consider the global behavior of the  $|u_{i,j}|$ 's, and we prove (Theorem 2.3) that for Wigner matrices whose entries have moments of all orders, the process  $(B_{s,t}^n)_{(s,t) \in [0,1]^2}$  has a limit in a weaker sense than for the Skorokhod topology and that this weak limit is the bivariate Brownian bridge if and only if the off-diagonal entries of the matrix have the same fourth moment as the GOE or GUE matrix (quite surprisingly, no hypothesis on the third moment is necessary). Under some additional hypotheses on the atom distributions (more coinciding moments and continuity), we prove the convergence for the Skorokhod topology (Theorem 2.6).

The paper is organized as follows. The main results are stated in Section 2, where we also make some comments on their hypotheses, an outline of their proofs and their formal proofs are given in Section 3 and, at last, Section 4 is devoted to the definitions of the functional spaces and their topologies and to the proofs of several technical results needed in Section 3.

**Acknowledgements:** It is a pleasure for the author to thank Djalil Chafaï for having generously introduced him to this interesting question and for the nice discussions we had about it. We also would like to thank Alice Guionnet, for her availability and her precious advices, and Terry Tao, who kindly answered several naive questions asked by the author on his blog.

## 2. MAIN RESULTS

For each  $n$ , let us consider a real symmetric or Hermitian random matrix

$$X_n := \frac{1}{\sqrt{n}} [x_{i,j}^{(n)}]_{i,j=1}^n.$$

For notational brevity,  $x_{i,j}^{(n)}$  will be denoted by  $x_{i,j}$ . We make the following hypotheses.

Let us denote by  $\lambda_1 \leq \dots \leq \lambda_n$  the eigenvalues of  $X_n$  and consider an orthogonal or unitary matrix  $U_n = [u_{i,j}]_{i,j=1}^n$  such that

$$X_n = U_n \text{diag}(\lambda_1, \dots, \lambda_n) U_n^*.$$

Note that  $U_n$  is not uniquely defined. However, one can choose it in any measurable way.

We define the bivariate càdlàg process

$$\left( B_{s,t}^n := \sqrt{\frac{2}{\beta}} \sum_{\substack{1 \leq i \leq ns, \\ 1 \leq j \leq nt}} (|u_{i,j}|^2 - 1/n) \right)_{(s,t) \in [0,1]^2},$$

---

<sup>1</sup>A *Wigner matrix* is a real symmetric or Hermitian random matrix with independent, centered entries whose variance is one. Its *atom distributions* are the distributions of its entries.

where  $\beta = 1$  in the real case and  $\beta = 2$  in the complex case.

**Assumption 2.1.** *For each  $n$ , the random variables  $x_{i,j}$ 's are independent (up to the symmetry), with the same distribution on the diagonal and the same distribution above the diagonal.*

**Assumption 2.2.** *For each  $k \geq 1$ ,  $\sup_n \mathbb{E}[|x_{1,1}|^k + |x_{1,2}|^k] < \infty$ . Moreover,*

$$(2) \quad \mathbb{E}[x_{1,1}] = \mathbb{E}[x_{1,2}] = 0, \quad \mathbb{E}[|x_{1,2}|^2] = 1$$

and  $\mathbb{E}[|x_{1,2}|^4]$  has a limit as  $n \rightarrow \infty$ .

The definitions of the functional spaces and their topologies can be found in Section 4.1 and the bivariate Brownian bridge has been defined in the introduction.

**Theorem 2.3.** *Suppose that Assumptions 2.1 and 2.2 are satisfied. Then the sequence*

$$(\text{distribution}(B^n))$$

*has a unique possible accumulation point supported by  $C([0,1]^2)$ . This accumulation point is the distribution of a centered Gaussian process which depends on the distributions of the  $x_{i,j}$ 's only through  $\lim_{n \rightarrow \infty} \mathbb{E}[|x_{1,2}|^4]$ , and which is the bivariate Brownian bridge if and only if  $\lim_{n \rightarrow \infty} \mathbb{E}[|x_{1,2}|^4] = 4 - \beta$ .*

To get a stronger statement where the convergence in distribution to the bivariate Brownian bridge is actually stated, one needs stronger hypotheses.

**Assumption 2.4.** *For each  $k \geq 1$ ,  $\sup_n \mathbb{E}[|x_{1,1}|^k + |x_{1,2}|^k] < \infty$ . Moreover, for each  $r, s \geq 0$ ,*

$$(3) \quad r + s \leq 10 \implies \mathbb{E}[\Re(x_{1,1})^r \Im(x_{1,1})^s] = \mathbb{E}[\Re(g_{1,1})^r \Im(g_{1,1})^s]$$

and

$$(4) \quad r + s \leq 12 \implies \mathbb{E}[\Re(x_{1,2})^r \Im(x_{1,2})^s] = \mathbb{E}[\Re(g_{1,2})^r \Im(g_{1,2})^s].$$

where the  $g_{i,j}$ 's are the entries of a standard GOE or GUE matrix.

**Assumption 2.5.** *The distributions of the entries of  $X_n$  are absolutely continuous with respect to the Lebesgue measure.*

**Theorem 2.6.** *Suppose that Assumptions 2.1, 2.4 and 2.5 are satisfied. Then as  $n \rightarrow \infty$ , bivariate process  $B^n$  converges in distribution, for the Skorokhod topology in  $D([0,1]^2)$ , to the bivariate Brownian bridge.*

**Remark 2.7. Weakening of the assumptions.** As explained above, the distributions of the entries  $x_{i,j} = x_{i,j}^{(n)}$  are allowed to depend on  $n$ . In order to remove Assumption 2.5 below (which we did not manage to do yet), it might be useful to weaken Assumptions 2.2 and 2.4 in the following way: one can easily see that the proofs still work if in Equations (2), (3) and (4), one replaces the identities by the same identities up to an error which is  $O(n^{-\alpha})$  for all  $\alpha > 0$ .

**Remark 2.8. Comments on the hypotheses.** In order to prove the convergence in the Skorokhod topology, we had to suppose the continuity of the atom distributions and the coincidence of their 10 (on the diagonal) and 12 (above the diagonal) first moments with the ones of a GOE or GUE matrix. The reason is that we needed these assumptions to control the discontinuities of the process  $B^n$ . Even though these hypotheses might not be optimal, a bound on the tails of the atom distributions seems to be necessary to avoid too large variations of the process  $B^n$ . Indeed, as shown by Figure 1, while for a GOE matrix (*left picture*),  $|u_{i,j}|^2$  is close to  $1/n$  for all  $i, j$  with high probability, when the atom distributions have not more than a second moment (*right picture*), the matrix  $X_n$  looks more like a *sparse matrix*, and so does  $U_n$ , which implies that for certain  $(i, j)$ 's,  $(|u_{i,j}|^2 - 1/n) \gg 0$ .




FIGURE 1. **Influence of the tails of the atom distributions of  $X_n$  on the  $|u_{i,j}|^2$ 's:** Plot of the map  $(i/n, j/n) \mapsto ||u_{i,j}|^2 - 1/n|$  for two different choices of atom distributions. *Left:* GOE matrix. *Right:* Wigner matrix with atom distribution admitting moments only up to order  $2 + \varepsilon$  for a small  $\varepsilon$ . For both pictures,  $n = 50$ .

### 3. PROOFS OF THEOREMS 2.3 AND 2.6

**3.1. Outline of the proofs.** Firstly, Theorem 2.6 can be deduced from Theorem 2.3 by proving that the sequence  $(\text{distribution}(B^n))$  is tight and only has  $C([0, 1]^2)$ -supported accumulation points. This can be done via some upper bounds on the fourth moment of the increments of  $B^n$  and on its jumps (or discontinuities). These bounds are given in the proof of Lemma 3.1 below, and rely on the existing bounds in the case where  $X_n$  is a GOE or GUE matrix and on the “one-by-one entries replacement method” developed by Terence Tao and Van Vu in recent papers, such as [18, 19].

Secondly, the proof of Theorem 2.3 relies on the following remark, inspired by some ideas of Jack Silverstein (see [2, Chap. 10] and [17]): even though we do not have any “direct access” to the eigenvectors of  $X_n$ , we have access to the process  $(B_{s, F_{\mu_{X_n}}^n}^n)_{s \in [0, 1], u \in \mathbb{R}}$ , for  $F_{\mu_{X_n}}(u) := \frac{1}{n} \#\{i; \lambda_i \leq u\}$ . Indeed,

$$B_{s, F_{\mu_{X_n}}^n}^n = \sqrt{\frac{2}{\beta}} \sum_{1 \leq i \leq ns} \sum_{\substack{1 \leq j \leq n \\ \text{s.t. } \lambda_j \leq u}} (|u_{i,j}|^2 - 1/n),$$

hence for all fixed  $s \in [0, 1]$ , the function  $u \in \mathbb{R} \mapsto B_{s, F_{\mu_{X_n}}^n}^n$  is the cumulative distribution function of the signed measure

$$(5) \quad \sqrt{\frac{2}{\beta}} \sum_{1 \leq i \leq ns} \sum_{j=1}^n (|u_{i,j}|^2 - 1/n) \delta_{\lambda_j},$$

which can be studied via its moments

$$\sum_{1 \leq i \leq ns} \left( e_i^* X_n^k e_i - \frac{1}{n} \operatorname{Tr} X_n^k \right) \quad (k \geq 1),$$

where the  $e_i$ 's are the vectors of the canonical basis. From the asymptotic behavior of the *moments* of the signed measure of (5), one can find out the asymptotic behavior of its *cumulative distribution function*.

Then, once the asymptotic distribution of the process  $(B_{s, F_{\mu_{X_n}}(u)}^n)_{s \in [0,1], u \in \mathbb{R}}$  identified, one can obtain the asymptotic distribution of the process  $(B_{s,t}^n)_{s \in [0,1], t \in [0,1]}$  because the function  $F_{\mu_{X_n}}$  tends to the (non random) cumulative distribution function  $F_{\text{semicircle}}$  of the semicircle law.

**3.2. Formal proofs.** By a standard *tightness + uniqueness of the accumulation point* argument, Theorem 2.6 will follow from the following lemma and Theorem 2.3. The proof of the lemma is postponed to Section 4.4.

**Lemma 3.1.** *Under Assumptions 2.1, 2.4 and 2.5, the sequence (distribution( $B^n$ )) is  $C$ -tight, i.e. is tight and has only  $C([0, 1]^2)$ -supported accumulation points.*

So let us now prove Theorem 2.3 under Assumptions 2.1 and 2.2. Note that it suffices to prove that the sequence (distribution( $B^n$ )) has a unique possible accumulation point supported by  $C([0, 1]^2)$  and that this accumulation point is the distribution of a centered Gaussian process which depends on the distributions of the  $x_{i,j}$ 's only through  $\lim_{n \rightarrow \infty} \mathbb{E}[|x_{1,2}|^4]$  (and actually does depend on  $\lim_{n \rightarrow \infty} \mathbb{E}[|x_{1,2}|^4]$ ). Indeed, in this case, by Theorem 1.1 of [9], this limit distribution is the bivariate Brownian bridge if and only if  $\lim_{n \rightarrow \infty} \mathbb{E}[|x_{1,2}|^4]$  is the same as for  $X_n$  a GOE or GUE matrix, i.e. equal to  $4 - \beta$ .

The following proposition is the key of the proof, since it allows to transfer the problem from the eigenvectors to some more accessible objects: the weighted spectral distributions of  $X_n$ .

**Proposition 3.2.** *Let  $e_i$  denote the  $i$ th vector of the canonical basis. To prove Theorem 2.3, it suffices to prove that each finite-dimensional marginal distribution of the process*

$$\left( \sum_{1 \leq i \leq ns} \left( e_i^* X_n^k e_i - \frac{1}{n} \operatorname{Tr} X_n^k \right) \right)_{s \in [0,1], k \geq 1}$$

*converges to a centered Gaussian measure and that the covariance of the limit process depends on the distributions of the  $x_{i,j}$ 's only through  $\lim_{n \rightarrow \infty} \mathbb{E}[|x_{1,2}|^4]$  and actually does depend on  $\lim_{n \rightarrow \infty} \mathbb{E}[|x_{1,2}|^4]$ .*

**Proof.** Let  $\mu_{X_n} := \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i}$  be the empirical spectral law of  $X_n$  and  $F_{\mu_{X_n}}(u) := \frac{1}{n} \#\{i; \lambda_i \leq u\}$  be its cumulative distribution function. We know that  $F_{\mu_{X_n}}$  converges in probability, as  $n$  tends to infinity, to the cumulative distribution function  $F_{\text{semicircle}}$  of the semicircle law in the space  $D_c(\mathbb{R}, [0, 1])$  (see Section 4.1 for the definitions of the functional spaces and their topologies). Moreover, the map

$$\begin{aligned} D_0([0, 1]^2) \times D_c(\mathbb{R}, [0, 1]) &\longrightarrow D_c([0, 1] \times \mathbb{R}) \\ ((G_{s,t})_{s,t \in [0,1]}, (g(u))_{u \in \mathbb{R}}) &\longmapsto (G_{s,g(u)})_{(s,u) \in [0,1] \times \mathbb{R}} \end{aligned}$$

is continuous at any pair of continuous functions. Hence for any continuous process  $(B_{s,t})_{s,t \in [0,1]}$  whose distribution is an accumulation point of the sequence (distribution( $B^n$ )) for the Skorokhod

topology in  $D([0, 1]^2)$ , the process

$$(B_{s, F_{\mu_{X_n}}^n}(u))_{(s,u) \in [0,1] \times \mathbb{R}}$$

converges in distribution (up to the extraction of a subsequence) to the process

$$(B_{s, F_{\text{semicircle}}}(u))_{(s,u) \in [0,1] \times \mathbb{R}}$$

(for the reader who is not used to weak convergence of probability measures, this assertion relies on Theorem 4.4 and Corollary 1 of Theorem 5.1 in Chapter 1 of [5]). Now, note that  $F_{\text{semicircle}} : \mathbb{R} \rightarrow [0, 1]$  admits a right inverse, so the distribution of the process  $(B_{s,t})_{s,t \in [0,1]}$  is entirely determined by the one of the process  $(B_{s, F_{\text{semicircle}}}(u))_{(s,u) \in [0,1] \times \mathbb{R}}$ .

As a consequence, to prove Theorem 2.3, it suffices to prove that the distribution of the process

$$(B_{s, F_{\mu_{X_n}}^n}(u))_{(s,u) \in [0,1] \times \mathbb{R}}$$

admits a unique  $C_c([0, 1] \times \mathbb{R})$ -supported accumulation point as  $n \rightarrow \infty$ .

Now, note that any  $f \in C_c([0, 1] \times \mathbb{R})$  is entirely determined by the collection of real numbers  $(\int_{u \in \mathbb{R}} u^k f(s, u) du)_{s \in [0,1], k \geq 0}$  in a way that preserves “the fact of being Gaussian and centered” if  $f$  is random. Indeed, let us fix  $(s, u_0) \in [0, 1] \times \mathbb{R}$ , let  $M > 0$  be such that the support of  $f(s, \cdot)$  is contained in  $[-M, M]$  and let, for each  $p \geq 1$ ,  $(P_{p,q})_{q \geq 1}$  be a sequence of polynomials that is uniformly bounded on  $[-M, M]$  and that converges pointwise to  $\mathbb{1}_{[u_0, u_0+1/p]}$  on  $[-M, M]$ . Then one has

$$f(s, u_0) = \lim_{p \rightarrow \infty} p \int_{u_0}^{u_0+1/p} f(s, u) du = \lim_{p \rightarrow \infty} \lim_{q \rightarrow \infty} p \int_{u \in \mathbb{R}} P_{p,q}(u) f(s, u) du.$$

As a consequence, to prove Theorem 2.3, it suffices to prove that each finite-dimensional marginal distribution of the process

$$\left( \int_{u \in \mathbb{R}} u^k B_{s, F_{\mu_{X_n}}^n}(u) du \right)_{s \in [0,1], k \geq 0}$$

converges to a centered Gaussian measure and that the covariance of the limit process depends on the distributions of the  $x_{i,j}$ 's only through  $\lim_{n \rightarrow \infty} \mathbb{E}[|x_{1,2}|^4]$  and actually does depend on  $\lim_{n \rightarrow \infty} \mathbb{E}[|x_{1,2}|^4]$ .

But for all  $(s, u) \in [0, 1] \times \mathbb{R}$ ,

$$B_{s, F_{\mu_{X_n}}^n}(u) = \sqrt{\frac{2}{\beta}} \sum_{1 \leq i \leq ns} \sum_{\substack{1 \leq j \leq n \\ \text{s.t. } \lambda_j \leq u}} (|u_{i,j}|^2 - 1/n),$$

hence

$$(6) \quad B_{s, F_{\mu_{X_n}}^n}(u) = \sqrt{\frac{2}{\beta}} \sum_{1 \leq i \leq ns} F_{\mu_{X_n, e_i} - \mu_{X_n}}(u),$$

where:

- $\mu_{X_n}$  is still the empirical spectral law of  $X_n$ ,
- $\mu_{X_n, e_i}$  is the weighted spectral law of  $X_n$ , defined by  $\mu_{X_n, e_i} := \sum_{j=1}^n |u_{i,j}|^2 \delta_{\lambda_j}$ ,
- $F_{\mu_{X_n, e_i} - \mu_{X_n}}$  is the cumulative distribution function of the null-mass signed measure  $\mu_{X_n, e_i} - \mu_{X_n}$ .

The following lemma will allow to conclude the proof of Proposition 3.2.

**Lemma 3.3.** *Let  $\mu$  be a compactly supported null-mass signed measure and set  $F_\mu(u) := \mu((-\infty, u])$ . Then for all  $k \geq 0$ ,*

$$\int_{u \in \mathbb{R}} u^k F_\mu(u) du = - \int_{x \in \mathbb{R}} \frac{x^{k+1}}{k+1} d\mu(x).$$

**Proof.** Let  $a < b$  be such that the support of  $\mu$  is contained in the open interval  $(a, b)$ .  $F_\mu$  is null out of  $(a, b)$  and satisfies  $F_\mu(u) = -\mu((u, b))$ , so by Fubini's Theorem,

$$\int_{u \in \mathbb{R}} u^k F_\mu(u) du = \int_{x=a}^b \int_{u=a}^x -u^k du d\mu(x) = \int_{x=a}^b \frac{-x^{k+1}}{k+1} d\mu(x) + \frac{a^{k+1}}{k+1} \mu((a, b)).$$

□

It follows from this lemma and from (6) that for all  $s \in [0, 1]$ ,  $k \geq 0$ ,

$$\int_{u \in \mathbb{R}} u^k B_{s, F_\mu X_n}^n(u) du = \frac{-1}{k+1} \sqrt{\frac{2}{\beta}} \sum_{1 \leq i \leq ns} \left( e_i^* X_n^{k+1} e_i - \frac{1}{n} \text{Tr} X_n^{k+1} \right),$$

which proves Proposition 3.2. □

Now, Theorem 2.3 is a direct consequence of the following proposition, whose proof is postponed below to Section 4.2.

**Proposition 3.4.** *Under Assumptions 2.1 and 2.2, each finite-dimensional marginal distribution of the process*

$$\left( \sum_{1 \leq i \leq ns} (e_i^* X_n^k e_i - \frac{1}{n} \text{Tr}(X_n^k)) \right)_{s \in [0,1], k \geq 1}$$

*converges to a centered Gaussian measure. The covariance of the limit distribution, denoted by*

$$(\text{Cov}_{s_1, s_2}(k_1, k_2))_{s_1, s_2 \in [0,1], k_1, k_2 \geq 1},$$

*depends on the distributions of the  $x_{i,j}$ 's only through  $\mathbb{E}[|x_{1,2}|^4]$ . Moreover, we have*

$$\text{Cov}_{s_1, s_2}(2, 2) = (\mathbb{E}[|x_{1,2}|^4] - 1)(\min\{s_1, s_2\} - s_1 s_2).$$

## 4. PROOFS OF THE TECHNICAL RESULTS

**4.1. Functional spaces and associated topologies.** In this paper, we use several functional spaces:

- $C([0, 1]^2)$  (resp.  $C_c([0, 1] \times \mathbb{R})$ ) is the set of continuous functions on  $[0, 1]^2$  (resp. *compactly supported* continuous function on  $[0, 1] \times \mathbb{R}$ ), endowed with the uniform convergence topology.
- $D_c(\mathbb{R}, [0, 1])$  is the set of *compactly supported* càdlàg functions on  $\mathbb{R}$  taking values in  $[0, 1]$ , endowed with the topology defined by  $f_n \rightarrow f$  if and only if the bounds of the support of  $f_n$  tend to the ones of the support of  $f$  and for all  $M > 0$ , after restriction to  $[-M, M]$ ,  $f_n \rightarrow f$  (the topology of  $D([-M, M])$  being deduced from the one of  $D([0, 1])$  defined in [5]).



- $D([0, 1]^2)$  (resp.  $D_c([0, 1] \times \mathbb{R})$ ) is the set of functions  $f : [0, 1]^2 \rightarrow \mathbb{R}$  (resp. of *compactly supported* functions  $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ ) admitting limits in all “orthants”, more precisely such that for each  $s_0, t_0$ , for each pair of symbols  $\diamond, \diamond' \in \{<, \geq\}$ ,

$$\lim_{\substack{s \diamond s_0 \\ t \diamond' t_0}} f(s, t)$$

exists, and is equal to  $f(s_0, t_0)$  if both  $\diamond$  and  $\diamond'$  are  $\geq$ . The space  $D([0, 1]^2)$  is endowed with the Skorokhod topology defined in [4] and the space  $D_c([0, 1] \times \mathbb{R})$  is endowed with the topology defined by  $f_n \rightarrow f$  if and only if for all  $M > 0$ , after restriction to  $[0, 1] \times [-M, M]$ ,  $f_n \rightarrow f$  (the topology of  $D([0, 1] \times [-M, M])$  being deduced from the one of  $D([0, 1]^2)$ ).

- $D_0([0, 1]^2)$  is the set of functions in  $D([0, 1]^2)$  vanishing at the border of  $[0, 1]^2$ , endowed with the induced topology.

**4.2. Proof of Proposition 3.4.** Note that by invariance of the law of  $X_n$  under conjugation by any permutation matrix, the expectation of the random law  $\mu_{X_n, e_i}$  does not depend on  $i$ . So for all  $s \in [0, 1]$ ,  $k \geq 1$ ,

$$(7) \quad \sum_{1 \leq i \leq ns} (e_i^* X_n^k e_i - \frac{1}{n} \text{Tr}(X_n^k)) = \sum_{1 \leq i \leq ns} (e_i^* X_n^k e_i - \mathbb{E}[e_i^* X_n^k e_i]) - \frac{\lfloor ns \rfloor}{n} \sum_{1 \leq i \leq n} (e_i^* X_n^k e_i - \mathbb{E}[e_i^* X_n^k e_i]).$$

Hence we are led to study the limit, as  $n \rightarrow \infty$ , of the finite-dimensional marginal distributions of the process

$$(8) \quad \left( \sum_{1 \leq i \leq ns} (e_i^* X_n^k e_i - \mathbb{E}[e_i^* X_n^k e_i]) \right)_{s \in [0, 1], k \geq 1}$$

The random variables of (8) are going to be studied via their joint moments. So let us fix  $p \geq 1$ ,  $s_1, \dots, s_p \in [0, 1]$  and  $k_1, \dots, k_p \geq 1$ . We shall study the limit, as  $n$  tends to infinity, of

$$(9) \quad \mathbb{E} \left[ \prod_{\ell=1}^p \sum_{1 \leq i \leq ns_\ell} (e_i^* X_n^{k_\ell} e_i - \mathbb{E}[e_i^* X_n^{k_\ell} e_i]) \right].$$

We introduce the set

$$(10) \quad \mathcal{E} := \{\overline{0}^1, \dots, \overline{k}_1^1\} \cup \dots \cup \{\overline{0}^p, \dots, \overline{k}_p^p\},$$

where the sets  $\{\overline{0}^1, \overline{1}^1, \dots\}$ ,  $\{\overline{0}^2, \overline{1}^2, \dots\}$ ,  $\dots$ ,  $\{\overline{0}^p, \overline{1}^p, \dots\}$  are  $p$  disjoint copies of the set of nonnegative integers. The set  $\mathcal{E}$  is ordered as presented in (10). For each partition  $\pi$  of  $\mathcal{E}$ , for each  $x \in \mathcal{E}$ , we denote by  $\pi(x)$  the index of the class of  $x$ , after having ordered the classes according to the order of their first element (for example,  $\pi(\overline{1}^1) = 1$ ;  $\pi(\overline{2}^1) = 1$  if  $\overline{1}^1 \preceq \overline{2}^1$  and  $\pi(\overline{2}^1) = 2$  if  $\overline{1}^1 \succ \overline{2}^1$ ).

By Assumption 2.1, the expectation of (9) can be expanded and expressed as a sum on the set  $\text{Part}(\mathcal{E})$  of partitions of the set  $\mathcal{E}$  introduced above. We get

$$(11) \quad \mathbb{E} \left[ \prod_{\ell=1}^p \sum_{1 \leq i \leq ns_\ell} (e_i^* X_n^{k_\ell} e_i - \mathbb{E}[e_i^* X_n^{k_\ell} e_i]) \right] = n^{-\frac{k_1 + \dots + k_p}{2}} \sum_{\pi \in \text{Part}(\mathcal{E})} A(n, \pi) \mathbb{E} \left[ \prod_{\ell=1}^p \left( x_{\pi(\overline{0}^\ell), \pi(\overline{1}^\ell)} \cdots x_{\pi(\overline{k_\ell-1}^\ell), \pi(\overline{k_\ell}^\ell)} - \mathbb{E} \left[ x_{\pi(\overline{0}^\ell), \pi(\overline{1}^\ell)} \cdots x_{\pi(\overline{k_\ell-1}^\ell), \pi(\overline{k_\ell}^\ell)} \right] \right) \right],$$

where for each  $\pi \in \text{Part}(\mathcal{E})$ ,  $A(n, \pi)$  is the number of families of indices of

$$(i_{\overline{0}^1}, \dots, i_{\overline{k_1}^1}, i_{\overline{0}^2}, \dots, i_{\overline{k_2}^2}, \dots, i_{\overline{0}^p}, \dots, i_{\overline{k_p}^p}) \in \{1, \dots, n\}^{\mathcal{E}}$$

whose level sets partition is  $\pi$  and who satisfy, for each  $\ell = 1, \dots, p$ ,

$$(12) \quad 1 \leq i_{\overline{0}^\ell} = i_{\overline{k_\ell}^\ell} \leq ns_\ell.$$

For any  $\pi \in \text{Part}(\mathcal{E})$ , let us define  $G_\pi$  to be the graph with vertex set the set  $\{\pi(x); x \in \mathcal{E}\}$  and edge set

$$E_\pi := \{ \{ \pi(\overline{m-1}^\ell), \pi(\overline{m}^\ell) \} ; 1 \leq \ell \leq p, m \in \{1, \dots, k_\ell\} \}.$$

For the term associated to  $\pi$  in (11) to be non zero, we need to have:

- (i)  $\pi(\overline{0}^\ell) = \pi(\overline{k_\ell}^\ell)$  for each  $\ell = 1, \dots, p$ ,
- (ii) each edge of  $G_\pi$  is visited at least twice by the union of the  $p$  paths  $(\pi(\overline{0}^\ell), \dots, \pi(\overline{k_\ell}^\ell))$  ( $\ell = 1, \dots, p$ ),
- (iii) for each  $\ell = 1, \dots, p$ , there exists  $\ell' \neq \ell$  such that at least one edge of  $G_\pi$  is visited by both paths  $(\pi(\overline{0}^\ell), \dots, \pi(\overline{k_\ell}^\ell))$  and  $(\pi(\overline{0}^{\ell'}), \dots, \pi(\overline{k_{\ell'}}^{\ell'}))$ .

Indeed, (i) is due to (12), (ii) is due to the fact that the  $x_{i,j}$ 's are centered and (iii) is due to the fact the the  $x_{i,j}$ 's are independent.

For each  $\ell = 1, \dots, p$  and each  $m = 0, \dots, k_\ell$ , set

$$s(\overline{m}^\ell) = \begin{cases} s_\ell & \text{if } m = 0 \text{ or } m = k_\ell, \\ 1 & \text{otherwise,} \end{cases}$$

and

$$(13) \quad s_\pi := \prod_{B \text{ bloc of } \pi} \min_{x \in B} s(x).$$

Then one can easily see that, as  $n \rightarrow \infty$ ,

$$(14) \quad A(n, \pi) \sim s_\pi n^{|\pi|}.$$

Thus for  $\pi$  to have a non zero asymptotic contribution to (11), we need to have moreover that

$$(iv) \quad \frac{k_1 + \dots + k_p}{2} \leq |\pi|.$$

The following lemma is a generalization of [1, Lem. 2.1.34]. Its proof goes along the same lines as the proof of the former.

**Lemma 4.1.** *Let  $\pi \in \text{Part}(\mathcal{E})$  satisfy (i), (ii), (iii). The the number  $c_\pi$  of connected components of  $G_\pi$  is such that  $c_\pi \leq p/2$  and*

$$|\pi| \leq c_\pi - \frac{p}{2} + \frac{k_1 + \dots + k_p}{2}.$$

As a consequence, if  $\pi$  also satisfies (iv), we have

- (a)  $c_\pi = p/2$ ,
- (b)  $p$  is even,
- (c)  $|\pi| = \frac{k_1 + \dots + k_p}{2}$  (so that  $k_1 + \dots + k_p$  is also even).

Note also that by (ii), we also have

$$(d) |E_\pi| \leq \frac{k_1 + \dots + k_p}{2}.$$

To sum up, by (11) and (14), we have

$$(15) \quad \lim_{n \rightarrow \infty} \mathbb{E} \left[ \prod_{\ell=1}^p \sum_{1 \leq i \leq ns_\ell} (e_i^* X_n^{k_\ell} e_i - \mathbb{E}[e_i^* X_n^{k_\ell} e_i]) \right] = \\ \sum_{\pi} s_\pi \lim_{n \rightarrow \infty} \mathbb{E} \left[ \prod_{\ell=1}^p \left( x_{\pi(\overline{0}^\ell), \pi(\overline{1}^\ell)} \cdots x_{\pi(\overline{k_\ell-1}^\ell), \pi(\overline{k_\ell}^\ell)} - \mathbb{E}[x_{\pi(\overline{0}^\ell), \pi(\overline{1}^\ell)} \cdots x_{\pi(\overline{k_\ell-1}^\ell), \pi(\overline{k_\ell}^\ell)}] \right) \right],$$

where the sum is taken over the partitions  $\pi$  of  $\mathcal{E}$  which satisfy (i), (ii), (iii) and (iv) above (such partitions also satisfy (a), (b), (c) and (d) above).

**Case where  $p$  is odd:** By (b), we know that when  $p$  is odd, there is no partition  $\pi$  satisfying the above conditions. So by (15),

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[ \prod_{\ell=1}^p \sum_{1 \leq i \leq ns_\ell} (e_i^* X_n^{k_\ell} e_i - \mathbb{E}[e_i^* X_n^{k_\ell} e_i]) \right] = 0.$$

**Case where  $p = 2$ :** In this case, by (a), for each partition  $\pi$  satisfying (i), (ii), (iii) and (iv),  $G_\pi$  is connected, so that  $|\pi| - 1 \leq |E_\pi|$ . Hence by (c) and (d),  $|E_\pi|$  is either equal to  $\frac{k_1+k_2}{2}$  or to  $\frac{k_1+k_2}{2} - 1$ .

In the case where  $|E_\pi| = \frac{k_1+k_2}{2} - 1$ , the graph  $G_\pi$  has exactly one more vertex than edges, hence is a tree. As a consequence, the paths  $(\pi(\overline{0}^1), \dots, \pi(\overline{k_1}^1))$  and  $(\pi(\overline{0}^2), \dots, \pi(\overline{k_2}^2))$ , which have the same beginning and ending vertices, have the property to visit an even number of times each edge they visit. By an obvious cardinality argument, only one edge is visited more than twice, and it is visited four times (twice in each sense). The other edges are visited once in each sense. It follows that the expectation associated to  $\pi$  in (15) is equal to  $\mathbb{E}[|x_{1,2}|^4] - 1$ .

In the case where  $|E_\pi| = \frac{k_1+k_2}{2}$ , by a cardinality argument again, we see that each edge of  $G_\pi$  is visited exactly twice (in fact, the configuration is the one described in [1, Sect. 2.1.7], where  $G_\pi$  is a *bracelet*). It follows that the expectation associated to  $\pi$  in (15) is equal to 1.

As a consequence, as  $n$  tends to infinity,

$$\mathbb{E} \left[ \sum_{1 \leq i \leq ns_1} (e_i^* X_n^{k_1} e_i - \mathbb{E}[e_i^* X_n^{k_1} e_i]) \times \sum_{1 \leq i \leq ns_2} (e_i^* X_n^{k_2} e_i - \mathbb{E}[e_i^* X_n^{k_2} e_i]) \right]$$

converges to a number that we shall denote by

$$(16) \quad \text{Cov}_{s_1, s_2}^{\text{centered}}(k_1, k_2)$$

and which depends on the distributions of the  $x_{i,j}$ 's only through  $\lim_{n \rightarrow \infty} \mathbb{E}[|x_{1,2}|^4]$ .

**Case where  $p$  is  $> 2$  and even:** By (a) above, for each partition  $\pi$  satisfying (i), (ii), (iii) and (iv),  $G_\pi$  has exactly  $p/2$  connected components. By (iii), each of them contains the support of exactly two of the  $p$  paths

$$(17) \quad (\pi(\overline{0}^\ell), \dots, \pi(\overline{k_\ell}^\ell)) \quad (\ell = 1, \dots, p).$$

Let us define  $\sigma_\pi$  to be the *matching* (i.e. a permutation all of whose cycles have length two) of  $\{1, \dots, p\}$  such that for all  $\ell = 1, \dots, p$ , the paths with indices  $\ell$  and  $\sigma_\pi(\ell)$  of the collection (17) above are supported by the same connected component of  $G_\pi$ .

We shall now partition the sum of (15) according to the value of the matching  $\sigma_\pi$  defined by  $\pi$ . We get

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[ \prod_{\ell=1}^p \sum_{1 \leq i \leq n_{s_\ell}} (e_i^* X_n^{k_\ell} e_i - \mathbb{E}[e_i^* X_n^{k_\ell} e_i]) \right] = \sum_{\sigma} \sum_{\pi} s_{\pi} \lim_{n \rightarrow \infty} \mathbb{E} \left[ \prod_{\ell=1}^p \left( x_{\pi(\bar{0}^\ell), \pi(\bar{1}^\ell)} \cdots x_{\pi(\bar{k}_\ell-1)^\ell, \pi(\bar{k}_\ell)^\ell} - \mathbb{E}[x_{\pi(\bar{0}^\ell), \pi(\bar{1}^\ell)} \cdots x_{\pi(\bar{k}_\ell-1)^\ell, \pi(\bar{k}_\ell)^\ell}] \right) \right],$$

where the first sum is over the matchings  $\sigma$  of  $\{1, \dots, p\}$  and the second sum is over the partitions  $\pi$  satisfying (i), (ii), (iii) and (iv) such that  $\sigma_\pi = \sigma$ .

Note that for each matching  $\sigma$  of  $\{1, \dots, p\}$ , the set of partitions  $\pi$  of  $\mathcal{E}$  such that  $\sigma_\pi = \sigma$  can be identified with the cartesian product, indexed by the set of cycles  $\{\ell, \ell'\}$  of  $\sigma$ , of the set of partitions  $\tau$  of the set

$$\mathcal{E}_{\ell, \ell'} := \{\bar{0}^\ell, \dots, \bar{k}_\ell^\ell\} \cup \{\bar{0}^{\ell'}, \dots, \bar{k}_{\ell'}^{\ell'}\} \quad (\text{subset of } \mathcal{E})$$

satisfying the following conditions

- (i.2)  $\tau(\bar{0}^\ell) = \tau(\bar{k}_\ell^\ell)$  and  $\tau(\bar{0}^{\ell'}) = \tau(\bar{k}_{\ell'}^{\ell'})$ ,
- (ii.2) each edge of the graph  $G_\tau$  is visited at least twice by the union of the 2 paths  $(\tau(\bar{0}^\ell), \dots, \tau(\bar{k}_\ell^\ell))$  and  $(\tau(\bar{0}^{\ell'}), \dots, \tau(\bar{k}_{\ell'}^{\ell'}))$ ,
- (iii.2) at least one edge of  $G_\tau$  is visited by both previous paths,
- (iv.2)  $\frac{k_\ell + k_{\ell'}}{2} \leq |\tau|$ .

Moreover, by independence of the random variables  $x_{i,j}$ 's, the expectation

$$\mathbb{E} \left[ \prod_{\ell=1}^p \left( x_{\pi(\bar{0}^\ell), \pi(\bar{1}^\ell)} \cdots x_{\pi(\bar{k}_\ell-1)^\ell, \pi(\bar{k}_\ell)^\ell} - \mathbb{E}[x_{\pi(\bar{0}^\ell), \pi(\bar{1}^\ell)} \cdots x_{\pi(\bar{k}_\ell-1)^\ell, \pi(\bar{k}_\ell)^\ell}] \right) \right]$$

factorizes along the connected components of  $G_\pi$ . The factor  $s_\pi$ , defined in (13), also factorizes along the connected components of  $G_\pi$ . It follows that we have

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[ \prod_{\ell=1}^p \sum_{1 \leq i \leq n_{s_\ell}} (e_i^* X_n^{k_\ell} e_i - \mathbb{E}[e_i^* X_n^{k_\ell} e_i]) \right] = \sum_{\sigma} \prod_{\{\ell, \ell'\}} \lim_{n \rightarrow \infty} \mathbb{E} \left[ \sum_{1 \leq i \leq n_{s_\ell}} (e_i^* X_n^{k_\ell} e_i - \mathbb{E}[e_i^* X_n^{k_\ell} e_i]) \times \sum_{1 \leq i \leq n_{s_{\ell'}}} (e_i^* X_n^{k_{\ell'}} e_i - \mathbb{E}[e_i^* X_n^{k_{\ell'}} e_i]) \right],$$

where the sum is over the matchings  $\sigma$  of  $\{1, \dots, p\}$  and for each such  $\sigma$ , the product is over the cycles  $\{\ell, \ell'\}$  of  $\sigma$ .

By the definition of  $\text{Cov}_{s_1, s_2}^{\text{centered}}(k_1, k_2)$  in (16), we get

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[ \prod_{\ell=1}^p \sum_{1 \leq i \leq n_{s_\ell}} (e_i^* X_n^{k_\ell} e_i - \mathbb{E}[e_i^* X_n^{k_\ell} e_i]) \right] = \sum_{\sigma \text{ matching}} \prod_{\{\ell, \ell'\} \text{ cycle of } \sigma} \text{Cov}_{s_\ell, s_{\ell'}}^{\text{centered}}(k_\ell, k_{\ell'}).$$

By Vick's formula and Equation (7), we have proved the first part of Proposition 3.4.

**Computation of  $\text{Cov}_{s_1, s_2}(2, 2)$ :** We have, by the paragraph devoted to the case  $p = 2$  above,

$$\text{Cov}_{s_1, s_2}^{\text{centered}}(2, 2) = \sum_{\pi, G_\pi \text{ is a tree}} s_\pi \times (\mathbb{E}[|x_{1,2}|^4] - 1) + \sum_{\pi, G_\pi \text{ is a bracelet}} s_\pi.$$

There are exactly one tree and zero bracelet with 2 vertices. We represent this tree in the following way:  $\bullet - \circ$ . There are two associated partitions  $\pi$ :

- the first one is defined by

$$(\pi(\bar{0}^1), \pi(\bar{1}^1), \pi(\bar{2}^1)) = (\pi(\bar{0}^2), \pi(\bar{1}^2), \pi(\bar{2}^2)) = (\bullet, \circ, \bullet),$$

and satisfies  $s_\pi = \min\{s_1, s_2\}$ ,

- the second one is defined by

$$(\pi(\bar{0}^1), \pi(\bar{1}^1), \pi(\bar{2}^1)) = (\bullet, \circ, \bullet), \quad (\pi(\bar{0}^2), \pi(\bar{1}^2), \pi(\bar{2}^2)) = (\circ, \bullet, \circ),$$

and satisfies  $s_\pi = s_1 s_2$ .

As a consequence,  $\text{Cov}_{s_1, s_2}^{\text{centered}}(2, 2) = (\mathbb{E}[|x_{1,2}|^4] - 1)(\min\{s_1, s_2\} + s_1 s_2)$ .

From this formula and Equation (7), we easily deduce the formula of  $\text{Cov}_{s_1, s_2}(2, 2)$ . It concludes the proof of Proposition 3.4.  $\square$

**4.3. A preliminary result for the proof of Lemma 3.1.** Proposition 4.3 below will allow to prove Lemma 3.1. Let us now consider two independent Wigner matrices  $X_n, X'_n$ . Let us introduce a modified version of Assumption 2.4, depending on an integer  $m \geq 2$ :

**Assumption 4.2.** For each  $k \geq 1$ ,  $\sup_n \mathbb{E}[|x_{1,1}|^k + |x_{1,2}|^k + |x'_{1,1}|^k + |x'_{1,2}|^k] < \infty$ . Moreover, there exists  $\varepsilon_0 > 0$  such that for each  $r, s \geq 0$ ,

$$(18) \quad r + s \leq m - 2 \implies \mathbb{E}[\Re(x_{1,1})^r \Im(x_{1,1})^s] - \mathbb{E}[\Re(x'_{1,1})^r \Im(x'_{1,1})^s] = O(n^{-\varepsilon_0 + 1 + \frac{r+s-m}{2}})$$

and

$$(19) \quad r + s \leq m \implies \mathbb{E}[\Re(x_{1,2})^r \Im(x_{1,2})^s] - \mathbb{E}[\Re(x'_{1,2})^r \Im(x'_{1,2})^s] = O(n^{-\varepsilon_0 + \frac{r+s-m}{2}}).$$

Let  $U'_n = [u'_{i,j}]_{i,j=1}^n$  be the associated eigenvectors matrix of  $X'_n$  (like  $U_n = [u_{i,j}]_{i,j=1}^n$  for  $X_n$ ).

**Proposition 4.3.** We suppose that  $X_n, X'_n$  both satisfy Assumptions 2.1, 2.5 and 4.2. Let us fix a positive integer  $k$  and a polynomial function  $G$  on  $\mathbb{C}^k$ . For each  $n$ , let us consider a collection  $(i_1, p_1, q_1), \dots, (i_k, p_k, q_k) \in \{1, \dots, n\}^3$  (this collection might depend on  $n$ ). Then for a certain constant  $C$  independent of  $n$  and of the choices of  $(i_1, p_1, q_1), \dots, (i_k, p_k, q_k)$  (but depending on  $k$  and on  $G$ )

$$(20) \quad |\mathbb{E}[G(n u_{p_1, i_1} \bar{u}_{q_1, i_1}, \dots, n u_{p_k, i_k} \bar{u}_{q_k, i_k})] - \mathbb{E}[G(n u'_{p_1, i_1} \bar{u}'_{q_1, i_1}, \dots, n u'_{p_k, i_k} \bar{u}'_{q_k, i_k})]| \leq C n^{2 - \frac{m}{2}}.$$

**Proof.** The proof follows the ideas developed by Terence Tao and Van Vu in recent papers, such as [18, 19]. More precisely, it makes an intensive use of the strategy and of some estimations given in [19].

An event  $E = E^{(n)}$  depending on the parameter  $n$  will be said to be true with *overwhelming probability* if for all  $c > 0$ ,  $1 - \mathbb{P}(E) = O(n^{-c})$ . One can neglect any such event in the proof of (20) because  $G$  has polynomial growth and the entries of unitary matrices are bounded by 1.

By Assumption 2.4 and Chebyshev's inequality, that for each  $k$ , as  $t \rightarrow \infty$ ,

$$\mathbb{P}(|x_{i,j}| \geq t) = O(t^{-k}) \quad (\text{uniformly in } i, j \text{ and in } n),$$

so that by the union bound,  $\mathbb{P}(\max_{1 \leq i, j \leq n} |x_{i,j}| \geq n^\varepsilon) = O(n^{2-k\varepsilon})$  for any  $\varepsilon, k$ . Hence by what precedes, for a fixed  $\varepsilon_1$  (a parameter fixed later), one can suppose that for each  $n$ , the  $x_{i,j}$ 's satisfy  $|x_{i,j}| \leq n^{\varepsilon_1}$  for all  $i, j$ . Of course, one can also suppose that  $|x'_{i,j}| \leq n^{\varepsilon_1}$  for all  $i, j$ .

To prove (20), it suffices to prove that we can replace the  $x_{i,j}$ 's by the  $x'_{i,j}$ 's one by one up to an error in the considered expectation which is  $O(n^{1 - \frac{m}{2}})$  for each diagonal replacement and

$O(n^{-\frac{m}{2}})$  for each off-diagonal replacement (and, of course, that these bounds on the errors are uniform in the  $\frac{n(n+1)}{2}$  replacements).

So let us fix  $1 \leq p \leq q \leq n$  and define, for each  $z \in \mathbb{C}$  (or  $z \in \mathbb{R}$  if  $p = q$ ),  $A_{p,q}(z)$  be the matrix obtained by "mixing" the (rescaled) matrices  $X_n$  and  $X'_n$  in the following way:

- for each  $1 \leq i \leq j \leq n$  such that  $(i, j)$  is below  $(p, q)$  for the lexicographic order, the  $i, j$  entry of  $A_{p,q}(z)$  and its symmetric one are the corresponding ones in the ones of  $nX'_n$ ,
- the  $p, q$  entry of  $A_{p,q}(z)$  and its symmetric one are  $z$  and  $\bar{z}$ ,
- for each  $1 \leq i \leq j \leq n$  such that  $(i, j)$  is above  $(p, q)$  for the lexicographic order, the  $i, j$  entry of  $A_{p,q}(z)$  and its symmetric one are the corresponding ones in the ones of  $nX_n$ .

We set  $U(z) = [u_{i,j}(z)]_{i,j=1}^n$  to be the eigenvectors matrix of  $A_{p,q}(z)$  and

$$F(z) := G(n u_{p_1, i_1}(z) \bar{u}_{q_1, i_1}(z), \dots, n u_{p_k, i_k}(z) \bar{u}_{q_k, i_k}(z)).$$

Let  $\mathbb{E}[\cdot | A_{p,q}(0)]$  denote the conditional expectation with respect to the  $\sigma$ -algebra generated by the entries of  $A_{p,q}(0)$ . We have to prove that with overwhelming probability, we have, uniformly on the probability space and on  $p, q$ ,

$$(21) \quad \mathbb{E}[F(\sqrt{n}x_{p,q}) | A_{p,q}(0)] = \mathbb{E}[F(\sqrt{n}x'_{p,q}) | A_{p,q}(0)] + O(n^{\mathbb{1}_{p=q} - \frac{m}{2}}).$$

To prove it, we shall use the Taylor expansion of  $F$  around zero. Let us for example treat the case  $p \neq q$ . For  $|x| \leq n^{\varepsilon_1}$ , we have

$$F(\sqrt{n}x) = \sum_{r+s \leq m} \frac{n^{\frac{r+s}{2}}}{r!s!} \partial_{\Re(z)}^r \partial_{\Im(z)}^s F(0) \Re(x)^s \Im(x)^s + O \left( n^{(m+1)(\varepsilon_1+1/2)} \sup_{\substack{r+s=m+1 \\ |z| \leq n^{\varepsilon_1+1/2}}} |\partial_{\Re(z)}^r \partial_{\Im(z)}^s F(z)| \right).$$

Moreover, by [19, Propo. 20], we know that with overwhelming probability,  $A_{p,q}(0)$  is a *good configuration* (in the sense of [19, Def. 18]). By Equation (27) of [19, Lem. 24], it implies that with overwhelming probability, for all  $l = 1, \dots, k$ , for all  $r, s$  such that  $r + s \leq 10$ ,

$$\sup_{|z| \leq n^{1/2+\varepsilon_1}} |\partial_{\Re(z)}^r \partial_{\Im(z)}^s (n u_{p_l, i_l}(z) \bar{u}_{q_l, i_l}(z))| = O(n^{\varepsilon_1 - (r+s)})$$

(in the statement of [19, Lem. 24],  $m$  must be bounded above by 10, but the bound 10 can be replaced by any finite one). The function  $G$  is a polynomial (let  $d$  be its degree). It follows that with overwhelming probability, for all  $r, s$ , we have

$$\sup_{|z| \leq n^{1/2+\varepsilon_1}} |\partial_{\Re(z)}^r \partial_{\Im(z)}^s F(z)| = O(n^{d\varepsilon_1 - (r+s)}).$$

As a consequence, by the bounds of (19),

$$\mathbb{E}[F(\sqrt{n}x_{p,q}) | A_{p,q}(0)] - \mathbb{E}[F(\sqrt{n}x'_{p,q}) | A_{p,q}(0)] = \sum_{r+s \leq m} O(n^{d\varepsilon_1 - \frac{m}{2} - \varepsilon_0}) + O(n^{(m+1+d)\varepsilon_1 - \frac{m+1}{2}})$$

Now, choosing  $\varepsilon_1$  small enough so that  $d\varepsilon_1 < \varepsilon_0$  and  $(m+1+d)\varepsilon_1 < 1/2$ , we get (21).  $\square$

**4.4. Proof of Lemma 3.1.** To prove Lemma 3.1, we shall use the following proposition, which is the obvious multidimensional generalization of Proposition 3.26 of Chapter VI of [13] (note also that the result of [13] is devoted to processes on  $[0, \infty)$  and not on  $[0, 1]$ , so that the  $N$  of [13] means nothing to us).

For  $f \in D([0, 1]^2)$  and  $(s_0, t_0) \in [0, 1]^2$ , we define  $\Delta_{s_0, t_0} f$  to be the *maximal discontinuity* of  $f$  at  $(s_0, t_0)$ , i.e.

$$\Delta_{s_0, t_0} f := \max_{\diamond, \diamond' \in \{<, \geq\}} \left| f(s_0, t_0) - \lim_{\substack{s \diamond s_0 \\ t \diamond' t_0}} f(s, t) \right|.$$

**Proposition 4.4.** *If the sequence (distribution( $B^n$ )) is tight and satisfies*

$$(22) \quad \forall \varepsilon > 0, \quad \mathbb{P}\left(\sup_{(s, t) \in [0, 1]^2} \Delta_{s, t} B^n > \varepsilon\right) \xrightarrow{n \rightarrow \infty} 0,$$

*then the sequence (distribution( $B^n$ )) is  $C$ -tight, i.e. is tight and can only have  $C([0, 1]^2)$ -supported accumulation points.*

So to prove Lemma 3.1, let us first prove that the sequence (distribution( $B^n$ )) is tight. Note that the process  $B^n$  vanishes at the border of  $[0, 1]^2$ . So according to [4, Th. 3] and to Cauchy-Schwartz inequality, it suffices to prove that there is  $C < \infty$  such that for  $n$  large enough, for all  $s < s', t < t' \in [0, 1]$ ,

$$(23) \quad \mathbb{E} \left[ \left( \sum_{ns < i \leq ns'} \sum_{nt < j \leq nt'} (|u_{i, j}|^2 - 1/n) \right)^4 \right] \leq C(s' - s)^2(t' - t)^2.$$

Set  $I = \{i = 1, \dots, n; ns < i \leq ns'\}$  and  $J = \{j = 1, \dots, n; nt < j \leq nt'\}$ . We have

$$\mathbb{E} \left[ \left( \sum_{i \in I, j \in J} (|u_{i, j}|^2 - 1/n) \right)^4 \right] = n^{-4} \sum_{(i_\ell, j_\ell)_{\ell=1}^4 \in (I \times J)^4} \mathbb{E} \left[ \prod_{\ell=1}^4 (n|u_{i_\ell, j_\ell}|^2 - 1) \right].$$

Let us now apply Proposition 4.3 for  $X'_n$  a GOE or GUE matrix and  $m = 12$  (indeed, Assumption 2.4 implies Assumption 4.2 for  $X'_n$  a GOE or GUE matrix and  $m = 12$ ). Let  $U'_n = [u'_{i, j}]_{i, j=1}^n$  be a Haar-distributed orthogonal or unitary matrix. By Proposition 4.3, we have

$$\mathbb{E} \left[ \left( \sum_{i \in I, j \in J} (|u_{i, j}|^2 - 1/n) \right)^4 \right] = O(n^{-8} (\#I \times J)^4) + n^{-4} \sum_{(i_\ell, j_\ell)_{\ell=1}^4 \in (I \times J)^4} \mathbb{E} \left[ \prod_{\ell=1}^4 (n|u'_{i_\ell, j_\ell}|^2 - 1) \right],$$

where the  $O(n^{-8} (\#I \times J)^4)$  is uniform in  $s, s', t, t'$ , hence is bounded (for some constants  $C, C'$ ) by

$$C \frac{(\#I \times J)^4}{n^8} \leq 2C'(s' - s)^4(t' - t)^4 \leq 2C'(s' - s)^2(t' - t)^2.$$

As a consequence, it suffices to prove the result for  $U'_n$  instead of  $U_n$ . But this job is already done in [9]: the unitary case is treated in Section 3.4.1 (see specifically Equation (3.25)) and the orthogonal case is treated, more elliptically, in Section 4.5 (to recover the details of the proof, join Equations (3.26), (4.5) and (4.17)).

So the proof of (23) is complete.

Let us now prove (22). Note that  $\sup_{(s, t) \in [0, 1]^2} \Delta_{s, t} B^n = \max_{1 \leq i, j \leq n} ||u_{i, j}|^2 - 1/n|$ . As a consequence, by the union bound, it suffices to prove that for each  $\varepsilon > 0$ , there is  $C < \infty$  independent of  $i, j$  and  $n$  such that for all  $i, j$ ,

$$\mathbb{P}(|u_{i, j}|^2 - 1/n| > \varepsilon) \leq Cn^{-4}.$$

By Chebyshev's inequality,

$$\mathbb{P}(|u_{i, j}|^2 - 1/n| > \varepsilon) = \mathbb{P}(|n|u_{i, j}|^2 - 1| > n\varepsilon) \leq \frac{1}{(n\varepsilon)^4} \mathbb{E}[(n|u_{i, j}|^2 - 1)^4].$$



So to prove (22), one only needs to prove that  $\mathbb{E}[(n|u_{i,j}|^2 - 1)^4]$  is bounded independently of  $i, j$  and  $n$ . To do so, let us again apply Proposition 4.3 for  $X'_n$  a GOE or GUE matrix and  $m = 4$ : it allows to reduce the problem to the case where the  $u_{i,j}$ 's are the entries of a Haar-distributed matrix. In this case, the distribution of  $u_{i,j}$  does not depend on  $i, j$  and it is well known that the moments of  $\sqrt{n}u_{1,1}$  converge to the ones of a standard real or complex Gaussian variable. We did not find any concise enough reference for this moments convergence, but in the unitary case, Proposition 3.4 of [3] allows to compute  $\mathbb{E}[(n|u_{1,1}|^2 - 1)^4]$  and to verify its boundedness. To treat the orthogonal case, the most direct way to compute  $\mathbb{E}[(n|u_{1,1}|^2 - 1)^4]$  is to use the fact that  $u_{1,1}^2$  has the same distribution as  $Z_1^2/(Z_1^2 + \dots + Z_n^2)$  for  $Z_i$  independent standard Gaussian variables (see e.g. [15, Lem. 2.1]) and then to use [15, Lem. 2.4].  $\square$

## REFERENCES

- [1] G. Anderson, A. Guionnet, O. Zeitouni *An Introduction to Random Matrices*. Cambridge studies in advanced mathematics, 118 (2009).
- [2] Z. D. Bai, J. W. Silverstein *Spectral analysis of large dimensional random matrices*, Second Edition, Springer, New York, 2009.
- [3] F. Benaych-Georges *Classical and free infinitely divisible distributions and random matrices*. Annals of Probability. Vol. 33, no. 3 (2005) 1134–1170.
- [4] P.J. Bickel, M.J. Wichura *Convergence criteria for multiparameter stochastic processes and some applications*, Ann. Math. Statist., 42(5):1656–1670, 1971.
- [5] P. Billingsley *Convergence of Probability Measures*. New York, John Wiley and Sons (1968).
- [6] S. Chatterjee, E. Meckes *Multivariate normal approximation using exchangeable pairs*, ALEA 4, 257–283 (2008).
- [7] B. Collins *Moments and Cumulants of Polynomial Random Variables on Unitary Groups, the Itzykson-Zuber Integral, and Free Probability*. Int. Math. Res. Not., No. 17, 2003.
- [8] B. Collins, P. Sniady *Integration with respect to the Haar measure on unitary, orthogonal and symplectic group*. Comm. Math. Phys., 264, 773–795, 2006.
- [9] C. Donati-Martin, A. Rouault *Truncations of Haar unitary matrices, traces and bivariate Brownian bridge*, arXiv.
- [10] L. Erdős, B. Schlein, H.-T. Yau *Semicircle law on short scales and delocalization of eigenvectors for Wigner random matrices*. Ann. Probab. 37 (2009), no. 3, 815–852.
- [11] L. Erdős, B. Schlein, H.-T. Yau *Local semicircle law and complete delocalization for Wigner random matrices*. Comm. Math. Phys. 287 (2009), no. 2, 641–655.
- [12] L. Erdős, B. Schlein, H.-T. Yau *Wegner estimate and level repulsion for Wigner random matrices*. To appear in Int. Math. Res. Not.
- [13] J. Jacod, A.N. Shiryaev *Limit theorems for stochastic processes*, Berlin, Springer-Verlag (1987).
- [14] T. Jiang *How many entries of a typical orthogonal matrix can be approximated by independent normals?* Ann. Probab. 34(4), 1497–1529. 2006.
- [15] T. Jiang *A variance formula related to a quantum conductance problem*. Physics Letters A, vol. 373 (25), 2117–2121, 2009.
- [16] A. Knowles, J. Yin *Eigenvector Distribution of Wigner Matrices*, arXiv:1102.0057.
- [17] J. W. Silverstein *Weak convergence of random functions defined by the eigenvectors of sample covariance matrices*. Ann. Probab. 18 (1990), no. 3, 1174–1194.
- [18] T. Tao, V. Vu *Random matrices: universality of local eigenvalue statistics*, arXiv 0906.0510.
- [19] T. Tao, V. Vu *Random matrices: universality of eigenvectors*, arXiv 1103.2801.

FLORENT BENAYCH-GEORGES, LPMA, UPMC UNIV PARIS 6, CASE COURIER 188, 4, PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE AND CMAP, ÉCOLE POLYTECHNIQUE, ROUTE DE SACLAY, 91128 PALAISEAU CEDEX, FRANCE. URL: <http://www.cmapx.polytechnique.fr/~benaych/>

*E-mail address:* florent.benaych@gmail.com