



**HAL**  
open science

# Convex inner approximations of nonconvex semialgebraic sets applied to fixed-order controller design

Didier Henrion, Christophe Louembet

► **To cite this version:**

Didier Henrion, Christophe Louembet. Convex inner approximations of nonconvex semialgebraic sets applied to fixed-order controller design. 2011. hal-00585633v1

**HAL Id: hal-00585633**

**<https://hal.science/hal-00585633v1>**

Submitted on 13 Apr 2011 (v1), last revised 10 Jan 2012 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Convex inner approximations of nonconvex semialgebraic sets applied to fixed-order controller design

Didier Henrion<sup>1,2,3</sup>, Christophe Louembet<sup>1,2</sup>

April 13, 2011

## Abstract

We describe an elementary algorithm to build convex inner approximations of nonconvex sets. Both input and output sets are basic semialgebraic sets given as lists of defining multivariate polynomials. Even though no optimality guarantees can be given (e.g. in terms of volume maximization for bounded sets), the algorithm is designed to preserve convex boundaries as much as possible, while removing regions with concave boundaries. In particular, the algorithm leaves invariant a given convex set. The algorithm is based on Gloptipoly 3, a public-domain Matlab package solving nonconvex polynomial optimization problems with the help of convex semidefinite programming (optimization over linear matrix inequalities, or LMIs). We illustrate how the algorithm can be used to design fixed-order controllers for linear systems, following a polynomial approach.

**Keywords:** polynomials; nonconvex optimization; LMI; fixed-order controller design

## 1 Introduction

The set of controllers stabilizing a linear system is generally nonconvex in the parameter space, and this is an essential difficulty faced by numerical algorithms of computer-aided control system design, see e.g. [4] and references therein. It follows from the derivation of the Routh-Hurwitz stability criterion (or its discrete-time counterpart) that the set of stabilizing controllers is real basic semialgebraic, i.e. it is the intersection of sublevel sets of given multivariate polynomials. A convex inner approximation of this nonconvex semialgebraic stability region was obtained in [4] in the form of linear matrix inequalities (LMI) obtained from univariate polynomial positivity conditions, see also [9]. Convex polytopic inner approximations were also obtained in [13], for discrete-time stability, using

---

<sup>1</sup>CNRS; LAAS; 7 avenue du colonel Roche, F-31077 Toulouse; France.

<sup>2</sup>Université de Toulouse; UPS, INSA, INP, ISAE; UT1, UTM, LAAS; F-31077 Toulouse; France

<sup>3</sup>Faculty of Electrical Engineering, Czech Technical University in Prague, Technická 2, CZ-16626 Prague, Czech Republic

reflection coefficients. Convex inner approximations make it possible to design stabilizing controllers with the help of convex optimization techniques, at the price of loosing optimality w.r.t. closed-loop performance criteria ( $H_2$  norm,  $H_\infty$  norm or alike).

Generally speaking, the technical literature abounds of convex *outer* approximations of nonconvex semialgebraic sets. In particular, such approximations form the basis of many branch-and-bound global optimization algorithms [12]. By construction, Lasserre's hierarchy of LMI relaxations for polynomial programming is a sequence of embedded convex outer approximations which are semidefinite representable, i.e. which are obtained by projecting affine sections of the convex cone of positive semidefinite matrices, at the price of introducing lifting variables [6].

After some literature search, we could not locate any systematic constructive procedure to generate convex *inner* approximations of nonconvex semialgebraic sets, contrasting sharply with the many convex outer approximations mentioned above. In the context of fixed-order controller design, inner approximations correspond to a guarantee of stability, at the price of loosing optimality. No such stability guarantee can be ensured with outer approximations.

The main contribution of this paper is therefore an elementary algorithm, readily implementable in Matlab, that generates convex inner approximations of nonconvex sets. Both input and output sets are basic semialgebraic sets given as lists of defining multivariate polynomials. Even though no optimality guarantees can be given in terms of volume maximization for bounded sets, the algorithm is designed to preserve convex boundaries as much as possible, while removing regions with concave boundaries. In particular, the algorithm leaves invariant a given convex set. The algorithm is based on Gloptipoly 3, a public-domain Matlab package solving nonconvex polynomial optimization problems with the help of convex LMIs [7]. We illustrate how the algorithm can be used to design fixed-order controllers for linear systems, following a polynomial approach.

## 2 Convex inner approximation

Given a basic closed semialgebraic set

$$S = \{x \in \mathbb{R}^n : p_1(x) \leq 0 \dots p_m(x) \leq 0\} \quad (1)$$

where  $p_i$  are multivariate polynomials, we are interested in computing another basic closed semialgebraic set

$$\bar{S} = \{x \in \mathbb{R}^n : \bar{p}_1(x) \leq 0 \dots \bar{p}_m(x) \leq 0\} \quad (2)$$

which is a valid inner approximation of  $S$ , in the sense that

$$\bar{S} \subset S.$$

Ideally, we would like to find the tightest possible approximation, in the sense that the complement set  $S \setminus \bar{S} = \{x \in S : x \notin \bar{S}\}$  is as small as possible. Mathematically we may formulate the problem as the volume minimization problem

$$\inf_{\bar{S}} \int_{S \setminus \bar{S}} dx$$

but since set  $S$  is not necessarily bounded we should make sure that this integral makes sense. Moreover, computing the volume of a given semialgebraic set is a difficult task in general [8], so we expect that optimizing such a quantity is as much as difficult. For these reasons, we have not been able to define a mathematically sound while tractable measure of tightness of the inner approximation. In practice we will content ourselves of an inner approximation that removes the nonconvex parts of the boundary and keeps the convex parts as much as possible.

### 3 Detecting nonconvexity

Before describing the method, let us recall some basics definitions on polynomials and differential geometry. Let  $x \in \mathbb{R}^n \mapsto p_i(x) \in \mathbb{R}[x]$  be a multivariate polynomial of total degree  $d$ . Let

$$g_i(x) = \left[ \frac{\partial p_i(x)}{\partial x_j} \right]_{j=1 \dots n} \in \mathbb{R}^n[x]$$

be its gradient vector and

$$H_i(x) = \left[ \frac{\partial^2 p_i(x)}{\partial x_j \partial x_k} \right]_{j,k=1 \dots n} \in \mathbb{R}^{n \times n}[x]$$

its (symmetric) Hessian polynomial matrix. Define the optimization problem

$$\begin{aligned} q_i &= \min_{x,y} y^T H_i(x, y) y \\ \text{s.t.} & \quad p_i(x) = 0 \\ & \quad p_j(x) \leq 0, j = 1 \dots m, j \neq i \\ & \quad y^T g_i(x) = 0 \\ & \quad y^T y = 1 \end{aligned} \tag{3}$$

with global minimizers  $\{x^1 \dots x^{k_i}\}$  and  $\{y^1 \dots y^{k_i}\}$ .

Let us make the following nondegeneracy assumption on defining polynomials  $p_i(x)$ :

**Assumption 1** *There is no point  $x$  such that  $p_i(x)$  and  $g_i(x)$  vanish simultaneously while satisfying  $p_j(x) \leq 0$  for  $j = 1, \dots, m, j \neq i$ .*

Since the polynomial system  $p_i(x) = 0, g_i(x) = 0$ , involves  $n + 1$  equations for  $n$  unknown, Assumption 1 is satisfied generically. In other words, in the Euclidean space of coefficients of polynomials  $p_i(x)$ , instances violating Assumption 1 belong to a variety of measure zero, and an arbitrarily small perturbation on the coefficients generates a perturbed set  $S_\epsilon$  satisfying Assumption 1.

**Theorem 1** *Under Assumption 1, polynomial level set (1) is convex if and only if  $q_i \geq 0$  for all  $i = 1, \dots, m$ .*

**Proof:** The boundary of set  $S$  consists of points  $x$  such that  $p_i(x) = 0$  for some  $i$ , and  $p_j(x) \leq 0$  for  $j \neq i$ . In the neighborhood of such a point, consider the Taylor series

$$p_i(x + y) = p_i(x) + y^T g_i(x) + y^T H_i(x) y + O(y^3) \quad (4)$$

where  $O(y^3)$  denotes terms of degree 3 or higher in entries of vector  $y$ , the local coordinates. By Assumption 1, the gradient  $g_i(x)$  does not vanish along the boundary, and hence convexity of the boundary is inferred from the quadratic term in expression (4). More specifically, when  $y^T g_i(x) = 0$ , vector  $y$  belongs to the hyperplane tangent to  $S$  at point  $x$ . Let  $V$  be a matrix spanning this linear subspace of dimension  $n - 1$  so that  $y = V\hat{y}$  for some  $\hat{y}$ . The quadratic form  $y^T H_i(x) y = \hat{y}^T V^T H_i V \hat{y}$  can be diagonalised with the congruence transformation  $\hat{y} = U\bar{y}$  (Schur decomposition), and hence  $y^T H_i(x) y = \bar{y}^T U^T V^T H_i V U \bar{y} = \sum_{i=1}^{n-1} h_i(x) \bar{y}_i^2$ . The eigenvalues  $h_i(x)$ ,  $i = 1, \dots, n - 1$  are reciprocals of the principal curvatures of the surface. Problem (3) then amounts to finding the minimum curvature, which is non-negative when the surface is locally convex around  $x$ .  $\square$

In the case of three-dimensional surfaces ( $n = 3$ ), the ideas of tangent plane, local coordinates and principal curvatures used in the proof of Theorem 1 are standard notions of differential geometry, see e.g. Section 3.3. in [2] and in particular Example 5 for connections between principal curvatures and eigenvalues of the local Hessian form (called the second fundamental form, once suitably normalized).




Figure 1: Hyperboloid of one sheet (white), with tangent plane (gray) at the origin, a saddle point with a tangent convex parabola (thick black) and a tangent concave hyperbola (thick black).

As an example illustrating the proof of Theorem 1, consider the hyperboloid of one sheet  $S = \{x \in \mathbb{R}^3 : p_1(x) = x_1^2 - x_2^2 - x_3 \leq 0\}$  with gradient and Hessian

$$g_1(x) = \begin{bmatrix} 2x_1 \\ -2x_2 \\ -1 \end{bmatrix}, \quad H_1(x) = \begin{bmatrix} 2 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

At the origin  $x = 0$ , the tangent plane is  $T = \{y \in \mathbb{R}^3 : y_3 = 0\}$  and  $p_1(y) = 2y_1^2 - 2y_2^2$  is a bivariate quadratic form with eigenvalues 2 and  $-2$ , corresponding respectively to the convex parabola  $\{x : x_2^2 + x_3 = 0\}$  (positive curvature) and concave hyperbola  $\{x : x_1^2 - x_3 = 0\}$  (negative curvature), see Figure 1.

Theorem 1 can be exploited in an algorithmic way to generate a convex inner approximation of a semialgebraic set.

### Algorithm 1 (Convex inner approximation)

*Input:* Polynomials  $p_i$ ,  $i = 1 \dots m$  defining set  $S$  as in (1). Small nonnegative scalar  $\epsilon$ .

*Output:* Polynomials  $\bar{p}_i$ ,  $i = 1 \dots \bar{m}$  defining set  $\bar{S}$  as in (2).

*Step 1:* Let  $i = 1$ .

*Step 2:* If  $\deg p_i \leq 1$  then go to Step 5.

*Step 3:* If  $p_i(x) \in S$ , solve optimization problem (3) for optimum  $q_i$  and minimizers  $\{x^1 \dots x^k\}$ . If  $p_i(x) \notin S$ , go to Step 5.

*Step 4:* If  $q_i < 0$ , then select one of the minimizers  $x^j$ ,  $j = 1 \dots k_i$ , let  $p_{m+1} = g_i(x^j)(x - x^j) + \epsilon$ . Then let  $m = m + 1$ , and go to step 3.

*Step 5:* Let  $i = i + 1$ . If  $i \leq m$  then go to Step 2.

*Step 6:* Return  $\bar{p}_i = p_i$ ,  $i = 1, \dots, m$ .

The idea behind the algorithm is as follows. At Step 3, by solving the polynomial optimization problem of Theorem 1 we identify a point of minimal curvature along algebraic varieties defining the boundary of  $S$ . If the minimal curvature is negative, then we separate the point from the set with a gradient hyperplane, and we iterate on the resulting semialgebraic set. At the end, we obtain a valid inner approximation.

Note that Step 2 checks if the boundary is affine, in which case the minimum curvature is zero and there is no optimization problem to be solved.

The key parameter of the algorithm is the small positive scalar  $\epsilon$  used at Step 4 for separating strictly a point of minimal curvature, so that the algorithm does not identify it again at the next iteration. Moreover, in Step 4, one must elect arbitrarily a minimizer. We will discuss this issue later in this paper.

## 4 Matlab code and examples

At each step of Algorithm 1 we have to solve a potentially nonconvex polynomial optimization problem. For that purpose, we use Gloptipoly 3, a public-domain Matlab package [7]. The methodology consists in building and solving a hierarchy of embedded linear matrix inequality (LMI) relaxations of the polynomial optimization problem, see the survey [11].

The LMI problems are solved numerically with the help of any semidefinite programming solver (by default Gloptipoly 3 uses SeDuMi). Under the assumption that our original semi-algebraic set is compact, the sequence of minimizers obtained by solving the LMI relaxations is ensured to converge monotonically to the global minimum. Under the additional assumption that the global optima live on a zero-dimensional variety (i.e. there is a finite number of them), Gloptipoly 3 eventually extracts some of them (not necessarily all, but at least one) using numerical linear algebra. The LMI problems in the hierarchy have a growing number of variables and constraints, and the main issue is that we cannot predict in advance how large has to be the LMI problem to guarantee global optimality. In practice however we observe that it is not necessary to go very deep in the hierarchy to have a numerical certificate of global optimality.

## 4.1 Hyperbola

Let us first with the elementary example of an unbounded nonconvex hyperbolic region  $S = \{x \in \mathbb{R}^2 : p_1(x) \leq 0\}$  with  $p_1(x) = -1 + x_1x_2$ , for which optimization problem (3) reads

$$\begin{aligned} \min \quad & 2y_1y_2 \\ \text{s.t.} \quad & x_2y_1 + x_1y_2 = 0 \\ & -1 + x_1x_2 = 0 \\ & y_1^2 + y_2^2 = 1. \end{aligned}$$

Necessary optimality conditions yield immediately  $k_1 = 2$  global minimizers  $x^1 = \frac{\sqrt{2}}{2}(1, 1)$ ,  $y^1 = \frac{\sqrt{2}}{2}(1, -1)$  and  $x^2 = \frac{\sqrt{2}}{2}(-1, -1)$ ,  $y^2 = \frac{\sqrt{2}}{2}(-1, 1)$ , and hence two additional (normalized) affine constraints  $p_2(x) = -2 + x_1 + x_2$  and  $p_3(x) = -2 - x_1 - x_2$  defining the slab  $\bar{S} = \{x : p_i(x) \leq 0, i = 1, 2, 3\} = \{x : -2 \leq x_1 + x_2 \leq 2\}$  which is indeed a valid inner approximation of  $S$ .

## 4.2 Egg quartic

Now we show that Algorithm 1 can be used to detect convexity of a semialgebraic set. Consider the smooth quartic sublevel set  $S = \{x \in \mathbb{R}^2 : p_1(x) = x_1^4 + x_2^4 + x_1^2 + x_2 \leq 0\}$  represented on Figure 2. Assumption 1 is ensured since the gradient  $g_1(x) = [2x_1(x_1^2 + 2) \ 4x_2^3 + 1]$  cannot vanish for real  $x$ .

A Matlab implementation of the first steps of the algorithm can be easily written using Gloptipoly 3:

```
% problem data
mpol x y 2
p1 = x(1)^4+x(2)^4+x(1)^2+x(2);
g1 = diff(p1,x); % gradient
H1 = diff(g1,x); % Hessian
% LMI relaxation order
order = 3;
% build LMI relaxation
```




Figure 2: Convex smooth quartic.

```
P = msdp(min(y'*H1*y), p1==0, ...
    g1*y==0, y'*y==1, order);
% solve LMI relaxation
[status,obj] = msol(P)
```

Notice that we solve the LMI relaxation of order 3 (e.g. moments of degree 6) of problem (3). Running the above script, Gloptipoly returns `obj = 2.0000` and `status = 1`, certifying that the minimal curvature is strictly positive, and hence that the polynomial sublevel set is convex.

Note that in this simple case, convexity of set  $S$  follows directly from positive semidefiniteness of the Hessian  $H_1(x) = \text{diag}(12x_1^2 + 2, 12x_2^2)$ , yet Algorithm 1 can systematically detect convexity in more complicated cases.

### 4.3 Waterdrop quartic

Consider the quartic  $S = \{x \in \mathbb{R}^2 : p_1(x) = x_1^4 + x_2^4 + x_1^2 + x_2^3 \leq 0\}$  which has a singular point at the origin, hence violating Assumption 1.

Applying Algorithm 1, the LMI relaxation of order 4 (moments of degree 8) yields a globally minimal curvature of  $-0.094159$  achieved at the 2 points  $x^1 = (-0.048892, -0.14076)$  and  $x^2 = (0.048896, -0.14076)$ . With the two additional affine constraints  $p_k(x) = g_1(x^k)(x - x^k) \leq 0$ ,  $k = 2, 3$ , the resulting set  $\bar{S}$  has a globally minimal curvature of 1 certified at the LMI relaxation of order 4, and therefore it is a valid convex inner approximation of  $S$ , see Figure 3.






Figure 3: Nonconvex waterdrop quartic (light gray) and its convex inner approximation (dark gray) obtained by adding affine constraints at two points  $x^1$  and  $x^2$  of minimal curvature.

This example illustrates that Algorithm 1 can work even when Assumption 1 is violated. Here the singularity is removed by the additional affine constraints. This example also shows that symmetry of the problem can be exploited, since two global minimizers are found (distinct points with the same minimal curvature) to remove two nonconvex parts of the boundary simultaneously.

#### 4.4 Singular quartic

Consider the quartic  $S = \{x \in \mathbb{R}^2 : p_1(x) = x_1^4 + x_2^4 + x_2^3 \leq 0\}$  which has a singular point at the origin, hence violating Assumption 1.

Running Algorithm 1, we obtain the following sequence of bounds on the minimum curvature, for increasing LMI relaxation orders:

order	2	3	4	5
obj	$-7.5000 \cdot 10^{-1}$	$-7.7502 \cdot 10^{-2}$	$-8.5855 \cdot 10^{-3}$	$-4.9525 \cdot 10^{-3}$

GloptiPoly is not able to certify global optimality, so we can only speculate that the global minimum is zero and hence that set  $S$  is convex, see Figure 4. We may say that set  $S$  is numerically convex.

Indeed if we strengthen the constraint  $p_1(x) \leq 0$  into  $p_1(x) + \epsilon \leq 0$  for a small positive  $\epsilon$ , say  $10^{-3}$ , then GloptiPoly 3 certifies global optimality and convexity with  $\text{obj} = -4.0627\text{e-}7$  at the 4th LMI relaxation. On the other hand, if we relax the constraint into  $p_1(x) + \epsilon \leq 0$




Figure 4: Numerically convex singular quartic.

with a negative  $\epsilon = -10^{-3}$ , then GloptiPoly 3 certifies global optimality and nonconvexity with  $\text{obj} = -0.22313$  at the 4th LMI relaxation. We can conclude that the optimum of problem 3 is sensitive, or ill-conditioned, with respect to the problem data, the coefficients of  $p_1(x)$ . The reason behind this ill-conditioning is the singularity of  $S$  at the origin, see Figure 5 which represents the effect of perturbing the constraint  $p_1(x) \leq 0$  around the singularity.

## 5 Control applications

In this section we focus on control applications of Algorithm 1, which is used to generate convex inner approximation of stability regions in the parameter space.

### 5.1 Fourth-order discrete-time system

Consider now the fourth degree polynomial  $x_2 + x_1z - (x_1 + x_2)z^3 + z^4$ . Its roots are in the open unit disk if and only if  $x = (x_1, x_2)$  belongs to the interior of stability region  $S = \{x \in \mathbb{R}^2 : p_1(x) = 2x_1^2x_2 + 3x_1x_2^2 + 2x_1^2 + x_1x_2 + x_2^2 + x_2 - 1 \leq 0, p_2(x) = -2x_2 - 1 \leq 0, p_3(x) = -2x_1 + x_2 - 2 \leq 0\}$ .

With the following GloptiPoly 3 implementation of Steps 1-3 of Algorithm 1:

```
% problem data
mpol x y 2
```




Figure 5: Perturbed quartic  $p_1(x) + \epsilon \leq 0$  (bold line) can be convex ( $\epsilon = 10^{-3}$ ) or nonconvex ( $\epsilon = -10^{-3}$ ) near singularity of original quartic level set  $p_1(x) = 0$  (light line).

```

p1 = 2*x(1)^2*x(2)+3*x(1)*x(2)^2+2*x(1)^2+x(1)*x(2)+x(2)^2+x(2)-1;
p2 = -2*x(2)-1;
p3 = -2*x(1)+x(2)-2;
g1 = diff(p1,x); % gradient
H1 = diff(g1,x); % Hessian
% LMI relaxation order
order = 4;
% build LMI relaxation
P = msdp(min(y'*H1*y), p1==0, p2<=0, p3<=0, ...
        g1*y==0, y'*y==1, order);
% solve LMI relaxation
[status,obj] = msol(P)

```

we obtain along the cubic boundary  $p_1(x) = 0$  a strictly negative minimum curvature of  $\text{obj} = -6.0344$  at the LMI relaxation of order 4 (moments of degree 8), certified by  $\text{status}=1$ . Region  $S$  is therefore nonconvex. GloptiPoly 3 extracts two solutions  $x^1 = -(0.9784, 0.3499)$ ,  $y^1 = (0.7931, 0.6090)$  and  $x^2 = -(0.9784, 0.3499)$ ,  $y^2 = -(0.7931, 0.6090)$ .

We follow Step 4 of Algorithm 1 add we add the affine constraint  $p_4(x) = g_1^T(x^1)(x - x^1) + \epsilon \leq 0$  with  $\epsilon$  a small positive real, say  $10^{-3}$ , and we run Step 3 again on the new problem:

```

% select one solution
x1 = double(x); x1 = x1(:, :, 1);
y1 = double(y); y1 = y1(:, :, 1);
g11 = double(g1); g11 = g11(:, :, 1);
% new constraint
p4 = g11*(x-x1)+1e-3;
% LMI relaxation order
order = 4;
% build LMI relaxation
P = msdp(min(y'*H1*y), p1==0, p2<=0, p3<=0, p4<=0, ...
    g1*y==0, y'*y==1, order);
% solve LMI relaxation
[status,obj] = msol(P)

```

GloptiPoly 3 now returns a positive minimum curvature  $\text{obj}=0.2663$  certified by  $\text{status}=1$  and hence the resulting semialgebraic set  $\bar{S} = \{x : p_1(x) \leq 0, \dots, p_4(x) \leq 0\}$  is a valid convex inner approximation of nonconvex stability region  $S$ , see Figure 6.




Figure 6: Convex inner approximation (dark gray) of nonconvex fourth-order discrete-time stability region (light gray).

Let us remark that set  $S$  actually does not satisfy Assumption 1 since the cubic curve  $p_3(x) = 0$  has a singular point at  $x = (-1/2, 1)$  denoted by  $s$  on Figure 6. However, since the point of minimum curvature  $x^1$  is found elsewhere and the corresponding affine cut removes  $P$  from  $\bar{S}$ , this is not troublesome.

Finally, from Figure 6 we see that the choice of point of minimum curvature  $x^1$  is not

necessarily optimal in terms of maximizing the surface of  $\bar{S}$ . A point chosen between  $x^1$  and  $s$  along the boundary would be likely to generate a larger convex inner approximation.

## 5.2 Third-order discrete-time stability region

Algorithm 1 can lend insight into the (nonconvex) geometry of the stability region. Consider the simplest non-trivial case of a third-order discrete-time polynomial  $x_1 + x_2z + x_3z^2 + z^3$  which is stable (roots within the open unit disk) if and only if parameter  $x = (x_1, x_2, x_3)$  lies within the interior of compact region  $S = \{x \in \mathbb{R}^3 : p_1(x) = -x_1 - x_2 - x_3 - 1 \leq 0, p_2(x) = x_1 - x_2 + x_3 - 1 \leq 0, p_3(x) = x_1^2 - x_1x_3 + x_2 - 1 \leq 0\}$ . Stability region  $S$  is nonconvex, delimited by two planes  $p_1(x) = 0, p_2(x) = 0$  and a hyperbolic paraboloid  $p_3(x) = 0$  see e.g. [1, Example 11.4].

Optimization problem (3) corresponding to convexity check of the hyperbolic paraboloid reads as follows:

$$\begin{aligned}
\min \quad & -2y_1^2 + 2y_1y_3 \\
\text{s.t.} \quad & x_1^2 - x_1x_3 + x_2 - 1 = 0 \\
& -x_1 - x_2 - x_3 - 1 \leq 0 \\
& x_1 - x_2 + x_3 - 1 \leq 0 \\
& (2x_1 - x_3)y_1 + y_2 + x_3y_3 = 0 \\
& y_1^2 + y_2^2 + y_3^2 = 1.
\end{aligned} \tag{5}$$

The objective function and the last constraint depend only on  $y$ , and necessary optimality conditions obtained by differentiating the Lagrangian  $-2y_1^2 + 2y_1y_3 + t(y_1^2 + y_2^2 + y_3^2 - 1)$  with respect to  $y$  yield the symmetric pencil equation

$$\begin{bmatrix} -4 + 2t & 0 & 2 \\ 0 & 2t & 0 \\ 2 & 0 & 2t \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = 0$$

From the determinant of the above 3-by-3 matrix, equal to  $t(t^2 - 2t - 1)$ , we conclude that multiplier  $t$  can be equal to  $1 - \sqrt{2}, 0$  or  $1 + \sqrt{2}$ . The choice  $t = 0$  implies  $y_1 = 0, y_2 = 1, y_3 = 0$  which is inconsistent with the last but one constraint in (5). The choice  $t = 1 - \sqrt{2}$  yields  $y_1 = \pm(1 + \sqrt{2})\alpha, y_2 = 0, y_3 = \pm\alpha$  with  $\alpha = 1/\sqrt{4 - 2\sqrt{2}}$  and the objective function  $-2y_1^2 + 2y_1y_3 = -1 + \sqrt{2}$ . The choice  $t = 1 + \sqrt{2}$  yields  $y_1 = \pm\alpha, y_2 = 0, y_3 = \pm(-1 - \sqrt{2})\alpha$  and the objective function  $-1 - \sqrt{2}$ , a negative minimum curvature. Therefore region  $S$  is indeed nonconvex.

From the remaining constraints in (5), we conclude that the minimal curvature points  $x$  can be found along the portion of parabola  $\sqrt{2}x_1^2 - x_2 + 1 = 0$  included in the half-planes  $(2 + \sqrt{2})x_1 + x_2 + 1 \geq 0$  and  $-(2 + \sqrt{2})x_1 + x_2 + 1 \geq 0$ . Any plane tangent to the hyperbolic paraboloid  $p_3(x) = 0$  at a point along the parabola  $\sqrt{2}x_1^2 - x_2 + 1 = 0$  can be used to generate a valid inner approximation of the stability region. For example, with the choice  $x^1 = (0, 1, 0)$ , we generate the gradient half-plane  $p_4(x) = g_3(x^1)(x - x^1) = x_2 - 1 \leq 0$ .

More generally, for discrete-time polynomials of degree  $n \geq 3$ , stability region  $S$  is the image of the box  $B = [-1, 1]^n$  (of so-called reflection coefficients) though a multiaffine mapping, see e.g. [13] and references therein. The boundary of  $S$  consists of ruled

surfaces, and the convex hull of  $S$  is generated by the images of the vertices of  $B$  through the multi-affine mapping. It would be interesting to investigate whether this particular geometry can be exploited to generate systematically a convex inner approximation of maximum volume of the stability region  $S$ .

## 6 Conclusion

We have presented a general-purpose computational algorithm to generate a convex inner approximation of a given basic semialgebraic set. The inner approximation is not guaranteed to be of maximum volume, but the algorithm has the favorable features of leaving invariant a convex set, and preserving convex boundaries while removing nonconvex regions by enforcing linear constraints at points of minimum curvature.

Each step of the algorithm consists in solving a potentially nonconvex polynomial optimization problem with the help of a hierarchy of convex LMI relaxations. For this we use Gloptipoly 3, unfortunately with no guarantee of a priori computational burden, even though in practice it is observed that global optimality is ensured at a moderate cost, as soon as the dimension of the ambient space is small. Numerical experiments indicate that the approach may be practical for ambient dimensions up to 4 or 5. For larger problems, we can rely on more sophisticated nonlinear or global optimization codes [12], even though this possibility has not been investigated in this paper. Indeed, our main driving force is to contribute with a readily available Matlab implementation.

Generally speaking, one may question the relevance of applying a relatively complex algorithm to obtain a convex inner approximation in the form of a list of defining polynomials which are not necessary individually convex. Indeed, if convexity of the inner approximation is guaranteed in the presented work, convexity of the defining polynomials would allow the use of constant multipliers to certificate optimality in a nonlinear optimization framework. Instead, with no guarantee of convexity of the defining polynomials, the geometric property of convexity of the sets is more delicate to exploit efficiently by optimization algorithms. However, it is conjectured that all convex semialgebraic sets are semidefinite representable in [3], see also [10]. It may then become possible to fully exploit the geometric convexity of our inner convex through an explicit representation as a projection of an affine section of the semidefinite cone. For example, in our target application domain, this would allow to use semidefinite programming to find a suboptimal stabilizing fixed-order controller.

## Acknowledgements

The first author is grateful to J. W. Helton for many discussions and ideas leading to Theorem 1.

## References

- [1] J. Ackermann et al. Robust control: the parameter space approach. Springer, 2nd edition, 2002.
- [2] M. P. do Carmo. Differential geometry of curves and surfaces. Prentice Hall, 1976.
- [3] J. W. Helton, J. Nie. Sufficient and necessary conditions for semidefinite representability of convex hulls and sets, *SIAM J. Optimization*, 20(2):759-791, 2009.
- [4] D. Henrion, M. Šebek, V. Kučera. Positive polynomials and robust stabilization with fixed-order controllers, *IEEE Trans. Autom. Control*, 48(7):1178-1186, 2003.
- [5] D. Henrion, D. Peaucelle, D. Arzelier, M. Šebek. Ellipsoidal approximation of the stability domain of a polynomial, *IEEE Trans. Autom. Control*, 48(12):2255-2259, 2003.
- [6] D. Henrion, J. B. Lasserre. Solving nonconvex optimization problems - How GloptiPoly is applied to problems in robust and nonlinear control, *IEEE Control Systems Magazine*, 24(3):72-83, 2004.
- [7] D. Henrion, J. B. Lasserre, J. Löfberg, GloptiPoly 3: moments, optimization and semidefinite programming, *Optimization Methods and Software*, 24(4-5):761-779, 2009.
- [8] D. Henrion, J. B. Lasserre, C. Savorgnan. Approximate volume and integration for basic semialgebraic sets, *SIAM Review*, 51(4):722-743, 2009.
- [9] A. Karimi, H. Khatibi, R. Longchamp. Robust control of polytopic systems by convex optimization. *Automatica*, 43(6):1395-1402, 2007.
- [10] J. B. Lasserre. Convex sets with semidefinite representation, *Math. Programming*, 120:457-477, 2009.
- [11] M. Laurent. Sums of squares, moment matrices and optimization over polynomials, in: M. Putinar, S. Sullivant (Eds.), *Emerging applications of algebraic geometry*, IMA Vol. Math. Appli., 149:157-270, Springer, 2009.
- [12] A. Neumaier. Complete search in continuous global optimization and constraint satisfaction, in: A. Iserles (Ed.), *Acta Numerica*, Cambridge Univ. Press, 271-369, 2004.
- [13] U. Nurges. Robust pole assignment via reflection coefficients of polynomials, *Automatica*, 42(7):1223-1230, 2006.