



HAL
open science

Seismic stratigraphy and deformational styles of the offshore Cyrenaica (Libya) and bordering Mediterranean Ridge

Yem L. Mbida, L. Camera, J. Mascle, A. Ribodetti

► **To cite this version:**

Yem L. Mbida, L. Camera, J. Mascle, A. Ribodetti. Seismic stratigraphy and deformational styles of the offshore Cyrenaica (Libya) and bordering Mediterranean Ridge. *Geophysical Journal International*, 2011, 185 (1), pp.65-77. 10.1111/j.1365-246X.2011.04928.x . hal-00585749

HAL Id: hal-00585749

<https://hal.science/hal-00585749>

Submitted on 2 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Seismic stratigraphy and deformational styles of the offshore Cyrenaica (Libya) and bordering Mediterranean Ridge

Lionel Mbida Yem,^{1,2} Laurent Camera,³ Jean Mascle³ and Alessandra Ribodetti¹

¹Université de Nice Sophia-Antipolis, IRD, Observatoire de la Côte d'Azur, Géoazur F-06235, BP 48, Villefranche-sur-mer, France.

E-mail: ribodetti@geoazur.obs-vlfr.fr

²Applied Geology Laboratory UYI, Yaoundé, Cameroon

³Université de Nice Sophia-Antipolis, CNRS, Observatoire de la Côte d'Azur, Géoazur F-06235, BP 48, Villefranche-sur-mer, France

Accepted 2010 December 21. Received 2010 December 13; in original form 2009 June 4

SUMMARY

Off northwest Libya the Cyrenaica foreland basin domain and its Pan-African continental crust, which constitute the African promontory, are overthrust by the Mediterranean Ridge Complex. The thrust belt contact and its seismic stratigraphy have been analysed using pre-stack depth-migrated multichannel seismic (MCS) lines recorded during the MEDISIS survey (2002). The geometry and sedimentary distribution analysis through the wedge-top depocentres allow reconstruction of schematic cross-sections of the tectono-sedimentary wedge that includes two major thrust sequences separated by an apparently poorly deformed transition zone. Based on time–space variation of several piggyback basins, we propose that these thrust sequences relate to distinct phases of shortening. (1) A first event, which probably occurred just prior to the Messinian crisis in latest Miocene (Tortonian times?) and (2) A second event, that has finally led to the present-day overthrusting of the Mediterranean Ridge over the Libyan continental slope.

Key words: Image processing; Continental margins: convergent; Continental margins: divergent; Africa; Europe.

1 INTRODUCTION

Extending off Libya, the north Cyrenaica foreland basin (Fig. 1) results from a diachronous geological evolution including a passive margin stage and its subsequent tectonic inversion (and uplift), which originated from a still ongoing convergence between the African and Eurasian plates. According to DeCelles & Giles (1996) this area, extending along the southern Neo-Tethys border, can be regarded as part of a foreland-type basin which includes (1) a foredeep depositional system, the present-day Dernah-Tubruq and Benghazi basins and (2) a wedge-top depozone, the Mediterranean Ridge (Fig. 1).

Up to now the foredeep region remains still poorly studied and it therefore remains difficult to properly locate, through the foreland basin, the seaward extension of the passive African continental margin. While considered for a long time as enigmatic, the geology of the Central Mediterranean Ridge has recently been clarified through the use of detailed swath bathymetric data (Huguen & Mascle 2005; Huguen *et al.* 2005), various academic seismic reflection data (Cita *et al.* 1989; Camerlenghi *et al.* 1992, 1995; De Voogd *et al.* 1992; Zitter 2004; Chamot-Rooke *et al.* 2005) including several multichannel seismic (MCS) lines (Finetti 1982; Chaumillon 1995; Kopf *et al.* 1998; Mascle & Chaumillon 1998; Costa *et al.* 2004) and ODP leg 160 drilling results (Emeis *et al.* 1996). Some of these studies have shown that the wedge complex includes strati-

graphic mélanges made of Mesozoic to Plio-Quaternary sediments (Finetti 1982; Ahlbrandt 2001) emplaced on the subducting African basement. According to Huguen & Mascle (2005), the wedge can be divided into three different morphostructural provinces: (1) a southern front overthrusting the passive African continental margin, (2) a central, rather flat, domain and (3) a northern province in backthrust contact to the North with the Hellenic active margin. Whether this morphologic feature expresses only relatively shallow thrust sequences within the wedge-top depozone, or deeply rooted structures of the subducting African crust, remains an open debate.

The few available geophysical data have always faced difficulties providing reliable images of the deep structure of the wedge-top depozone, mainly due to the presence, in the shallow section of the wedge, of thick Messinian salt deposits (Hsü *et al.* 1973; Finetti 1982; Mascle & Chaumillon 1998), and to its overall and intense tectonic deformation. In this study we have applied pre-stack depth migration (PSDM) techniques on five regional MCS lines to tentatively obtain better depth images of the geological structures and of the structural relationships of this accreted wedge with its foredeep domain, the passive Libyan continental margin.

Bosworth *et al.* (2008) proposed that a 'strain shadow' (see their Fig. 1) affects the shore area just south and southeast of Cyrenaica where only little shortening is evidenced. However, we can wonder whether some deformation may occur on the opposite side of




Figure 1. Main morphostructural elements of the studied area, including the Cyrenaica foreland basin and the bordering Mediterranean Ridge (modified from Chamot-Rooke *et al.*, 2005) are shown on a shaded bathymetric map (from Gebco, 2003). Location of MCS lines MD11, 13, 17, 18 and 19, illustrated and discussed in this paper, are indicated in blue lines. In red, main thrust zones (frontal thrust and back thrust of the Mediterranean Ridge) and main strike-slip fault zones across the Mediterranean Ridge and its backstop. In upper left corner: plates, main kinematic, tectonic elements and location of the study area (after Huguen *et al.*, 2005).


the ‘strain shadow’ zone (offshore Cyrenaica), which has not been explored and evidenced so far. In this case, how this shortening is expressed and what its timing is are questions we are tentatively trying to answer.

2 REGIONAL GEODYNAMIC SETTING

According to Janssen *et al.* (1995), the Apulian domain was, until Jurassic times, part of the African Plate, and was then bounded to the North by the Early Triassic Tethyan Ocean (Makhlouf 2006). Palaeogeographic reconstructions of the Tethyan-Mediterranean domain which extended from Northern Libya to Southern Aegean sea (e.g. Guiraud & Bosworth 1999; Stampfli & Borel 2000) show that the onset of the closure of the Tethyan oceanic space, generated by a left-lateral translation of the East Saharian domain with respect to the Hellenides-Rhodopes superterrane, probably initiated in Barremian times (Dercourt *et al.* 1986; Janssen *et al.* 1995). The northward drift of Africa relative to Europe, emphasized by the opening of the South and Equatorial Atlantic Oceans (Craig *et al.* 2008), led to (1) the collision, and subsequent suture, of different cratonic blocks across the remnants of the Tethys ocean, (2) the rifting of Apulia away from Africa (Janssen *et al.* 1995) and

finally (3) the opening of the Mesogean Sea (Dercourt *et al.* 1986; Ricou *et al.* 1986; Le Pichon *et al.* 1988; Ziegler 1988). Geophysical and field data indicate that the late Mesozoic rifting phase of the North Africa basement (Guiraud & Bosworth 1999) was accompanied by a series of aborted rifts south of the actual continental margin (Tawadros 2001; Craig *et al.* 2008). A late Cretaceous regional high subsidence rate, possibly associated with the Sirte collapse (Rusk 2001; Carr 2003), is effectively recognized in the Atlas mountains (Zouaghi 2005) and in Libya’s Kufrah, Murzuq, Ghadami and Tripolitania basins. Through Cenozoic times, the progressive shortening between the African continental margin (Cyrenaica) and the Hellenic arc (Kempfer & Garfunkel 1994; Rosenbaum *et al.* 2002) has led to the creation of a tectono-sedimentary wedge, the Mediterranean Ridge complex (Fig. 1; McKenzie 1970), and even to some tectonic inversion of the Cyrenaica shore basins (Röhlich 1980; El Hawat & Abdulsamad, 2004). According to Bosworth *et al.* (2008), the stratigraphy of the Tubruq offshore basin includes Palaeozoic silico-clastic sediments, unconformably overlain by Triassic to Early-Cretaceous pre-inversion package, topped by foreland sequences, including Late Cretaceous to Eocene and Oligocene to Holocene sediments (Table 1). Seismic reflection profiles shown in this paper provide images of this thick sedimentary section within both the foredeep system and the wedge-top area.

Table 1. Synthetic stratigraphy of Mesozoic sequences (modified from Carr (2003)). NA, The Mesozoic succession of North Africa is divided into two sequences (NA 6, Early Triassic to Middle Cretaceous; and NA 7, Middle to Late Cretaceous); LST, lowstand systems tract; TST, transgressive systems tract; HST, highstand systems tract; MFS, maximum flooding surface; SB, sequence boundary; AP, second-order sequence stratigraphic framework for the Arabian Plate [Sequence AP 6 (Late Permian to Toarcian); Sequence AP 7 (Toarcian to Tithonian); Sequence AP 8 (Tithonian to Turonian); and Sequence AP 9 (Turonian to Palaeocene)].



Downloaded from https://academic.oup.com/gji/article/185/1/65/605919 by guest on 02 October 2021

3 DATA PROCESSING

During the MEDISIS survey (run in 2002) several MCS profiles were recorded across the Libya foreland basin to tentatively image, beneath the Mediterranean Ridge, the northward extension of this segment of the Mesozoic African passive margin and to evaluate the structure and nature of the sedimentary cover of both domains. These lines were recorded using a 4515 cubic inches air gun array, with shot spacing of 75 m, and a 360 channel, 4.5-km-long

streamer; the source and receivers configuration resulted in a 30 common depth point (CDP) fold coverage. Before performing a pre-stack depth migration, a more conventional seismic processing sequence was carried out using Geovector software to improve the data signal-to-noise ratio. This sequence included minimum phase signal conversion, predictive deconvolution, Butterworth filtering (3–6 and 65–75 Hz), *F–K* antimultiple attenuation and normal moveout correction. Amplitudes of the data were preserved as much as possible. The velocity analysis was performed by both

semblance panel techniques and variant stack corrections at every 100 or 200 CDP, depending on the length of each line and of the structural complexity of the area. After conversion into interval velocities, using Dix equations, the inferred velocity model was interpolated and subsequently smoothed for dynamic ray tracing (Lambaré *et al.* 1992) and for better estimate of traveltimes, amplitude and ray-related parameters. PSDM (Rai–Born inversion tomography) was then applied on data sorted in shotpoint gathers. During the PSDM process (Thierry *et al.* 1999), common image gathers (CIG) and semblance panels are stored for each kilometre. These CIG were used for quality control of the final migrated image and to further refine the PSDM accuracy by iterative velocity analysis in the depth domain. The redundant image of reflectors for each offset, or diffraction angle in the CIG, must be flat and at the same depth position to guarantee optimal processing of the final section (Fig. 2). To improve the accuracy of the PSDM method used to migrate the MEDISIS data, we have performed two iterative migration-velocity analyses for each line shown and discussed in this paper.

4 THE DIFFERENT SEISMIC SEQUENCES

4.1 Seismic sequences

Analysis of seismic sections obtained after PSDM processing has allowed us to distinguish through the study area three major seismic sequences respectively labeled A, B and C in Figs 4–8. These sequences correlate with a well-stratified sedimentary cover whose thickness varies between 2500 and 5000 m along the foredeep basin and may reach 6000 m in the distal domain of the wedge-top depozone.

The strategy used to define these reflection packages is based on (1) their seismic stratigraphy, (2) some sequential stratigraphy analysis, as defined by Vail *et al.* (1977) and Van Wagoner *et al.* (1988); this technique recommends interpreting the depositional sequences on seismic sections by identifying discontinuities at the base of reflection terminations and (3) stratigraphy as derived from on-shore observations and drilling (Duronio *et al.* 1991), and regional geological synthesis (Carr 2003; Bosworth *et al.* 2008). Based on these various approaches, three major bounding discontinuities were recognized: (1) a probable maximum flooding surface (MFS) lies between A1 and A2 seismic facies units; (2) a sequence boundary (SB1), which truncates the depositional sequence A2 and (3) a second sequence boundary (SB2) capping seismic sequence B. In the following section, seismic sequences are described from the oldest to the youngest one, and from the foredeep towards the distal wedge-top depozone area to facilitate stratigraphic correlation.

4.1.1 Seismic sequence A

This sequence represents the deepest observed seismic unit. Its base is not clearly imaged through the seismic data. However, a banded package of high amplitude seismic reflectors, seen at the base of a structural high on line MD18 (Fig. 3), possibly delineates sequence A basement. A can be divided into two distinct seismic facies units (A1, A2) with different characters, separated by a probable MFS. The lower interval (A1) is made of discontinuous parallel, fair and high amplitude seismic reflections, sometimes affected by extensional faulting (Fig. 4; line MD17). In addition, this unit displays local mound-shaped strata (shale? salt?) and lenses of chaotic reflection

bodies (Figs 4 and 5) possibly indicating channel fills (Mitchum 1977).

Towards the distal foreland, A1 presents staircase structures associated with low angle northward dipping oblique faults (Fig. 4). Through the Benghazi wedge-top inner domain (Fig. 6; line MD11) the A1 seismic expression is less obvious. The poor quality of seismic information in this area may be due to fluid circulation along in-sequence thrust and released on the seafloor as mud volcanoes. Further north, a banded reflection package, similar to the one observed in the Tubruq foredeep basin, is found between 3.5 and 8.2 km depth on lines MD11 and MD13 (Figs 6 and 7). According to Bally (1987), sedimentary lenses are common in prograding facies units and in near deltaic environments. We thus interpret A1 as a probable parasequence set including a late low stands system tract and sediments deposited from the onset of sea level rise until the time of maximum transgression on the coast. Additional supports for such interpretation are the presence of potential failing stage salt deposits through the sequence and the presence of an overlying MFS (Fig. 3).

An upper sequence, A2, highlights aggrading (Fig. 3) and prograding (Fig. 4) seismic reflections with a bank-like external geometry. This package is characterized by more continuous, parallel and high amplitude seismic reflections (Fig. 3). Local disruptions of the A2 seismic facies unit, together with dispersed reflections, are observed in the area of deformation (i.e. high angle normal fault) or in dome-like rising features above the general level of surrounding strata, or in areas of potential fluid migration (Figs 3 and 4). Following Wilson (1975), the A2 layer-cake high amplitude reflectors (Fig. 4) can be interpreted as an aggrading shelf edge complex that builds out in deep water during sea level high stand (Vail *et al.* 1977; Sarg *et al.* 1999).

Among this high-stand systems tract (Vail 1987; Posamentier *et al.* 1988; Posamentier & Allen 1999), the eroded unconformity (SB1) displays a linear shaped seismic reflection pattern that extends north of the Cyrenaica foredeep basins and may define a major global sea level fall after the deposition of A2 sequence facies units.

4.1.2 Seismic sequence B

This seismic sequence shows a base-concordant relation with SB1 sequence boundary along the shelf edge (Fig. 3), and down-dip succession of its lower strata against MFS in a basin-ward direction (Fig. 4). These bounding patterns suggest a parasequence set with bank-like external geometry (Kendall *et al.* 2003).

B is widespread on the study area and characterized by two different types of seismic reflections. A first group is seen along the present-day shelf edge and shows stratification patterns with good continuity, exhibiting high amplitude locally interrupted by discontinuous low amplitudes. A second type, observed in the deep water basin and through the wedge-top depozone, includes chaotic and blank massive shape sequences with intermittent internal reflections. This deep-water seismic facies unit displays strong variations in thickness associated with post-deposition deep normal fault reactivation and piggyback-basin development (Fig. 4).

Following Fontaine & Cussey (1987), the up-dip B made of parallel and high frequency reflections may indicate pelagic carbonates accumulation during the transgressive stage. We therefore interpret the B sequence as a potential carbonate parasequence (Van Wagoner *et al.* 1988; changing in the basin-ward direction to downslope mass flows) and dolomitic limestone associated with a post-deposition basin restriction. According to Deelman (1999) and Sarg (2002),




Figure 2. Examples of quality and accuracy controls of depth-migrated seismic line. (A): Common image gathers (in grey and white); iso-X-panels (in colour) show maximum coherence (green-blue), around one for the main reflectors on semblance panels; blue dots and arrows indicate the locations of main seismic discontinuities; SR, seabed reflector; SB, sequences boundaries (1, 2 and 3). (B): Example of pre-stack depth-migrated line MD17, from the Libyan continental slope to the Mediterranean Ridge.




Figure 3. Seismic stratigraphy and internal structure of the Tubruq foredeep basin from PSDM line MD18. A, B and C indicate the different seismic sequences as indicated in the text. MFS, maximum flooding surface; SB1 and SB2, sequence boundaries.




Figure 4. Seismic stratigraphy and structural style at the contact between the Mediterranean Ridge complex and the Tubruq foredeep sequences from PSDM line MD17. A, B and C indicate the different seismic sequences as indicated in the text. MFS: maximum flooding surface; SB1 and SB2: sequence boundaries. Blue arrows underline terminations against sequence boundaries. Bold line: thrust or extensional faults; regular lines: boundaries of main seismic reflectors.




Figure 5. Seismic stratigraphy and structural patterns of the Tubruq foredeep basin/Cyrenaica thrust sheets and seaward extension of the foredeep sequence from PSDM image MD 19. A, B and C indicate the different seismic sequences as indicated in the text, SB1 and SB2: sequence boundaries.




Figure 6. Seismic sequences and structural style of the inner Mediterranean complex as illustrated on PSDM line MD11. A, B and C indicate seismic sequences. Bold lines: thrust systems; regular lines: boundaries of main seismic reflectors.




Figure 7. Seismic sequences and structural patterns of Neo-Tethys Duplex, as interpreted from PSDM lines MD13. Black bold lines: thrust or detachment level; regular lines: boundaries of main seismic reflectors. SB1 and SB2: sequences boundaries. A, B and C: seismic sequences.

tectonic settings prone to hydrographic restriction (such as foreland basin) provide favourable conditions for carbonate to evaporite transition during sea level low stands. A large-scale shelf unconformity surface (SB2), seen at the top of **B** seismic sequence, attests for a post-date sea level drawdown associated with uplift and thrusting (Fig. 4). This combined tectonic–eustatic event probably accommodated restricted conditions leading to **B** potential carbonate dolomitization. Our hypothesis is supported by the presence of diapiric-type structures observed within **B** in the distal foreland region (Fig. 7).

4.1.3 Seismic sequence C

C lies between the regional SB2 seismic sequence boundary and the seafloor everywhere in the foreland basin. Its internal pattern is also characterized by two seismic facies units: **C1** and **C2**. A lower package (**C1**), well developed in the Tubruq foredeep depozone, shows subparallel and high amplitude banded reflections (Figs 3 and 4). An upper package (**C2**) is characterized by transparent weak seismic reflection. Both units form a stratigraphic melange with fill configuration in the toe of the Tubruq accretionary complex (Fig. 4) and linear shape along the distal part of the wedge-top depozone (Figs 6 and 7). This stratigraphic melange is characterized by interbedded, fine and high frequency continuous seismic reflections with which some weak transparent facies can be recognized. In addition, it exhibits evidence of active salt welts, created by salt flowage in the lowermost part of the sequence (Fig. 4).

In summary, the distribution patterns and internal reflections of **C** seismic sequence lead to interpret a shelf margin wedge (Homewood *et al.* 2002) including forced regressive salt deposits (likely Messinian) developed during stepped sea level falls.

4.2 Tentative correlation with the regional chronostratigraphy

An integrated approach, based on seismic stratigraphy analysis and previous geological synthesis (Duronio *et al.* 1991; Carr 2003; Bosworth *et al.* 2008), leads us to document three mega sequences within the Cyrenaica foreland depozone. (1) A lowermost one overlies an irregular acoustic basement expressed by series of tilted blocks along the shelf edge (Figs 3 and 4) and thick-skinned thrusting within the distal foreland; (2) an intermediate sequence is widespread and displays syn and post-deposition tectonic features along the wedge-top depozone and (3) a last depositional sequence reflects tectonic reactivation of foredeep extensional faults coeval with piggyback-basin development.

The sedimentary sequences deduced from the acoustic facies analysis are related to two parasequences (low stands and lower transgressive system tract) **A1**, topped by a high stand system tract (**A2**) and an upper transgressive system tract (**B**) overlain by a shelf margin wedge (**C**). Along the Eastern Mediterranean domain, as well as in Libyan shore basins, comparable seismic sequences and lithofacies interpretation have been proposed by several authors (Finetti 1982; Biju-Duval *et al.* 1987; Duronio *et al.* 1991; Shegawi 1992; Chaumillon 1995; Emeis *et al.* 1996; Gumati *et al.* 1996; Mascle & Chaumillon 1998; Ahlbrandt 2001; Rusk 2001; Tay *et al.* 2002; Carr 2003; El Hawat & Abdulsamad 2004; Bosworth *et al.* 2008). The regional history stemming from these studies includes Triassic-to-Early-Cretaceous crustal extension, followed by uplift and folding during Eocene–Oligocene times (Gumati & Nairn 1991). According to Gumati *et al.* (1996) and Bosworth *et al.*

(2008), the stratigraphic succession of the Cyrenaica platform includes a Palaeozoic basement unconformably overlain by Triassic to Cretaceous sediments. Carr (2003) divided the Mesozoic succession across North Africa (including Sirte and Cyrenaica) into two sequences: (1) an Early Triassic to middle Cretaceous sequence, unconformably overlain by (2) a middle to late Cretaceous sedimentary cover. Despite a lack of control by well data, a comparison of its second-order sequences stratigraphy framework (Table 1) with our interpretation lead us to respectively correlate MFS to the Oxfordian maximum transgression, A2 to the Late Jurassic Early Cretaceous and SB1 to the top of the Albian unconformity. From this attempt, B can be assigned to the Cenomanian–Eocene sequence (Bosworth *et al.* 2008), SB2 to the top of an Eocene unconformity (Ceriani *et al.* 2002) and C to an Oligocene–Holocene sequence (Ahlbrandt 2001; Bosworth *et al.* 2008).

5 TECTONIC FEATURES OF THE CYRENAICA WEDGE-TOP

Interpretation of MD11, 13, 17 and 19 lines tends to indicate that several phases of tectonic activity have affected the Cyrenaica basement; they include the formation of NE–SW trending grabens during Triassic to Early Cretaceous (Fig. 3) and the development of regional highs and lows associated with Eocene–Holocene thrusting (Fig. 4). In the following section, we will focus on the structural analysis of the wedge-top depozone to tentatively reconstruct the post-rift deformation of the Cyrenaica foreland basin. As can be seen in Fig. 1, the Central Mediterranean Ridge complex includes two distinct mega thrust sequences; the two sequences are separated by an area characterized by thick-slip sliding structures (Fig. 4), and a high velocity (6.2 km s^{-1}) according to PSDM iterative velocity analysis. We have named the northern thrust wedge, Neo-Tethyan-Duplex and the frontal prism, Cyrenaica thrust-sheet.

5.1 Neo-Tethyan Duplex

MCS sections MD18, 17, 19, 11 and 13 (Figs 3–7, respectively) indicate that the Neo-Tethyan Duplex extends between 70 and 130 km away from the base of the Cyrenaica continental slope (Fig. 1). These lines show that the Neo-Tethyan Duplex may include several ramps potentially at a crustal scale (Figs 6 and 7). According to the hypothesis of Bosworth *et al.* (2008) and Carr (2003), folds and thrusting in this area may involve sediments up to Jurassic to middle Cretaceous. Moreover, when considering the seismic features and thickness variations, as illustrated on lines MD11 and MD13, the Neo-Tethyan Duplex seems to be characterized by hinterland-direct shortening migration attesting probable back stepping of wedge accretion. This observation leads us to interpret a wedge over a critical profile (Dahlen *et al.* 1984; Muños *et al.* 1985; Zoetemeijer *et al.* 1992). This hypothesis is supported by the northward passive duplex (roof backthrust) that marked the structural front of the Neo-Tethyan Duplex. The flat bedding unconformity of likely Early Pliocene age, that caps the top of the Miocene fan shape sequence, leads to believe that the basal detachment (D1) of the Neo-Tethyan Duplex reached its critical dip position at that time, or just prior to the onset of Cyrenaica thrust-sheet accretion.

5.2 Cyrenaica thrust-sheet

Structural analysis of seismic profiles MD17 and MD19 shows that the Cyrenaica thrust-sheet spreads over nearly 40 km (Fig. 5) above

late Cretaceous and Tertiary fore deep sequence, directly above the passive African continental margin. Unlike the Neo-Tethyan Duplex, the Cyrenaica thrust-sheet displays typical thin-skinned thrust system (Fig. 4) that may involve Upper Cretaceous to Holocene sedimentary units. This package is detached from its Jurassic to Early Cretaceous sole (the sedimentary cover of the African margin) along fault plane (D2) with ramp-flat-geometry. Its hinterland zone is characterized by tectonic denudation of foredeep sequence where exhumation of deep-seated rocks was accomplished by ramp-decollement geometry. The presence of a wide and thick Messinian piggyback basin behind the passive duplex indicates that the onset of Cyrenaica thrust-sheet accretion is most likely coeval with the latest Miocene deformation and uplift.

6 DISCUSSION AND CONCLUSIONS

6.1 The Cyrenaica margin

Tentative chronostratigraphic correlation tends to indicate that our interpretation is, in general, in good agreement with usually proposed Mesozoic evolution of the southern Tethyan margin. According to the MCS data interpretation the sedimentary pile of this margin segment may include Triassic salt, and evaporite accumulated during the Messinian crisis. Up to now no evidence of evaporite older than the Messinian ones has been reported offshore Eastern Libya; in western Libya however, salt deposits (Late Triassic to Middle Jurassic in age) constitute one of the major hydrocarbon plays within the Pelagian domain (Sabratat subbasin in Libya, Asstart subbasin in Tunisia), where they generated domes-like structures that later affected both sedimentation and oil entrapment (Bosworth, 2009, personal communication). The lack of published well data needed to calibrate our seismic data makes uncertain the Triassic age we have tentatively attributed to deep diapirs-like and folded features observed beneath the Tubruq shelf basin (Fig. 3). Our hypothesis is however supported by several lines of reasoning. First, the mother layers of the postulated salt features, which can be seen in the deepest part of the cross-section, do not appear affected by superficial compression and do not show any structural relationship with identified Messinian sequences. Secondly, the North Tubruq Triassic salt basin shows some NE–SW sag similar to the early Mesozoic evaporite through known in the central Atlas of Tunisia (Bédir *et al.* 2001; Zouaghi *et al.* 2005). Finally, assuming an early stage, in late Palaeozoic, for the opening of the Tethyan ocean (Sengör 1979; Stampfli *et al.* 2004; Garfunkel 2004), it may not be surprising to observe salt deposits in confined depressions, as is usually the case during early rifting phases.

On our seismic lines we have tentatively recognized within the post-Jurassic cover of the eastern Libyan margin two main regional unconformities: a first one on top of the Albian, and a second on top of the Eocene. These unconformities and their erosional surfaces (see Figs 3 and 4) indicate that along the Tubruq region foredeep the depocentres shifted from south to north prior to Tertiary times and maybe before a Santonian event (Guiraud *et al.* 2005) indicating a tectonic inversion of the Neotethyan basins. Such a scenario is supported by a continuous uplift and erosion time and shift during Campanian and Maastrichtian of the southern Neo-Tethyan margin, which is supported by the deposition of thick Palaeocene pile directly overlying late Jurassic strata (Fig. 3). This observation is in good agreement with the hypothesis of Bosworth *et al.* (2008), which indicates a global change in the kinematic of the North African terrains in Santonian times. The sedimentary




Figure 8. Schematic North-South regional cross-section across the Cyrenaica wedge-top depozone illustrating its main structural characteristics.

distribution and observed tectonic features indicate that the Cyrenaica foreland basin can be divided into domains: (1) the Dernah-Tubruq basin and (2) Benghazi (Jabal al Akhdar) foredeep, separated by a master, NNE-SSW trending, strike-slip fault zone (Fig. 1).

6.2 The northern bordering Mediterranean Ridge

From the tentative reconstructed cross-section shown in Fig. 8 we may distinguish two different thrust sequences: (1) a northern one, or Neo-Tethyan Duplex, characterized by ramp anticlines and (2) a southern one, or Cyrenaica thrust sheets, overthrusting the foredeep sequences. The identification of Messinian salt sequences, the discrimination between regional seismic sequence boundaries and the architectures of both thrust domains allow us to identify two main and successive phases of shortening, respectively during late Miocene time (possibly the Tortonian event of Guiraud *et al.* (2005)) and just after the Messinian time. Our results are somewhat consistent with what is known from the evolution of the Southern Apennines (Ori & Friend 1984; Bonini *et al.* 1998; Piedilato & Prosser 2005) and of southwestern Turkey margin (Alçiçek & Veen 2008) where comparable events have been described, including a relative out-of-sequence thrusting during the Miocene (post-Aquitania). This seems to support the occurrence of a regional shortening event sometimes during upper Miocene (Tortonian?) times affecting most of the Eastern Mediterranean domain. An important issue concerns the interpretation of the inner domain of the Mediterranean ridge (Figs 3–7 and 8); due to its relatively elevated position with respect to the southern thrust-sheet, we suspect that this thick-slip sliding domain correlates with a faulted acoustic basement bulge. Accordingly, the specific acoustic response of the overlying sediments may result from fluid circulations along in-sequence thrusts or along normal faults reactivated by compression events. If correct, this would indicate that mud expulsion and gas chimneys observed through the inner part of Mediterranean ridge Complex are controlled by deep-seated tectonic features such as thrusts or reverse faults as already proposed by Huguen *et al.* (2001, 2005).

Besides their scientific impact, the results presented in this paper may also provide interesting issues for petroleum exploration along the Cyrenaica foreland basin. Further investigations, involving wide-angle seismic data, should be undertaken especially within the central domain of the Mediterranean ridge to improve our knowledge of its internal structure and lithology.

ACKNOWLEDGMENTS

This work is part of the L. Mbida Yem' Ph.D., supported by a grant from Institut de Recherche pour le Developpement (IRD). The authors gratefully acknowledge the chief scientist and staff onboard the N/O NADIR during the Medisis cruise. We express our thanks to P. Thierry, G. Lambaré and S. Operto, who developed the preserved amplitude pre-stack depth migration software used for this study; special thanks to G. Lambaré for providing his dynamic ray tracing code. The first author is also grateful to F. Sage-Khadir for her assistance during pre-processing of seismic data on Geovector, and to E. Lopez for stimulating discussions. We are grateful to Ben Yates for his proofreading and English comments.

REFERENCES

- Ahlbrandt, T., 2001. Sirte Basin Province: Sirte-Zelten Total Petroleum System, *U.S. Geol. Surv. Bull.*, 2202–F, 29pp., <http://geology.cr.usgs.gov/pub/bulletins/b2202-f>.

- Alçiçek, M.C. & ten Veen, J.H., 2008. The late early Miocene Acipayam piggy-back basin: refining the last stages of Lycian nappe emplacement in the SW Turkey, *Sediment. Geol.*, **208**, 101–113.
- Bally, A.W., 1987. Atlas of seismic stratigraphy, *AAPG Stud. Geol.*, **1**(27), 1–10.
- Bédir, M. et al., 2001. Subsurface Mesozoic basins in the central Atlas of Tunisia: tectonics, sequence deposit distribution, and hydrocarbon potential, *AAPG Bull.*, **85**(5), 885–907.
- Biju-Duval, B., Morel, Y., Burolet, P.F., Winnock, E., Ravenne, C., Mascle, G. & Charier S., 1987. Le plateau Cyrénien: promontoire Africain sur la marge ionienne, *Revue de l'institut français du pétrole.*, **42**(4), 419–447.
- Bonini M, Moratti G. & Sani F., 1998. Evolution and decollement migration in thrust-top basins: inferences from the Messinian Velona Basin (Northern Apennines, Italy), *Tectonophysics*, **304**, 95–108.
- Bosworth, W., El-Hawat, A.S., Helgeson, D.E. & Burke, K., 2008. Cyrenaica “shock absorber” and associated inversion strain shadow in the collision zone of northeast Africa, *Geology*, **36**(9), 695–698.
- Camerlenghi, A., Cita, M.B., Hieke, W. & Ricchiuto T., 1992. Geological evidence for mud diapirism on the Mediterranean Ridge accretionary complex: Fluids in convergent margins, *Earth planet. Sci. Lett.*, **109**(3–4), 493–504.
- Camerlenghi, A., Cita, M.B., Della Vedova, B., Fusi, N., Mirabile, L. & Pellis G., 1995. Geophysical evidence of mud diapirism on the Mediterranean Ridge accretionary complex, *Mar. geophys. Res.*, **17**(2), 115–141.
- Carr, I.D., 2003. A sequence stratigraphy synthesis of the North African Mesozoic, *J. Petrol. Geol.*, **26**, 133–152.
- Ceriani, A., Di Giulio, A., Goldstein, R.H. & Rossi C., 2002. Diagenesis associated with cooling during burial: and example for Lower Cretaceous reservoir sandstones (Sirt basin, Libya), *AAPG Bull.*, **86**(9), 1573–1591.
- Chamot-Rooke N., Rangin, C. Le Pichon, X. & Dotmed Working Group, 2005. DOTMED Deep Offshore Tectonics of the Mediterranean: a synthesis of deep marine data in Eastern Mediterranean, *Mémoire Société géologique de France and American Association of Petroleum Geologists*, N. numero special, 177, 64 pp.
- Chaumillon, E., 1995. Structure de la ride méditerranéenne: apports de la sismique multitraçage, *PhD thesis*, Peirre & Marie Curie University, Paris.
- Cita, M.B. et al., 1989. Discovery of mud diapirism in the Mediterranean Ridge. A preliminary report, *Bull. Soc. Geol. It.*, **108**, 537–543.
- Costa, E. et al., 2004. Modeling deformation and salt tectonics in the eastern Mediterranean Ridge accretionary wedge, *GSA Bull.*, **116**, 880–894.
- Craig, J. et al., 2008. Structural styles and prospectivity in the Precambrian and Palaeozoic Hydrocarbon Systems of North Africa, *Geology of East Libya*, **4**, 51–122.
- Dahlen, F.A., Suppe, J. & Davis, D., 1984. Mechanics of fold and thrust belts and accretionary wedges: cohesive Coulomb theory, *J. geophys. Res.*, **89**, 10 087–10 101.
- DeCelles, P.G. & Giles, K.A., 1996. Foreland basin systems, *Basin Res.*, **8**, 105–123.
- Deelman, J.C., 1999. Low-temperature nucleation of magnesite and dolomite., *N. Jb. Miner. Mh.*, **7**, 289–302.
- Dercourt, J. et al., 1986. Geological evolution of the Tethys belt from the Atlantic to the Pamirs since the Lias, *Tectonophysics*, **123**, 241–315.
- De Voogd B., Truffert, C., Chamot-Rooke, N., Huchon, P., Lallemand, S. & Le Pichon, X., 1992. Two-ship deep seismic soundings in the basins of the Eastern Mediterranean Sea (Pasiphae cruise), *Geophys. J. Int.*, **109**, 536–552.
- Duronio, P., Dakshi A. & Bellini E., 1991. Stratigraphy of the offshore Cyrenaica, Libya, in *The Geology of Libya*, Vol. IV, pp. 1589–1620, eds Salem, M.J., Hammuda, O.S. & Eliagoubi, B.A., Elsevier, Amsterdam.
- El-Hawat, A.S. & Abdulsamad, E.O., 2004. *The Geology of Cyrenaica: A Field Seminar*. Special publication, 130pp., Earth Sciences Society of Libya (ESSL), Tripoli.
- Emsis et al., 1996. Proceeding of the Ocean Drilling Program Initial Reports, **160**, College Station, TX.
- Finetti, I., 1982. Structure, stratigraphy and evolution of central Mediterranean, *Boll. Geof. Teor. Appl.*, **24**(96), 247–312.
- Fontaine, J.M. & Cussey, R., 1987. Seismic interpretation of carbonate depositional environments, *AAPG Bull.*, **71**(3), 281–297.
- Garfunkel, Z., 2004. Origin of the Eastern Mediterranean basin: a re-evaluation, *Tectonophysics*, **391**, 11–34.
- GEBCO, 2003, updated: 2008, General bathymetric chart of the Oceans.
- Guiraud, R. & Bosworth, W., 1999. Phanerozoic geodynamic evolution of northeastern Africa and the northwestern Arabian platform, *Tectonophysics*, **315**, 73–108.
- Guiraud R., Bosworth W., Thierry J. & Delplanque A., 2005. Phanerozoic geological evolution of Northern and Central Africa: an overview, *J. Afric. Earth. Sci.*, **43**, 83–143.
- Gumati, Y.D. & Nairn, A.E.M., 1991. Tectonic subsidence of the Sirte Basin, Libya. *J. Petrol. Geol.*, **14**, 93–102.
- Gumati, Y.D., Kanes, W.H. & Schamel, S., 1996. An evaluation of the hydrocarbon potential of the sedimentary basins of Libya, *J. Petrol. Geol.*, **19**, 95–112.
- Hamed, S. & El Hawat 2005. Dynamic stratigraphy of Cyrenaica: an implication of the Mediterranean Tectonics, in: *Proceedings of the Cyrenaica Workshop*, Bremen, 2005.
- Homewood, P.W., Mauriaud, P. & Lafont, F., 2002. Best practices in sequence stratigraphy for explorationists and reservoir engineers, *Bull. Centre Rech. Elf Explor. Prod. Mem.*, **25**, 81pp.
- Hsü, K.J., Cita, M.B. & Ryan W.B.F., 1973. The origin of the Mediterranean evaporates, in *Initial Reports of the Deep Sea Drilling Project*, Vol. 13, pp. 1203–1231, eds Ryan, W.B.F., et al., US Govt. Printing Office, Washington DC.
- Huguen, C. & Mascle, J., 2005. La ride méditerranéenne: apport de la cartographie multifaisceaux à l'analyse morphologique d'un prisme d'accrétion-collision, *Géomorphologie: relief, processus, environnement*, **2**, 91–104.
- Huguen, C., Mascle, J., Chaumillon, E., Woodside, J.M., Benkheilil, J., Kopf, A. & Wolkonskaia, A., 2001. Deformation styles of the eastern Mediterranean Ridge and surroundings from combined swath mapping and seismic reflection profiling, *Tectonophysics*, **343**, 21–47.
- Huguen, C., Mascle, J., Woodside, J.M., Zitter, T. & Foucher, J.P., 2005. Mud volcanoes and mud domes of central Mediterranean Ridge: near-bottom and *in situ* observations, *Deep Sea Res. Part I: Oceanographic Res. Papers*, **52**, 1911–1931.
- Janssen, M.E., Stephenson, R.A. & Cloetingh, S., 1995. Temporal and spatial correlations between changes in plate motions and the evolutions of rifted basins in Africa, *Geol. Soc. Am. Bull.*, **107**(11), 1317–1332.
- Kempler, D. & Garfunkel, Z., 1994. Structures and kinematics in the north-eastern Mediterranean: a study of an unusual plate boundary, *Tectonophysics*, **234**, 19–32.
- Kendall, C.G.St.C., Lake, P., Weathers III, Dallon H., Lakshmi, Venkat, Althausen, John & Alsharhan, Abdulrahman S., 2003. Evidence of rain shadow in the geologic record: repeated evaporite accumulation at extensional and compressional plate margins, in: *The Desertification in the Third Millennium*, pp. 45–52, eds Sharhan, A.S., Wood, W.W., Goudie, A.S., Fowler, A. and Abdellatif, E.M., Swets & Zeitlinger Publishers (Balkema), Lisse, The Netherlands.
- Kopf, A., Robertson, A.H.F., Clennell, M.B. & Flecker, R., 1998. Mechanisms of mud extrusion on the Mediterranean Ridge Accretionary Complex, *Geo-Mar. Lett.*, **18**, 97–114, doi:10.1007/s003670050058.
- Lambare, G., Virieux, J., Madariaga, R. & Jin, S., 1992. Iterative asymptotic inversion in the acoustic approximation, *Geophysics*, **57**, 1138–1154.
- Le Pichon, X., Bergerat, F. & Roulet, M.J., 1988. Plate kinematics and tectonics leading to the alpine belt formation. A new analysis, *Geol. Soc. Am. Special Paper*, **218**, 111–131.
- Makhlouf, I.M., 2006. Late Triassic Early Jurassic Neotethyan evolution at Northern Gondwana (Jordan and Libya), Mediterranean Region, *Geologica Acta*, **4** (3), 371–376.
- Mascle, J. & Chaumillon, E., 1998. An overview of Mediterranean ridge collisional accretionary complex as deduced from multichannel seismic data, *Mar. Geol.*, **18**, 81–89.
- McKenzie, D.P., 1970. Plate tectonics of the Mediterranean region, *Nature*, **226**, 239–243.
- Mitchum, R.M., Jr, 1977. Seismic stratigraphy and global changes of sea level. Part 11: glossary of terms used in seismic stratigraphy, *Am. Assoc. Petrol. Geol. Mem.*, **26**, 205–212.

- Muños, J.A. Martinez, A. & Verges, J., 1985. Thrust sequences in the eastern Spanish Pyrenees, *J. Struct. Geol.*, **8**(3–4), 399–405.
- Ori, G.G. & Friend, P.F., 1984. Sedimentary basins formed and carried piggyback on active thrust sheets, *Geology*, **12**, 475–478.
- Piedilato, S. & Prosser, G., 2005. Thrust sequences and evolution of the external sector of a fold and thrust belt: an example from the southern Apennines (Italy), *J. Geodyn.*, **39**, 386–402.
- Posamentier, H.W. & Allen, G.P., 1999. Siliciclastic sequence stratigraphy: concepts and applications, in *SEPM Concepts in Sedimentology and Paleontology*, 7, Society of Economic Paleontologists and Mineralogists, Tulsa, OK, 204pp., doi:10.1017/S0016756801216082.
- Posamentier, H.W., Jervey, M.T. & Vail, P.R., 1988. Eustatic controls on clastic deposition: I. Conceptual framework, in: *Sea Level Changes—An Integrated Approach*, SEPM Special Pub. Vol. 42, pp. 110–124, eds Wilgus, C.K., Hastings, B.S., Kendall, C.G.St.C., Posamentier, H.W., Ross, C.A. & Van Wagoner, J.C., Society of Economic Paleontologists and Mineralogists, Tulsa, OK.
- Ricou, L.E., Dercourt, J., Geysant, J., Grandjacquet, C., Lepvrier, C. & Biju-Duval, B., 1986. Geological constraints on the alpine evolution of the Mediterranean Tethys, *Tectonophysics*, **123**, 235–256.
- Robertson, A.H.F. & Shipboard Scientific Party 1996. Mud volcanism on the Mediterranean Ridge: initial results of Ocean Drilling Program, Leg 160, *Geology*, **24**, 239–242.
- Röhlich, P., 1980. Tectonic development of Jabal al Akhdar, in: *The Geology of Libya*, Vol. 3, pp 923–931, eds Salem, M.J. and Buserwil, M.T., Academic Press, London.
- Rosenbaum, G., Lister, G.S. & Duboz, C., 2002. Relative motions of Africa, Iberia and Europe during Alpine orogeny, *Tectonophysics*, **359**, 117–129.
- Rusk, D.C., 2001. Libya: petroleum potential of the underexplored basins—a twenty-first-century challenge, *Am. Assoc. Petrol. Geol. Mem.*, **74**, 429–452.
- Sarg, J.F., 2002. The Saline Giants—Sedimentary Extremes that are part of the Earth's Rhythms. AAPG Conf. Abstract, session 1704.
- Sarg, J.F., Markello, J.R. & Weber, L.J., 1999. The second-order cycle, carbonate-platform growth, and reservoir, source, and trap prediction, in *Advances in Carbonate Sequence Stratigraphy: Application to Reservoirs, Outcrops and Models*, SEPM Special Pub. Vol. 63, pp 11–34, Society of Economic Paleontologists and Mineralogists, Tulsa, OK.
- Sengör, A.M.C., 1979. Mid-Mesozoic closure of Permo-Triassic Tethys and its implications, *Nature*, **279**, 590–593.
- Shegawi, O.M., 1992. *Onshore and Offshore Basins of Northeast Libya: Their Origin and Hydrocarbon Potential*. University of South Carolina, Columbia, SC, 607pp.
- Stampfli, G.M. & Borel, G., 2000. *Mesozoic Plate Tectonic of the Western Tethyan Domain*, 6th Int. Cretaceous Symp, Vienna.
- Tawadros, E.E., 2001. *Geology of Egypt and Libya*. Taylor & Francis, London, 468pp.
- Tay, P.L., Lonergan, L., Warner, M. & Jones, K.A., 2002. Seismic investigation of thick evaporate deposits on the central and inner unit of the Mediterranean Ridge accretionary complex, *Mar. Geol.*, **186**, 167–194.
- Thierry, P., Operto, S. & Lambaré, G., 1999. Fast 2-D ray+born migration/inversion in complex media, *Geophysics*, **64**, 162–181.
- Vail, P.R., 1987. Seismic stratigraphy interpretation using sequence stratigraphy. Part I: seismic stratigraphy interpretation procedure. in: *Atlas of Seismic Stratigraphy*, AAPG Studies in Geology No. 27, pp. 1–10, ed. Bally, A.W., American Association of Petroleum Geologists, Tulsa, OK.
- Vail, P.R., Todd, R.G. & Sangree, J.B., 1977. Seismic Stratigraphy and Global Changes of Sea Level. Part 5: Chronostratigraphic Significance of Seismic Reflections: Section 2, Application of Seismic Reflection Configuration to Stratigraphic Interpretation, *AAPG Special Volumes*, **26**, 99–116.
- Van Wagoner, J.C., Posamentier, H.W., Mitchum, R.M., Vail, P.R., Sarg, J.F., Loutit, T.S. & Hardenbol, J., 1988. An overview of sequence stratigraphy and key definitions. in: *Sea Level Changes—An Integrated Approach*, SEPM Special Pub. Vol. 42, pp. 39–45, eds Wilgus, C.K., Hastings, B.S., Kendall, C.G.St.C., Posamentier, H.W., Ross, C.A., Van Wagoner, J.C., Society of Economic Paleontologists and Mineralogists, Tulsa, OK.
- Wilson, J.L., 1975. *Carbonate Facies in Geologic History*, Springer Verlag, New York, 471pp.
- Zeigler, P.A., 1988. Evolution of the Arctic-North Atlantic and the Western Tethys, *Am. Assoc. Petrol. Geol. Mem.*, **43**, 198.
- Zitter, T.A.C., 2004. Mud volcanism and fluid migrations in the Eastern Mediterranean neotectonic zones. *PhD thesis*. Vrije University.
- Zoetemeijer, R., Sassi, W., Roure, F. & Cloetingh, S., 1992. Stratigraphy and kinematic modelling of thrust evolution, northern Apennines, Italy, *Geology*, **20**, 1035–1038.
- Zouaghi, T., Bédir, M. & Inoubli, M. H., 2005. 2D seismic interpretation of strike-slip faulting, salt tectonics, and Cretaceous unconformities, Atlas Mountains, central Tunisia, *J. Africa Earth Sci.*, **43**, 464–486.