



HAL
open science

Early and treatment-related deaths in childhood acute myeloid leukaemia in the Nordic countries: 1984-2003

Lene Molgaard-Hansen, Merja Möttönen, Heidi Glosli, Guðmundur K. Jónmundsson, Jonas Abrahamsson, Henrik Hasle

► **To cite this version:**

Lene Molgaard-Hansen, Merja Möttönen, Heidi Glosli, Guðmundur K. Jónmundsson, Jonas Abrahamsson, et al.. Early and treatment-related deaths in childhood acute myeloid leukaemia in the Nordic countries: 1984-2003. *British Journal of Haematology*, 2010, 151 (5), pp.447. 10.1111/j.1365-2141.2010.08389.x . hal-00586990

HAL Id: hal-00586990

<https://hal.science/hal-00586990>

Submitted on 19 Apr 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Early and treatment-related deaths in childhood acute myeloid leukaemia in the Nordic countries: 1984-2003

Journal:	<i>British Journal of Haematology</i>
Manuscript ID:	BJH-2010-00926.R1
Manuscript Type:	Ordinary Papers
Date Submitted by the Author:	19-Jul-2010
Complete List of Authors:	Molgaard-Hansen, Lene; Aarhus University Hospital Skejby, Department of Paediatrics Möttönen, Merja; University Hospital, Oulu, Department of Paediatrics Glosli, Heidi; Oslo University Hospital, Rikshospitalet, Department of Paediatrics Jónmundsson, Guðmundur; Landspítalinn University Hospital, Reykjavik, Department of Paediatrics Abrahamsson, Jonas; The Queen Silvia Children's Hospital, Department of Paediatric Oncology Hasle, Henrik; Aarhus University Hospital Skejby, Department of Paediatrics
Key Words:	ACUTE MYELOID LEUKAEMIA, CHILDREN, EARLY DEATHS, TREATMENT-RELATED MORTALITY , INFECTION

1
2
3
4
5
6
7
8
9

Early and treatment-related deaths in childhood acute myeloid leukaemia in the Nordic countries: 1984-2003

10
11
12
13
14
15
16
17
18
19

Lene Molgaard-Hansen¹, Merja Möttönen², Heidi Glosli³, Guðmundur K. Jónmundsson⁴, Jonas Abrahamsson⁵, Henrik Hasle¹

20
21
22
23

On behalf of the Nordic Society of Paediatric Haematology and Oncology (NOPHO)

24
25
26
27

1) Department of Paediatrics, Aarhus University Hospital Skejby, Aarhus, Denmark.

28
29
30
31
32
33
34
35
36
37

2) Department of Paediatrics, University Hospital, Oulu, Finland. 3) Department of Paediatrics, Oslo University Hospital, Rikshospitalet, Oslo, Norway. 4) Department of Paediatrics, Landspítalinn University Hospital, Reykjavik, Iceland. 5) Department of Paediatric Oncology, The Queen Silvia Children's Hospital, Gothenburg, Sweden.

38
39
40
41

Disclaimers: none

42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements of research support (alphabetically):

The A. P. Møller Foundation for the Advancement of Medical Science. The Aarhus University Hospital Research Initiative Foundation. The Aase and Ejnar Danielsen Foundation. The Anders Hasselbalch Foundation. The Bent Bøgh and wife Inge Bøgh Foundation. The Carl J. Beckers Foundation. The Dagmar Marshall Foundation. The Danish Cancer Society. The Danish Childhood Cancer Foundation. The Danish Graduate School in Clinical Oncology. The Eva and Henry Frænkel Foundation. The Frode V. Nyegaard and Wife Foundation. The Johannes Fogh-Nielsen and wife Ella Fogh-Nielsen Grant. **The Karen Elise Jensen Foundation.**

Early and treatment-related deaths in childhood AML

2

1
2
3
4 The Kurt Bønnelycke and wife Grethe Bønnelycke Foundation. The M. Brogaard and
5
6 Wife Foundation. The Max and Anna Friedmann Grant. The Meta and Haakon
7
8 Bagger Foundation. The Oticon Foundation. The Otto Christensen Foundation. The
9
10 Simon Fougner Hartmann and Family Foundation. The Sophus Jacobsen and wife
11
12 Astrid Jacobsen Foundation. The Thora and Viggo Grove Grant. The University of
13
14 Aarhus.
15
16
17
18
19
20

21 Correspondence:

22
23 Lene Molgaard-Hansen. Department of Paediatrics, Aarhus University Hospital
24
25 Skejby, DK-8200 Aarhus, Denmark. Telephone: +45-20959325. Fax: +45-89496023.
26
27 E-mail: lene.molgaard@dadlnet.dk
28
29
30
31
32

33 Presentation of the study: none

34
35
36
37 Conflict of interest: none
38
39
40
41

42 Word count summary: **175**

43
44
45 Word count text: **3861**
46
47

48
49 References: **39**
50
51

52 Figures: 1
53
54

55 Tables: **8**
56
57
58
59
60

Summary

Despite major improvements in the cure rate of childhood acute myeloid leukaemia (AML), 5-15% of patients still die from treatment-related complications. In a historical prospective cohort study, we analyzed frequency, clinical features and risk factors for early deaths (ED) and treatment-related deaths (TRD) in 525 children **included in the Nordic protocols NOPHO-AML-84, -88 and -93.**

Seventy patients (13%) died before starting treatment or from treatment-related complications. The death rate rose from 11% in NOPHO-AML-84 to 29% in -88, but then fell to 8% in -93. Sixteen patients (3%) died within the first two weeks, mainly from bleeding or leukostasis. Hyperleukocytosis, age <2 or ≥ 10 years were risk factors. **After day 15, 10% of patients died from treatment-related complications with infection as the main cause of death. Risk factors were age <2 or ≥ 10 years and treatment according to the NOPHO-AML-88 protocol.**

The number of EDs and TRDs in AML is high. Therefore optimal antifungal prophylaxis is essential, and studies on the benefit of antibacterial prophylaxis and individual risk factors for ED and TRD are needed.

Keywords

Acute myeloid leukaemia, children, early deaths, treatment-related mortality, infection

Introduction

The last two decades have seen a considerable improvement in the prognosis for children with acute myeloid leukaemia (AML) owing to the introduction of more intensive chemotherapy, better risk-group stratification and improved supportive care.

The overall survival rate in childhood AML has reached 60-65% (Kaspers & Creutzig, 2005). Most deaths are due to progressive disease, but 5-15% die from toxic effects of treatment (Riley *et al*, 1999; Creutzig *et al*, 2004; Rubnitz *et al*, 2004; Slats *et al*, 2005). Further improvement in AML survival rates will probably require introduction of individualized therapy where targeted and more leukaemia-specific drugs are used in an effort to arrest progressive disease in combination with better supportive care to prevent early deaths (ED) and treatment-related deaths (TRD).

The aim of the present study was to identify the frequency, the clinical features and possible risk factors for EDs and TRDs in childhood AML in the Nordic countries.

Patients and methods

Eligibility and methods

Since July 1984, all children diagnosed with AML in the Nordic countries (Denmark, Finland, Iceland, Norway and Sweden) have been registered and treated on common protocols conducted by the NOPHO-AML Study Group (Nordic Society of Paediatric Haematology and Oncology). **The study** is population-based for patients younger than 15 years of age, whereas 15-18-years-old patients are enrolled according to local practice. Patients diagnosed from 1 July 1984 to 31 December 2003 were identified in the database and included in the present historical prospective cohort study. Excluded were patients treated according to other protocols than NOPHO-AML-84/88/93 or pre-treated with cytostatic drugs for more than 14 days, patients with myeloid leukaemia of Down syndrome, Fanconi anaemia, Kostmann syndrome, extramedullary myeloid tumour without significant bone marrow involvement, preceding myelodysplastic syndrome or therapy-related AML. A total of 525 patients fulfilled the inclusion criteria. Among these 489 were younger than 15 years at the time of diagnosis and 36 were 15-18 years old. The patients were diagnosed in 21 hospitals in Denmark (n=114), Finland (n=97), Iceland (n=6), Norway (n=106) and Sweden (n=202). The allogeneic transplants were performed in eight centres.

The Nordic AML database contains information about the time and cause of death, and 10 children dying before starting treatment, and **78** dying from treatment-related complications were identified. **Information about patients registered as dying from treatment-related complications in second or third complete remission is not reported. Additional data concerning the remaining 70 patients were collected by reviewing each patient's file using a specific registration form. The**

1
2
3
4 files of 29 patients registered as non-responders were reviewed to verify the
5
6 correctness of the data. Leukaemia-related data such as white blood cell count
7
8 (WBC), platelet count and presence of CNS leukaemia at diagnosis, FAB-type and
9
10 cytogenetics were collected from the database. **Performance status at diagnosis**
11
12 **was not available.**
13
14
15
16
17

18 **NOPHO-AML-84/88/93 treatment**

19
20 The treatment regimens of the NOPHO-AML trials are shown in Figure I and the
21
22 treatment elements including cumulative drug doses are summarised in Tables 1 and
23
24 2. Details concerning patient characteristics and clinical outcome have been reported
25
26 elsewhere (Lie *et al*, 2005).
27
28
29

30 The NOPHO-AML-84 protocol was of moderate intensity with an induction of three
31
32 courses of cytarabine, 6-thioguanine and doxorubicin followed by four consolidation
33
34 courses with high-dose cytarabine. To reduce the frequency of non-response and
35
36 relapse in the NOPHO-AML-88 study, etoposide and mitoxantrone were added in
37
38 selected courses during induction and consolidation, and the interval between
39
40 induction courses was reduced. The event-free survival rose from 29% to 41%, but
41
42 the NOPHO-AML-88 protocol suffered from an unacceptably high frequency of toxic
43
44 deaths during induction and in early consolidation. The NOPHO-AML-93 used the
45
46 same induction blocks as the NOPHO-AML-88, but the dose intensity was modified:
47
48 children with a good response after the first course were given their second course
49
50 only following haematological recovery. Consolidation was identical with NOPHO-
51
52 **AML-88. The event-free survival at five years was 48% in the NOPHO-AML-93**
53
54 **protocol.**
55
56
57
58
59
60

Intrathecal methotrexate was given on the first day of all courses.

1
2
3
4 In the NOPHO-AML-84/88/93 studies, allogeneic haematopoietic stem cell
5 transplantation (HSCT) in first complete remission (CR1) was recommended to all
6 patients with a HLA-matched family donor. For patients without a donor, autologous
7 HSCT was optional in the NOPHO-AML-84, -88 and the first part of -93, and
8 consequently, autologous HSCT was performed on a non-randomised basis decided
9 by the responsible physician. HSCT conditioning regimens and supportive care
10 followed local guidelines.
11
12
13
14
15
16
17
18
19

20
21 Recommendations for supportive care measures in the NOPHO-AML-84/88/93
22 protocols included trimethoprim/sulfamethoxazole prophylaxis for *Pneumocystis*
23 *jiroveci*. Systemic antifungal prophylaxis and granulocyte-colony-stimulating factor
24 (G-CSF) were not recommended. Cyto-reduction prior to chemotherapy was
25 performed at the discretion of the attending physician. All-trans-retinoic acid (ATRA)
26 was recommended to patients with acute promyelocytic leukaemia (APL) as from
27 1999.
28
29
30
31
32
33
34
35
36
37
38
39

40 **Definitions**

41
42 The *diagnosis of AML* was established by morphological analysis of bone marrow
43 (BM) aspirates according to the FAB and WHO classifications.
44
45

46
47 *Central nervous system (CNS) involvement* was defined as more than five leucocytes
48 per microlitre in the cerebrospinal fluid (CSF) in combination with detectable
49 leukaemic cells in the cytospin and/or presence of neurological symptoms such as
50 cranial nerve palsy.
51
52
53
54
55

56
57 *Complete remission (CR)* was defined according to the CALGB criteria (Cheson *et al*,
58 1990) with minor deviations: $\leq 5\%$ leukaemic blasts in the BM with signs of normal
59 haematopoiesis in the BM and with clear signs of regeneration of normal blood cell
60

1
2
3
4 production in the peripheral blood (PB) (platelets $>80 \times 10^9/l$ without transfusions and
5
6 neutrophils $>1.0 \times 10^9/l$), and no leukaemic cells in the PB or in the CSF. Response
7
8 after induction was evaluated close to day 15; a blast cell count in the BM less than
9
10 5% was considered as good response.
11

12
13 *Non-responders (NR)* included all patients surviving the first 42 days of treatment
14
15 without achieving CR. Patients with CR criteria lasting less than four weeks were also
16
17 classified as NR. NRs at day 42 (n=29) were censored in the analysis of TRD.
18

19
20 *Early death (ED)* was defined as a fatal event occurring before or within the first 42
21
22 days of treatment. ED was subclassified into: a) ED₀₋₁₄ before or during and after the
23
24 first therapy course (<15 days of treatment) and b) ED₁₅₋₄₂ in aplasia between days
25
26 15-42 from treatment start. This classification reflects the ED rate and its association
27
28 with initial problems (hyperleukocytosis, leukostasis) or aplasia after induction
29
30 therapy.
31
32

33
34 ***Treatment-related death (TRD)*** was defined as death unrelated to progressive
35
36 disease occurring after day 42 of treatment start in patients who achieved
37
38 complete remission. In this paper, only TRDs in CR1 are reported.
39
40

41
42 ***Hyperleukocytosis*** was defined as WBC $\geq 100 \times 10^9/l$ at diagnosis.
43

44
45 *Infections* were categorized as clinically, radiologically or microbiologically
46
47 documented. Viral disease was registered when both relevant disease symptoms
48
49 and virus were detected. Fungal infections were defined as proven invasive infection
50
51 (positive histopathological findings or positive culture) (Ascioglu *et al*, 2002).
52

53
54 *Cytogenetics*: Patients with t(8;21)(q22;q22), t(9;11)(p22;q23), t(15;17)(q22;q12-21),
55
56 inv16(p13q22) and t(16;16)(p13;q22) were classified as having a favourable
57
58 karyotype (Grimwade *et al*, 1998).
59
60

Statistics

Subgroups of patients were compared by Chi-square or Fischer's exact tests. Kaplan-Meier plots were used for survival analysis, including estimation of cumulative incidence of EDs and TRDs. The main event in the analysis of possible risk factors was composite endpoints consisting of ED₁₅₋₄₂ or TRD. Time was defined as time from day 15 after diagnosis to date of death. Patients who experienced a relapse (n=203) or a second malignant neoplasm (n=2) were censored at the time of this event. Patients without events were censored at the time of last follow-up. Cox proportional hazard regression analysis was performed with the composite endpoints as events and time to this as outcome measure. The following covariates were included: age at diagnosis (<2, 2-9, ≥10 years), WBC at diagnosis (<20, 20-99, ≥100 x 10⁹/l) and protocol (NOPHO-AML-84 vs. NOPHO-AML-88 or NOPHO-AML-93). For patients who underwent HSCT, a time-dependent covariate was defined as zero before, and one after the date of HSCT, and for all other patients, this time-dependent covariate was defined as zero.

A stepwise, multiregression analysis according to Cox was also performed for composite endpoints consisting of **ED₁₅₋₄₂ or TRD** caused by infection. The analysis was adjusted for age at diagnosis and protocol.

The p-values reported are two-tailed, and p-values less than 0.05 were considered significant. **The statistical analyses were performed using Stata Statistical Software, Release 10 (College Station, TX: StataCorp 2007).**

Results

In the Nordic countries, 13% (70/525) of children diagnosed with AML died before starting treatment or from treatment-related complications (Table 3). ED and TRD rose from 11% in the NOPHO-AML-84 protocol to 29% in -88 and then declined to 8% in -93. The frequency of EDs and TRDs in the first half of the protocol period compared with the second half of the protocol was 9% vs. 14% in NOPHO-AML-84 ($p=0.5$), 35% vs. 24% in NOPHO-AML-88 ($p=0.2$), and 9% vs. 8% in NOPHO-AML-93 ($p=0.8$).

The analysis was performed in February 2010 with a median follow-up of 11.8 years (range, 2.8-25.1 years) for patients registered as alive.

Early death day 0-14

The main causes of ED₀₋₁₄ and characteristics of patients appear from Table 4. The death rate was largely unchanged over time with 3%, 6% and 2% in the NOPHO-AML-84, -88 and -93 protocols, respectively ($p=0.4$ and $p=0.05$ when comparing NOPHO-AML-93 with -84 and -88, respectively). Ten (63%) patients died before protocol treatment was begun and three shortly after the first induction course. Among the 16 patients (3%) who died early, 11 (69%) died from intracranial haemorrhage, three (19%) from leukostasis and two (12%) from sepsis. Age was a risk factor for ED₀₋₁₄: 87% ($n=14$) of the patients were <2 years or ≥ 10 years at diagnosis ($p=0.02$). Hyperleukocytosis was also a risk factor. It was present in 75% of the ED₀₋₁₄ patients compared with 15% of the AML cases in general ($p<0.001$). FAB M4 or M5 was not a significant risk factor for ED₀₋₁₄ compared with other FAB-types

($p=0.2$). However, 55% of the patients who died from a cerebral haemorrhage had FAB types M4 or M5, vs. 38% in the AML population ($p=0.1$).

Early death day 15-42

The principal cause of **death** among the 15 patients (3%), who died between days 15-42, was infection (93%) (Table 5). Four (29%) already had infection signs at the time of their last course of chemotherapy, which was the first induction course in one case. One patient died from a gastrointestinal bleeding after the second induction course.

Treatment-related death (> day 42)

TRD occurred in connection with *chemotherapy* in 26 (5%) patients. The main cause of death was infection ($n=22$, 85%) (Table 5). Eight (36%) of these patients had signs of infection (fever and/or respiratory symptoms) when the last course of chemotherapy before death was being administered. One patient with FAB M4 died from a cerebral bleeding. Two patients died from acute respiratory distress syndrome (ARDS). A 13-year-old girl died unexpectedly one day after the first consolidation course. She had been complaining of chest pain, but the autopsy revealed no signs of heart disease or other causes of death.

TRD occurred in ten (7%) of 150 patients *transplanted in CR1*. Eight (80%) of these deaths were caused by a severe infection. One patient died from veno-occlusive disease (VOD) and another from graft-versus-host-disease (GVHD) with bronchiolitis obliterans.

Three patients, all treated with chemotherapy only, died from TRD *more than one year after end of AML treatment*. A 13-year-old female died from cardiac failure 11

1
2
3
4 years after end of treatment. She had received a heart transplant seven years prior to
5
6 her death due to anthracycline-induced cardiomyopathy. A 17-year-old boy had
7
8 candidiasis in the liver and spleen during induction. Later, liver cirrhosis with
9
10 oesophageal varices developed. He died from multiorgan failure seven years after
11
12 end of treatment, following a month with symptoms of pulmonary infection. A seven
13
14 months old girl with AML FAB M4, WBC $170 \times 10^9/l$ and platelets $75 \times 10^9/l$ presented
15
16 at diagnosis with cerebral bleeding and died twelve years later as a consequence of
17
18 the severe brain damage.
19
20
21
22
23
24
25

26 **Risk factors for ED₁₅₋₄₂ or TRD**

27
28 Both univariate and multiple Cox regression analyses of the composite endpoints
29
30 consisting of ED₁₅₋₄₂ or TRD showed that age at diagnosis was a significant
31
32 prognostic factor (Table 6). Eighteen (13%) of 141 patients <2 years of age ($p=0.03$)
33
34 and 23 (14%) of 169 patients ≥ 10 years ($p=0.01$) died from treatment-related
35
36 complications as compared with 13 (6%) of 215 children between 2 and 9 years of
37
38 age. Another risk factor was treatment according to the NOPHO-AML-88 protocol
39
40 ($p=0.01$). The number of ED₁₅₋₄₂ or TRDs was 26 (23%) in the NOPHO-AML-88 as
41
42 compared with 8 (8%) and 20 (6%) in the NOPHO-AML-84 and -93, respectively.
43
44 WBC $20-99 \times 10^9/l$ at diagnosis seemed to be a risk factor as well, though this
45
46 endpoint only reached statistical significance in the multiple regression analysis
47
48 ($p=0.04$). FAB M3 was a risk factor for ED₁₅₋₄₂ or TRD in the multiple Cox regression
49
50 analysis (11% vs. 5%, $p=0.05$), but only few patients had this FAB type. Allogeneic
51
52 HSCT was also associated with a slight excess risk of death from treatment-related
53
54 complications (HR, 2.0; 95% confidence interval (CI), 0.8-4.9; $p=0.1$). Sex, platelet
55
56
57
58
59
60

count and presence of CNS leukaemia at diagnosis, cytogenetics and size of the treating department were not statistically significant risk factors.

Death in FAB M3

Eight (29%) of 28 patients with FAB type M3 suffered ED or TRD, two (25%) within the first two weeks of diagnosis both from a CNS bleeding before starting chemotherapy or ATRA (Table 4). The other six all died from infections. FAB M3 was not a significant risk factor compared with other FAB types for ED₀₋₁₄ (7% vs. 3%, $p=0.2$), but it was a significant risk factor for ED₁₅₋₄₂ or TRD (11% vs. 5%, $p=0.05$).

Cause of death

Death from infection:

Infection was the main reason for ED or TRD ($n=46$; **66%**) (Table 5). After day 15, infection was considered the cause of death in **44 (81%) of 54** deaths. Infection was the cause of death in eight (80%) of ten patients dying from treatment-related causes during HSCT.

In **fourteen** patients, fatal complications caused by infections occurred as a result of bacterial pathogens, in **six** cases a gram-positive isolate and in **eight** a gram-negative isolate. *Pseudomonas aeruginosa* ($n=7$) were predominant among gram-negative pathogens, *Klebsiella* was isolated in a single case. The gram-positive pathogens were *Streptococci* ($n=2$), *Staphylococci* ($n=2$), *Bacillus cereus* ($n=1$) and *Gemella* ($n=1$). **Eleven patients died of fungal infection. This was caused by *Aspergillus* in six patients, by *Candida* in four patients and in one case no specific microorganism could be identified. Nine (82%) of 11 cases of fungal infection were only diagnosed by autopsy.** Viruses causing TRD were CMV ($n=1$)

Early and treatment-related deaths in childhood AML

14

1
2
3
4 and RSV (n=1). In four cases, the infection was polybacterial or –microbial. Fifteen
5
6 patients died of clinically/radiologically documented infections without having a
7
8 pathogen identified.
9

10
11 **Age <2 (p=0.02) or ≥10 years (p=0.01) at diagnosis were the only statistically**
12
13 **significant risk factors for infection-related ED₁₅₋₄₂ or TRD (Table 7).**
14
15
16
17

18
19 Death from cardiac failure:

20
21 Only one patient receiving a cumulative dose of doxorubicin of 150 mg/m² and
22
23 mitoxantrone 60 mg/m² died from cardiac failure. Six patients dying from infection
24
25 had cardiac insufficiency a few days preceding their death but in all cases, cardiac
26
27 failure was considered part of septic shock.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

The three consecutive NOPHO-AML protocols have shown a successive improvement in outcome from a 5-year overall survival of 38% in the NOPHO-AML-84 to 64% in the NOPHO-AML-93 (Lie *et al*, 2005). **This analysis shows that 13% of children diagnosed with AML in the Nordic countries died before starting treatment or from treatment-related complications.** The rates of ED and TRD rose from 11% in NOPHO-AML-84 to 29% in -88, but then fell to 8% in -93. A total of 88% of the patients with ED₀₋₁₄ died due to haemorrhage or leukostasis. The risk factors for death before day 15 were hyperleukocytosis, age <2 years or ≥10 years. After day 15, infection caused 81% of ED₁₅₋₄₂ and TRDs. Risk factors were age (<2 years or ≥10 years at diagnosis), WBC 20-99 x 10⁹/l, FAB M3 and treatment according to the NOPHO-AML-88 protocol.

The number of EDs and TRDs has been estimated in different AML trials as shown in **Table 8** (Tsukimoto *et al*, 2009; Riley *et al*, 1999; Creutzig *et al*, 2004; Rubnitz *et al*, 2004; Slats *et al*, 2005; Ribeiro *et al*, 2005; Gibson *et al*, 2005; Creutzig *et al*, 2006; Kardos *et al*, 2005; Ravindranath *et al*, 2005; Kaspers & Creutzig, 2005; Smith *et al*, 2005; Entz-Werle *et al*, 2005; Perel *et al*, 2005; Dluzniewska *et al*, 2005; Pession *et al*, 2005). ED occurred in 2-10% and TRD in 2-16%. The percentage was higher with a high rate of HSCT. With a 5-year overall survival of 60% or above, our ED and TRD results resemble those reported by others (Riley *et al*, 1999; Creutzig *et al*, 2004). Several groups have reported a decrease in toxic death rates over time without concomitant changes in therapy or even with more intensive therapy. However in the 1982-1997 period, the Dutch childhood oncology group observed that chemotherapy-related mortality rose from 3% to 8% during consolidation, which was perhaps related

1
2
3
4 to the scheduling of intensification blocks (Slats *et al*, 2005). This illustrates the
5
6 delicate balance between minor changes in treatment and the risk of treatment-
7
8 related mortality. **Within the different NOPHO-AML protocols, no significant**
9
10 **reduction in the number of EDs and TRDs occurred over time in contrast to the**
11
12 **learning curve demonstrated in the CCG-2961 study (Lange *et al*, 2008).**

13
14
15
16 The causes of ED and TRD were similar in the different AML studies despite
17
18 variations in supportive care practices (Lehrnbecher *et al*, 2009). **In contrast to the**
19
20 **excess mortality rate due to cardiac causes (standardized mortality rate=5.0) in**
21
22 **the Childhood Cancer Survivor Study (CCSS) (Armstrong *et al*, 2009), the**
23
24 **frequency of cardiac mortality in our population was low.**

25
26
27
28 **Three patients died 7-12 years after end of treatment from late effects of**
29
30 **treatment or complications in relation to initial AML manifestations, which**
31
32 **underlines the importance of continued long-term follow-up.** In the CCSS, 4% of
33
34 272 5-year AML survivors treated without HSCT died from other causes than relapse
35
36 8-26 years after diagnosis (Mulrooney *et al*, 2008).

37
38
39
40 In the AML-BFM-93 and -98 trials, risk factors for death before day 15 were low
41
42 performance status at presentation, initial CNS involvement, hyperleukocytosis and
43
44 FAB M5 (Creutzig *et al*, 2004). After day 15, age above 10 years and FAB M0 were
45
46 risk factors. **Further studies are needed to determine whether pharmacokinetic**
47
48 **differences account for the increased death rate among children <2 years and**
49
50 **≥10 years of age (Palle *et al*, 2006; Palle *et al*, 2009).** We had no data on BMI at
51
52 diagnosis, but the Children's Cancer Group (CCG)-2961 showed that overweight and
53
54 underweight children and adolescents with AML were more likely to experience
55
56 treatment-related mortality (Lange *et al*, 2005). **The high rate of ED and TRD in**
57
58 **APL patients in the NOPHO-AML studies may be due to the low incidence of**
59
60

FAB M3 in Northern Europe and therefore limited experience and late introduction of ATRA.

In the AML-BFM-78 and -83 studies, the risk of ED due to leukostasis and/or haemorrhage was significantly increased in children with FAB M5, hyperleukocytosis, and extramedullary organ involvement (Creutzig *et al*, 1987). These findings are in line with the results from our studies. We also found a higher proportion of EDs from leukostasis/haemorrhage in children <2 years than in children between 2 and 9 years of age. We observed a small decline in the incidence of hyperleukocytosis over time from 18% in the NOPHO-AML-84, 19% in the NOPHO-AML-88 to 13% in the NOPHO-AML-93 protocol. The low number of EDs in the NOPHO-AML-84 (3%) and -93 (2%) precludes any conclusion as to whether the lower frequency of hyperleukocytosis reduced ED as shown in the study from St. Jude Children's Research Hospital (Inaba *et al*, 2008).

Next to relapses, infection remains the most important cause of morbidity and mortality among children with AML. Infections caused **85%** of all EDs and TRDs when deaths before starting treatment were excluded. The infection-related mortality of 9% (46/525) is comparable to that reported in other recent AML studies quoting frequencies of 5-11% (Riley *et al*, 1999; Lehrnbecher *et al*, 2004; Brunet *et al*, 2006; Sung *et al*, 2007). However, it is difficult to compare results and draw conclusions about trends in infection-related mortality due to the use of different definitions and underestimation of the true incidence of these infections. We had a large amount of infection-related deaths where no microorganism could be identified. Like in other studies, a significant number of invasive fungal infections were diagnosed only at autopsy (Groll *et al*, 1996). Risk factors for infection-related deaths in the CCG 2961 trial were underweight, non-white race and age above 16 years (Sung *et al*, 2007). In

1
2
3
4 **our study, age <2 or \geq 10 years at diagnosis were risk factors for death due to**
5 **infection.** The NOPHO-AML-88 used the principle of timed sequential induction
6
7 which caused an unacceptable increase in ED and TRD. This is similar to the results
8
9 from the CCG 2891 trial, where fatal infections were more common during intensive
10
11 than during standard timed induction (Sung *et al*, 2009).
12
13

14
15
16 To allow comparison between cooperative AML groups and to improve our
17
18 understanding of infection outcomes, standard infection definitions should be used
19
20 (Sung *et al*, 2008). To reduce infection-related mortality, it is important to improve
21
22 both prevention and treatment through prophylaxis, early diagnosis and aggressive
23
24 management. **The current NOPHO-AML 2004 protocol includes prophylactic**
25
26 **anti-fungal therapy with fluconazole but no prophylactic G-CSF or anti-bacterial**
27
28 **therapy since there is no solid evidence that these measures reduce mortality**
29
30 **in children with AML. However, a recent study from St. Jude Children's**
31
32 **Research Hospital reduced the odds of bacterial sepsis by 93% using**
33
34 **intravenous vancomycin with oral ciprofloxacin or a cephalosporin (Kurt *et al*,**
35
36 **2008). A relatively small number of patients were studied and larger multi-**
37
38 **institutional studies are needed before recommendations on these**
39
40 **prophylactic antibiotic regimens can be made.** The development of targeted and
41
42 more leukaemia-specific drugs may reduce the toxic effect on normal cells and
43
44 thereby reduce infection-related morbidity. Identification of possible immunogenetic
45
46 risk profiles at the start of therapy could potentially be helpful for risk-adapted and
47
48 individualized supportive care (Hartel *et al*, 2007). Initial data have demonstrated that
49
50 IL-6 and chitotriosidase polymorphisms may modify infection risk (Lehrnbecher *et al*,
51
52 2005). In a study from the CCG, polymorphisms in the gene encoding cytidin
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

deaminase were associated with an increased risk of treatment-related mortality with cytarabine-based therapy for AML (Bhatla *et al*, 2009).

Our study has some limitations. AML is rare in children and the relatively low number of patients therefore hampers statistical analysis of risk factors at subgroup level. Limited information in the medical records sometimes made it difficult to categorize the cause of death. In some cases, it was difficult to assess whether the patient was in complete remission, and to determine whether the cause of death was treatment- or disease-related.

The number of EDs and TRDs in childhood AML in the Nordic countries remains substantial. Further studies should be conducted to explore the pathogenesis behind leukostasis/haemorrhage and prospective studies of the effect of cytoreduction should be performed. **Optimal antifungal prophylaxis is essential, and studies on the benefit of antibacterial prophylaxis and individual risk factors for ED and TRD are needed.**

Acknowledgements

We are grateful to professor M.Vaeth, Department of Biostatistics, University of Aarhus for excellent statistical assistance.

We thank all principal investigators of the NOPHO-AML-84/88/93 studies and the Nordic paediatric oncology centres for providing patient files for this study.

Investigators in Denmark: S.Rosthøj/E.Oestergaard, Aalborg; N.Clausen/H.Hasle/H.Schroeder, Aarhus; K.Schmiegelow/M.Yssing/C.Rechnitzer/B.Lausen, Copenhagen; N.Carlsen/M.Hejl, Odense. *Finland:* L.Hovi/M.A.Siimes/K.Vettenranta/U.Pihkala, Helsinki; M.Perkkiö/P.Riikonen, Kuopio; A.Mäkipernaal/M.Arrola/K.Parto, Tampere; T.Salmi, Turku; M.Lanning/M.Möttönen, Oulu. *Iceland:* G.Jónmundsson/J.Kristinsson, Reykjavik. *Norway:* S.Lie/A.Glomstein/B.Zeller/H.Glosli, Oslo; M.Hellebostad, Ullevaal; J.Helgestad/B.Moström, Bergen; S.Kolmannskog/R.Nygaard/B.Lund, Trondheim; T.Stockland/T.Flaegstad, Tromsø. *Sweden:* L.Mellander/J.Abrahamsson, Gothenburg; M.Behrendtz, Linköping; J.Heldrup/S.Garwicz/L.Hjorth, Lund; G.Gustafsson/S.Soderhall/M.Heyman, Stockholm; E.Forestier/K.Johansson/ P.Sandström, Umeå; A.Kreuger/G.Lönnerholm/B.M.Frost/J.Palle, Uppsala.

Authorship

Lene Molgaard-Hansen participated in the design of the study, provided clinical data, handled the data, performed the statistical analysis and drafted the manuscript.

Henrik Hasle participated in the design of the study, participated in the interpretation of the data and the drafting of the manuscript.

Merja Möttönen, Heidi Glosli, Guðmundur K Jónmundsson and Jonas Abrahamsson provided clinical data in their capacity as national coordinators of this study and participated in the interpretation of the data.

All authors read and approved the final manuscript.

For Peer Review

References

- Armstrong,G.T., Liu,Q., Yasui,Y., Neglia,J.P., Leisenring,W., Robison,L.L., & Mertens,A.C. (2009) Late mortality among 5-year survivors of childhood cancer: a summary from the Childhood Cancer Survivor Study. *Journal of Clinical Oncology*, **27**, 2328-2338.
- Ascioglu,S., Rex,J.H., de,P.B., Bennett,J.E., Bille,J., Crokaert,F., Denning,D.W., Donnelly,J.P., Edwards,J.E., Erjavec,Z., Fiere,D., Lortholary,O., Maertens,J., Meis,J.F., Patterson,T.F., Ritter,J., Selleslag,D., Shah,P.M., Stevens,D.A., & Walsh,T.J. (2002) Defining opportunistic invasive fungal infections in immunocompromised patients with cancer and hematopoietic stem cell transplants: an international consensus. *Clinical Infectious Diseases*, **34**, 7-14.
- Bhatla,D., Gerbing,R.B., Alonzo,T.A., Conner,H., Ross,J.A., Meshinchi,S., Zhai,X.W., Zamzow,T., Mehta,P.A., Geiger,H., Perentesis,J., & Davies,S.M. (2009) Cytidine deaminase genotype and toxicity of cytosine arabinoside therapy in children with acute myeloid leukemia. *British Journal of Haematology*, **144**, 388-394.
- Brunet,A.S., Ploton,C., Galambrun,C., Pondarre,C., Pages,M.P., Bleyzac,N., Freydiere,A.M., Barbe,G., & Bertrand,Y. (2006) Low incidence of sepsis due to viridans streptococci in a ten-year retrospective study of pediatric acute myeloid leukemia. *Pediatric Blood and Cancer*, **47**, 765-772.
- Cheson,B.D., Cassileth,P.A., Head,D.R., Schiffer,C.A., Bennett,J.M., Bloomfield,C.D., Brunning,R., Gale,R.P., Grever,M.R., Keating,M.J., & . (1990)

1
2
3
4 Report of the National Cancer Institute-sponsored workshop on definitions of
5 diagnosis and response in acute myeloid leukemia. *Journal of Clinical*
6
7 *Oncology*, **8**, 813-819.
8
9

10
11
12 Creutzig,U., Ritter,J., Budde,M., Sutor,A., & Schellong,G. (1987) Early deaths due to
13 hemorrhage and leukostasis in childhood acute myelogenous leukemia.
14
15 Associations with hyperleukocytosis and acute monocytic leukemia. *Cancer*,
16
17 **60**, 3071-3079.
18
19

20
21
22 Creutzig,U., Zimmermann,M., Lehrnbecher,T., Graf,N., Hermann,J., Niemyer,C.M.,
23
24 Reiter,A., Ritter,J., Dworzak,M., Stry,J., & Reinhardt,D. (2006) Less toxicity
25
26 by optimizing chemotherapy, but not by addition of granulocyte colony-
27
28 stimulating factor in children and adolescents with acute myeloid leukemia:
29
30 results of AML-BFM 98. *Journal of Clinical Oncology*, **24**, 4499-4506.
31
32
33

34
35
36 Creutzig,U., Zimmermann,M., Reinhardt,D., Dworzak,M., Stry,J., & Lehrnbecher,T.
37
38 (2004) Early deaths and treatment-related mortality in children undergoing
39
40 therapy for acute myeloid leukemia: analysis of the multicenter clinical trials
41
42 AML-BFM 93 and AML-BFM 98. *Journal of Clinical Oncology*, **22**, 4384-4393.
43
44
45

46
47 Dluzniewska,A., Balwierz,W., Armata,J., Balcerska,A., Chybicka,A., Kowalczyk,J.,
48
49 Matysiak,M., Ochocka,M., Radwanska,U., Rokicka-Milewska,R., Sonta-
50
51 Jakimczyk,D., Wachowiak,J., & Wysocki,M. (2005) Twenty years of Polish
52
53 experience with three consecutive protocols for treatment of childhood acute
54
55 myelogenous leukemia. *Leukemia*, **19**, 2117-2124.
56
57

58
59 Entz-Werle,N., Suci,S., van der Werff ten Bosch, Vilmer,E., Bertrand,Y., Benoit,Y.,
60
Margueritte,G., Plouvier,E., Boutard,P., Vandecruys,E., Ferster,A., Lutz,P.,

Uyttebroeck,A., Hoyoux,C., Thyss,A., Rialland,X., Norton,L., Pages,M.P., Philippe,N., Otten,J., & Behar,C. (2005) Results of 58872 and 58921 trials in acute myeloblastic leukemia and relative value of chemotherapy vs allogeneic bone marrow transplantation in first complete remission: the EORTC Children Leukemia Group report. *Leukemia*, **19**, 2072-2081.

Gibson,B.E., Wheatley,K., Hann,I.M., Stevens,R.F., Webb,D., Hills,R.K., De Graaf,S.S., & Harrison,C.J. (2005) Treatment strategy and long-term results in paediatric patients treated in consecutive UK AML trials. *Leukemia*, **19**, 2130-2138.

Grimwade,D., Walker,H., Oliver,F., Wheatley,K., Harrison,C., Harrison,G., Rees,J., Hann,I., Stevens,R., Burnett,A., & Goldstone,A. (1998) The importance of diagnostic cytogenetics on outcome in AML: analysis of 1,612 patients entered into the MRC AML 10 trial. The Medical Research Council Adult and Children's Leukaemia Working Parties. *Blood*, **92**, 2322-2333.

Groll,A.H., Shah,P.M., Mentzel,C., Schneider,M., Just-Nuebling,G., & Huebner,K. (1996) Trends in the postmortem epidemiology of invasive fungal infections at a university hospital. *Journal of Infection*, **33**, 23-32.

Hartel,C., Deuster,M., Lehrnbecher,T., & Schultz,C. (2007) Current approaches for risk stratification of infectious complications in pediatric oncology. *Pediatric Blood and Cancer*, **49**, 767-773.

Inaba,H., Fan,Y., Pounds,S., Geiger,T.L., Rubnitz,J.E., Ribeiro,R.C., Pui,C.H., & Razzouk,B.I. (2008) Clinical and biologic features and treatment outcome of

1
2
3
4 children with newly diagnosed acute myeloid leukemia and hyperleukocytosis.

5
6
7 *Cancer*, **113**, 522-529.

8
9
10 Kardos,G., Zwaan,C.M., Kaspers,G.J., de-Graaf,S.S., de Bont,E.S., Postma,A.,
11 Bokkerink,J.P., Weening,R.S., van der Does-van den Berg, van Wering,E.R.,
12 Korbijn,C., & Hahlen,K. (2005) Treatment strategy and results in children
13 treated on three Dutch Childhood Oncology Group acute myeloid leukemia
14 trials. *Leukemia*, **19**, 2063-2071.

15
16
17
18
19
20
21
22
23 Kaspers,G.J. & Creutzig,U. (2005) Pediatric acute myeloid leukemia: international
24 progress and future directions. *Leukemia*, **19**, 2025-2029.

25
26
27
28
29 Kurt,B., Flynn,P., Shenep,J.L., Pounds,S., Lensing,S., Ribeiro,R.C., Pui,C.H.,
30 Razzouk,B.I., & Rubnitz,J.E. (2008) Prophylactic antibiotics reduce morbidity
31 due to septicemia during intensive treatment for pediatric acute myeloid
32 leukemia. *Cancer*, **113**, 376-382.

33
34
35
36
37
38
39 Lange,B.J., Gerbing,R.B., Feusner,J., Skolnik,J., Sacks,N., Smith,F.O., &
40 Alonzo,T.A. (2005) Mortality in overweight and underweight children with
41 acute myeloid leukemia. *JAMA*, **293**, 203-211.

42
43
44
45
46
47
48 Lange,B.J., Smith,F.O., Feusner,J., Barnard,D.R., Dinndorf,P., Feig,S.,
49 Heerema,N.A., Arndt,C., Arceci,R.J., Seibel,N., Weiman,M., Dusenbery,K.,
50 Shannon,K., Luna-Fineman,S., Gerbing,R.B., & Alonzo,T.A. (2008) Outcomes
51 in CCG-2961, a children's oncology group phase 3 trial for untreated pediatric
52 acute myeloid leukemia: a report from the children's oncology group. *Blood*,
53
54
55
56
57
58
59 **111**, 1044-1053.
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Lehrnbecher,T., Bernig,T., Hanisch,M., Koehl,U., Behl,M., Reinhardt,D., Creutzig,U., Klingebiel,T., Chanock,S.J., & Schwabe,D. (2005) Common genetic variants in the interleukin-6 and chitotriosidase genes are associated with the risk for serious infection in children undergoing therapy for acute myeloid leukemia. *Leukemia*, **19**, 1745-1750.

Lehrnbecher,T., Ethier,M.C., Zaoutis,T., Creutzig,U., Gamis,A., Reinhardt,D., Aplenc,R., & Sung,L. (2009) International variations in infection supportive care practices for paediatric patients with acute myeloid leukaemia. *British Journal of Haematology*, **147**, 125-128.

Lehrnbecher,T., Varwig,D., Kaiser,J., Reinhardt,D., Klingebiel,T., & Creutzig,U. (2004) Infectious complications in pediatric acute myeloid leukemia: analysis of the prospective multi-institutional clinical trial AML-BFM 93. *Leukemia*, **18**, 72-77.

Lie,S.O., Abrahamsson,J., Clausen,N., Forestier,E., Hasle,H., Hovi,L., Jonmundsson,G., Mellander,L., Siimes,M.A., Yssing,M., Zeller,B., & Gustafsson,G. (2005) Long-term results in children with AML: NOPHO-AML Study Group--report of three consecutive trials. *Leukemia*, **19**, 2090-2100.

Mulrooney,D.A., Dover,D.C., Li,S., Yasui,Y., Ness,K.K., Mertens,A.C., Neglia,J.P., Sklar,C.A., Robison,L.L., & Davies,S.M. (2008) Twenty years of follow-up among survivors of childhood and young adult acute myeloid leukemia: a report from the Childhood Cancer Survivor Study. *Cancer*, **112**, 2071-2079.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Palle,J., Frost,B.M., Gustafsson,G., Hellebostad,M., Kanerva,J., Liliemark,E., Schmiegelow,K., & Lonnerholm,G. (2006) Etoposide pharmacokinetics in children treated for acute myeloid leukemia. *Anticancer Drugs*, **17**, 1087-1094.

Palle,J., Frost,B.M., Petersson,C., Hasle,H., Hellebostad,M., Kanerva,J., Schmiegelow,K., & Lonnerholm,G. (2009) Thioguanine pharmacokinetics in induction therapy of children with acute myeloid leukemia. *Anticancer Drugs*, **20**, 7-14.

Perel,Y., Auvrignon,A., Leblanc,T., Michel,G., Reguerre,Y., Vannier,J.P., Dalle,J.H., Gandemer,V., Schmitt,C., Mechinaud,F., Lejars,O., Piguet,C., Couillaud,G., Pautard,B., Landman-Parker,J., Thuret,I., Aladjidi,N., Baruchel,A., & Leverger,G. (2005) Treatment of childhood acute myeloblastic leukemia: dose intensification improves outcome and maintenance therapy is of no benefit-- multicenter studies of the French LAME (Leucemie Aigue Myeloblastique Enfant) Cooperative Group. *Leukemia*, **19**, 2082-2089.

Pession,A., Rondelli,R., Basso,G., Rizzari,C., Testi,A.M., Fagioli,F., De,S.P., & Locatelli,F. (2005) Treatment and long-term results in children with acute myeloid leukaemia treated according to the AIEOP AML protocols. *Leukemia*, **19**, 2043-2053.

Ravindranath,Y., Chang,M., Steuber,C.P., Becton,D., Dahl,G., Civin,C., Camitta,B., Carroll,A., Raimondi,S.C., & Weinstein,H.J. (2005) Pediatric Oncology Group (POG) studies of acute myeloid leukemia (AML): a review of four consecutive childhood AML trials conducted between 1981 and 2000. *Leukemia*, **19**, 2101-2116.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Ribeiro,R.C., Razzouk,B.I., Pounds,S., Hijiya,N., Pui,C.H., & Rubnitz,J.E. (2005) Successive clinical trials for childhood acute myeloid leukemia at St Jude Children's Research Hospital, from 1980 to 2000. *Leukemia*, **19**, 2125-2129.

Riley,L.C., Hann,I.M., Wheatley,K., & Stevens,R.F. (1999) Treatment-related deaths during induction and first remission of acute myeloid leukaemia in children treated on the Tenth Medical Research Council acute myeloid leukaemia trial (MRC AML10). The MCR Childhood Leukaemia Working Party. *British Journal of Haematology*, **106**, 436-444.

Rubnitz,J.E., Lensing,S., Zhou,Y., Sandlund,J.T., Razzouk,B.I., Ribeiro,R.C., & Pui,C.H. (2004) Death during induction therapy and first remission of acute leukemia in childhood: the St. Jude experience. *Cancer*, **101**, 1677-1684.

Slats,A.M., Egeler,R.M., van der Does-van den Berg, Korbijn,C., Hahlen,K., Kamps,W.A., Veerman,A.J., & Zwaan,C.M. (2005) Causes of death--other than progressive leukemia--in childhood acute lymphoblastic (ALL) and myeloid leukemia (AML): the Dutch Childhood Oncology Group experience. *Leukemia*, **19**, 537-544.

Smith,F.O., Alonzo,T.A., Gerbing,R.B., Woods,W.G., & Arceci,R.J. (2005) Long-term results of children with acute myeloid leukemia: a report of three consecutive Phase III trials by the Children's Cancer Group: CCG 251, CCG 213 and CCG 2891. *Leukemia*, **19**, 2054-2062.

Sung,L., Aplenc,R., Zaoutis,T., Groll,A.H., Gibson,B., & Lehrnbecher,T. (2008) Infections in pediatric acute myeloid leukemia: Lessons learned and unresolved questions. *Pediatric Blood and Cancer*, **51**, 458-460.

1
2
3
4 Sung,L., Gami,A., Alonzo,T.A., Buxton,A., Britton,K., Deswarte-Wallace,J., &
5
6 Woods,W.G. (2009) Infections and association with different intensity of
7
8 chemotherapy in children with acute myeloid leukemia. *Cancer*, **115**, 1100-
9
10 1108.
11

12
13
14
15 Sung,L., Lange,B.J., Gerbing,R.B., Alonzo,T.A., & Feusner,J. (2007)
16
17 Microbiologically documented infections and infection-related mortality in
18
19 children with acute myeloid leukemia. *Blood*, **110**, 3532-3539.
20
21

22
23 Tsukimoto,I., Tawa,A., Horibe,K., Tabuchi,K., Kigasawa,H., Tsuchida,M., Yabe,H.,
24
25 Nakayama,H., Kudo,K., Kobayashi,R., Hamamoto,K., Imaizumi,M.,
26
27 Morimoto,A., Tsuchiya,S., & Hanada,R. (2009) Risk-stratified therapy and the
28
29 intensive use of cytarabine improves the outcome in childhood acute myeloid
30
31 leukemia: the AML99 trial from the Japanese Childhood AML Cooperative
32
33 Study Group. *Journal of Clinical Oncology*, **27**, 4007-4013.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure legend

Figure I

Flow diagram of the NOPHO-AML-84/88/93 protocols with indication † of early deaths and treatment-related deaths (n=70). The treatment elements are described in Tables 1 and 2.

Among patients who suffered ED day 0-14; ten died before starting treatment, three during the first induction, and three after finishing the first induction course. ED day 15-42 included seven patients dying after the first induction course and eight after the second.

Abbreviations: A, cytarabine; T, 6-thioguanine; Dox, doxorubicin; CR, complete remission; n, number; HA, high-dose cytarabine; E, etoposide; M, mitoxantrone; BM, bone marrow; ED, early death; TRD, treatment-related death.

Table 1. NOPHO-AML-84/88/93 treatment elements

ATDox	Cytarabine 100 mg/m ² /day continuous infusion day 1-4 6-Thioguanine 100 mg/m ² orally every 12 h day 1-4 Doxorubicin 75 mg/m ² 8-hour infusion day 5
HA₂	Cytarabine 2 g/m ² 2-hour infusion every 12 h day 1-3
ATEDox	Cytarabine 200 mg/m ² /day continuous infusion day 1-4 6-Thioguanine 100 mg/m ² orally every 12 h day 1-4 Etoposide 100 mg/m ² /day continuous infusion day 1-4 Doxorubicin 75 mg/m ² 8-hour infusion day 5
AM	Cytarabine 100 mg/m ² /day continuous infusion day 1-5 Mitoxantrone 10 mg/m ² 30-minute infusion day 1-3
HA₁M	Cytarabine 1 g/m ² 2-hour infusion every 12 h day 1-3 Mitoxantrone 10 mg/m ² 30-minute infusion day 3-5
HA₂E	Cytarabine 2 g/m ² 2-hour infusion every 12 h day 1-3 Etoposide 100 mg/m ² /day 1-hour infusion day 2-5
HA₃	Cytarabine 3 g/m ² 2-hour infusion every 12 h day 1-3
CNS therapy	Methotrexate age-adjusted 6-12 mg intrathecal on the first day of all courses
For children less than 2 years of age, drug doses were calculated according to body weight with 1 m ² equalling 30 kg	

Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. NOPHO-AML-84/88/93 cumulative doses of cytarabine, etoposide and anthracyclines

	Cytarabine	Etoposide	Doxorubicin / Mitoxantrone
	(g/m ²)	(mg/m ²)	Cumulative dose of anthracyclines* (mg/m ²)
NOPHO-AML-84	50.4	0	225 / 0 Cum. dose = 225
NOPHO-AML-88	50.1	1600	150 / 60 Cum. dose = 450
NOPHO-AML-93	GR: 49.6 PR: 61.3	GR: 1600 PR: 1600	GR: 150 / 30 PR: 75 / 60 Cum. dose GR = 300, PR = 375

Abbreviations: GR, good responders; PR, poor responders

* Calculated applying the following conversion factors to daunorubicin-equivalents: doxorubicin 1 x, mitoxantrone 5 x

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Number of early deaths, treatment-related deaths, non-responders and relapses in NOPHO-AML-84/88/93 protocols

Status	Period	NOPHO AML-84		NOPHO AML-88		NOPHO AML-93		Total	
		No. of patients	%	No. of patients	%	No. of patients	%	No. of patients	%
No. of patients		96		114		315		525	
Early death (day 0 – 42)	ED day 0 - 14	3	3	7	6	6	2	16	3
	ED day 15 – 42	3	3	9	8	3	1	15	3
Treatment-related death (> day 42)	During chemotherapy:								
	1) Induction	2	2	8	7	5	2	15	3
	2) Consolidation	1	1	6	5	4	1	11	2
	Transplant-related	2	2	1	1	7	2	10	2
	> 1 year after end of treatment	0	0	2	2	1	0.3	3	1
Total ED/TRD		11	11	33	29	26	8	70	13
Death without CR (non-responders)	NR > day 42	14	15	1	1	14	4	29	6
Relapse (cumulative incidence)		44	46	39	34	120	38	203	39

Abbreviations: ED, early death; TRD, treatment-related death; CR, complete remission; NR, non-responders

Table 4. Characteristics of patients who suffered early death day 0 - 14 in NOPHO-AML-84/88/93 protocols

Patient no.	Survival after diagnosis (days)	Age at diagnosis (months/years)	Sex	WBC at diagnosis ($\times 10^9/l$)	Platelets at diagnosis ($\times 10^9/l$)	Cytogenetics	FAB	Treatment	Causes of death
1	0	9 y	F	550	69	Normal	M1	No	Bleeding (CNS)
2	0	11 y	M	290	45	Normal	M4	No	Bleeding (CNS)
3	1	0 mo	F	450	20	Other	M4	No chemotherapy. Exchange transfusion	Leukostasis
4	1	15 y	F	41	15	No data	M3	No	Bleeding (CNS)
5	2	11 y	M	338	65	No data	M4	No chemotherapy. Neurosurgery	Bleeding (CNS)
6	2	14 y	M	173	76	t(15;17)(q22;q21)	M3	Started 1st induction	Bleeding (CNS)
7	2	15 y	F	365	12	No data	M1	No (Jehovah's Witness)	Bleeding (CNS)
8	3	0 mo	F	400	No data	No data	M5	No	Leukostasis
9	3	7 mo	F	442	35	Other MLL than t(9;11)	M0	Started 1st induction. Neurosurgery	Bleeding (CNS)
10	3	12 y	M	317	35	Normal	M4	Started 1st induction	Bleeding (CNS)
11	3	14 y	F	245	89	Normal	M5a	Cytoreduction (cytarabine, hydroxy-urea)	Bleeding (CNS)
12	5	8 mo	F	458	158	t(9;11)(p22;q23)	M5a	No chemotherapy. Neurosurgery	Bleeding (CNS)
13	9	8 y	M	247	17	Normal	M1	No (Jehovah's Witness)	Leukostasis
14	11	4 mo	F	10	10	No data	RAEB-T	1st induction. Antibiotics for 3 days before death, no antifungal treatment	Sepsis (Pseudomonas aeruginosa)
15	11	11 y	F	7	108	Normal	M4	1st induction. Antibiotics for 7 days, antifungals for 1 day before death	Sepsis (no positive culture), bleeding (lungs)
16	14	3 mo	M	30	22	No data	M5	1st induction	Bleeding (CNS)

Abbreviations: WBC, white blood count; F, female; M, male

Table 5. Causes of early deaths and treatment-related deaths in NOPHO-AML-84/88/93 protocols

Status	Period	No. of patients	Bleeding / Leukostasis	Infection				Cardiac	Other	Unknown	
				Bacteria	Fungal	Viral	Mixed				Clinically / radiologically documented, no positive culture
Early death (day 0 - 42)	ED day 0 - 14	16	14	1	0	0	0	0	0	0	
	ED day 15 - 42	15	1	3	4	0	1	6	0	0	
Treatment-related death (> day 42)	During chemotherapy:										
	1) Induction	15	0	5	4	1	1	2	0	2	0
	2) Consolidation	11	1	3	1	0	1	4	0	0	1
	Transplant-related	10	0	2	2	1	1	2	0	2	0
	> 1 year after end of treatment	3	0	0	0	0	0	0	1	1	1
Total ED/TRD		70	16	14	11	2	4	15	1	5	2

Abbreviations: ED, early death; TRD, treatment-related death

Early and treatment-related deaths in childhood AML

Table 6. Risk factors for early death day 15 - 42 or treatment-related death in NOPHO-AML-84/88/93 protocols

Risk factors	All patients		ED ₁₅₋₄₂ or TRD		HR (95% CI)	Adjusted HR* (95% CI)
	n=525	%	n=54	%	Simple	Multiple
Sex						
Female	272	52	30	56	1.2 (0.7-1.0)	1.2 (0.7-2.0)
Male	253	48	24	44	1.0	1.0
Age at diagnosis (years)						
<2	141	27	18	33	2.3 (1.1-4.6)	2.3 (1.1-4.7)
2-9	215	41	13	24	1.0	1.0
≥10 (10-14)	169	32	23 (14)	43	2.4 (1.2-4.7)	2.4 (1.2-4.8)
Leukocytes at diagnosis (x10 ⁹ /l)						
<20	285	54	25	46	1.0	1.0
20-99	159	30	22	41	1.7 (0.9-3.0)	1.8 (1.0-3.2)
≥100	81	16	7	13	1.2 (0.5-2.7)	1.0 (0.4-2.3)
Platelets at diagnosis (x10 ⁹ /l)						
<20	108	20	12	22	1.3 (0.6-2.8)	1.4 (0.6-3.0)
20-99	273	52	29	54	1.2 (0.6-2.3)	1.2 (0.6-2.3)
≥100	140	27	13	24	1.0	1.0
No data	4	1	0	-	-	-
CNS leukaemia at diagnosis						
Yes	29	6	4	7	1.6 (0.6-4.4)	1.3 (0.5-3.6)
No	485	92	48	89	1.0	1.0
No data	11	2	2	4	-	-
FAB-type						
M0	19	4	1	2	0.5 (0.1-4.3)	1.0 (0.1-8.3)
M1	81	15	8	15	1.0	1.0
M2	131	25	14	26	1.0 (0.4-2.5)	1.2 (0.5-2.9)
M3	28	5	6	11	2.3 (0.8-6.5)	2.9 (1.0-8.4)
M4	106	20	9	17	0.8 (0.3-2.2)	0.9 (0.4-2.4)
M5	95	18	11	20	1.2 (0.5-2.9)	1.5 (0.6-3.7)
M6	14	3	2	4	1.4 (0.3-6.4)	2.8 (0.6-13.6)
M7	21	4	0	-	-	-
Others / No data	30	6	3	5	1.0 (0.3-3.7)	1.0 (0.3-3.7)
Cytogenetics						
Normal	93	18	14	26	1.4 (0.7-3.1)	1.4 (0.7-3.1)
Cytogenetic favourable†	100	19	12	22	1.0	1.0
Other MLL than t(9;11)	32	6	4	8	1.2 (0.4-3.6)	2.4 (0.7-7.9)
Others	144	27	12	22	0.7 (0.3-1.6)	0.7 (0.3-1.5)
No data	156	30	12	22	-	-
Allogeneic HSCT						
Yes	150	28	10	19	2.1 (0.9-4.9)	2.0 (0.8-4.9)
No	374	71	44	81	1.0	1.0
No data	1	1	0	-	-	-
Protocol						
NOPHO-AML-84	96	18	8	15	1.0	1.0
NOPHO-AML-88	114	22	26	48	2.8 (1.3-6.3)	2.8 (1.3-6.1)
NOPHO-AML-93	315	60	20	37	0.7 (0.3-1.6)	0.7 (0.3-1.6)
Size of treating department						
<2 AML patients per year	287	55	33	61	1.0	1.0
≥2 AML patients per year	238	45	21	39	0.8 (0.5-1.4)	0.7 (0.4-1.3)

Abbreviations: ED₁₅₋₄₂, early death day 15-42; TRD, treatment-related death; HR, hazard ratio; 95% CI, 95% confidence interval; HSCT, haematopoietic stem cell transplantation
* Adjusted for age (2-9 vs. <2 or ≥10 years), leukocytes at diagnosis (<20 vs. 20-99 or ≥100 x 10⁹/l) and protocol (NOPHO-AML-84 vs. NOPHO-AML-88 or NOPHO-AML-93)
† Favourable cytogenetics: t(8;21)(q22;q22), t(9;11)(p22;q23), t(15;17)(q22;q12-21), inv16(p13q22) and t(16;16)(p13;q22)

Table 7. Risk factors for early death day 15 - 42 or treatment-related death caused by infection in NOPHO-AML-84/88/93 protocols

Risk factors	All patients		ED ₁₅₋₄₂ or TRD caused by infection		HR (95% CI)	Adjusted HR* (95% CI)
	n=525	%	n=46	%	Simple	Multiple
Sex						
Female	272	52	26	57	1.2 (0.7-2.2)	1.2 (0.7-2.2)
Male	253	48	20	43	1.0	1.0
Age (years)						
<2	141	27	16	35	2.9 (1.3-6.6)	2.8 (1.2-6.3)
2-9	215	41	9	19	1.0	1.0
≥10 (10-14)	169	32	21 (21)	46	3.1 (1.4-6.9)	3.1 (1.4-6.7)
Leukocytes at diagnosis (x10 ⁹ /l)						
<20	285	54	23	50	1.0	1.0
20-99	159	30	18	39	1.5 (0.8-2.7)	1.6 (0.9-2.9)
≥100	81	16	5	11	0.9 (0.4-2.4)	0.7 (0.3-2.0)
Protocol						
NOPHO-AML-84	96	18	8	17	1.0	1.0
NOPHO-AML-88	114	22	20	44	2.2 (1.0-5.1)	2.1 (0.9-4.7)
NOPHO-AML-93	315	60	18	39	0.6 (0.3-1.5)	0.6 (0.3-1.4)

Abbreviations: ED₁₅₋₄₂, early death day 15-42; TRD, treatment-related death; HR, hazard ratio; 95% CI, 95% confidence interval

* Adjusted for age (2-9 vs. <2 or ≥10 years) and protocol (NOPHO-AML-84 vs. NOPHO-AML-88 or NOPHO-AML-93)

Table 8. Number of early deaths and treatment-related deaths in recent paediatric AML studies from major groups

Study group	Years of enrolment	No. of patients	Age range (years)	Early death rate (%)	TRD rate (%)	Overall survival rate (5 years) (%)
AIEOP-92 (Kaspers et al, 2005 and Pession et al, 2005)	1992-2001	160	0-14	6	7	60
AML-BFM 98 (Creutzig et al, 2004 and 2006)	1998-2002	430	0-18	3	4*	62†
AML-PLLSG 98 (Dluzniewska et al, 2005)	1998-2002	104	0-14	8	10	50
AML99 (Tsukimoto et al, 2009)	2000-2002	240	0-18	2‡	4	76
CCG 2891 (Smith et al, 2005)	1989-1995	750	0-14	4	15	47
DCOG AML-92/94 (Kardos et al, 2005)	1992-1998	78	0-14	10	16	50
EORTC-CLG 58921 (Entz-Werle et al, 2005)	1993-2000	166	0-14	2	6	62
LAME 91 (Perel et al, 2005)	1991-1998	247	0-14	4	6	61
MRC AML 12 (Gibson et al, 2005)	1995-2002	455	0-14	4	6	66
NOPHO-AML-93	1993-2003	292	0-14	5	2	65
NOPHO-AML-93	1993-2003	315	0-18	3	5	64
POG 8821 (Ravindranath et al, 2005)	1988-1993	511	0-14	4	8	42
St. Jude-AML91 (Ribeiro et al, 2005)	1991-1997	62	0-14	3	Not reported	57

Abbreviations: TRD, treatment-related death; AIEOP, Associazione Italiana di Ematologia e Oncologia Pediatrica; AML99, Japanese Leukemia/Lymphoma Study Group; BFM, Berlin-Frankfurt-Münster Group; PLLSG, the Polish Pediatric Leukemia/Lymphoma Study Group; CCG, the Children's Cancer Group; DCOG, Dutch Childhood Oncology Group; EORTC, European Organization of Research and Treatment of Cancer; LAME, Leucémie Aiguë Myéloblastique Enfant; MRC, Medical Research Council; POG, Pediatric Oncology Group

* Without non-responders who are included in the AML-BFM study on early death and treatment-related death

† Is from the paper (Creutzig et al, 2006) which included 473 patients in the period from 1998 to 2003

‡ Patients dying before initiation therapy were excluded (n=3)