

Controversial results of the genetic analysis of a canine distemper vaccine strain

Zoltán Demeter, Elena Alina Palade, Ákos Hornyák, Miklós Rusvai

► To cite this version:

Zoltán Demeter, Elena Alina Palade, Ákos Hornyák, Miklós Rusvai. Controversial results of the genetic analysis of a canine distemper vaccine strain. Veterinary Microbiology, 2010, 142 (3-4), pp.420. 10.1016/j.vetmic.2009.10.017. hal-00587286

HAL Id: hal-00587286 https://hal.science/hal-00587286

Submitted on 20 Apr 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Controversial results of the genetic analysis of a canine distemper vaccine strain

Authors: Zoltán Demeter, Elena Alina Palade, Ákos Hornyák, Miklós Rusvai





Please cite this article as: Demeter, Z., Palade, E.A., Hornyák, Á., Rusvai, M., Controversial results of the genetic analysis of a canine distemper vaccine strain, *Veterinary Microbiology* (2008), doi:10.1016/j.vetmic.2009.10.017

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1	Controversial results of the genetic analysis of a canine distemper vaccine strain
2	
3	Zoltán Demeter ¹ *, Elena Alina Palade ¹ , Ákos Hornyák ² , Miklós Rusvai ¹
4	
5	¹ Department of Pathology and Forensic Veterinary Medicine, Szent István University, Faculty
6	of Veterinary Science, H-1078 Budapest, István u. 2, Hungary
7	² Department of Microbiology and Infectious Diseases, Szent István University, Faculty of
8	Veterinary Science, H-1143 Budapest, Hungária krt. 23-25, Hungary
9	
10	*Corresponding author. Tel: +36-1-478-4178; +36-70-222-3365. Fax: +36-1-478-4284.
11	E-mail address: Demeter.Zoltan@aotk.szie.hu
12	

13 Abstract

Canine distemper (CD) is a highly contagious, often fatal, multisystemic viral disease of 14 15 receptive carnivores. The presence of a PsiI cleavage site on a specific location of the 16 hemagglutinin (H) gene was found to be a hallmark of vaccine strains, thus, a previously 17 published restriction fragment length polymorphism (RFLP) test using PsiI theoretically 18 allows the distinction between all currently used vaccine strains and virulent field strains. The 19 RFLP test was carried out on all brands of CD vaccines available in Hungary. The present 20 work describes the extensive sequencing and phylogenetic study of the strain present in 21 Vanguard (Pfizer Animal Health) vaccines, which following the PsiI based RFLP test reacted 22 as a wild-type strain. Based on the product description provided by the manufacturer, all 23 batches should have contained a virus strain (Snyder Hill) belonging to the group of vaccine 24 strains (America-1). Extensive genetic analysis involving the full nucleic acid sequence of four other genes (N, M, P and F) of the CDV genome revealed that the incriminated virus 25

strain showed a higher level of genetic identity to wild-type strains from the America-2 group than to any of the strains belonging to America-1 group, therefore the vaccine does not contain the virus strain stated by the manufacturer in its product description and has not been containing it since at least 1992.

5

6 **Keywords:** canine distemper, vaccine strain, phylogenetic analysis

7

8 **Running title:** Canine distemper vaccine strain ambiguity

9

10 **1. Introduction**

11 Canine distemper (CD) is a highly contagious viral disease that has a wide range of receptive 12 carnivore animal families, such as the Canidae, Mustelidae, Procyonidae, Felidae and many 13 others. It is caused by the canine distemper virus (CDV) which belongs to the Morbillivirus 14 genus of the Paramyxoviridae virus family (Appel, 1987; Kabakci et al., 2004; Lednicky et 15 al., 2004). Although live attenuated vaccines have been used for many years to control 16 distemper (Chappuis, 1997; Barrett, 1996; Patel & Heldens, 2009), CDV continues to cause 17 outbreaks in receptive domestic carnivores all around the world, in particular young dogs in breeding units and shelter facilities which most often suffer fatal disease (Appel, 1987; 18 19 Blixenkrone-Møller et al., 1993; Barrett, 1996; Gemma et al., 1996; Demeter et al., 2006; 20 Martella et al., 2008). Previous genetic and phylogenetic studies have revealed that the virus strains used in the currently applied vaccines form a distinct group (America-1) from the large 21 22 group of wild-type strains that were reported worldwide in the previous decades (Martella et 23 al., 2007; Demeter, 2009). The most frequently used strains in the manufacturing process of 24 CDV vaccines are the Onderstepoort, Snyder Hill, Rockborn, Lederle, and Convac. The Onderstepoort strain, used worldwide as an attenuated live vaccine, dates back to a disease 25

1 outbreak among North American ranched foxes in the 1930's (Haig, 1956; Martella et al., 2 2007). The Snyder Hill strain was isolated in Ithaca, USA in the 1950's from the brain of a dog. The virus was then passaged in vivo in dogs before being adapted to cell growth in NL-3 4 DKC cells (Brown et al., 1972; Martella et al., 2007). The Rockborn strain is undistinguishable from the Snyder Hill strain (Greene & Appel, 2006), while all the other 5 6 vaccine strains (Lederle, Convac etc.) also belong to the America-1 group. Nowadays it is not 7 clear whether these strains are still circulating in the field, since they have not been detected 8 over the last five decades (Martella et al., 2007).

9 In recent years the number of vaccinated dogs that got infected with CDV and 10 developed characteristic clinical signs of CD has increased (Appel, 1987; Blixenkrone-Møller 11 et al., 1993; Kai et al., 1993; Shin et al., 1995; Gemma et al., 1996; Lan et al., 2006). The 12 molecular analysis of the isolates from the vaccinated yet infected canines demonstrated that 13 the causative agent in these cases was not closely related to any of the currently used vaccine 14 strains (Pardo et al., 2005; Lan et al., 2006; Calderon et al., 2007; Martella et al., 2007). On 15 the other hand, cases when vaccinated animals shed the vaccine strain for a limited period of 16 time were also described (Greene & Appel, 2006). Considering the fact that in case of 17 recently vaccinated animals the virus strain present in the vaccine can interfere with 18 polymerase chain reaction (PCR) based diagnostic tests, leading to diagnostically false 19 positive results, in order to rule out the mentioned misleading factors, a PCR coupled with 20 restriction fragment length polymorphism (RFLP) assay using a PsiI cleavage site of the 21 hemagglutinin (H) gene of the viral genome was designed (Demeter et al., 2007). According 22 to the nucleotide sequences deposited in the publicly accessible GenBank, the presence of 23 PsiI cleavage site is characteristic only for vaccine strains, thus, PsiI RFLP was expected to 24 allow a reliable differentiation of all CDV vaccine strains from the wild-type ones. During the examination of CDV vaccines marketed in Hungary by PsiI RFLP, the vaccine strain present 25

3

in a Vanguard vaccine (Pfizer Animal Health, USA) exhibited wild-type genetic feature
remaining undigested by *Psi*I. This study presents the results of the extensive sequencing and
phylogenetic investigation of batches of this vaccine brand purchased in different countries
(Hungary, Malta, Israel, and USA), as well as batches dating back to 1992 and 1994.

5

6 **2. Materials and methods**

7 2.1. Vaccine batches

8 Products of four different manufacturers were purchased and tested during this study. A total 9 number of ten vaccines were investigated, of which six were of the same brand (vaccines 10 number 1 to 6), but purchased in different countries (Hungary, Israel, Malta and United States 11 of America) or were from batches dated back to 1992, 1994 and 2006 (Table 1). All modified 12 live CD vaccines were stored at the recommended temperature of 4 °C.

13

14 2.2. RNA purification

Vaccines were prepared according to the protocol provided by each manufacturer. Following dilution of the lyophilized component with the solution recommended and supplied together with the vaccine, viral RNA was isolated from each vaccine using QIAamp viral RNA Mini Kit (Qiagen, Germany) according to the manufacturer's instructions. After purification, the RNA samples were stored at -80 °C until the following steps of the investigations.

20

21 2.3. Primers

The primers used in the present study were designed using the Primer Designer 4 software (Scientific & Educational Software, USA). Sequences of different vaccine and wild-type CDV strains were retrieved from GenBank and were aligned to ensure the attaching of the

newly designed primers to the vaccine strains and wild-type strains as well. Primers were
 synthesized by BIOMI Kft. (Gödöllő, Hungary).

3

4 2.4. RT-PCR assays, RFLP and gel electrophoresis

5 Reverse transcription and amplifications were performed in one-step RT-PCR method using 6 the QIAGEN OneStep RT-PCR Kit (Qiagen, Germany), according to the manufacturer's instructions. Reverse transcription was carried out at 50 °C for 30 min. Following an initial 7 8 denaturation at 95 °C for 15 min, the reaction mixture was subjected to 40 cycles of heat 9 denaturation at 94 °C for 45 sec, the corresponding annealing temperature for each primer pair for 1 min, and DNA extension at 72 °C for 2 min, followed by a final extension of 10 min 10 11 at 72 °C. The reactions were performed in a SPRT001 PCR Sprint Thermal Cycler (Hybaid 12 Ltd, UK).

Following RT-PCR 7.5µl of amplicons was electrophoresed at 80 V for 80 min in a 1.2% Tris acetate-EDTA-agarose gel (Merck, Germany), stained with ethidium bromide. The bands were visualized by UV transillumination at 312 nm. Product sizes were determined with reference to 100 bp and 1 kb molecular weight ladders (Fermentas, Lithuania). The *Psi*I (SibEnzyme, Russia) based RFLP was carried out according to the previously described protocol (Demeter et al., 2007).

19

20 2.5. Sequence analysis and phylogeny

DNA was extracted from the gel with the QiaQuick Gel Extraction Kit (Qiagen, Germany).
Fluorescence-based direct sequencings were performed in both directions on the amplicons by
BIOMI Kft. (Gödöllő, Hungary) using an ABI 3100 genetic analyzer (Applied Biosystems,
USA). Nucleotide sequences were identified by BLAST search against gene bank databases
(http://www.ncbi.nlm.nih.gov/BLAST/). The complete sequences of each gene were compiled

1 and aligned to the complete genome sequence of CDV using the Align Plus 4 software 2 (Scientific & Educational Software, USA). The nucleotide sequences of the Vanguard vaccine 3 strain were deposited in the GenBank under the following accession numbers: EF095750 (H 4 gene), EU072198 (F gene), EU072199 (M gene), EU072200 (N gene) and EU072201 (P 5 gene). Phylogenetic trees of the nucleic acid and putative amino acid (aa) sequences were 6 established by using sequence data of the analyzed vaccine strains and the ones retrieved from 7 the GenBank (most of the available complete sequences of the analyzed genes of vaccine and 8 wild-type strains as well). In case of the Vanguard vaccine, each gene of the CDV genome 9 was analyzed, except the highly conserved large polymerase (L) gene. Phylogenetic analyses 10 were performed with the help of the ClustalX program and the robustness of the groupings in 11 the neighbor-joining analysis was assessed with 1000 bootstrap resampling. Results of the 12 phylogenetic analysis were verified using Bayesian and maximum likelihood (ML) 13 approaches (Ronquist, F. & Huelsenbeck, 2003) with the MrBayes 3.1.2 program. In order to 14 identify recombination recombination any the detection program (RDP: 15 http://darwin.uvigo.es/rdp/rdp.html) was used (Martin & Rybicki, 2000).

16

17 **3. Results**

18 3.1. RT-PCRs and RFLP analysis

19 The RT-PCRs resulted amplicons of the expected size (Table 1) in case of every pair of 20 primers used. The RFLP resulted two clearly differentiable bands in case of the currently used 21 vaccine strains at the predicted heights of 294 bp and 816 bp, while the amplicons obtained 22 from field viruses and vaccines 1 to 6 (Table 1) remained undigested (1110 bp).

Different batches of Vanguard vaccines (Pfizer Animal Health, USA) dating back to 1992, 1994 and 2006 were also involved in the study. Furthermore, batches of the same brand were purchased in different countries as well, such as Hungary, Israel, Malta and the United

- States of America (Table 1). The RFLP test unequivocally resulted in undigested amplicons in
 case of all batches from different time and geographical origin.
- 3

4 *3.2. Genetic analyses*

5 To identify the strain present in the Vanguard vaccines, primers were designed to cover more 6 than half of the viral genome. The nucleotide sequences of all CDV genes, except the highly 7 conserved L gene were determined. The genetic analyses of all investigated genes of the 8 Vanguard vaccines (Table 1) have revealed that the virus strain present in all of them is more 9 closely related to different wild-type strains from the America-2 group, than to the one stated 10 by the manufacturer (Snyder Hill strain) or any of the viruses from the group of vaccine 11 strains (Table 2). As a next step the full nucleotide sequences of the highly variable H gene of 12 all vaccine brands currently used in Hungary were included in the study (Table 1). Since the 13 obtained nucleotide sequences of vaccines 1 to 6 turned out to be 100% similar in case of all 14 batches, the subsequent investigations performed on other domains of the virus genome were 15 carried out using only vaccine number 6. According to the investigation of the complete 16 sequence of the H gene, the investigated virus strain was not positioned anywhere close to the 17 Snyder Hill strain, but it showed a significantly higher similarity to a lesser panda isolate 18 (99.45%) and a wild-type strain (98.90%) isolated from a naturally infected dog (strain 25259, 19 accession number: AY964114) in North America (Figure 1). As expected following the initial 20 RFLP, the virus strains present in vaccines 7-10 were positioned in the group of vaccine 21 strains (America-1). The findings of the neighbour-joining method-based phylogenetic 22 analysis were confirmed by the results of the Bayesian and ML methods as well: the 23 Vanguard vaccine strain showed a considerably higher level of identity with wild-type viruses 24 from the America-2 group (data not shown). RDP could not identify any recombination in 25 case of any of the analyzed genes of the Vanguard vaccine strain (data not shown).

1 The genetic analyses of the full nucleic acid sequences of the F, N, M and P genes 2 only confirmed the initial observations of the H gene analysis: in each case the virus strain 3 present in the Vanguard vaccine was not positioned in the group of vaccine strains (America-4 1). Unfortunately the nucleic acid sequence of the Snyder Hill strain is not publicly available for these genomic segments, except for a partial, 335 bps long F gene sequence: AY288312 5 6 and a 390 bps long partial P gene segment: AY286481. Even tough it cannot be considered as 7 reliable as complete gene sequence analyses, the genetic analysis of such small segments 8 yielded results identical with the initial investigations performed on the H gene.

9

10 **4. Discussion**

11 In the present study, we have applied a previously described PsiI-based RFLP test (Demeter 12 et al., 2007) that theoretically allows the differentiation of all wild-type strains with known 13 nucleotide sequence from all vaccine CDV strains. The differentiating RFLP analysis carried 14 out on CD vaccine brands currently used in Hungary revealed that the CDV strain present in 15 the Vanguard (Pfizer Animal Health, USA) vaccine reacted as a wild-type strain. According 16 to the statement of the manufacturer, the European Vanguard products used to immunize dogs 17 against CD contain the Snyder Hill strain. Based on the nucleotide sequence available in the 18 GenBank (complete sequence of the H gene of the Snyder Hill strain: accession number: 19 AF259552), the amplicons of the virus strain present in the Vanguard vaccine should have 20 been digested by the *Psi*I enzyme, but they remained undigested, hence reacted as a wild-type 21 virus. In order to elucidate the dilemma, an extensive genetic and phylogenetic analysis of the 22 incriminated virus strain was conducted. Primer pairs that enabled the amplification of 23 complete genes of the CDV genome were designed and the nucleotide sequences of all CDV 24 genes (H, N, M, P and F), except the highly conserved L gene were determined. For phylogenetic purposes the H gene was chosen, because of its vital role in the infectious 25

1 process (Murphy et al., 1999) this gene is considered as one of the most variable one in the 2 CDV genome (Haas et al., 1997; Harder & Osterhaus, 1997; Iwastsuki et al., 1997; 3 Mochizuki et al., 1999; Uema et al., 2005; Martella et al., 2006, McCarthy et al., 2007), 4 therefore it is profoundly suitable for phylogenetic analysis. H gene analysis further emphasized the previous findings: the investigated virus strain was not positioned anywhere 5 close to the Snyder Hill strain, but turned out to be most closely related to a wild-type virus 6 strain isolated from a lesser panda (AF178039). Similar observations were reported by 7 8 another research group (Pardo et al., 2005), but the investigated segment was too small to 9 obtain relevant results (979 bases). Based on the findings of this research group the virus 10 strain present in the Vanguard vaccine was most closely related to the A75/17 wild-type strain 11 isolated from a dog, but the investigation of the complete sequence of the H gene has revealed 12 that it is more closely related to the lesser panda isolate (99.45 %) and a wild-type strain 13 (98.90 %) isolated in North America from a naturally infected dog (strain 25259, accession 14 number: AY964114) by the same research group (Pardo et al., 2005). The analysis of other 15 genes of CDV has only confirmed the findings of the H gene analysis.

16 In order to extend our investigations in time and geographical respect, Vanguard 17 vaccine vials produced in different time (1992 and 1994) and different geographical origin 18 (Malta, Israel, and USA) were purchased and the same RFLP test was applied to determine 19 whether these vaccines contain the same virus strain as the vaccine purchased in Hungary in 20 2006. The result of the RFLP test and the complete sequencing of the H gene of the virus 21 strains present in all these Vanguard vaccine batches revealed that all of them contain exactly 22 the same virus strain, showing not even one nucleotide difference on this relatively large and 23 highly variable gene. Due to these findings, the vaccine batches dated back to 1992 and 1994 24 and the ones purchased in other countries were not investigated on other segments of the CDV genome, since those results most probably would have only confirmed the previous 25

observations. On the other hand, these findings led to the conclusion that the Vanguard
 vaccines have been containing the analyzed virus strain since at least 1992.

3 There could be several possible explanations for the findings of the present study, such 4 as (1) the incorrect labelling of the Snyder Hill strain deposited in the GenBank (AF259552) contradicted by its positioning among the group of vaccine strains in case of the phylogenetic 5 6 trees constructed based on the nucleic acid sequences of the full length of the H gene (Figure 7 1) and partial segment of the F gene (data not shown); (2) a possible recombination of the 8 seed virus with a wild-type strain – this theory is eliminated by the simultaneous RDP-based 9 analysis of several different genes; (3) due to a taxonomical and technical error (i.e. erroneous 10 laboratory registries) the strain was incorrectly labelled when the vaccine was registered and 11 finally (4) contamination of the seed virus stock used in the vaccine production, as also 12 suggested by Pardo et al. (2005). At this point the last two explanations cannot be 13 demonstrated neither eliminated. The significance of these findings is inconclusive without 14 further studies, as phylogeny can not reveal whether or not a virus is virulent, and there may 15 be little genetic distance between virus passages with dramatic changes in their phenotypes, 16 including virulence. At this point it can only be declared that the vaccine does not contain the 17 virus strain erroneously stated by the manufacturer in its product description and has not been 18 containing it since at least as 1992. Surely the vaccine strain underwent all the required tests 19 before being licensed and proved to be safe to be used in animals, and no claims of inefficacy 20 of the Vanguard vaccines have been reported. There are only a few reports of animals 21 vaccinated with Vanguard vaccines that still developed clinical disease (McInnes et al., 1992; 22 Pardo et al., 2005), but the exact cause of the lack of protection and the exact role of the 23 vaccine strain in the infection was not elucidated, therefore no relevant conclusions can be 24 drawn regarding the efficacy of the Vanguard vaccines, as there are numerous aspects that can 25 interfere with the successful immunization of an animal (Povey, 1986; Greene & Appel,

2006). As no documented concerns regarding the safety and efficacy of the Vanguard strains
 have been raised since at least 1992, it is possible that this recently and apparently
 successfully attenuated wild-type strain could provide, at least theoretically, a better
 protection against the currently circulating wild-type CDV strains.

5 On the other hand, these findings suggest that the previously described *Psi*I-based 6 identification of America-1 (vaccine) strains could lead to diagnostically false results in case 7 of animals recently vaccinated with any Vanguard vaccine, or in case of animals with 8 uncertain vaccination history that could have been vaccinated with a Vanguard vaccine.

9

10 **5. Acknowledgements**

11 The authors would like to thank Pfizer Animal Health division of Pfizer Inc. for the support 12 and assistance during the investigations described in the present study.

13

14 **6. Follow-up**

Following the notification of the research group, the manufacturer has performed the 15 16 sequencing of the vaccine master-seed strain used globally for the production of the vaccine. 17 The work has resulted in a regulatory review of, and a change in, the strain designation in the 18 product literature globally in early 2009. This work demonstrated that the distemper strain 19 used in vaccine production was misidentified back in 1956 when it was first acquired. 20 Documentary evidence of these findings was supplied to the European regulatory agencies, 21 and they have accepted the evidence of the misclassification, and the integrity of the 22 registration study data and subsequent production batches. The strain is now identified in the product descriptions as N-CDV. 23

24

1	6. References				
2					
3	Appel, M.J.G., 1987. Canine distemper virus. In: Horzinek MC, editor. Virus infections of				
4	carnivores, Vol. 1. Amsterdam, Elsevier, pp. 133-159.				
5					
6	Barrett, T., 1999. Morbillivirus infections, with special emphasis on morbilliviruses of				
7	carnivores. Vet. Microbiol. 69, 3-13.				
8					
9	Blixenkrone-Møller, M., Svansson, V., Have, P., Orvell, C., Appel, M., Pedersen, I.R., Dietz,				
10	H.H., Henriksen, P., 1993. Studies on manifestations of canine distemper virus infection in an				
11	urban dog population. Vet. Microbiol. 37, 163-173.				
12					
13	Brown, A.L., Vitamvas, J.A., Merry, D.L. Jr., Beckenhauer, W.H., 1972. Immune response of				
14	pups to modified live-virus canine distemper-measles vaccine. Am. J. Vet. Res. 33, 1447-				
15	1456.				
16					
17	Calderon, M.G., Remorini, P., Periolo, O., Iglesias, M., Mattion, N., La Torre, J., 2007.				
18	Detection by RT-PCR and genetic characterization of canine distemper virus from vaccinated				
19	and non-vaccinated dogs in Argentina. Vet. Microbiol. 125, 341-349.				
20					
21	Chappuis, G., Control of canine distemper. Veterinary Microbiol 1997; 44: 351-358.				
22					
23	Demeter, Z., 2009. Polymerase chain reaction based investigations of canine distemper and				
24	parvovirus strains from Hungary. Doctoral Thesis. Postgraduate School of Veterinary				
25	Science, Budapest, Hungary				

1	
2	Demeter, Z., Palade, E.A., Lakatos, B., Kozma, T., Rusvai, M., 2006. Experiences on the
3	control of endemic distemper at the flaying-house of Budapest. Magyar Állatorvosok Lapja
4	128, 665-673. [in Hungarian]
5	
6	Demeter, Z., Lakatos, B., Palade, E.A., Kozma, T., Forgách, P., Rusvai, M., 2007. Genetic
7	diversity of Hungarian canine distemper virus strains. Vet. Microbiol. 122, 258-269.
8	
9	Gemma, T., Watari, T., Akiyama, K., Miyashita, N., Shin, Y.S., Iwatsuki, K., Kai, C.,
10	Mikami, T., 1996. Epidemiological observations on recent outbreaks of canine distemper in
11	Tokyo area. J. Vet. Med. Sci. 58, 547-550.
12	
13	Greene, C.E., Appel, M.J., 2006. Canine distemper. In: Greene CE, editor: Infectious diseases
14	of the dog and cat, 3 rd edition. Saunders-Elsevier, pp. 25-41.
15	
16	Haas, L., Martens, W., Greiser-Wilke, I., Mamaev, L., Butina, T., Maack, D., Barrett, T.,
17	1997. Analysis of the haemagglutinin gene of current wild-type canine distemper virus
18	isolates from Germany. Virus Res. 48, 165-171.
19	
20	Haig, D.A., 1956. Canine distemper: immunization with avianized virus. Onderstepoort
21	Journal of Veterinary Research 17, 19-53.
22	
23	Han, G.Z., Liu, X.P., Li, S.S., 2008. Cross-species recombination in the haemagglutinin gene
24	of canine distemper virus. Virus Res. 136, 198-201.
25	

- Harder, T.C., Osterhaus, A.D.M.E., 1997. Canine distemper virus A morbillivirus in search
 of new hosts? Trends Microbiol. 5, 120-124.
- 3

Iwatsuki, K., Miyashita, N., Yoshida, E., Gemma, T., Shin, Y.S., Mori, T., Hirayama, N., Kai,
C., Mikami, T., 1997. Molecular and phylogenetic analyses of the haemagglutinin (H)
proteins of field isolates of canine distemper virus from naturally infected dogs. J. Gen. Virol.
78, 373-280.

8

9 Kabakci, N., Yarim, M., Karahan, S., Guvenc, T., Yagci, B.B., Gurcan, I.S., 2004.
10 Immunohistochemical investigation of cerebellum in dogs infected with canine distemper
11 virus. Acta Vet. Hung. 52, 327-337.

12

Kai, C., Ochikubo, F., Okita, M., Iinuma, T., Mikami, T., Kobune, F., Yamanouchi, K., 1993.
Use of B95a cells for isolation of canine distemper virus from clinical cases. J. Vet. Med. Sci.
55, 1067-1070.

16

Lan, N.T., Yamaguchi, R., Inomata, A., Furuya, Y., Uchida, K., Sugano, S., Tateyama, S.,
2006. Comparative analyses of canine distemper viral isolates from clinical cases of canine
distemper in vaccinated dogs. Vet. Microbiol. 115, 32-42.

20

Lednicky, J.A., Dubach, J., Kinsel, M.J., Meehan, T.P., Bocchetta, M., Hungerford, L.L.,
Sarich, N.A., Witecki, K.E., Braid, M.D., Pedrak, C., Houde, C.M., 2004. Genetically distant
American canine distemper virus lineages have recently caused epizootics with somewhat
different characteristics in raccoons living around a large suburban zoo in the USA. Virology
I-2, 1-14.

1	
2	Martella, V., Cirone, F., Elia, G., Lorusso, E., Decaro, N., Campolo, M., Desario, C., Lucente,
3	M.S., Bellacicco, A.L., Blixenkrone-Møller, M., Carmichael, L.E., Buonavoglia, C., 2006.
4	Heterogeneity within the hemagglutinin genes of canine distemper virus (CDV) strains
5	detected in Italy. Vet. Microbiol. 116: 301-309.
6	
7	Martella, V., Elia, G., Buonavoglia, C., 2008. Canine distemper virus. Vet. Clin. N. Am.
8	Small 38, 787-797.
9	
10	Martella, V., Elia, G., Lucente, M.S., Decaro, N., Lorusso, E., Bányai, K., Blixenkrone-
11	Møller, M., Lan, N.T., Yamaguchi, R., Cirone, F., Carmichael, L.E., Buonavoglia, C., 2007.
12	Genotyping canine distemper virus (CDV) by a hemi-nested multiplex PCR provides a rapid
13	approach for investigation of CDV outbreaks. Vet. Microbiol. 122, 32-42.
14	
15	Martin, D., Rybicki, E., 2000. RDP: detection of recombination amongst aligned sequences.
16	Bioinformatics 16, 562-563.
17	
18	McCarthy, A.J., Shaw, M.A., Goodman, S.J., 2007. Pathogen evolution and disease
19	emergence in carnivores. Proc. R. Soc. B. 274, 3165-3174.
20	
21	McInnes, E.F., Burroughs, R.E.J., Duncan, N.M., 1992. Possible vaccine-induced canine
22	distemper in a South American bush dog (Speothos venaticus). J. Wildlife Dis. 28, 614-617.

1	Mochizuki, M., Hashimoto, M., Hagiwara, S., Yoshida, Y., Ishiguro, S., 1999. Genotypes of
2	canine distemper virus determined by analysis of the hemagglutinin genes of recent isolates
3	from dogs in Japan. J. Clin. Microbiol. 37, 2936-2942.
4	
5	Murphy, F., Gibbs, J., Horzinek, M., Studdert, J., 1999. Veterinary Virology, 3rd edition, San
6	Diego, CA, USA, Academic Press
7	
8	Pardo, I.D., Johnson, G.C., Kleiboeker, S.B., 2005. Phylogenetic characterization of canine
9	distemper viruses detected in naturally infected dogs in North America. J. Clin. Microbiol. 43,
10	5009-5017.
11	
12	Patel, J.R., Heldens, J.G.M., 2009. Review of companion animal viral diseases and
13	prophylaxis. Vaccine 27, 491-504.
14	
15	Povey, R.C., 1986. Distemper vaccination of dogs: factors which could cause vaccine failure.
16	Can. Vet. J. 27, 321-323.
17	
18	Ronquist, F., Huelsenbeck, J. P., 2003. MrBayes 3: Bayesian phylogenetic inference under
19	mixed models. Bioinformatics 19, 1572-1574.
20	
21	Shin, Y.S., Mori, T., Okita, M., Gemma, T., Kai, C., Mikami, T., 1995. Detection of canine
22	distemper virus nucleocapsid protein gene in canine peripheral blood mononuclear cells by
23	RT-PCR. J. Vet. Med. Sci. 57, 439-445.
24	

- 1 Uema, M., Ohashi, K., Wakasa, C., Kai, C., 2005. Phylogenetic and restriction fragment
- 2 length polymorphism analyses of hemagglutinin (H) protein of canine distemper virus isolates
- 3 from domestic dogs in Japan. Virus Res. 109, 59-63.
- 4

S

- 1 **Figure 1:** Phylogenetic tree constructed based on a total number of 165 complete nucleotide
- 2 sequences (1824 bp) of the H gene. In case of each strain the accession number along with the
- 3 species and country of origin is presented (when applicable).
- 4 * result of a recombination between an "Asia-1" and an "European wildlife" strain (Han et al.,
- 5 2008)



	Vaccine brand	Vaccine brand Manufacturer		Year of production	Country of origin		
1	Vanguard	Norden Laboratories	NAª	1992	Hungary		
2	Vanguard	Smith Kline NA ^a		1994	Hungary		
3	Vanguard Plus 5	Pfizer Animal Health	L53665	2006	Hungary		
4	Vanguard Plus 5	Pfizer Animal Health	A602088C	2006	Israel		
5	Vanguard Plus 7	Pfizer Animal Health	L60065	2006	Malta		
6	Vanguard Plus 5	Pfizer Animal Health	A602620B	2006	USA		
7	Canigen DH(A2)PPi	Ceva Sante Animale	12TU	2006	Hungary		
8	Eurican	Merial	L246779	2006	Hungary		
9	Nobivac DHP	Intervet	A030C01	2006	Hungary		
10	Canvac 8	Dyntec	040407	2007	Hungary		
not available							

Table 1: Description of the vaccines used in the present study

^a not available

Table 2: The level of identity between the nucleotide and amino acid sequence of the H gene and H protein of the strain present in the Vanguard vaccines and those of the viruses from the group of vaccine strains (shaded) and of the wild-type strains, which showed the highest level of identity and the Snyder Hill strain (bolded). The values in italic represent the level of identity at the amino acid level.

	Vanguard	Panda	Dog	Snyder Hill	Convac	Canigen	Onder.	Eurican	Japan
Vanguard	-	98.84	98.35	91.77	90.95	90.95	91.44	90.13	91.44
Panda	99.45	-	97.86	91.61	91.44	90.78	91.28	89.96	91.28
Dog	98.90	98.57	-	90.78	89.96	89.96	90.46	89.14	90.46
Snyder Hill	92.92	92.70	92.32	-	95.39	95.23	95.72	93.91	95.72
Convac	93.09	93.31	92.48	97.09		96.87	97.53	95.55	97.36
Canigen	93.03	92.81	92.43	96.87	98.46	-	99.01	98.19	99.17
Onder.	93.25	93.03	92.65	97.09	98.68	99.39	-	97.69	99.50
Eurican	92.59	92.37	91.99	96.32	97.91	99.23	98.84	-	97.69
Japan	93.25	93.03	92.65	97.09	98.68	99.45	99.61	98.90	-

Vanguard: EF095750; Snyder Hill: AF259552; Convac: Z35493; Canigen: DQ903854; Onder: AF378705, Eurican: AF014953; Japan: AB212966; Panda (Lesser Panda): AF178039; Dog (25259): AY964114