



**HAL**  
open science

# Lieb-Thirring inequalities for radial magnetic bottles in the disk

Francoise Truc

► **To cite this version:**

Francoise Truc. Lieb-Thirring inequalities for radial magnetic bottles in the disk. 2011. hal-00588164v2

**HAL Id: hal-00588164**

**<https://hal.science/hal-00588164v2>**

Preprint submitted on 25 May 2011 (v2), last revised 9 Sep 2011 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Lieb-Thirring inequalities for radial magnetic bottles in the disk

Françoise Truc\*

May 25, 2011

## Abstract

We consider a Schrödinger operator  $H_A$  with a non vanishing radial magnetic field  $B = dA$  and Dirichlet boundary conditions on the unit disk. We assume growth conditions on  $B$  near the boundary which guarantee in particular the compactness of the resolvent of this operator. Under some assumptions on an additional radial potential  $V$  the operator  $H_B + V$  has a discrete negative spectrum and we prove a Lieb-Thirring inequality on these negative eigenvalues. As a consequence we get an explicit upperbound of the number  $N(H_A, \lambda)$  of eigenvalues of  $H_A$  less than any positive value  $\lambda$ , which depends on the minimum of  $B$  and on the integral of the square of any gauge associated to  $B$ .

## 1 Introduction

Let us consider a particle in a domain  $\Omega$  in  $\mathbb{R}^2$  in the presence of a *magnetic field*  $B$ . We define the 2-dimensional magnetic Laplacian associated to this particle as follows:

Let  $A$  a *magnetic potential* associated to  $B$ ; it means that  $A$  is a smooth real one-form on  $\Omega \subset \mathbb{R}^2$ , given by  $A = \sum_{j=1}^2 a_j dx_j$ , and that the *magnetic field*  $B$  is the two-form  $B = dA$ . We have  $B(x) = \mathbf{b}(x) dx_1 \wedge dx_2$  with  $\mathbf{b}(x) = \partial_1 a_2(x) - \partial_2 a_1(x)$ . The magnetic connection  $\nabla = (\nabla_j)$  is the differential operator defined by

$$\nabla_j = \frac{\partial}{\partial x_j} - ia_j .$$

The 2-dimensional magnetic Schrödinger operator  $H_A$  is defined by

$$H_A = - \sum_{j=1}^2 \nabla_j^2 .$$

---

\*Institut Fourier, francoise.truc@ujf-grenoble.fr Unité mixte de recherche CNRS-UJF 5582, BP 74, 38402-Saint Martin d'Hères Cedex (France)

The magnetic Dirichlet integral  $h_A = \langle H_A \cdot | \cdot \rangle$  is given, for  $u \in C_0^\infty(\Omega)$ , by

$$h_A(u) = \int_{\Omega} \sum_{j=1}^2 |\nabla_j u|^2 |dx|. \quad (1.1)$$

From the previous definitions and the fact that the formal adjoint of  $\nabla_j$  is  $-\nabla_j$ , it is clear that the operator  $H_A$  is symmetric on  $C_0^\infty(\Omega)$ .

In [5] we discuss the essential self-adjointness of this operator. The result in dimension 2 is the following

**Theorem 1.1** *Assume that  $\partial\Omega$  is compact and that  $B(x)$  satisfies near  $\partial\Omega$*

$$\mathbf{b}(x) \geq (D(x))^{-2}, \quad (1.2)$$

*then the Schrödinger operator  $H_A$  is essentially self-adjoint. ( $D(x)$  denotes the distance to the boundary). This still holds true for any gauge  $A'$  such that  $dA' = dA = B$ .*

We have, using Cauchy-Schwarz inequality,

$$|\langle \mathbf{b}(x)u, u \rangle| = |\langle [\nabla_1, \nabla_2]u, u \rangle| \leq \|\nabla_1 u\|^2 + \|\nabla_2 u\|^2 \quad u \in C_o^\infty(\Omega).$$

This gives the well-known lower bound

$$\forall u \in C_o^\infty(\Omega), \quad h_A(u) \geq \left| \int_{\Omega} \mathbf{b}(x)|u|^2 |dx| \right|. \quad (1.3)$$

In this paper, we do not use the conditions (1.2) but we assume nevertheless that  $\mathbf{b}(x)$  grows to infinity as  $x$  approaches the boundary. The operator  $H_A^D$  defined by Friedrich's extension of the quadratic form  $h_A$  has a compact resolvent. We call such an operator a magnetic bottle, by similarity with magnetic bottles in the whole space ([1], [4], [21]). We add a suitable negative potential in order to have a discrete negative spectrum and we address the question of the existence of Lieb-Thirring inequalities ([17], [15]) in the radially symmetric case.

## Aknowledgements

It is a pleasure to thank A. Laptev for fruitful discussions about the question discussed here and Y. Colin de Verdière for useful comments.

## 2 Inequalities: the main results

We consider a magnetic field  $B = \mathbf{b}(x)dx_1 \wedge dx_2$  and a scalar potential  $V$  on the unit disk  $\Omega = \{x = (x_1, x_2) \in \mathbb{R}^2 \mid x_1^2 + x_2^2 = r^2 < 1\}$  so that

- $(H_1)$   $K = \inf_{x \in \Omega} \mathbf{b}(x) > 0$  and  $\mathbf{b}(x) \rightarrow +\infty$  as  $D(x) \rightarrow 0$  (i.e as  $x$  approaches the boundary.)
- $(H_2)$   $B$  is radially symmetric,
- $(H_3)$   $V \in L^1(\Omega)$ ,  $V$  radial and non negative,  $V$  bounded from above .

From assumption  $(H_1)$  and from inequality (1.3) we deduce that for any gauge  $A$  associated to  $B$ , the operator  $H_A$  has a compact resolvent, and assumption  $(H_3)$  entails that the negative spectrum of  $H_A - V$  is discrete, where  $H_A - V$  denotes the operator defined by Friedrich's extension of the quadratic form  $h_A - V$ . Using assumption  $(H_2)$  we can write any vector potential as  $A = A(r)d\theta$ .

The first theorem deals with the number  $N(A, V)$  of negative eigenvalues of the operator  $H_A - V$ . Noticing that we have  $N(A, V) = N(A', V)$  for any gauge  $A'$  so that  $dA' = dA = B$ , we will prove that

**Theorem 2.1** *If assumptions  $(H_1)(H_2)(H_3)$  are verified and if moreover*

$$\mathbf{b}(x) \leq (D(x))^{-\beta}, \quad \beta < \frac{3}{2} \quad (2.1)$$

then

$$N(A, V) \leq \frac{1}{\sqrt{1-\alpha}} \int_{[0,1[} [(\frac{1}{\alpha}-1)A'^2(r)+V(r)]rdr + 2 \int_0^1 [1 + |\log[r\sqrt{K}]|] V(r)rdr$$

for any  $\alpha \in ]0, 1[$  and any radial gauge  $A'$  such that  $dA' = dA = B$ .

The second theorem is a consequence of the first one and provides an explicit upper-bound of the number  $N(H_A, \lambda)$  of the eigenvalues of  $H_A$  less than any positive value  $\lambda$  :

**Theorem 2.2** *If assumptions  $(H_1)$  and  $(H_2)$  are verified and if moreover*

$$\mathbf{b}(x) \leq (D(x))^{-\beta}, \quad \beta < \frac{3}{2}$$

then the number of eigenvalues of the operator  $H_A$  less than  $\lambda$  satisfies the following inequality

$$N(A, \lambda) \leq c_K \lambda + \frac{\lambda}{2\sqrt{1-\alpha}} + \frac{\sqrt{1-\alpha}}{\alpha} \int_{[0,1[} r A'^2(r) dr \quad (2.2)$$

with

- $c_K = \frac{3 - \log K}{2}$  if  $K \leq 1$
- and
- $c_K = \frac{1 + \log K}{2} + \frac{1}{K}$  if  $K > 1$ ,

for any  $\alpha \in ]0, 1[$  and any radial gauge  $A'$  such that  $dA' = dA = B$ .

**Remark 2.3** The minimum of the righthandside is obtained by choosing the radial gauge  $A'$  so that  $\int_0^1 A'(r)rdr = 0$ , and then by taking  $\alpha_\lambda = \frac{-6I + \sqrt{I^2 + 4I\lambda}}{\lambda - 2I}$  with  $I := \int_{]0,1[} r A'^2(r)dr$ .

**Remark 2.4** The inequality of Theorem 2.1 is a "magnetic" version of the Cwikel-Lieb-Rosenblum inequality [6] [16] [18]. CLR inequalities apply to Schrödinger operators in  $\mathbb{R}^d$  for  $d \geq 3$  and  $A \equiv 0$  and are a particular case of Lieb-Thirring inequalities. In the case of dim 2 (and  $A \neq 0$ ), analogues of CLR inequalities can be found in [3] and [14] (for a Aharanov-Bohm magnetic field) and more recently in [12] (for a large class of magnetic fields, in a weighted version). Let us emphasize that the bounds in [12] in the radial case do not depend on the magnetic field and are obtained only for bounded magnetic potentials  $A$ , assumption which we do not need (example 2.5, with  $1 < \beta < 3/2$ ). Moreover the constants in our results are explicit. This implies that our theorems can not be derived from [12].

Concerning general magnetic Lieb-Thirring inequalities we refer to [10] for Lieb-Thirring inequalities for constant magnetic fields in dim 2 and 3 which depend on the field strength, to [7] and [8] for magnetic Lieb-Thirring inequalities related to Pauli operators, and to [9] for links between magnetic and non magnetic Lieb-Thirring inequalities.

**Example 2.5** Consider a magnetic field  $B$  as in the definition (3.1) below, and assume  $b(r) \equiv 1$  and  $\beta \neq 1$ . Then  $c_K = \frac{3}{2}$ , the optimal gauge is  $A' = A_\beta(r)d\theta$  with

$$A_\beta = \frac{1}{1 - \beta} \left[ \frac{1}{(1 - r)^{\beta-1}} - \frac{1}{(2 - \beta)(3 - \beta)} \right], \quad (2.3)$$

and the corresponding minimal value of  $I$  is

$$I_\beta = \int_0^1 A_\beta^2(r)rdr = \frac{1}{(1 - \beta)^2(3 - 2\beta)(4 - 2\beta)}. \quad (2.4)$$

### 3 Proofs

#### 3.1 Proof of Theorem 2.1

Let us introduce the polar coordinates  $x = (r, \theta)$ ,  $r \in \mathbb{R}^+$ ,  $\theta \in [0, 2\pi[$ . Due to assumption (2.1) the magnetic field we have to consider is of the type

$$B(r) = \frac{b(r)}{(1-r)^\beta} dr \wedge d\theta, \quad \text{with} \quad \max_{[0,1[} b(r) \leq M \quad \text{and} \quad \beta < \frac{3}{2}. \quad (3.1)$$

We first prove the following

**Lemma 3.1** *If  $B$  satisfies (3.1), then, for any radial magnetic potential  $A$  associated to  $B$ , there exists a constant  $K$  such that  $A$  writes*

- if  $\beta \neq 1$   $A = A(r)d\theta = \frac{a(r)}{(1-r)^{\beta-1}}d\theta$   
with  $a(r) = K(1-r)^{\beta-1} + \tilde{a}$ ,  $\max_{[0,1[} \tilde{a}(r) \leq C$ .
- if  $\beta = 1$   $A = A(r)d\theta = a(r) \ln(1-r)d\theta$   
with  $a(r) = \frac{K}{\ln(1-r)} + \tilde{a}$ ,  $\max_{[0,1[} \tilde{a}(r) \leq C$ .

In particular  $\int_{[0,1[} r A^2(r) dr < \infty$ .

*Proof.*–

Let us explain the case  $\beta \neq 1$ . The method for the case  $\beta = 1$  is the same. The function  $a(r)$  satisfies the equation

$$(\beta - 1)a(r) - (1-r)a'(r) = b(r).$$

This implies that

$$a(r) = k(r)(1-r)^{\beta-1}, \quad \text{with} \quad k(r) = \int_r^1 b(t)(1-t)^{-\beta} dt + K. \quad (3.2)$$

From (3.1) we get

$|\int_r^1 b(t)(1-t)^{-\beta} dt| \leq M \int_r^1 (1-t)^{-\beta} dt \leq M \frac{(1-r)^{-\beta+1}}{1-\beta}$  and the result follows. □

We come now to the proof of Theorem 2.1, following the method of [13]. The quadratic form associated to  $H_A^D - V$  can be rewritten as

$$h_{A,V}(u) = \int_0^1 \int_0^{2\pi} \left[ \left| \frac{\partial u}{\partial r} \right|^2 - V(r)|u|^2 + r^{-2} \left[ \left[ \frac{\partial}{\partial \theta} - iA(r) \right] u \right]^2 \right] r dr d\theta \quad (3.3)$$

for any  $u \in C_0^\infty([0, 1[ \times [0, 2\pi[)$ . Changing variables  $r = e^t$  and denoting  $w(t, \theta) = u(e^t, \theta)$  for  $t \in ]-\infty, 0[$  and  $\theta \in [0, 2\pi[$  we transfer the form  $h_{A,V}(u)$  to

$$\tilde{h}_{A,V}(w) = \int_{-\infty}^0 \int_0^{2\pi} \left[ \left| \frac{\partial w}{\partial t} \right|^2 - \tilde{V}(t)|w|^2 + \left[ \frac{\partial}{\partial \theta} - if(t) \right] w \right]^2 dt d\theta \quad (3.4)$$

with

$$\tilde{V}(t) = e^{2t}V(e^t), \quad f(t) = e^tA(e^t).$$

By expanding a given function  $w \in C_0^\infty([-\infty, 0[ \times [0, 2\pi[)$  into a Fourier series we obtain that  $\tilde{h}_{A,V}(w) = \oplus_{\ell \in \mathbb{Z}} h_{\ell,V}(w_\ell)$  with

$$h_{\ell,V}(v) = \int_{-\infty}^0 \left| \frac{\partial v}{\partial t} \right|^2 + \left[ (\ell - f(t))^2 - \tilde{V}(t) \right] |v|^2 dt,$$

and  $w_\ell = \Pi_\ell(w)$  where  $\Pi_\ell$  is the projector acting as

$$\Pi_\ell(w)(r, \theta) = \frac{1}{2\pi} \int_0^{2\pi} e^{i\ell(\theta - \theta')} w(r, \theta') d\theta'.$$

We write, for any  $\alpha \in ]0, 1[$  and any  $\ell \in \mathbb{Z}^*$

$$h_{\ell,V}(v) \geq \int_{-\infty}^0 \left| \frac{\partial v}{\partial t} \right|^2 + \left[ \left(1 - \frac{1}{\alpha}\right) f^2(t) - \tilde{V}(t) + (1 - \alpha)\ell^2 \right] |v|^2 dt.$$

Let us denote by  $L_\alpha$  the operator associated via Friedrich's extension to the quadratic form

$$q_\alpha(v) = \int_{-\infty}^0 \left| \frac{\partial v}{\partial t} \right|^2 D_t^2 + \left[ \left(1 - \frac{1}{\alpha}\right) f^2(t) - \tilde{V}(t) \right] |v|^2 dt.$$

$L_\alpha$  and  $q_\alpha$  depend on  $V$  but we skip the reference to  $V$  in notations for the sake of simplicity. Since

$$h_{\ell,V} \geq q_\alpha + (1 - \alpha)\ell^2,$$

the number  $N(h_{\ell,V})$  of negative eigenvalues of  $h_{\ell,V}$  is less than the number of negative eigenvalues of  $L_\alpha + (1 - \alpha)\ell^2$ . So denoting by  $\{-\mu_k^\alpha\}$  the negative eigenvalues of  $L_\alpha$  and by  $I_\ell$  the set  $\{k \in \mathbb{N}; -\mu_k^\alpha + (1 - \alpha)\ell^2 < 0\}$  for any  $\ell \in \mathbb{Z}^*$ , we get

$$N(A, V) \leq \sum_{\ell \in \mathbb{Z}^*} \sum_{k \in I_\ell} 1 + N(h_{0,V}).$$

Noticing that the sum in the righthandside is taken over the  $(\ell, k)$  so that  $0 < |\ell| \leq \frac{1}{\sqrt{1-\alpha}} \sqrt{\mu_k^\alpha}$  we write

$$N(A, V) \leq \frac{2}{\sqrt{1-\alpha}} \sum_{k \in \mathbb{N}} \sqrt{\mu_k^\alpha} + N(h_{0,V}). \quad (3.5)$$

Let us extend the functions  $f$  and  $\tilde{V}$  to  $\mathbb{R}$  by zero and denote respectively by  $f_1$  and  $\tilde{V}_1$  these extensions.

Since  $C_0^\infty([-\infty, 0]) \subset C_0^\infty(\mathbb{R})$ , the negative eigenvalues  $\{-\nu_k^\alpha\}$  of the operator  $L_1^\alpha$  associated via Friedrich's extension to the quadratic form

$$q_1^\alpha(v) = \int_{-\infty}^0 \left| \frac{\partial v}{\partial t} \right|^2 D_t^2 + \left[ \left(1 - \frac{1}{\alpha}\right) f_1^2(t) - \tilde{V}_1(t) \right] |v|^2 dt$$

verify

$$\sum_{k \in \mathbb{N}} \sqrt{\mu_k^\alpha} \leq \sum_{k \in \mathbb{N}} \sqrt{\nu_k^\alpha}. \quad (3.6)$$

Applying the sharp inequality of Hundertmarkt-Lieb-Thomas [11] (see Appendix) to the operator  $L_1^\alpha$  we get

$$\begin{aligned} \sum_{k \in \mathbb{N}} \sqrt{\nu_k^\alpha} &\leq \frac{1}{2} \int_{-\infty}^{+\infty} \left[ \left(\frac{1}{\alpha} - 1\right) f_1^2(t) + \tilde{V}_1(t) \right] dt \\ &\leq \frac{1}{2} \int_{-\infty}^0 \left[ \left(\frac{1}{\alpha} - 1\right) f^2(t) + \tilde{V}(t) \right] dt \\ &\leq \frac{1}{2} \int_0^1 \left[ \left(\frac{1}{\alpha} - 1\right) A^2(r) + V(r) \right] r dr. \end{aligned} \quad (3.7)$$

To conclude we need the following

**Lemma 3.2** *Assume that  $K = \inf_{x \in \Omega} \mathbf{b}(x) > 0$ . Then for any  $\varepsilon \in ]0, 1[$*

$$N(h_{0,V}) = N(h_{0,0} - V) \leq \frac{1}{\varepsilon} \int_0^1 \left[ 1 + \left| \log \left( \sqrt{\frac{(1-\varepsilon)K}{\varepsilon}} r \right) \right| \right] V(r) r dr, \quad (3.8)$$

*In particular*

$$N(h_{0,V}) \leq 2 \int_0^1 \left[ 1 + \left| \log(\sqrt{K}r) \right| \right] V(r) r dr. \quad (3.9)$$

*Proof.*–

**Step 1** :From (1.3) we get that  $h_A(u) \geq K \int_\Omega |u|^2 dx \quad \forall u \in C_0^\infty(\Omega)$ , which implies for  $h_{0,0}$  (returning to the variable  $r$  and considering  $V \equiv 0$ ),

$$\begin{aligned} h_{0,0}(w) &= \int_0^1 \left[ \left| \frac{\partial w}{\partial r} \right|^2 + r^{-2} A^2(r) |w|^2 \right] r dr \\ &\geq K \int_0^1 |w|^2 r dr \quad \forall w \in C_0^\infty([0, 1]). \end{aligned}$$

We write for any  $\varepsilon \in ]0, 1[$

$$N(h_{0,0} - V) \leq N(\varepsilon h_{0,0} + (1 - \varepsilon)K - V) \leq N\left(h_{0,0} + \frac{(1 - \varepsilon)K}{\varepsilon} - \frac{V}{\varepsilon}\right), \quad (3.10)$$

where we have used the fact that multiplying an operator by a positive constant does not change the number of its negative eigenvalues.

Step 2 : We establish the following upperbound :

$$N(h_{0,0} + 1 - V) = N(h_{0,V} + 1) \leq \int_0^1 [1 + |\log r|] V(r) r dr. \quad (3.11)$$

We have

$$\begin{aligned} h_{0,V}(w) &= \int_0^1 \left[ \left| \frac{\partial w}{\partial r} \right|^2 + [r^{-2}A^2(r) - V(r)] |w^2| \right] r dr \\ &\geq \int_0^1 \left[ \left| \frac{\partial w}{\partial r} \right|^2 - V(r) |w^2| \right] r dr \quad \forall w \in C_o^\infty([0, 1]). \end{aligned}$$

By the variational principle,

$$N(h_{0,V} + 1) \leq N(P_0 + 1 - V), \quad (3.12)$$

where  $P_0$  is the operator generated by the closure, in  $L^2([0, 1], r dr)$  of the quadratic form

$$\int_0^1 \left| \frac{\partial w}{\partial r} \right|^2 r dr, \quad w \in C_o^\infty([0, 1]).$$

Considering the mapping  $U : L^2([0, 1], r dr) \rightarrow L^2([0, 1], dr)$  defined by  $(Uf)(r) = r^{1/2}f(r)$  we get that

$$N(P_0 + 1 - V) \leq N(T_0 + 1 - V) \quad (3.13)$$

where the operator  $T_0 = UP_0U^{-1}$  is the Sturm Liouville operator on  $L^2([0, 1], dr)$  acting on its domain by

$$(T_0u)(r) = -u''(r) - \frac{u(r)}{4r^2}, \quad u(0) = u(1) = 0. \quad (3.14)$$

The upperbound (3.11) will follow from the properties of  $G(r, r, 1)$ , the diagonal element of the integral kernel of  $(T_0 + 1)^{-1}$ . Precisely we have

$$G(r, r, 1) \leq r(1 + |\log r|), \quad r \in [0, 1[. \quad (3.15)$$

The proof of (3.15) is given in Appendix B. The Birman-Swinger principle then yields

$$N(T_0 + 1 - V) \leq \int_0^1 G(r, r, 1) V(r) dr \leq \int_0^1 [1 + |\log r|] V(r) r dr. \quad (3.16)$$

This ends the proof of (3.11), together with the inequalities (3.12) and (3.13).

Step 3 : We mimick the previous method to get, for any strictly positive number  $k$

$$N(h_{0,0} + k^2 - V) \leq \int_0^1 [1 + |\log(kr)|] V(r) r dr . \quad (3.17)$$

Due to the Birman-Swinger principle it suffices to prove that, for any strictly positive number  $k$

$$G(r, r, k^2) \leq r(1 + |\log(kr)|), \quad r \in [0, 1[ . \quad (3.18)$$

This is done in Appendix C.

Step 4 :Returning to (3.10) and applying (3.17) with  $k^2 = \frac{(1-\varepsilon)K}{\varepsilon}$  and  $\frac{V}{\varepsilon}$  instead of  $V$  we get, for any  $\varepsilon \in ]0, 1[$

$$N(h_{0,0} - V) \leq N\left(h_{0,0} + \frac{(1-\varepsilon)K}{\varepsilon} - \frac{V}{\varepsilon}\right) \quad (3.19)$$

$$\leq \frac{1}{\varepsilon} \int_0^1 \left[1 + \left|\log\left(\sqrt{\frac{(1-\varepsilon)K}{\varepsilon}} r\right)\right|\right] V(r) r dr , \quad (3.20)$$

and taking  $\varepsilon = \frac{1}{2}$  we obtain Lemma 3.2. □

Theorem 2.1 follows from Lemma 3.2 together with inequalities (3.5), (3.6), and (3.7).

### 3.2 Proof of Theorem 2.2

Noticing that for any  $\lambda > 0$  the constant potential  $V(x) \equiv \lambda$  is in  $L^1(\Omega)$ , and that  $N(A, \lambda)$  denotes the number of eigenvalues of the operator  $H_A^D$  less than  $\lambda$ , we apply Theorem 2.1 to  $V(x) \equiv \lambda$ . To get the result it suffices to compute  $\int_0^1 [1 + |\log(kr)|] r dr$ . We get after computation that

$$\int_0^1 [1 + |\log(kr)|] r dr = \gamma_k, \quad (3.21)$$

with

- $\gamma_k = \frac{3 - 2 \log k}{4}$  if  $k \leq 1$
- $\gamma_k = \frac{1 + 2 \log k}{4} + \frac{1}{2k^2}$  if  $k > 1$ .

### 3.3 Proof of Remark 2.3

The choice of  $A' = A + c$  is obtained by taking the minimum over the constants  $c$  of the function  $F(c) = \int_0^1 (A + c)^2(r) r dr$ . To get the minimum over the values of  $\alpha$  we study the sign of the expression, for any  $\alpha \in ]0, 1[$ , of

$$g_\lambda(\alpha) := \frac{\lambda}{2\sqrt{1-\alpha}} + \frac{\sqrt{1-\alpha}}{\alpha} I .$$

A direct computation shows that the value  $\alpha_\lambda$  which realizes the minimum of  $g_\lambda(\alpha)$  is the positive solution of

$$\alpha^2(\lambda - 2I) + 6\alpha I - 4I = 0 . \quad (3.22)$$

## 4 An eigenvalue asymptotic upperbound

From Theorem 2.2 we get easily an asymptotic estimate for the righthandside of (2.2) when  $\lambda$  tends to  $\infty$  :

**Corollary 4.1** *If assumptions  $(H_1)$  and  $(H_2)$  are satisfied and if moreover*

$$\mathbf{b}(x) \leq (D(x))^{-\beta}, \quad \beta < \frac{3}{2}$$

*then the number of eigenvalues of the operator  $H_A$  less than  $\lambda$  satisfies, as  $\lambda$  tends to  $\infty$*

$$N(A, \lambda) \leq \left(\frac{1}{2} + c_K\right)\lambda + \sqrt{\lambda}\sqrt{I} + O(1), \quad (4.1)$$

where

$$I = \int_0^1 A'^2(r) r dr ,$$

and

- $c_K = \frac{3 - \log K}{2}$  if  $K \leq 1$

and

- $c_K = \frac{1 + \log K}{2} + \frac{1}{K}$  if  $K > 1$ ,

*This holds for any radial gauge  $A'$  associated to  $B$ , and the minimum of the righthand-side is obtained by choosing  $A'$  so that  $\int_0^1 A'(r) r dr = 0$ .*

**Example 4.2** *Assume  $b(r) \equiv 1$  in (3.1) and  $\beta \neq 1$ . Then  $c_K = \frac{3}{2}$  and the minimum is obtained for  $I = I_\beta = \int_0^1 A_\beta^2(r) r dr$ , where  $A_\beta$  is defined as in (2.3), so that*

$$I_\beta = \frac{1}{(1-\beta)^2(3-2\beta)(4-2\beta)} . \quad (4.2)$$

*Proof.*–

We define as previously, for any  $\alpha \in ]0, 1[$ ,

$$g_\lambda(\alpha) := \frac{\lambda}{2\sqrt{1-\alpha}} + \frac{\sqrt{1-\alpha}}{\alpha} I$$

and we want to determine the asymptotic behavior as  $\lambda$  tends to  $\infty$  of  $g_\lambda(\alpha_\lambda)$ , where  $\alpha_\lambda$  is the minimum of  $g_\lambda(\alpha)$ .

From (3.22) we compute the following asymptotics

$$\alpha_\lambda = \frac{2\sqrt{I}}{\sqrt{\lambda}} + O\left(\frac{1}{\lambda}\right)$$

$$\sqrt{1-\alpha_\lambda} = 1 - \frac{\sqrt{I}}{\sqrt{\lambda}} + O\left(\frac{1}{\lambda}\right),$$

and this gives the result.

The minimal value is obtained as previously by taking the minimum over the constants  $c$  of the function  $F(c) = \int_0^1 (A+c)^2(r) r dr$ .

□

**Remark 4.3** *The leading term in the estimate (4.1) is of the same order than the leading term in the Weyl formula for the Dirichlet Laplacian (corresponding to the case  $A \equiv 0$ ) in the unit disk.*

## 5 Appendix A

We recall the sharp inequality of Hundertmarkt-Lieb-Thomas

**Theorem 5.1** *Let*

$$Lv(t) = -v''(t) - W(t)v(t), \quad W \geq 0 \quad W \in L^1(\mathbb{R})$$

*be defined in the sense of quadratic forms on  $\mathbb{R}$ , and assume that the negative spectrum of  $L$  is discrete. Denote by  $\{-\nu_k, k \in \mathbb{N}\}$  the negative eigenvalues of  $L$ . Then*

$$\sum_{k \in \mathbb{N}} \sqrt{\nu_k} \leq \frac{1}{2} \int_{-\infty}^{+\infty} W(t) dt .$$

## 6 Appendix B

Let us compute the diagonal element for the Green function  $G(r, r', 1)$  of the operator  $T_0$  defined by (3.14).  $G(r, r', 1)$  is the solution of

$$((T_0 + 1)u)(r) = \delta_{r'}(r), \quad u(0) = u(1) = 0. \quad (6.1)$$

We have

$$\begin{aligned} G(r, r', 1) &= A_1 u_1(r) + A_2 u_2(r) \quad r \leq r' \\ G(r, r', 1) &= B_1 u_1(r) + B_2 u_2(r) \quad r > r', \end{aligned}$$

where  $u_1(r) = \sqrt{r}I_0(r)$  and  $u_2(r) = \sqrt{r}K_0(r)$  are independent solutions of the related homogeneous equation, ( $I_0$  and  $K_0$  are the modified Bessel functions).

The coefficients depend of  $r'$  but we omit the indices for the sake of clarity. Due to the boundary conditions and to the fact that the derivative (with respect to  $r$ ) of  $G(r, r', 1)$  has the discontinuity in  $r'$  of a Heaviside function, they satisfy :

$$\begin{aligned} A_1 u_1(0) + A_2 u_2(0) &= 0 & B_1 u_1(1) + B_2 u_2(1) &= 0 \\ B_1 - A_1 &= \frac{-u_2(r')}{W(r')} & B_2 - A_2 &= \frac{u_1(r')}{W(r')} \end{aligned}$$

where  $W(r')$  is the value of the Wronskian of  $u_1$  and  $u_2$  taken at the point  $r'$ .

The first equation is always satisfied since  $u_1(0) = u_2(0) = 0$ . Let us set  $A_2 = 0$ . We have  $W(r') = u_1'(r')u_2(r') - u_1(r')u_2'(r') = r'\hat{W}(r')$  where  $\hat{W}(r')$  is the Wronskian of the modified Bessel functions  $I_0$  and  $K_0$ . As  $r'\hat{W}(r') = 1$  (see [2]), we get after solving the above system, and doing  $r = r'$  :

$$\begin{aligned} G(r, r, 1) &= u_1(r) \left[ -u_1(r) \frac{u_2(1)}{u_1(1)} + u_2(r) \right] \\ &= rI_0(r) \left[ -I_0(r) \frac{K_0(1)}{I_0(1)} + K_0(r) \right]. \end{aligned}$$

Using again the properties of the modified Bessel functions we can write

$$G(r, r, 1) \leq rI_0(r)K_0(r)$$

and observe (see figure 1) that the function

$$g(r) = \frac{I_0(r)K_0(r)}{1 + |\log r|}$$

is decreasing on  $]0, \infty[$  and has a limit at  $r = 0$  equal to 1, so we get

$$G(r, r, 1) \leq r(1 + |\log r|), \quad r \in [0, 1[.$$




Figure 1: The function  $g$

## 7 Appendix C

We now compute the diagonal element for the Green function  $G(r, r', k^2)$  of the operator  $T_0$  defined by (3.14).  $G(r, r', k^2)$  is the solution of

$$((T_0 + k^2)u)(r) = \delta_{r'}(r), \quad u(0) = u(1) = 0. \quad (7.1)$$

We have, as previously

$$G(r, r, k^2) = u_1(r) \left[ -u_1(r) \frac{u_2(1)}{u_1(1)} + u_2(r) \right]$$

where  $u_1(r) = \sqrt{r}I_0(kr)$  and  $u_2(r) = \sqrt{r}K_0(kr)$  are independent solutions of the related homogeneous equation. This leads to

$$G(r, r, k^2) = rI_0(kr) \left[ -I_0(kr) \frac{K_0(k)}{I_0(k)} + K_0(kr) \right] \leq rI_0(kr)K_0(kr) \leq r(1 + |\log(kr)|).$$

## References

- [1] J. Avron, I. Herbst & B. Simon. *Schrödinger operators with magnetic fields*, Duke. Math. J. **45** (1978), 847-883.
- [2] M. Abramovitz, I. Stegun. *Handbook of mathematical functions*, National Bureau of Standards (1964)
- [3] A. Balinski, W. Evans & R. Lewis. *On the number of negative eigenvalues of Schrödinger operators with an Aharonov-Bohm magnetic field*, R. Soc. Lon. Proc. Ser. A Math. Phys. Eng. Sci. **457** (2001), 2481-2489.

- [4] Y. Colin de Verdière. *L'asymptotique de Weyl pour les bouteilles magnétiques*, Commun. Math. Phys. **105** (1986), 327-335.
- [5] Y. Colin de Verdière & F. Truc *Confining particles in a purely magnetic field*, Ann Inst Fourier (to appear).
- [6] M. Cwickel. *Weak type estimates for singular values and the number of bound states of Schrödinger operators*, Ann. of Math. **106** (1977), 93-100.
- [7] L. Erdős & J.P. Solovej. *Magnetic Lieb-Thirring inequalities with optimal dependence on the field strength*, Journ. Stat. Phys. **116** (2004), 475-506.
- [8] L. Erdős & J.P. Solovej. *Magnetic Lieb-Thirring inequalities for the 3-dimensional Pauli operator with a strong nonhomogeneous magnetic fields*, Ann. Henri Poincaré **5** (2004), 671-745.
- [9] R. Frank *Remarks on eigenvalue estimates and semi-group domination*, Spectral and Scattering Theory for Quantum Magnetic Systems, P. Briet et al (eds), 63-86, Contemp. Math. 500, Amer. Soc., Providence, RI, 2009
- [10] R. Frank & R. Olofsson. *Eigenvalue bounds for Schrödinger operators with a homogeneous magnetic field*, arXiv 1102.0329
- [11] D.Hundertmarkt, E.H.Lieb & L.E.Thomas. *A sharp bound for an eigenvalue moment of the one-dimensional Schrödinger operator*, Adv. Theor. Math. Phys. **2** (1998), 719-731.
- [12] H. Kovarik. *Eigenvalue bounds for two-dimensional magnetic Schrödinger operators*, arXiv 1103.5194
- [13] A. Laptev. *The negative spectrum of a class of two-dimensional Schrödinger operators with spherically symmetric potentials*, Func. Anal. Appl. **34** (2000), 305-307.
- [14] A. Laptev. *Spectral Inequalities for Partial Differential Equations and their applications* to appear in Proceedings of the 5th ICCM Beijing 2010
- [15] A. Laptev, T. Weidl *Recent results on Lieb-Thirring inequalities*, Journées Equations aux dérivées partielles, St Jean de Monts (2000).
- [16] E.H. Lieb. *Bound states of the Laplace and Schrödinger operators with spherically symmetric potentials*, Bull. Amer. Math. Soc. **82** (1976), 751-753.
- [17] E.H. Lieb, W. Thirring, *Inequalities for the moments of the eigenvalues of the Schrödinger Hamiltonian and their relation to Sobolev inequalities*, Studies in Mathematical Physics, Princeton Univ. Press, Princeton, NJ (1976), 269-303.

- [18] G.V. Rozenblum. *Distribution of the discrete spectrum of singular differential operators*, Soviet. Math. Soc. **20** (1976), 63-71.
- [19] B. Simon. *Essential Self-adjointness for Schrödinger operators with singular potentials: A generalized Kalf-Walter-Schminke theorem*, Arch. Rational Mech. Analysis, **52** (1973), 44-48.
- [20] B. Simon. *Schrödinger operators with singular magnetic vector potentials*. Math. Z. **131** (1973), 361–370.
- [21] F. Truc. *Semi-classical asymptotics for magnetic bottles*, Asympt. Anal. **15** (1997), 385-395.