HAL
open science

The Mediterranean oak forest: a new habitat for the Capercaillie?

Manuel A. González, Pedro P. Olea, Luis Robles, Vicente Ena

To cite this version:

Manuel A. González, Pedro P. Olea, Luis Robles, Vicente Ena. The Mediterranean oak forest: a new habitat for the Capercaillie?. Journal für Ornithologie = Journal of Ornithology, 2010, 151 (4), pp.901-906. 10.1007/s10336-010-0528-6 . hal-00588626

HAL Id: hal-00588626

https://hal.science/hal-00588626v1

Submitted on 25 Apr 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Mediterranean *Quercus pyrenaica* oak forest: a new habitat for the Capercaillie?

Manuel A. González[1]*, Pedro P. Olea[2], Luis Robles[3], Vicente Ena[1].

[1]Dpt. of Biodiversity and Environmental Management, University of León, Campus de Vegazana s/n, 24071 León, Spain.

[2]School of Biology, IE University, Campus Santa Cruz la Real, 40003 Segovia, Spain.

[3]Asociación para la Conservación del Urogallo (ACU), C/ Unic s/n, 24850 Baix Pallars (Buseu), Lleida, Spain.

*Corresponding author. Email: magong@unileon.es

Phone: +34 987 291907

ABSTRACT

We described an extension of the known distribution range of capercaillie into an atypical area and habitat for the species. Nine Cantabrian capercaillie *Tetrao urogallus cantabricus* leks and 14 cocks were registered in Mediterranean *Quercus pyrenaica* forests in an area of 1,500 km^2, of which 4,500 forest hectares were surveyed. At present this population represents both the southern-most distribution for capercaillie and the only one inhabiting Mediterranean *Q. pyrenaica* forests, what suggests a wider adaptation of this (sub)species than previously thought. This capercaillie population and its habitat need to be better studied, as well as to be considered in conservation planning for Cantabrian capercaillie.

KEY WORDS*:* conservation, leks, Mediterranean, *Quercus pyrenaica*, *Tetrao urogallus cantabricus*, plasticity.

INTRODUCTION

Populations across distribution range show different adaptations to local conditions from the core areas to the periphery of their distribution (Guo et al. 2003). In the case of capercaillie, the vast majority of populations inhabit boreal and montane coniferous forests with abundant bilberry *Vaccinium myrtillus* understory (Storch 1993; 2000a; 2000b, Storch 2001). However, the Cantabrian subspecies *Tetrao urogallus cantabricus* (Cantabrian Mountains, NW Spain) resides mainly in pure deciduous forests of beech *Fagus sylvatica*, birch *Betula pubescens* and sessile oak *Quercus petraea*, as well as mixed forests of beech and oaks (*Quercus robur*, *Q. petraea*, and *Q. pyrenaica*) (Bañuelos & Quevedo 2008). In these forests holly *Ilex aquifolium* provides a key, evergreen winter food resource for Cantabrian capercaillie (Storch 2001; Blanco-Fontao et al. 2009) whereas it relies on bilberry in summer-autumn (Blanco-Fontao et al. 2009). Previous studies showed, however, a negative selection to *Quercus pyrenaica* forests by Cantabrian capercaillie at the northern slope of the Cantabrian range (Quevedo et al. 2006b).

This endemic capercaillie subspecies is isolated and separated more than 300km from the nearest capercaillie population in the Pyrenees. After severe population decline in the last three decades (from 2,000 to 400 adults), only 108 leks remain occupied in the northern watershed and 85 leks in the southern, corresponding to an overall occupancy rate of 32% of all known leks (Bañuelos & Quevedo 2008). Consequently, Cantabrian capercaillie subspecies qualifies to be listed as ”Endangered” according to the IUCN criteria (Storch et al. 2006).

Capercaillie populations have so far been generally considered to be distributed in the Eurosiberian biogeographic region, outside Mediterranean

region (but see Handrinos and Akriotis 1997, Storch 2001; discussion section). Unlike the Eurosiberian region, the Mediterranean region is characterized by a period of drought during hot summer which determines the existing vegetation (e.g., Penas 1995; Blondel & Aronson 1999; Rivas-Martínez et al. 2005). This Mediterranean environment is thought to be unsuitable for a species primarily adapted to boreal environments. However, there are historical records suggesting capercaillie presence during 17th-19th centuries in Mediterranean areas in Spain (Madoz 1848; Castroviejo 1975; Martínez 1993), but the first sighting record in *Q. pyrenaica* forests in this Mediterranean area was unnoticed until May 1998 (Pollo et al. 2004) (No. 1 in Fig. 1).

In this study we present information on the number of cocks and characteristics of leks located in an atypical area and habitat for the capercaillie, i.e., Mediterranean *Q. pyrenaica* forests with virtual lack of bilberry and holly. This study is the first showing capercaillie widely present in this forest type, suggesting a wider ecological plasticity of this (sub)species than previously thought. We also discuss implications that this new capercaillie site may have in the conservation of Cantabrian capercaillie.

MATERIAL & METHODS

Study area

The study area is located on the southern slope of the Cantabrian Mountains in León province (NW Spain). It is centred at 42° 39'N and located in the Mediterranean biogeographical region bordering the Eurosiberian region (Atlantic) (Rivas-Martínez et al. 2004, European Environment Agency, 2008) (Fig. 1). The study area covers approximately 1,500km² below the putative line

separating the two biogeographical regions in a landscape slightly mountainous (elevation ranges from 800 to 1,700 m.a.s.l.). It belongs to the biogeographical unit Carpetan-Leonese Subprovince from the Mediterranean region with supramediterranean bioclimate (Penas 1995, Rivas-Martínez et al. 2005, del Río et al 2007). A meteorological station from our study area (Villameca) clearly shows two dry months ($P<2T$; Blondel & Aronson 1999) in summer: July (P=26; T=19.9) and August (P=19; T=19.1) (data based on 37 years).

Dominant forests are supramediterranean natural (more than 50 years old) and post-fire *Q. pyrenaica* forests frequently occurring interspersed with Scots pine *Pinus sylvestris* plantations (less than 50 years old). Both forest types are largely monoespecific in the study area. *Q. pyrenaica* forests cover 27,000ha (19% of the study area). Scots pine plantations occupy 11,000ha (8%).

Q. pyrenaica (Willd.) is a marcescent oak tree distributed widely in Spain as well as the western part of France and the northwest of Marocco. This tree develops under siliceous soils in Eurosiberian region as well as in the Mediterranean region with an extensive representation in this biogeographic region. In Spain, *Q. pyrenaica* represents the dominant species of different vegetation series (del Río et al 2007).

Forest fragments of the study area are embedded in a matrix mainly composed of heather *Erica australis* 1,560ha (1%), brooms *Genista florida* and Cy*tisus scoparius* (hereafter *C. scoparius*) 3,640ha (3%), meadows 3,460ha (3%) and riparian lowland forest of *Populus nigra*, *Fraxinus excelsior* and *Alnus glutinosa* 5,340ha (4%). Understory cover is mainly dominated by heath *Erica arborea* (hereafter *E. arborea*) and broom *C. scoparius* while bilberry and holly is completely absent or very scarce (<0.5% of the forest ground cover) in the study area. The remaining 62% surface corresponds with anthropogenic landscapes such

as villages, roads, opencast mines, valley bottoms, which are not suitable for capercaillie habitat. Human population is approximately 1,000 people (0.6 people/km²) and it has declined by 80% since the 1950's. Livestock rearing is the main economic activity in the study area, followed by coal mining industry, agriculture, forestry and hunting.

Methods

No systematic sampling of the species was performed for the whole study area. We gathered information on capercaillie presence from 2002-2009 from two sources: i) questionnaires and reports sent by forest wardens, hunters and local people to the León environmental agency from the regional government (Consejería de Medio Ambiente of the Junta de Castilla y León), and ii) field surveys. Samples were recorded over the whole year. Data from the first source were validated within two weeks after receipt of the sample through field surveys carried out by an experienced observer (L. Robles). Field surveys consisted of a 3-4 hours systematic zig-zag walk looking for presence signs of capercaillie (e.g., direct sightings, footprints, droppings or feathers), especially focusing on sites known to be highly used by the species (e.g., forest paths, dead trunks, big stones, oldest forest sites, etc.).

In forest patches bigger than 1km² (i.e., more than 3-4 hours of sampling required) we conducted surveys over consecutive days until covering the whole forest patch (i.e., a continuous forest isolated by a non-forest matrix). Direct sightings, footprints, droppings or feathers recorded in a site were considered as signs of capercaillie presence. Each forest patch was completely surveyed 6-10 times throughout one year, 1-2 months each between. If no sign of capercaillie

presence was found, the forest patch was labelled unoccupied and no further surveys were conducted in that specific patch.

Additionally, we surveyed the occupied patches in order to find leks. Selected sites were locations where survey after survey we found capercaillie signs. The visits were performed throughout April and May (i.e., display season) after 12 a.m. to avoid disturbing birds. A lek is defined as a site where one or more cocks consistently display for hens, plus the adjacent surrounding forest habitat. It has traditionally been used as capercaillie occurrence measure in the Cantabrian range (see Obeso & Bañuelos 2003).

During April-May where fresh capercaillie signs were found, 2-3 observers (forest wardens and L. Robles) visited the site at night as many as four times until the display finished, or until well past dawn. A site was considered as an occupied lek when at least one cock was heard calling or seen displaying.

Every detected lek was surveyed each year to label it as occupied or unoccupied just looking for presence signs after 12 a.m. (see above). In April-May 2009 every occupied lek was surveyed at night to get the number of displaying cocks.

Nine forest structure variables (Table 1) were measured in each lek consisting of twelve circular plots with 5m radius. Three plots separated by 10 meters among them were arranged in each cardinal direction from the lek centroid (i.e., the point where previously here a cock was recorded displaying).

RESULTS

A total of nine leks, including that discovered in 1998 (Pollo et al 2004), were registered in the study area, seven of them remained occupied in 2009 (Fig. 2). The highest occupancy was in 2006 with eight occupied leks (Fig. 2). Since 2006 no new lek was found, despite an increase of around 2,000 additional hectares surveyed (60% of the total surveyed surface, Fig. 2). Leks were grouped into three subareas according to the watersheds: one subarea had three leks, other had four and the last subarea had two (Fig. 1). Mean nearest neighbour distance between leks was 2.07 ± 0.74 kilometers (mean ± SE).

Leks were in *Q. pyrenaica* forests ranging from 1,132 to 1,398 m.a.s.l. (mean ± SE: 1275 ± 100.93, n=9) and mainly faced in a northerly direction with two exceptions facing south (Table 1). They showed heterogeneity in the forest structure, ranging from sites with young trees (perimeter 0.19 ± 0.005 meters) with high canopy cover (0.81 ± 0.06%) to mature trees (perimeter 0.90 ± 0.62 meters) with low canopy cover (0.28 ± 0.07%). The maximum tree height in leks varied from 7.75 ± 0.86 meters to 16.50 ± 3.20 meters (see Fig. 3).

Understory cover ranged from low (0.004 ± 0.86%) to high (0.62 ± 0.27%). In six of the nine leks the dominant understory species was *E. arborea* while *C. scoparius* dominated the remaining one (Table 1).

In 2009 twelve cocks were recorded at the seven occupied leks. Moreover, two additional cocks not referred to any lek were detected during the spring 2009 fieldwork. Two nests with five and seven eggs each (both in *Q. pyrenaica* forest) were found in early June 2009 (Fig. 1, No 5 and 8).

DISCUSSION

So far, capercaillie always occurred in the Eurosiberian biogeographical region with the exceptions of the probably introduced birds of Mount Athos (Greece; Handrinos and Akriotis 1997), currently extinct (S. Xirouchakis per. comm.), and two other unclear data in Mount Grammos and Lailias Forest Reserve in Greece (Storch 2001). Therefore, the capercaillie population we described here represents both the southern-most distribution for capercaillie and the only one inhabiting Mediterranean *Q. pyrenaica* forests. These forests are different from the nearest ones occupied by capercaillie leks located north of the putative line separating Eurosiberian and Mediterranean regions (Fig 1, Table 1 and 2). These northern leks are mainly located in birch and sessile oak forests with understory dominated by bilberry (see Table 2) in a Eurosiberian bioclimatic environment (superior montane thermotype and inferior humid ombrotype) (Penas 1995). Nonetheless lek 4, although in *Q. pyrenaica* forest, has an eurosiberian bioclimatic environment (Penas 1995), probably because of its most northerly relative position.

Despite an increase in the surveyed area, no new leks were found after 2006. However, the fact that we only surveyed 12% of the forest surface within our study area (Fig. 2), suggests new leks may be found in the future. Although the number of capercaillie in the study area is small, it represents approximately 7% of the global Cantabrian population, highlighting the conservation value of this site.

All the nine leks were located in *Q. pyrenaica* forests. Conversely no lek was located in pure Scots pine plantations, although these were available in the

study area. In addition, two of four radio-tracked (one hen and one cock) capers used all year round *Q. pyrenaica* forests of the studied area (authors, unpublished data) and two capercaillie nests were discovered in these forests. All of this suggests that *Q. pyrenaica* forests are suitable year-round habitat for the Cantabrian capercaillie. This is particularly important because the potential habitat for this endangered subspecies could thus be greatly increased, as *Q. pyrenaica* forests are highly represented with > 100,000 hectares at the southern slope of the Cantabrian Mountains (Gil-Sanchez & Torre-Antón 2007). Additionally, despite the idea that *Q. pyrenaica* forests are not considered as suitable habitat (Quevedo et al. 2006b) and neither are included in the recovery plans for the Cantabrian capercaillie, our study highlights the need to take into account these forests for the conservation of this subspecies. Nonetheless, the role of the pine plantations for capercaillie in the study area should be further assessed.

Finally, this study supports what is suggested by Quevedo et al. (2006a; 2006b) and Blanco-Fontao et al. (2009) that many general considerations for the species overall are not fully useful for the Cantabrian capercaillie, and suggests a wider adaptation and plasticity of this species than previously thought. For example, the near absence of bilberry and holly in the studied area suggest that capercaillie can maintain viable populations without this important summer and winter food resource for Cantabrian population and elsewhere (Storch 2001; Blanco-Fontao et al. 2009). Future research could address a detailed comparison of capercaillie suitable habitat between the two different biogeographical regions. Studying the ecology and genetic variability of this local population living in this atypical habitat would be of a great value to further understand and conserve this subspecies in particular and *Tetrao urogallus* in general.

Zusammenfassung

Der Mediterrane Pyrenäen-Eichenwald: ein neuer Lebensraum für das Auerhuhn?

Wir beschreiben eine Ausdehnung des bekannten Verbreitungsgebietes des Auerhuhns in ein atypisches Gebiet und Habitat für diese Art. In einem 1,500 km^2 großen Pyrenäen-Eichenwald Gebiet, wovon 4,500 ha Wald untersucht wurden, konnten neun Balzarenen des Kantabrischen Auerhuhns *Tetrao urogallus cantabricus* und 14 Hähne festgestellt werden. Aktuell repräsentiert diese Population die sowohl südlichste Verbreitung von Auerhühnern als auch die einzige, die mediterrane Pyrenäen-Eichenwälder bewohnt. Dieses weist auf eine größere Anpassung dieser (Unter-)Art hin, als vorher vermutet. Diese Auerhuhn-Population und ihr Lebensraum müssen jedoch noch besser untersucht werden und sie sollte auch in Schutzstrategien für das Kantabrische Auerhuhn berücksichtigt werden.

ACKNOWLEDGEMENTS

The forest wardens Fernando Gonzalo, Álvaro Ortiz and Ramón Balaguer showed us many little known aspects of the *Q. pyrenaica* forests. We wish to thank to María Angeles Osorio and César Pollo from the Consejería de Medio Ambiente de León of the Junta de Castilla y León. Particular thanks are due to Beatriz Blanco-Fontao, Patricia Mateo, Andy Dean and two anonymous reviewers whose revisions largely improved the manuscript. Manuel A. González is supported by a PhD scholarship of the Universidad de León.

LITERATURE CITED

Bañuelos M J, Quevedo M (2008) Update of the situation of the Cantabrian capercaillie *Tetrao urogallus cantabricus*: an ongoing decline. Grouse News 25:5-7.

Blanco-Fontao B, Fernández-Gil A, Obeso J R, Quevedo M (2009) Diet and habitat selection in Cantabrian capercaillie: ecological differentiation of a rear-edge population Journal of Ornithology *in press*.

Blondel, J., y Aronson, J. 1999. Biology and Wildlife of the Mediterranean Region. Oxford University Press, Oxford, UK.

Castroviejo J (1975) El urogallo en España. Monografías de la Estación Biológica de Doñana, CSIC.

Del Hoyo J, Elliot A, Sargatal J (ed) (1994) Handbook of the Birds of the World: New world vultures to guineafowl. Lynx Editions, Barcelona.

Del Rio S, Herrero L, Penas A (2007) Bioclimatic analysis of the Quercus pyrenaica forests in Spain. Phytocoenologia 37:541-560.

European Environment Agency (EEA). Biogeographical Regions, Europe 2008. http://www.eea.europa.eu/data-and-maps/data/biogeographical-regions-europe-2008/ (accessed 15 October 2009).

Gil-Sánchez L, Torre-Antón M (ed). 2007. Atlas forestal de Castilla y León, Valladolid.

Guo Q, Taper M, Schoenberger M, Brandle J (2005) Spatial-temporal population dynamics across species range: from centre to margin. Oikos 108:47-57.

Handrinos G, Akriotis T (1997) The birds of Greece. Christopher Helm, London.

Helle P, Kurki S, Lindén H (1999) Change in the sex ratio of the Finnish capercaillie *Tetrao urogallus* population. Wildl Biol 5:25-31.

Madoz P (1848) Diccionario geográfico-estadístico-histórico de España y sus posesiones de ultramar. Madrid.

Martínez AM (1993) Contribución al conocimiento de la eco-etología del urogallo cantábrico. Universidad de León. León.

Obeso JR, Bañuelos MJ (2003) El urogallo (*Tetrao urogallus cantabricus*) en la Cordillera Cantábrica. Servicio de Publicaciones de Parques Nacionales. Ministerio de Medio Ambiente, Madrid.

Penas A 1995. Atlas del Medio Natural de la provincia de León. Instituto Geominero de España.

Pollo CJ, Robles L, García-Miranda A (2004) Aumentan los urogallos "locos" en la cordillera Cantábrica. La Garcilla 119:20-21.

Quevedo M, Bañuelos MJ, Obeso JR (2006a) The decline of Cantabrian capercaillie: How much does habitat configuration matter?. Biol Conserv 127:190-200.

Quevedo M, Bañuelos MJ, Saez O, Obeso JR (2006b) Habitat selection by Cantabrian capercaillie *Tetrao urogallus cantabricus* at the edge of the species distribution. Wildl Biol 12:267-276.

Rivas-Martínez S, Loidi J (1999) Bioclimatology of the Iberian Peninsula. Itinera Geobot. 13:41-48.

Rivas-Martínez S, Penas A, Díaz T E (2004) Biogeographic Map of Europe. Cartographic Service. University of León, Spain. http://www.ucm.es/info/cif/form/bg_med.htm (accessed 10 September 2009).

Rivas-Martínez S et al. (2005). Mapa de series, geoseries y geopermaseries de vegetación de España. Memoria del mapa de vegetación potencial de España. Parte 1. http://www.ucm.es/info/cif/book/mapa_series/mapa_series_01.pdf (accessed 10 September 2009).

Storch I (1993) Habitat selection by capercaillie in summer and autumn: Is bilberry important?. Oecologia 95:257-265.

Storch I (2000a) Conservation status and threats to grouse worldwide: an overview. Wildl Biol 6:195-204.

Storch I (2000b) Status Survey and Conservation Action Plan 2000-2004: Grouse. WPA/BirdLife/SSC Grouse Specialist Group. IUCN, Glan, Switzerland and Cambridge, UK, and the World Pheasant Association, Reading, UK.

Storch I (2001) Capercaillie. Pages 1-24 in O. U. Press, editor. BWP Update: The journal of birds of the Western Palearctic. Oxford, UK.

Storch I, Bañuelos MJ, Fernandez-Gil A, Obeso JR, Quevedo M Rodriguez-Muñoz R (2006) Subspecies Cantabrian capercaillie *Tetrao urogallus cantabricus* endangered according to IUCN criteria. J Ornithol 147:653-655.

APPENDIX

Appendix 1. Description of the leks (n=9) in the study area (see ● in the map of the study area). Mean ± SE are shown for forest structure variables of the leks.

Footnote: **Lek** = lek number by discovery date; **Date** = year when the lek was first-time recorded (April-May); **Cocks** = No. of capercaillie cocks recorded in each lek in 2009; **m. a. s. l.** = elevation (meters above the sea level); **Exp.** = exposure in the slope; **Tree Perimeter** = mean trees perimeter measured in DBH (diameter at breast height). **Tree Height** = mean trees height; **Cover** = mean canopy cover (%); **Species** = tree dominant species; **Height =** mean understory height (meters); **Cover** = mean understory cover (%); **Species** = understory dominant specie.

					Trees				Understory		
Lek.	Date	Cocks	m a. s. l.	Exp.	Perimeter (m)	Height (m)	Cover (%)	Species	Height(m)	Cover (%)	Species
1	1998	2	1215	NW	0.45 ± 0.17	9.42 ± 1.38	0.57 ± 0.21	*Q. pyrenaica*	1.12 ± 0.38	0.39 ± 0.25	*C. scoparius*
2	2002	1	1131	NW	0.90 ± 0.62	16.50 ± 3.20	0.58 ± 0.15	*Q. pyrenaica*	0.85 ± 0.07	0.25 ± 0.06	*C. scoparius*
3	2002	0	1154	NW	0.32 ± 0.15	9.17 ± 1.75	0.54 ± 0.26	*Q. pyrenaica*	0.75 ± 0.50	0.16 ± 0.17	*E. arborea*
4	2003	2	1230	NE	0.70 ± 0.13	7.75 ± 0.86	0.28 ± 0.07	*Q. pyrenaica*	1.33 ± 0.24	0.45 ± 0.25	*E. arborea*
5	2004	3	1387	N	0.70 ± 0.15	14.08 ± 2.81	0.64 ± 0.22	*Q. pyrenaica*	0.03 ± 0.08	0.004 ± 0.01	*C. scoparius*
6	2006	1	1253	N	0.20 ± 0.05	9.50 ± 0.42	0.60 ± 0.25	*Q. pyrenaica*	0.50 ± 0.42	0.51 ± 1.41	*E. arborea*
7	2006	0	1356	N	0.23 ± 0.06	11.17 ± 0.38	0.47 ± 0.24	*Q. pyrenaica*	0.79 ± 0.25	0.39 ± 0.57	*E. arborea*
8	2006	2	1351	SE	0.43 ± 0.10	10.33 ± 1.37	0.58 ± 0.15	*Q. pyrenaica*	1.37 ± 0.31	0.40 ± 0.17	*E. arborea*
9	2006	1	1398	SW	0.19 ± 0.05	10.16 ± 0.57	0.81 ± 0.06	*Q. pyrenaica*	1.42 ± 0.45	0.62 ± 0.27	*E. arborea*

Appendix 2. Description of the Eurosiberian leks (n=17) north of the study area (see Fig 1). Mixed in "Tree species" involves no tree species as dominant, but occurring mixed: *Betula pubescens*, *Q. petraea*, *Q. pyrenaica* and *Sorbus aucuparia*.

Lek	m.a.s.l.	Exposure	Tree species	Understory species
1	1440	N-W	*Q. petraea*	*Vaccinium myrtillus*
2	1520	N-W	Mixed	*Vaccinium myrtillus*
3	1450	N-W	Mixed	*Vaccinium myrtillus*
4	1550	N-E	Mixed	*Vaccinium myrtillus*
5	1490	N	*Q. petraea*	*Vaccinium myrtillus*
6	1620	N	*Betula pubescens*	*Erica arborea*
7	1650	N	*Betula pubescens*	*Erica arborea*
8	1550	N	*Betula pubescens*	*Erica arborea*
9	1680	N	*Betula pubescens*	*Erica arborea*
10	1590	N-E	*Betula pubescens*	*Vaccinium myrtillus*
11	1680	N	*Betula pubescens*	*Vaccinium myrtillus*
12	1620	N-W	Mixed	*Vaccinium myrtillus*
13	1540	N-W	Mixed	*Vaccinium myrtillus*
14	1510	N-W	*Betula pubescens*	*Erica arborea*
15	1590	N	*Betula pubescens*	*Erica arborea*
16	1620	N-E	*Betula pubescens*	*Vaccinium myrtillus*
17	1590	N-W	Mixed	*Vaccinium myrtillus*

FIGURE LEGENDS

FIG. 1. Area of occupancy of Cantabrian capercaillie in 1980 (Obeso & Bañuelos 2003) (black left) and study area (right). Numbers correspond to the leks according to the discovery date. Lek 1 was the first capercaillie breeding record in *Q. pyrenaica* forest in 1998. Solid line represents the border between the Eurosiberian and the Mediterranean biogeographical regions after European Environment Agency -EEA- 2008 and Rivas-Martínez et al (2004). (●) represent the recently discovered Mediterranean leks while (▲) represent the traditional Eurosiberian leks. According to the government restrictions, leks location is slightly moved for security. Light and dark grey dominating the area represents *Pinus sylvestris* and *Q. pyrenaica* forests, respectively.

FIG. 2. Occurrence of the Mediterranean leks since 2002. Bars show the total accumulated forest surface surveyed (ha) through the study years. Solid line represents the occupied leks and broken line the known leks.

FIG. 1


Spain
Eurosiberian Region
Mediterranean Region
(Study area)
1
2
3
4
5
6
7
8
9
Mediterranean leks
Eurosiberian leks
Betula pubescens
Quercus petraea
Pinus sylvestris
Quercus pyrenaica
N
0
5
10 Km

FIG. 2


surveyed area (ha)
occupied display grounds
known display grounds
Acumulated surveyed area (ha)
No. of leks
Year
0
500
1000
1500
2000
2500
3000
3500
4000
4500
5000
0
1
2
3
4
5
6
7
8
9
10
2002
2003
2004
2005
2006
2007
2008
2009