



**HAL**  
open science

## **The reciprocal influences of asthma and obesity on lung function testing, AHR and airway inflammation in prepubertal children.**

Nicola Pietro Consilvio, Sabrina Di Pillo, Marcello Verini, Tommaso de Giorgis, Anna Cingolani, Valentina Chiavaroli, Francesco Chiarelli, Angelika Mohn

### ► **To cite this version:**

Nicola Pietro Consilvio, Sabrina Di Pillo, Marcello Verini, Tommaso de Giorgis, Anna Cingolani, et al.. The reciprocal influences of asthma and obesity on lung function testing, AHR and airway inflammation in prepubertal children.. *Pediatric Pulmonology*, 2010, 45 (11), pp.1103. <10.1002/ppul.21295>. <hal-00588663>

**HAL Id: hal-00588663**

**<https://hal.science/hal-00588663v1>**

Submitted on 26 Apr 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



**The reciprocal influences of asthma and obesity on lung function testing, AHR and airway inflammation in prepubertal children.**

Journal:	<i>Pediatric Pulmonology</i>
Manuscript ID:	PPUL-09-0317.R1
Wiley - Manuscript type:	Original Article
Date Submitted by the Author:	06-Mar-2010
Complete List of Authors:	<p>Consilvio, Nicola; University G. D'Annunzio, Department of Paediatrics                  Di Pillo, Sabrina; University G. D'Annunzio, Department of Paediatrics                  Verini, Marcello; University G. D'Annunzio, Department of Paediatrics                  de Giorgis, Tommaso; University G. D'Annunzio, Department of Paediatrics                  Cingolani, Anna; University G. D'Annunzio, Department of Paediatrics                  Chiavaroli, Valentina; University G. D'Annunzio, Department of Paediatrics                  Chiarelli, Francesco; University G. D'Annunzio, Department of Paediatrics                  Mohn, Angelika; University G. D'Annunzio, Department of Paediatrics</p>
Keywords:	Childhood Asthma, Obesity, Fractional exhaled Nitric Oxide, Lung Function Testing, Airways Hyperresponsiveness



1  
2  
3 **The reciprocal influences of asthma and obesity on lung function**  
4 **testing, AHR and airway inflammation in prepubertal children.**  
5  
6  
7  
8

9  
10 *Nicola P. Consilvio, MD, Sabrina Di Pillo, MD, Marcello Verini, MD, Tommaso de Giorgis, MD,*  
11 *Anna Cingolani, MD; Valentina Chiavaroli, MD, Francesco Chiarelli, PhD; Angelika Mohn, MD.*  
12  
13

14  
15  
16  
17 Department of Pediatrics, University of Chieti, Chieti, Italy.  
18

19  
20  
21  
22 **Short title:** Impact of asthma and obesity on lung function testing.  
23  
24  
25  
26  
27

28  
29 **Word count:** 3223

30 **Funding:** none

31  
32 **Conflict of Interest:** none  
33  
34  
35  
36  
37  
38  
39  
40

41 **Keywords:** Obesity, Asthma, Fractional exhaled Nitric Oxide, Lung Function Testing, Airways  
42 Hyperresponsiveness.  
43  
44  
45  
46  
47  
48  
49

50 **Correspondence:**

51 Angelika Mohn  
52 Department of Pediatrics  
53 University of Chieti  
54 Via Dei Vestini 5  
55 66100 Chieti  
56 Tel +39 0871 358827  
57 Fax +39 0871 584731  
58 Email amohn@unich.it  
59  
60

## SUMMARY

Although asthma and obesity are among the major chronic disorders their reciprocal or independent influences on lung function testing, airways hyperresponsiveness (AHR) and bronchial inflammation has not been completely elucidated.

In 118 pre-pubertal Caucasian children anthropometric measurements functional respiratory parameters (flow/volume curves at baseline and after 6-Minute Walk Test [6MWT]) together with bronchial inflammatory index (FeNO) were assessed. The study population was divided into four groups according to BMI and the presence or absence of asthma: Obese asthmatic (*ObA*) Normal-weight asthmatic (*NwA*), Obese non asthmatic (*Ob*), non-asthmatic Normal-weight children (*Nw*).

Baseline PEF and MEF<sub>75</sub> (%-expected) were significantly different across the four groups with significantly lower values of MEF<sub>75</sub> in ObA and Ob children when compared to Nw children ( $p=0.004$  and  $p=0.0001$ , respectively) and this independent role of obesity on upper respiratory flows was confirmed by multiple analysis of covariance.

After 6 MWT respiratory parameters decreased only in ObA and NwA children and 12 children presented a positive fall in FEV<sub>1</sub>, in contrast no changes of respiratory function testing were detected in Ob and Nw children, and only 2 Ob children presented a significant fall in FEV<sub>1</sub>.

FeNO analysis demonstrated significantly higher values in ObA and NwA children when compared to Ob ( $p=0.008$  and  $p=0.0002$ , respectively) and Nw children ( $p=0.0001$  and  $p=0.0003$ , respectively), although a significant difference was found between Ob and Nw children ( $p=0.0004$ ).

Multiple analysis of covariance confirmed an independent role of asthma on this parameter.

In conclusion while AHR and airway inflammation are clearly associated with an asthmatic status, obesity seems to induce reduction of upper airways flows associated with a certain degree of pro-inflammatory changes.

## INTRODUCTION

Over several decades the prevalence of obesity has steadily increased reaching epidemic proportions. Current estimates of childhood obesity coming from the National Health and Nutrition Examination Survey (NHANES) revealed an increase from 13.8% in 1999-2000 to 16.0% in 2003-2004 for American females and from 14.0% to 18.2% in American males subjects.<sup>1</sup> In Europe, Italy together with other Mediterranean countries explicates the highest rate of obesity in children aged between 7 and 11 years, reaching almost 17%.<sup>2</sup> This has been accompanied by a similar rise in the rates of asthma. In fact, the CDC National Surveillance for Asthma revealed that its prevalence in children has risen from 3.5% to 7.5% over a period ranging from 2001-2003.<sup>3</sup> Similarly in developing countries undergoing their epidemiological transition phase comparable significant rises have been observed.<sup>4</sup> Accordingly, the World Health Organization (WHO) includes both obesity and asthma among the major chronic disorders representing world-wide public health priorities.<sup>5</sup>

Given this dramatic rise in the prevalence of obesity and asthma, it is not surprising that there has been an increasing body of literature on the association between Body Mass Index (BMI) and asthma.<sup>6</sup>

Obesity may have effects on both the upper and lower airway,<sup>7</sup> and this hypothesis is supported to a certain extent by the observation that BMI, the best single measure of body adiposity,<sup>8</sup> seems to be a reliable predictor of allergic respiratory symptoms and airways hyperresponsiveness (AHR).<sup>9</sup> In fact, recent literature emphasises the role of obesity in inducing a pro-inflammatory state due to increased levels of different pro-inflammatory cytokines. This might be implicated in the genesis of chronic inflammation of the airway system<sup>10</sup> similar to those observed in asthma which is characterized by persistent infiltration of eosinophils, T-lymphocytes and mast cells combined with different cytokines. In this respect exhaled Nitric Oxide (eNO) has been shown to tightly reflect this airway inflammation<sup>11</sup> and its fractional concentration in exhaled air (FeNO) has been recently accepted by the PRACTALL Consensus Report for the Diagnosis, Management and Treatment of Childhood Asthma as a logical complementary item in the follow-up of bronchial inflammation.<sup>12</sup>

1  
2  
3 The reported association between increased BMI and onset of AHR has been elegantly  
4 demonstrated by Litonjua et al.<sup>13</sup> in a longitudinal study of aging and has been further confirmed by  
5 two large cross-sectional studies performed in adults.<sup>14,15</sup> This is in contrast with other  
6 epidemiological studies performed in childhood where this association has not been universally  
7 observed.<sup>16,17,18</sup> This represents an important discordance of the recent literature as it might indicate  
8 that different factors such as duration of obesity and subsequent physiological changes of airway  
9 structures might play an important role in the development of obesity induced AHR.  
10  
11  
12  
13  
14  
15  
16  
17  
18

19 Pre-pubertal children are a special study population as the shorter duration combined with better  
20 control of both diseases eliminates confounding factors commonly present in older subjects. In fact,  
21 this kind of population offers the exclusive opportunity to tangle whether obesity and asthma  
22 explicate an independent effect on lung function testing or influence themselves in a reciprocal  
23 manner.  
24  
25  
26  
27  
28  
29  
30

31 Therefore, with the primary aim to establish the possible dependent or independent role of asthma  
32 and obesity on airway system function we evaluated AHR and bronchial inflammation expressed by  
33 FeNO in a group of obese and non obese asthmatic children in comparison to non-asthmatic obese  
34 and normal-weight pre-pubertal children.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## MATERIALS AND METHODS

### Study subjects

Clinical, anthropometric measurements and respiratory parameters of a group of 118 Caucasian pre-pubertal children referred to the Children Obesity Clinic, to the Paediatric Allergy and Respiratory Clinic and to the Department of Paediatrics, University of Chieti, Italy, were sequentially evaluated between May 2008 and August 2008.

Asthma was classified according to Global Initiative for Asthma (GINA) guidelines<sup>19</sup> as intermittent asthma, mild persistent, moderate persistent and severe persistent asthma. The diagnosis was established by a single paediatric respiratory physician on the basis of a clinical history of repeated episodes of coughing, dyspnoea and wheezing, according to American Thoracic Society - European Respiratory Society (ATS-ERS) criteria.<sup>20</sup> Only subjects who fulfilled the definition of intermittent/mild persistent asthma were recruited. Low dose inhaled corticosteroids (ICSs) therapy was withdrawn for not more than weeks before the study evaluation. Over this period all children were closely monitored by the medical staff of the Paediatric Allergy and Respiratory Clinic, none experienced any adverse event and at the time of the study evaluation all subjects were symptom-free.

Obesity was defined as Body Mass Index [BMI] above 2 Standard Deviations (SD) for the mean of age and sex.<sup>21</sup> In all subjects endocrine diseases as cause for obesity were excluded prior to study entry. None of the non-asthmatic obese children had a personal history for asthma and allergic diseases.

We recruited as controls a group of healthy children, comparable for age, gender, and pubertal stage, who were admitted to the Department of Paediatrics for minor diseases. None of the children had a personal history for asthma and allergic diseases, and all were otherwise in a good health. Anthropometric measurement and respiratory function were assessed only after complete recovery of the disease.

1  
2  
3 Exclusion criteria for all children were: presence of upper or lower respiratory infections including  
4  
5  
6 asthma during the previous 4 weeks, congenital abnormalities of cardio-respiratory system, chest or  
7  
8 skeletal deformities, or neuromuscular system disease, chronic and autoimmune disease, gastro-  
9  
10 oesophageal reflux, presence of medical condition in which exercise was forbidden. None of the  
11  
12 children was taking any medication.

13  
14  
15 Based on the presence or absence of asthma and according to BMI, the study population was  
16  
17 divided into four groups:

- 18  
19 1. *Obese Asthmatic Children [ObA]* (sixteen boys and 14 girls);
- 20  
21 2. *Normal-weight Asthmatic Children [NwA]* (sixteen boys and 14 girls);
- 22  
23 3. *Non-Asthmatic Obese Children [Ob]* (fourteen boys and 16 girls);
- 24  
25 4. *Non-Asthmatic Normal-weight Children [Nw]* (fourteen boys and 14 girls).

26  
27  
28 All children underwent a complete physical examination including anthropometric parameters  
29  
30 [height, weight]. Pubertal stage was defined on the basis of breast development in girls and genital  
31  
32 development in boys according to the criteria of Tanner (all subjects had pre-adolescent  
33  
34 characteristics corresponding to stage 1). Respiratory function was assessed by standard  
35  
36 Flow/Volume curves at baseline and after Six-Minute Walk Test (6MWT), and respiratory  
37  
38 inflammatory status was assessed by FeNO. Only asthmatic children performed allergy evaluation  
39  
40 by Skin Prick Test (SPT), and blood sample was taken for specific IgE measurement.

41  
42  
43 Written informed consent was obtained from all parents and oral consent from all children.

44  
45  
46 The study was approved by Ethical Committee of University of Chieti.

## 47 48 49 50 51 52 53 **Methods**

### 54 55 *Anthropometric measurements*

56  
57 Body weight was determined to the nearest 0.1 Kg, and height was measured in triplicate with  
58  
59 Harpenden stadiometer to the nearest 0.1 cm. As fatness indices, we used BMI (the weight in  
60 kilograms divided by the square of the height in meters). SDS-height and BMI-SDS were calculated

1  
2  
3 as difference between the individual observed value and the reference mean for age and sex, divided  
4  
5 by the corresponding standard deviation.<sup>22</sup>  
6  
7

### 8 *Spirometric and Flowmetric Measurements:*

#### 9 *Baseline Flow/Volume curves*

10 Lung function evaluation was performed by Flow/Volume curves (KitMicro Spirometer 7.3a –  
11 COSMED) in the standing position with nose-clip until two consecutive technically acceptable  
12 curves were achieved, according to ATS/ERS Guidelines.<sup>23</sup> The main spirometric parameters  
13 included: Forced Expiratory Volume in 1 sec (FEV<sub>1</sub>), Forced Vital Capacity (FVC), FEV<sub>1</sub>/FVC,  
14 and Forced Expiratory Flows between 25% and 75% of the FVC (FEF<sub>25-75</sub>) and the results are  
15 expressed as Z-scores according to Stanojevic S et al.<sup>24</sup> Peak Expiratory Flow (PEF) and Maximal  
16 Expiratory Flows at 75% of the FVC (MEF<sub>75</sub>) were expressed as a percentage of theoretical values  
17 for age, gender, and height according to Quanjer PH et al.<sup>25</sup>  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

#### 31 *Six-Minute Walk Test (6MWT)*

32 In order to estimate the bronchial hyperresponsiveness, after baseline lung function testing, a  
33 6MWT was performed according to standard protocol.<sup>26</sup> The target of the test was to achieve 6  
34 minutes of free walk until the hearth-rate was 80-90% of the predicted maximum (calculated as 220  
35 – age in years), according to ATS/ERS recommendations for exercise challenge test in older  
36 children.<sup>27</sup> Spirometry was performed in duplicate sets at 1, 3, 5, 10 and 20 minutes post-exercise.  
37 The better of the two curves was selected as the representative value at each interval, but  
38 differences between the two values of FEV<sub>1</sub> had to be <5%. Exercise response was defined as the  
39 greatest decrease in FEV<sub>1</sub> expressed as a percentage of the baseline values. A bronchial obstruction  
40 response to exercise was considered positive when the FEV<sub>1</sub> reduction from baseline was greater  
41 than 12%.<sup>28</sup>  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56

#### 57 *Allergic sensitization*

58 Skin Prick Test (SPT) were performed and serum specific IgE were measured for the most common  
59 respiratory allergens: Dust Mite (*Dermatophagoides Pteronyssinus* and *Farinae*), Grass, Parietaria,  
60

1  
2  
3 Artemisia Vulgaris, Olive, Cypress, Lime, Stone, Elm, Plane, Cat and Dog dander, Alternaria  
4  
5 Alternata and Aspergillus Fumigatus (moulds). Determination of Allergen-Specific IgE was made  
6  
7 by an Immunoenzymatic Allergo-sorbent Test (Cap test Pharmacia).<sup>29,30</sup>  
8  
9

#### 10 *Respiratory inflammation marker*

11  
12 FeNO was determined with an on-line method using a single breath exhalation and a sensitive  
13  
14 chemiluminescence assay (Ecomedics CLD 88), according to ATS-ERS.<sup>31</sup> Patients made an  
15  
16 inspiration of eNO-free air via a mouthpiece immediately followed by full exhalation at a constant  
17  
18 rate (50 ml/sec) for at least 5 seconds. The mean of three readings at the end of the expiration  
19  
20 (plateau phase) was taken as the representative value for each measurements. 12 ppb or more were  
21  
22 considered positive, according to ATS-ERS criteria.<sup>32</sup>  
23  
24  
25  
26  
27  
28

#### 29 **Statistical analysis**

30 All values were expressed as Median and Interquartile Range.

31  
32  
33 Kruskal Wallis Test was performed for evaluating differences between the four groups (ObA, NwA,  
34  
35 Ob, Nw). Statistical significance level was  $p < 0.05$ . Subsequently, differences within the single  
36  
37 groups were tested by Mann-Whitney Test, adjusting the significance level by  $p < 0.008$ . Differences  
38  
39 in functional respiratory baseline parameters and after 6MWT within each group were analyzed  
40  
41 using Wilcoxon Signed Ranks Test. Categorical variables were tested by Pearson's chi-square ( $X^2$ )  
42  
43 test.  
44  
45  
46  
47

48 A multiple analysis of covariance (MANCOVA) was performed to investigate the independent  
49  
50 effect of obesity and asthma on respiratory indices (PEF, MEF<sub>75</sub> and FEV<sub>1</sub>/FVC) and bronchial  
51  
52 inflammation (FeNO), after adjusting for gender, age and height.  
53  
54

55 Statistical significance level was  $p < 0.05$ .

56  
57 All calculations were made with the computer program SPSS (Statistical Package for the Social  
58  
59 Science), version 16.0 software for Windows.  
60

## RESULTS

Anthropometric measurements and respiratory functional data of the whole study population are reported in Table 1. The four groups were similar for age, gender, and SDS-height.

As expected BMI-SDS was similar between ObA and Ob children, but was significantly different between obese children and normal-weight children (ObA vs NwA  $p=0.0004$ ; ObA vs Nw:  $p=0.0008$ ; Ob vs NwA  $p=0.0001$ ; Ob vs Nw:  $p=0.0002$ ).

All the asthmatic children were sensitized to *Dermatophagoides Pteronyssinus*.

### *Baseline Respiratory Functional Parameters*

No significant differences were found for FVC, FEV<sub>1</sub>, and FEF<sub>25-75</sub> Z-scores between the four groups, whereas FEV<sub>1</sub>/FVC Z-score resulted significantly different (Tab 1). In fact, this parameter was significantly lower in the ObA group compared to the Nw group ( $p=0.005$ ). Moreover a significant difference was detected between the four groups in terms of PEF ( $p=0.03$ ) and MEF<sub>75</sub> ( $p=0.001$ ) (Fig 1). By comparing the four groups directly no further significances could be detected for PEF, whereas MEF<sub>75</sub> was significantly lower in the ObA group and the Ob group compared to the Nw group ( $p=0.004$  and  $p=0.0001$ , respectively).

### *Respiratory Functional Parameters after 6MWT*

After 6MWT no significant differences were found in post-FVC and post-FEF<sub>25-75</sub> Z-scores among the four groups (Tab 1). Although post-FEV<sub>1</sub>/FVC Z-scores resulted significant different between the four groups, no further differences were detected by comparing the four groups directly. In contrast, post-FEV<sub>1</sub> Z-score resulted significantly lower in NwA children when compared to Ob children and Nw children ( $p=0.006$  and  $p=0.006$ , respectively). Post-PEF and post-MEF<sub>75</sub> resulted significantly different between the four groups ( $p=0.01$ , both). In fact, post-PEF was significantly lower in ObA children when compared to Nw children ( $p=0.002$ ). This was associated with significantly lower values of post-MEF<sub>75</sub> in the ObA group and the Ob group when compared to the Nw group ( $p=0.003$  and  $p=0.007$ , respectively).

1  
2  
3 Changes in respiratory parameters after 6MWT within each group are shown in Table 2. Post-  
4 FEV<sub>1</sub>/FVC Z-score was significantly lower in the ObA group, while in NwA children a significant  
5 reduction was detected for post-FEV<sub>1</sub>, post-FVC and post-FEF<sub>25-75</sub> Z-scores. In contrast no changes  
6 were detected for the Ob and Nw groups.  
7  
8  
9

10  
11 A positive response to exercise challenge was found in 5 ObA children and 7 NwA children  
12 compared to only 2 Ob children. As expected none of the healthy control children experienced a fall  
13 in FEV<sub>1</sub> greater than 12%.  
14  
15  
16  
17

#### 18 19 *Respiratory inflammation marker*

20 FeNO values were significantly different across all four groups (p=0.0002) (Tab. 1).  
21

22 In fact FeNO values were significantly higher in the ObA group and the NwA group compared to  
23 the Ob group (p=0.008 and p=0.0002, respectively) and to the Nw group (p=0.0001 and p=0.0003,  
24 respectively).  
25  
26  
27  
28  
29

30 Furthermore FeNO was significantly higher in the Ob group when compared to the Nw group  
31 (p=0.0004). In contrast no differences were detected between the two asthmatic groups.  
32  
33

#### 34 35 *Effect of asthma and obesity on respiratory functional parameters and bronchial inflammation*

36 In order to investigate the role of asthma and obesity on resulted significant different baseline  
37 respiratory variables and FeNO, a multiple analysis of covariance was performed. Asthma was  
38 significantly and independently related to FeNO values, whereas no association was found for  
39 obesity. In contrast obesity was significantly and independently related to PEF and MEF<sub>75</sub>. In the  
40 same model no relationship was found for asthma and obesity on FEV<sub>1</sub>/FVC Z-score (Table 3).  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## DISCUSSION

Although the effect of obesity on pulmonary function tests has been extensively investigated in adults, the number of similar studies in children is limited and confirm the controversial results observed in adults between obesity and AHR. In this study we compare for the first time obese asthmatic with obese non-asthmatic pre-pubertal children and healthy controls with the aim to analyse the reciprocal influences of obesity and asthma on lung function and bronchial inflammation. Our findings clearly demonstrate that while an asthmatic status is tightly and directly involved in AHR and bronchial inflammation, obesity mainly affects the upper airways flows.

As obese children often experience dyspnoea and breathlessness during physical activities it is a crucial clinical question as to whether these symptoms are effectively related to AHR such as in asthmatic children. In order to explore this possibility a 6MWT was performed. While basal values of spirometric indices were normal in all examined groups, in ObA and NwA children exercise challenge induced AHR documented by the reduction of functional respiratory parameters. In contrast in Ob and Nw children no changes of any of the obtained variables were detected. This is confirmed by the number of children exhibiting a positive response to exercise defined as fall in FEV1 greater than 12%. In fact within the asthmatic children 12 subjects met this definition compared to only 2 obese children. In agreement with our data, epidemiological studies do not uniformly support the conclusions that obesity leads to airflow limitation, AHR or asthma. In this respect Schachter and co-workers showed in a group of 1971 adults, that although BMI was associated with a clinical diagnosis of asthma and subjective symptoms of dyspnoea and wheeze, airflow obstruction or AHR subsequent to histamine challenge could not be detected in the sole obese subjects.<sup>16</sup> Similar findings were obtained by Bibi et al.<sup>17</sup> who demonstrated in a large population of pre-pubertal children that bronchodilation could not be induced by salbutamol inhalation in non-asthmatic obese subjects.<sup>33</sup> These data demonstrate that obesity should not be considered as being a risk factor for AHR in pre-pubertal children. Furthermore asthma like symptoms especially during physical activity should not be overestimated by parents and medical

1  
2  
3 staff. This is of paramount relevance as it might induce inappropriate treatment and refrain children  
4  
5 from performing sports leading to a vicious circle impeding improvement of both obesity and  
6  
7 respiratory symptoms.  
8  
9

10 Interestingly in the ObA and Ob groups we documented lower basal values of  $MEF_{75}$  when  
11  
12 compared to healthy controls. Furthermore PEF values were significantly different across the four  
13  
14 groups with a tendency of lower values in the two obese groups. In fact, by performing a multiple  
15  
16 analysis of covariance obesity was significantly and independently related to  $MEF_{75}$  and to PEF,  
17  
18 while this association was not found for asthma. Similar data were documented by two previous  
19  
20 studies<sup>34,35</sup> where obesity was associated with a significant reduction in  $FEF_{25-75}$ . These data  
21  
22 suggests that obesity might play a primary role on upper airway flows, and despite being possibly  
23  
24 explained by fatty infiltration of muscles narrowing directly the upper airway, recent literature  
25  
26 suggest that this functional parameter might be used as an early predictor of sub-clinical asthma.<sup>36</sup>  
27  
28  
29

30 In order to obtain a complete evaluation of the bronchial status we performed also FeNO analysis.  
31  
32 In effect, recent studies revealed that FeNO is a potentially useful measure to evaluate the role of  
33  
34 airways inflammation in asthma, as it represents the forerunner of the most important event namely  
35  
36 remodelling of bronchial airways,<sup>37</sup> offering therefore a logical complementary item to the only use  
37  
38 of clinical and functional data. In our study, as expected, both groups of asthmatic children  
39  
40 presented highly elevated FeNO values when compared to Ob children and normal controls  
41  
42 reflecting the underlying bronchial inflammation process. However, data from animal and human  
43  
44 studies clearly demonstrate that the enhancement of normal adipose tissue in obesity leads to a  
45  
46 systemic pro-inflammatory state.<sup>38</sup> Adipose tissue from obese individuals expresses a number of  
47  
48 pro-inflammatory molecules, such as leptin, tumor necrosis factor- $\alpha$ , IL-6, transforming growth  
49  
50 factor- $\beta$ 1, and C-reactive protein, and there appears to be significant overlap between the immune  
51  
52 function of adipocytes and the function of T-lymphocytes and macrophages, particularly with  
53  
54 respect to the elaboration of inflammatory cytokines.<sup>39,40</sup> Although a conclusive relationship  
55  
56 between obesity and airway inflammation has not yet to been described, it could be argued that the  
57  
58  
59  
60

1  
2  
3 role of the pro-inflammatory state underlying the main metabolic and cardiovascular complications  
4  
5 in obesity might play a pivotal role in determining airways inflammation independently of any  
6  
7 allergic mechanisms. This might explain the findings of our study where FeNO values were  
8  
9 significantly higher in Ob children when compared to healthy controls, and the findings of  
10  
11 Santamaria et al.<sup>41</sup> who found a positive correlation between FeNO and BMI in healthy children.  
12  
13 However it needs to be acknowledged that in the multiple analysis of covariance obesity was not a  
14  
15 predictor for increased FeNO values as only asthmatic status remained as such. This apparent  
16  
17 contrasting findings might be partially explained by the age group of our study population as  
18  
19 obesity related inflammation might not have reached its full implication and a possible involvement  
20  
21 of the respiratory system might require a long standing history of obesity.  
22  
23  
24  
25

26  
27 In this respect an important limitation of this study is the cross sectional design which does not  
28  
29 allow to speculate on the full clinical implications of our findings. In fact, as puberty is a well  
30  
31 known risk factor for developing obesity related complications it might be of crucial interest to  
32  
33 obtain longitudinal data starting in pre-puberty and lasting through all pubertal stages. This would  
34  
35 allow to fully explore this intriguing relationship and to not underestimate the possible role of  
36  
37 obesity on the development of bronchial inflammation.  
38  
39

40  
41 In conclusion, while AHR and airways inflammation are clearly associated to an asthmatic status,  
42  
43 obesity seems to induce reduction of upper airway flows combined with a certain degree of early  
44  
45 pro-inflammatory airways changes. Longitudinal data are required in order to clarify whether these  
46  
47 changes might represent risk factor for asthma later on in life.  
48  
49

## 50 51 52 53 54 55 **Acknowledgements**

56  
57 We thank all nurses of the Department of Pediatrics, University of Chieti, in particular Di Santo D,  
58  
59 Di Luca D, Diomedi A, for their daily support in clinical research and care of children.  
60

1  
2  
3 **REFERENCES**  
4

- 5  
6 1. Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of  
7  
8 overweight and obesity in the United States, 1999-2004. *JAMA* 2006;295(13):1549-1555.  
9  
10 2. Binkin N, Fontana G, Lamberti A, Cattaneo C, Baglio G, Perra A, Spinelli A. A national  
11  
12 survey of the prevalence of childhood overweight and obesity in Italy. *Obes Rev* 2009 Sep  
13  
14 17.  
15  
16 3. Moorman JE, Rudd RA, Johnson CA, King M, Minor P, Bailey C, Scalia MR, Akinbami  
17  
18 LJ; Centers for Disease Control and Prevention (CDC). *MMWR Surveill Summ*  
19  
20 2007;56(8):1-54.  
21  
22 4. World Health Organization. Obesity: preventing and managing the global epidemic. WHO  
23  
24 Technical Report Series 894. Geneva, Switzerland, WHO, 2000.  
25  
26 5. World Health Organization. Preventing chronic diseases: a vital investment. Geneva,  
27  
28 Switzerland: WHO, 2005.  
29  
30 6. Figueroa-Munoz JI, Chinn S, Rona RJ. Association between obesity and asthma in 4-11  
31  
32 year old children in the UK. *Thorax* 2001;56:133-137.  
33  
34 7. Deane S, Thomson A. Obesity and the pulmonologist. *Arch Dis Child* 2006;91(2):188-191.  
35  
36 8. Power C, Lake JK, Cole TJ. Measurement and long term health risks of child and adolescent  
37  
38 fatness. *Int J Obes Relat Metab Disord* 1997;21:507-526.  
39  
40 9. Gennuso J, Epstein LH, Paluch RA, Cerny F. The relationship between asthma and obesity  
41  
42 in urban minority children and adolescent. *Arch Pediatr Adolesc Med* 1998;152(12):1197-  
43  
44 1200.  
45  
46 10. Visser M, Bouter LM, McQuillan GM, Wener MH, Harris TB. Elevated C-reactive protein  
47  
48 levels in overweight and obese adults. *JAMA* 1999;282:2131-5.  
49  
50 11. Payne DN, Adcock IM, Wilson NM, Oates T, Scallan M, Bush A. Relationship between  
51  
52 exhaled nitric oxide and mucosal eosinophilic inflammation in children with difficult  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 asthma, after treatment with oral prednisolone. *Am J Respir Crit Care Med* 2001;164:1376-  
4  
5 1381.  
6  
7  
8 12. Bacharier LB, Boner A, Carlsen K-H, Eigenmann PA, Frischer T, Götz M, Helms PJ, Hunt  
9  
10 J, Liu A, Papadopoulos N, Platts-Mills T, Pohunek P, Simons FE, Valovirta E, Wahn U,  
11  
12 Wildhaber J; European Pediatric Asthma Group. Diagnosis and treatment of asthma in  
13  
14 childhood: a PRACTALL Consensus Report. *Allergy* 2008;63:5-34.  
15  
16  
17 13. Litonjua AA, Sparrow D, Celedon JC, DeMolles D, Weiss ST. Association of body mass  
18  
19 index with the development of methacholine airway hyperresponsiveness in men: the  
20  
21 Normative Aging Study. *Thorax* 2002;57:581-585.  
22  
23  
24 14. Chinn S, Jarvis D, Burney P. Relation of bronchial responsiveness to body mass index in the  
25  
26 ECRHS. European Community Respiratory Health Survey. *Thorax* 2002;57:1028-1033.  
27  
28  
29 15. Celedon JC, Palmer LJ, Litonjua AA, Weiss ST, Wang B, Fang Z, Xu X. Body mass index  
30  
31 and asthma in adults in families of subjects with asthma in Anqing, China. *Am J Respir Crit*  
32  
33 *Care Med* 2001;164:1835-1840.  
34  
35  
36 16. Schachter LM, Salome CM, Peat JK, Woolcock AJ. Obesity is a risk for asthma and wheeze  
37  
38 but not airway hyperresponsiveness. *Thorax* 2001;56:4-8.  
39  
40  
41 17. Bibi H, Shoseyov D, Feigenbaum D, Genis M, Friger M, Peled R, Sharff S. The relationship  
42  
43 between asthma and obesity in children: is it real or a case of over diagnosis? *J Asthma*  
44  
45 2004;41(4):403-410.  
46  
47  
48 18. Wickens K, Barry D, Friezema A, Rhodius R, Bone N, Purdie G, Crane J. Obesity and  
49  
50 asthma in 11-12 year old New Zealand children in 1989 and 2000. *Thorax* 2005;60:7-12.  
51  
52  
53 19. Global Strategy for Asthma Management And Prevention, Global Initiative for Asthma  
54  
55 (GINA) World Report 2006. Available from: <http://www.ginasthma.org>.  
56  
57  
58 20. National Asthma Education and Prevention Program. NAEPP expert panel report guidelines  
59  
60 for the diagnosis and management of asthma – update on selected topics 2002. National  
Institutes for Health 2006.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
21. Kiess W, Reich A, Müller G, Galler A, Kapellen T, Raile K, Böttner A, Seidel B, Kratzsch J. Obesity in childhood and adolescence: clinical diagnosis and management. *J Pediatr Endocrinol Metab* 2001;14(6):1431-1440.
  22. Lindgren G, Strandell A, Cole T, Healy M, Tanner J. Swedish population standards for height, weight and body mass index attained at 6 to 16 years (girls) or 19 years (boy). *Acta Paediatr* 1995;84:1019-1028.
  23. Laszlo G. Standardisation of lung function testing: helpful guidance from the ATS/ERS Task Force. *Thorax* 2006;61(9):744-746.
  24. Stanojevic S, Wade A, Stocks J, Hankinson J, Coates AL, Pan H, Rosenthal M, Corey M, Lebecque P, Cole TJ. Reference Ranges for Spirometry Across All Ages: A New Approach. *Am J Respir Crit Care Med* 2008;177:253-260.
  25. Quanjer PH, Borsboom GJ, Brunekreef B, Zach M, Forche G, Cotes JE, Sanchis J, Paoletti P. Spirometric reference values for white European children and adolescents: Polgar revisited. *Pediatr Pulmonol* 1995;19(2):135-142.
  26. ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. ATS statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med* 2002;166:111-117.
  27. Crapo RO, Casaburi R, Coates AL, Enright PL, Hankinson JL, Irvin CG, MacIntyre NR, McKay RT, Wanger JS, Anderson SD, Cockcroft DW, Fish JE, Sterk PJ. Guidelines for Methacholine and Exercise Challenge Testing-1999. This official statements of the American Thoracic Society was adopted by the ATS Board of Directors, July 1999. *Am J Respir Crit Care Med* 2000;161:309-329.
  28. Godfrey S, Springer C, Bar-Yishay E, Avital A. Cut-off points defining normal and asthmatic bronchial reactivity to exercise and inhalation challenges in children and young adults. *Eur Respir J* 1999;14:659-666.

- 1  
2  
3 29. Terho E, Frew AJ. Type III allergy skin testing. Position statement for EAACI  
4  
5 Subcommittee on Skin Tests and Allergen Standardization. *Allergy* 1995;50(5):392-396.  
6  
7  
8 30. Wolkerstorfer A, Wahn U, Kjellman NI, Diepgen TL, De Longueville M, Oranje AP.  
9  
10 Natural course of sensitization to cow's milk and hen's egg in childhood atopic dermatitis:  
11  
12 ETAC Study Group. *Clin Exp Allergy* 2002;32(1):70-73.  
13  
14  
15 31. American Thoracic Society; European Respiratory Society. ATS/ERS recommendations for  
16  
17 standardized procedures for the online and offline measurement of exhaled lower respiratory  
18  
19 nitric oxide and nasal nitric oxide, 2005. *Am J Respir Crit Care Med* 2005;171(8):912-930.  
20  
21  
22 32. Baraldi E, de Jongste JC; European Respiratory Society; American Thoracic Society.  
23  
24 Measurement of exhaled nitric oxide in children, 2001. *Eur Respir J* 2002;20(1):223-237.  
25  
26  
27 33. Bibi H, Montgomery M, Pasterkamp H, Chernick V. Relationship between response to  
28  
29 inhaled salbutamol and methacholine bronchial provocation in children with suspected  
30  
31 asthma. *Pediatr Pulmonol* 1991;10:244-248.  
32  
33  
34 34. Spathopoulos D, Paraskakis E, Trypsianis G, Tsalkidis A, Arvanitidou V, Emporiadou M,  
35  
36 Bouros D, Chatzimichael A. The Effect of Obesity on Pulmonary Lung Function of School  
37  
38 Aged Children in Greece. *Pediatr Pulmonol* 2009;44:273-280.  
39  
40  
41 35. Ülger Z, Demir E, Tanaç R, Gökşen D, Gülen F, Darcan Ş, Can D, Coker M. The effect of  
42  
43 childhood obesity on respiratory function tests and airway hyperresponsiveness. *The*  
44  
45 *Turkish Journal of Pediatrics* 2006;48:43-50.  
46  
47  
48 36. Drewek R, Garber E, Stanclik S, Simpson P, Nugent M, Gershan W. The FEF<sub>25-75</sub> and its  
49  
50 decline as a predictor of methacholine responsiveness in children. *J Asthma* 2009;46(4):375-  
51  
52 381.  
53  
54  
55 37. van Veen IH, Sterk PJ, Schot R, Gauw SA, Rabe KF, Bel EH. Alveolar nitric oxide versus  
56  
57 measures of peripheral airway dysfunction in severe asthma. *Eur Respir J* 2006;27(5):951-  
58  
59 956.  
60

- 1  
2  
3 38. Fantuzzi G. Adipose tissue, adipokines, and inflammation. *J Allergy Clin Immunol*  
4 2005;115:911-919.  
5  
6  
7  
8 39. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW Jr. Obesity is  
9 associated with macrophage accumulation in adipose tissue. *J Clin Invest* 2003;112:1796-  
10 1808.  
11  
12  
13 40. Wellen KE, Hotamisligil GS. Obesity-induced inflammatory changes in adipose tissue. *J*  
14 *Clin Invest* 2003;112:1785-1788.  
15  
16  
17 41. Santamaria F, Montella S, De Stefano S, Sperli F, Barbarano F, Valerio G. Relationship  
18 between exhaled nitric oxide and body mass index in children and adolescent. *J Allergy Clin*  
19 *Immunol* 2005;116:1163-1164.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **The reciprocal influences of asthma and obesity on lung function in**  
4 **prepubertal children.**  
5  
6  
7

8  
9  
10 *Consilvio Nicola Pietro, MD; Di Pillo Sabrina, MD; Verini Marcello, MD; Tommaso de Giorgis,*  
11 *MD; Anna Cingolani, MD; **Fabio Masuccio, MD**; Francesco Chiarelli, PhD; Angelika Mohn, MD.*  
12  
13

14  
15  
16  
17 Department of Pediatrics, University of Chieti, Chieti, Italy.  
18

19  
20  
21  
22 **Short title:** *Asthma and obesity impact on lung function.*  
23  
24  
25  
26  
27

28  
29 **Word count:** 2919  
30

31 **Funding:** none  
32

33 **Conflict of Interest:** none  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 **Keywords:** *Body Mass Index-Standard Deviation Score*, obesity, asthma, Fractional exhaled Nitric  
44 Oxide, Lung Function Testing, Airways Hyperresponsiveness.  
45  
46  
47  
48  
49

50 **Correspondence:**

51 Angelika Mohn  
52 Department of Pediatrics  
53 University of Chieti  
54 Via Dei Vestini 5  
55 66100 Chieti  
56 Tel +39 0871 358827  
57 Fax +39 0871 584731  
58 Email amohn@unich.it  
59  
60

**ABSTRACT**

As the incidence of obesity and asthma increase consistently, the reciprocal influences of both on lung function, airways hyperresponsiveness (AHR) and bronchial inflammation were analysed in a group of 118 caucasian prepubertal children.

Anthropometric measurements and respiratory parameters (Flow/Volume curves, “free run” Exercise Challenge Test, FeNO) were evaluated and the study population was divided into four groups according to Body Mass Index and the presence or absence of asthma: *Obese Asthmatic, Normal-weight Asthmatic, Obese non-Asthmatic, Normal-weight non-Asthmatic* children.

MEF75% was significantly lower in obese asthmatic and non-asthmatic children compared to healthy controls and the influence of obesity on MEF75% was confirmed by multiple linear regression. After the “free-run” Exercise Challenge Test all analysed respiratory parameters decreased significantly in asthmatic obese and non-obese children, in contrast in non-asthmatic obese children no differences were found. FeNO analysis demonstrate significantly higher values in asthmatic children compared to obese non-asthmatic and healthy controls. Furthermore FeNO was significantly higher in obese non-asthmatic children when compared to healthy controls.

In conclusion, while AHR and airways inflammation are clearly associated to an asthmatic status, obesity seems to induce early proinflammatory airways changes combined with reduction of higher intrathoracic airway flows.

## INTRODUCTION

Over the past 20 years the incidence and prevalence of obesity have steadily increased. The most recent data from the Centers for Disease Control's (CDC) Third National Health and Nutrition Examination Survey (NHANES III) revealed a prevalence of obesity – defined as a body mass index (BMI) in excess of 95% of the age specific distribution – of 14% in children aged 6 – 11 surveyed between 1988 and 1994.<sup>1</sup> In comparison NHANES II, conducted between 1976 and 1980, found a prevalence of obesity of 7.6% for the same population. This has been accompanied by a similar rise in the association rates of asthma. The CDC self-reported prevalence of asthma in children aged 5 – 14 rose from 42.8 per 1000 in 1980 to 74.4 per 1000 in 1994.<sup>2</sup> Epidemiologists have reported that both obesity and asthma are “endemic” in developed countries<sup>3,4</sup> and “epidemic” in developing countries undergoing their epidemiological transition phase.<sup>5</sup> Accordingly, the World Health Organization (WHO) includes both obesity and asthma among the major chronic disorders representing world-wide public health priorities.<sup>6</sup>

Given the dramatic rise in the prevalence of obesity and asthma, it is not surprising that there has been an increasing body of literature on the association between BMI and asthma.<sup>7</sup>

Obesity may have effects on both the upper and lower airway;<sup>8</sup> and this hypothesis is supported to a certain extent by the observation that BMI, the best single measure of body adiposity,<sup>9</sup> seems to be a reliable predictor of allergic respiratory symptoms and airways hyperresponsiveness (AHR).<sup>10,11,12</sup>

Infiltration by eosinophils, mast cells and T lymphocytes is a key feature of airway inflammation in asthma<sup>13</sup> and an increased eosinophil number in induced sputum of asthmatic children is tightly linked to current asthma symptoms and AHR.<sup>14</sup> Recently, exhaled Nitric Oxide (eNO) has been shown to have a good correlation with airway eosinophilic inflammation<sup>15</sup> and its fractional concentration in exhaled air (FeNO) has been accepted to be a useful indicator of airway inflammation. On the other hand recent literature emphasises the role of obesity in inducing a proinflammatory state as obesity is associated with increased levels of leptin and proinflammatory

1  
2  
3 cytokines such as interleukin (IL)-6 and IL-1 which might be implicated in a possible up-regulate  
4  
5 inflammation of the airway system.<sup>16</sup>  
6  
7

8 The reported association between increased BMI and onset of AHR has been elegantly  
9  
10 demonstrated by Litonjua et al.<sup>17</sup> in a longitudinal study of aging; and has been further confirmed  
11  
12 by two large cross-sectional studies.<sup>18,19</sup> This is in contrast with epidemiologic studies where this  
13  
14 association has not been universally observed, indicating that different factors such as duration of  
15  
16 obesity, physiological changes of airway structures might play an important role on the  
17  
18 development of this relationship.  
19  
20  
21

22 Prepubertal children might represent a special study population in order to investigate the reciprocal  
23  
24 and independent influences of obesity and asthma on lung function testing, and bronchial indices of  
25  
26 flogosis as both diseases should not necessarily having caused physiological changes commonly  
27  
28 observed in older subjects.  
29  
30

31 We therefore evaluated AHR and bronchial inflammation expressed by FeNO in obese and non  
32  
33 obese asthmatic children in comparison with non asthmatic obese and normalweight prepubertal  
34  
35 children.  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## METHODS

### *Study subjects*

Clinical, anthropometric measurements and respiratory parameters of a group of 118 Caucasian pre-pubertal children referred to the Department of Pediatrics, Children Obesity Clinic and to the Allergological and Pneumological Unity of the Pediatric Department, University of Chieti, Italy, between May 2008 and August 2008, were evaluated.

According to Body Mass Index [BMI], and based on the presence or absence of asthma, the study population was divided into four groups:

1. *Obese Asthmatic Prepubertal Children [ObA] (16 M and 14 F);*
2. *Normal-weight Asthmatic Prepubertal Children [NwA] (16 M and 14 F);*
3. *Non-Asthmatic Obese Prepubertal Children [Ob] (14 M and 16 F);*
4. *Non-Asthmatic Normal-weight Prepubertal Children [Nw] (14 M and 14 F)*

- Pubertal stage was defined on the basis of breast development in girls and genital development in boys according to the criteria of Tanner (all subjects had pre-adolescent characteristics corresponding to stage 1).

- A complete physical examination including [height, height Standard Deviation Score (SDS), weight, weight-SDS, BMI, BMI-SDS]. Children with values between -2 and 2 SD were defined normal-weight, while children with BMI above 2 SD were defined obese.<sup>20,21</sup>

- Asthma was classified according to GINA guidelines<sup>22</sup> as intermittent asthma, mild persistent, moderate persistent and severe persistent asthma. All subjects had been admitted for intermittent/mild persistent asthma and the diagnosis was made by a pediatric respiratory physician on the basis of clinical history of repeated episodes of coughing, dyspnea and wheezing, according to ATS-ERS criteria.<sup>23</sup> To evaluate the bronchial reactivity was made an Exercise Challenge Test. Moreover, to evaluate the bronchial inflammation all children underwent FeNO analysis. To evaluate the allergic sensitization we made Skin Prick Test (SPT) and serum specific IgE measurements for the most common respiratory allergens.

1  
2  
3 Asthmatic children had received an inhaled corticosteroids (ICS) therapy, but were free from  
4  
5 therapy for at least 2 weeks before the study evaluation; and all obese children were therapy-free.  
6  
7

8 All the subjects were symptom-free and had no clinical evidence of cardiopulmonary disease in a  
9  
10 preceding evaluation. Exclusion criteria from the study for all groups were: presence of upper or  
11  
12 lower respiratory infection or exacerbation of asthma during the previous 4 weeks, not well  
13  
14 controlled asthma according to current Global Initiative for Asthma (GINA) classification of  
15  
16 asthma,<sup>24</sup> congenital abnormalities of cardiorespiratory system, chest or skeletal deformities, or  
17  
18 neuromuscular system disease, chronic and autoimmune disease, gastroesophageal reflux, presence  
19  
20 of medical condition in which exercise was forbidden.  
21  
22  
23

24 Written informed consent was obtained from all parents and oral consent from all children.  
25  
26

27 The study was approved by Ethical Committee of University of Chieti.  
28  
29  
30

## 31 **Methods**

### 32 *Anthropometric measurements*

33  
34 Body weight was determined to the nearest 0.1 Kg, and height was measured in triplicate with  
35  
36 Harpenden stadiometer to the nearest 0.1 cm.  
37  
38

39 We used BMI for fatness indexes (the weight in kilograms divided by the square of the height in  
40  
41 meters); and also BMI expressed as a continuous variable in standard deviation score (BMI-SDS)  
42  
43 units (z-score), determined as difference between the individual observed value and the reference  
44  
45 mean for age and sex, divided by the corresponding standard deviation.<sup>25</sup>  
46  
47  
48  
49  
50  
51  
52

### 53 *Functional respiratory data*

54  
55 Evaluations were made by Flow/Volume curves (KitMicro Spirometer 7.3a – COSMED), according  
56  
57 ATS/ERS Guidelines,<sup>26</sup> and the main spirometric tests were considered: Forced Expiratory Volume  
58  
59 in 1 sec (FEV1), Forced Vital Capacity (FVC), Peak Expiratory Flow (PEF) and Maximal  
60

1  
2  
3 Expiratory Flows at 75%, 50% and 25% of the Forced Vital Capacity (MEF75%, MEF50%,  
4 MEF25%). All the tests were expressed as a percentage of theoretical values for eight.  
5  
6

7  
8 To evaluate the bronchial hyperresponsiveness measurements are performed in the standing  
9 position, with nose-clip until two consecutive technically acceptable curves are achieved. After the  
10 free-run, spirometry was measured in duplicate sets at 1, 3, 5, 10 and 20 minutes post-exercise. The  
11 better of the two curves was selected as the representative value at each interval, but differences  
12 between the two values of FEV1 had to be <5%. The target was to achieve 6 minutes of "free run"  
13 to increase heart-rate to 80% in the maximum for at least 4 minutes according to ATS/ERS  
14 recommendations for exercise challenge test in older children.<sup>27,28</sup> Exercise response was defined as  
15 the greatest decrease in FEV1 expressed as a percentage of the baseline values. A  
16 bronchoconstriction response to exercise was considered as positive when the FEV1 decrease from  
17 baseline was greater than 13%.<sup>29</sup>  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

#### 34 *Allergic sensitization*

35  
36 Was evaluated by Skin Prick Test (SPT) and serum specific IgE measurements for the most  
37 common respiratory allergens: Dust Mite (*Dermatophagoides Pteronyssinus* and *Farinae*), Grass,  
38 Parietaria, *Artemisia Vulgaris*, Olive, Cypress, Lime, Stone, Elm, Plane, Cat and Dog dander,  
39 Alternaria Alternata and *Aspergillus Fumigatus* (moulds). Determination of Allergen-Specific IgE  
40 was made by an Immunoenzymatic Allergo-sorbent Test (Cap test Pharmacia).<sup>30,31</sup>  
41  
42  
43  
44  
45  
46  
47  
48  
49

#### 50 *Respiratory inflammation marker*

51  
52 FeNO was determined with an on-line method using a single breath exhalation and a sensitive  
53 chemiluminescence assay (Ecomedics CLD 88), according to ATS-ERS.<sup>32</sup> Patients made an  
54 inspiration of eNO-free air via a mouthpiece immediately followed by full exhalation at a constant  
55 rate (50 ml/sec) for at least 5 seconds. The mean of three readings at the end of the expiration  
56  
57  
58  
59  
60

1  
2  
3 (plateau phase) was taken as the representative value for each measurements. 12 ppb or more were  
4  
5 considered positive, according to ATS-ERS criteria.<sup>33</sup>  
6  
7  
8  
9

### 10 *Statistical analysis*

11  
12 All values were expressed as **Means and SD**. Kruskal Wallis Test was performed for evaluating  
13 differences between the four groups (ObA, NwA, Ob, Nw). Statistical significance level was  
14  
15  $p < 0.05$ . Subsequently, differences within the single groups were tested by Mann-Whitney Test,  
16  
17  
18 adjusting the significance level by  $p < 0.0125$ . Differences in functional respiratory parameters **pre-**  
19  
20 **and post-“free-run” exercise Challenge Test** were analysed using Wilcoxon Signed Ranks Test  
21  
22  
23 within each group.  
24  
25

26  
27 **A multiple linear regression analysis was performed to asses the independent contribution of**  
28  
29 **obesity (BMI-SDS) and asthma on respiratory inflammation and respiratory functional parameters.**  
30  
31 **FeNO, PEF% and MEF75% were the dependent variables and BMI-SDS and asthma were the**  
32  
33 **independent variables.** Statistical significance level was  $p < 0.05$ .  
34  
35

36 All calculations were made with the computer program SPSS (Statistical Package for the Social  
37  
38 Science), version 16.0 software for Windows.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## RESULTS

Anthropometric measurements and respiratory functional data are reported in **table 1**.

The four groups were similar for age, gender, and **height**.

All the asthmatic children were **allergic** to Dermatophagoides Pteronyssinus (**Df. pt**).

As expected BMI and BMI-SDS was similar between obese asthmatic and obese non-asthmatic children, but was significantly different between obese and normal-weight children (**ObA vs Ob:  $p>0.05$ ; ObA vs NwA:  $p= 0.0004$ ; ObA vs Nw:  $p=0.0008$ ; Ob vs NwA:  $p=0.0001$ ; Ob vs Nw:  $p=0.0002$** ).

### *Baseline Respiratory Functional Parameters*

**No significant differences were found for FVC%, FEV1%, MEF50% and MEF25% between the four groups.**

**Despite detecting a trend of difference in PEF% between the four groups, no differences were found when performing sub-analysis between ObA vs NwA, ObA vs Ob, ObA vs Nw, NwA vs Ob, NwA vs Nw, and Ob vs Nw.**

We found that MEF75% was significantly lower in OA group and OnA group compared to NnA group ( $p=0.004$  and  $p=0.0002$ , respectively) (**Table 1**).

### *Respiratory Functional Parameters after “free run” Exercise Challenge Test*

No significant differences were found in **FEV1%, FVC%, MEF50% and MEF25%** after the **exercise test** among the four groups.

After the **“free-run” exercise Challenge Test** we found statistically significant differences in PEF% and MEF75%. In fact NnA had significantly higher values of these two parameters compared to OA ( $p=0.001$  for PEF% and  $p=0.002$  for MEF75%); furthermore obese non-asthmatic children had significantly lower value of MEF75% compared to the NnA group ( $p=0.007$ ) (**Table 1**).

1  
2  
3 As shown in Table 2, analysing functional respiratory data within each group after the “free-run”  
4 exercise Challenge Test, we found that all analysed parameters, with the exception of FVC% for the  
5 OA group and of MEF25% for the NA group, decreased significantly in asthmatic obese and non-  
6 obese children (without clinical and functional evidence of obstruction – all indices were in the  
7 normal range for age and height). In contrast in non-asthmatic obese children we did not detect any  
8 statistical significant differences in all functional respiratory parameters after “free run” exercise  
9 Challenge Test.

### 20 21 22 *Respiratory inflammation marker*

23 FeNO values were significantly different across all four groups ( $p=0.0002$ ).

24  
25 In fact FeNO values were significantly higher in the ObA group compared to Ob group and to the  
26 Nw group ( $p=0.011$  and  $p=0.0001$ , respectively); and were significantly higher in the NwA group  
27 compared to Ob group and to the Nw group ( $p=0.0002$  and  $p=0.0003$ , respectively). We did not  
28 detect any statistical significant difference in FeNO values between the asthmatic children  
29 ( $p=0.201$ ). Furthermore FeNO was significantly higher in the Ob group when compared to the Nw  
30 group ( $p=0.0004$ ) (Figure 1).

### 31 32 33 34 35 36 37 38 39 40 41 42 43 *Independent contribution of BMI-SDS and asthma on respiratory inflammation and respiratory* 44 *functional parameters*

45  
46 In a multiple linear regression analysis with FeNO as the dependent variable and BMI-SDS and  
47 asthma (asthmatic vs non-asthmatic children) as the independent variables, asthma was significantly  
48 and independently related to FeNO values, whereas no association was found for obesity (BMI-  
49 SDS). In a second multiple linear regression model with PEF% as the dependent variable and BMI-  
50 SDS and asthma as the independent variables, BMI-SDS was significantly and independently  
51 related to PEF% values. A further multiple linear regression analysis with MEF75% as the  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

dependent variable and BMI-SDS and asthma as the independent variables, showed again that BMI-SDS was significantly and independently related to MEF75% values (Table 3).

For Peer Review

## DISCUSSION

This is the first study comparing obese asthmatic with obese non-asthmatic children and healthy controls with the aim to analyse the reciprocal influences of obesity and asthma on lung function in prepubertal children. Although the effect of obesity on pulmonary function tests has been extensively investigated in adults, the number of similar studies in children is limited and confirm the controversial results between obesity and AHR observed in adults.

In order to evaluate the relationship between obesity and AHR and to detect whether breathlessness experienced by obese children during physical activities is effectively due to the airway hyper-responsiveness we performed a “free-run” Exercise Challenge test. While basal values of spirometric indexes were normal in all examined groups, in asthmatic obese and non-obese children free run induced AHR documented by reduction of all functional respiratory parameters. In contrast in obese children no changes in these functional respiratory parameters were detected. In addition, in a multiple linear regression we found that asthma remained the only predictor for AHR. In agreement to our data, epidemiological studies do not uniformly support the conclusions that obesity leads to airflow limitation, AHR or asthma. In this respect Schachter and coworkers showed in a group of 1971 adults, that although BMI was associated with a clinical diagnosis of asthma and subjective symptoms of dyspnea and wheeze, airflow obstruction or AHR subsequent to histamine challenge could not be detected in the sole obese subjects.<sup>34</sup> Similar findings were obtained by Bibi et al.<sup>35</sup> who demonstrated in a large population of prepubertal children that broncodilation could not be induced by salbutamol inhalation in non-asthmatic obese subjects.<sup>36</sup> These data demonstrate that obesity should not be considered as being a risk factor for AHR in prepubertal children. Furthermore asthma like symptoms especially during physical activity should not be overestimated by parents and medical staff. This is of paramount relevance as it might induce inappropriate treatment and refrain children from performing sports leading to a vicious circle impeding improvement of both obesity and respiratory symptoms.

1  
2  
3 Interestingly in obese asthmatic children and obese non-asthmatic children we documented lower  
4 basal values of MEF75% when compared to healthy controls. Furthermore by performing a  
5 multiple linear logistic regression analysis obesity, defined as BMI-SDS, was significantly and  
6 independently related to MEF75%; while no association was found for asthma. Similar data were  
7 documented by two previous studies<sup>37,38</sup> where obesity was associated with a significant reduction  
8 in FEF<sub>25-75</sub>. These data suggests that obesity might play a primary role on higher intrathoracic  
9 airway flows and despite being possibly explained by fatty infiltration of muscles narrowing  
10 directly the upper airway, recent literature suggest that this functional parameter might be used as  
11 an early predictor of subclinical asthma. In order to obtain a complete evaluation of bronchial status  
12 in this study we performed also FeNO analysis. In fact recent studies revealed that FeNO is a  
13 potentially useful measure to evaluate the role of airways inflammation in asthma, as it represents  
14 the forerunner of the most important event namely “remodelling” of bronchial airways.<sup>39</sup> In fact  
15 FeNO, as recommended in the Consensus Report Practall for the Diagnosis, Management and  
16 Treatment of Childhood Asthma<sup>40</sup> offers a logical complementary item to the only use of clinical  
17 and functional data, evaluating the bronchial eosinophilic inflammation. In our study, as expected,  
18 both groups of asthmatic children presented highly elevated FeNO values when compared to normal  
19 controls reflecting the underlying bronchial inflammation process. Data from animal and human  
20 studies clearly demonstrate that the enhancement of normal adipose tissue functions in obesity leads  
21 to a systemic proinflammatory state.<sup>41</sup> Adipose tissue from obese individuals expresses a number of  
22 proinflammatory molecules, such as leptin, tumor necrosis factor- $\alpha$  (TNF-  $\alpha$ ), IL-6, transforming  
23 growth factor- $\beta$ 1 (TGF-  $\beta$ 1), and C-reactive protein (CRP), and there appears to be significant  
24 overlap between the immune function of adipocytes and the function of T-lymphocytes and  
25 macrophages, particularly with respect to the elaboration of inflammatory cytokines.<sup>42,43</sup> Although  
26 a conclusive relationship between obesity and airway inflammation has not yet to been described, it  
27 could be argued that the role of the proinflammatory state underlying the main metabolic and  
28 cardiovascular complications in obesity might play a pivotal role in determining airways

1  
2  
3 inflammation independently of any allergic mechanisms. This might explain the findings of our  
4  
5 study where FeNO values were significantly higher in obese non-asthmatic children compared to  
6  
7 healthy controls and the findings of Santamaria et al.<sup>44</sup> who found a positive correlation between  
8  
9 FeNO and BMI in healthy children. However it needs to be acknowledged that in the multiple  
10  
11 regression analysis obesity was not a predictor for increased FeNO values. This apparent  
12  
13 contrasting findings might be partially explained by the age group of our study population as  
14  
15 obesity related inflammation might not have reached its full implication and a possible involvement  
16  
17 of the respiratory system might require a long standing history of obesity. As puberty is a well  
18  
19 known risk factor for developing obesity related complications it might be of crucial interest to  
20  
21 obtain longitudinal data starting in prepuberty and lasting through all pubertal stages in order to not  
22  
23 underestimate the possible role of obesity on the development of bronchial inflammation and  
24  
25 therefore to fully explore this intriguing relationship.  
26  
27  
28  
29  
30

31 In conclusion, while AHR and airways inflammation are clearly associated to an asthmatic status,  
32  
33 obesity seems to induce early proinflammatory airways changes combined with reduction of higher  
34  
35 intrathoracic airway flows. Longitudinal data are required in order to clarify whether these changes  
36  
37 might represent risk factor for asthma later on in life.  
38  
39  
40  
41  
42

### 43 Acknowledgements

44  
45 We thank all nurses of the Department of Pediatrics, University of Chieti, in particular Di Santo D,  
46  
47 Di Luca D, Diomedi A, for their daily support in clinical research and care of children.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## REFERENCES

1. Division of Health Examination Statistics, National Center for Health Statistics, Division of Nutrition and Physical Activity, National Center for Chronic Disease Prevention and Health Promotion, CDC Update. Prevalence of overweight among children, adolescents, and adults – United States, 1988-1994. *MMWR* 1997; 46: 199-202.
2. Mannino DM, Homa DM, Petrowski CA, Nixon LL, Johnson CA, Ball LB, Jack E, Kang DS. Surveillance for asthma – United States, 1960-1995. *MMWR CDC Surveill Summ.* 1998 Apr 24;47(1):1-27.
3. Woolcock AJ, Peat JK. Evidence for the increase of asthma worldwide. *Ciba Found Symp* 1997; 206: 122-34.
4. World Health Organization. Obesity: preventing and managing the global epidemic. WHO Technical Report Series 894. Geneva, Switzerland, WHO, 2000.
5. World Health Organization. Risk factors. *World Health Report 2002*. Geneva, Switzerland: WHO, 2002.
6. World Health Organization. Preventing chronic diseases: a vital investment. Geneva, Switzerland: WHO, 2005.
7. Figueroa-Munoz JI, Chinn S, Rona RJ. Association between obesity and asthma in 4-11 year old children in the UK. *Thorax* 2001; 56: 133-7.
8. Deane S, Thomson A. Obesity and the pulmonologist. *Arch Dis Child* 2006; 91: 188-191.
9. Power C, Lake JK, Cole TJ. Measurement and long term health risks of child and adolescent fatness. *Int J Obes Relat Metab Disord* 1997; 21: 507-526.
10. Schwartz J, Gold D, Dockery DW, Weiss ST, Speizer FE. Predictors of asthma and persistent wheeze in a national sample of children in U.S. Association with social class, perinatal events and race. *Am Rev Respir Dis* 1990; 142 (3): 555-562.

- 1  
2  
3 11. Gennuso J, Epstein LH, Paluch RA, Cerny F. The relationship between asthma and obesity  
4  
5 in urban minority children and adolescent. Arch Pediatr Adolesc Med 1998; 152 (12): 1197-  
6  
7 1200.  
8  
9
- 10 12. Huang SL, Shiao G, Chou P. Association between body mass index and allergy in teenage  
11  
12 girls in Taiwan. Clin Exp Allergy 1999; 29 (3): 323-329.  
13
- 14 13. Busse WW, Lemanske RF Jr. Asthma. N Eng J Med 2001; 344: 350-2.  
15
- 16 14. Gibson PG, Wlodarczyk JW, Hensley MJ, Gleeson M, Henry RL, Cripps AW, Clancy RL.  
17  
18 Epidemiological association of airway inflammation with asthma symptoms and airway  
19  
20 hyperresponsiveness in childhood. Am J Respir Crit Care Med 1998; 158: 36-41.  
21  
22
- 23 15. Payne DN, Adcock IM, Wilson NM, Oates T, Scallan M, Bush A. Relationship between  
24  
25 exhaled nitric oxide and mucosal eosinophilic inflammation in children with difficult  
26  
27 asthma, after treatment with oral prednisolone. Am J Respir Crit Care Med 2001; 164: 1376-  
28  
29 81.  
30  
31
- 32 16. Visser M, Bouter LM, McQuillan GM, Wener MH, Harris TB. Elevated C-reactive protein  
33  
34 levels in overweight and obese adults. JAMA 1999; 282: 2131-5.  
35  
36
- 37 17. Litonjua AA, Sparrow D, Celedon JC, DeMolles D, Weiss ST. Association of body mass  
38  
39 index with the development of methacholine airway hyperresponsiveness in men: the  
40  
41 Normative Aging Study. Thorax 2002; 57: 581-5.  
42  
43
- 44 18. Chinn S, Jarvis D, Burney P. Relation of bronchial responsiveness to body mass index in the  
45  
46 ECRHS. European Community Respiratory Health Survey. Thorax 2002; 57: 1028-33.  
47  
48
- 49 19. Celedon JC, Palmer LJ, Litonjua AA, Weiss ST, Wang B, Fang Z, Xu X. Body mass index  
50  
51 and asthma in adults in families of subjects with asthma in Anqing, China. Am J Respir Crit  
52  
53 Care Med 2001; 164: 1835-40.  
54  
55
- 56 20. Kiess W, Reich A, Müller G, Galler A, Kapellen T, Raile K, Böttner A, Seidel B, Kratzsch  
57  
58 J. Obesity in childhood and adolescence: clinical diagnosis and management. J Pediatr  
59  
60 Endocrinol Metab. 2001;14 Suppl 6:1431-40.

- 1  
2  
3 21. Dietz WH, Bellizzi MC. Introduction: the use of body mass index to assess obesity in  
4 children. *Am J Clin Nutr.* 1999 Jul;70(1):123S-5S.  
5  
6  
7  
8 22. <http://www.ginasthma.it> or [www.ginasthma.org](http://www.ginasthma.org) - GINA World Report 2006 - GLOBAL  
9 STRATEGY FOR ASTHMA MANAGEMENT AND PREVENTION.  
10  
11  
12 23. National Asthma Education and Prevention Program. NAEPP expert panel report guidelines  
13 for the diagnosis and management of asthma – update on selected topics 2002. National  
14 Institutes for Health 2006.  
15  
16  
17  
18  
19 24. Global Strategy for Asthma Management and Prevention, Global Initiative for Asthma  
20 (GINA) 2007. Available from: <http://www.ginasthma.org>.  
21  
22  
23  
24  
25 25. Lindgren G, Strandell A, Cole T, Healy M, Tanner J. Swedish population standards for  
26 height, weight and body mass index attained at 6 to 16 years (girls) or 19 years (boy). *Acta*  
27 *Pediatr* 1995; 84: 1019-1028.  
28  
29  
30  
31 26. Laszlo G. Standardisation of lung function testing: helpful guidance from the ATS/ERS  
32 Task Force. *Thorax.* 2006 Sep; 61 (9): 744-6.  
33  
34  
35  
36 27. Crapo RO, Casaburi R, Coates AL, Enright PL, Hankinson JL, Irvin CG, MacIntyre NR,  
37 McKay RT, Wanger JS, Anderson SD, Cockcroft DW, Fish JE, Sterk PJ. Guidelines for  
38 Methacholine and Exercise Challenge Testing-1999. This official statements of the  
39 American Thoracic Society was adopted by the ATS Board of Directors, July 1999. *Am J*  
40 *Respir Crit. Care Med* 2000;161:309-329.  
41  
42  
43  
44  
45  
46  
47  
48 28. ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories.  
49 ATS statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med* 2002;  
50 166: 111–117.  
51  
52  
53  
54  
55 29. Godfrey S, Springer C, Bar-Yishay E, Avital A. Cut-off points defining normal and  
56 asthmatic bronchial reactivity to exercise and inhalation challenges in children and young  
57 adults. *Eur Respir J.* 1999; 14: 659-66.  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
30. Terho E, Frew AJ et al., "Type III allergy skin testing. Position statement for EAACI Subcommittee on Skin Tests and Allergen Standardization", *Allergy*, vol. 50, no. 5, pp. 392-6, 1995.
  31. Wolkerstorfer A, Wahn U, Kjellman NI, Diepgen TL, De Longueville M, Oranje AP, "Natural course of sensitization to cow's milk and hen's egg in childhood atopic dermatitis: ETAC Study Group", *Clinical & Experimental Allergy*, vol. 32, no. 1, pp. 70-73, 2002.
  32. American Thoracic Society; European Respiratory Society. ATS/ERS recommendations for standardized procedures for the online and offline measurement of exhaled lower respiratory nitric oxide and nasal nitric oxide, 2005. *Am J Respir Crit Care Med*. 2005 Apr 15;171(8):912-30.
  33. Baraldi E, de Jongste JC; European Respiratory Society; American Thoracic Society. Measurement of exhaled nitric oxide in children, 2001. *Eur Respir J* 2002; 20 (1): 223-237.
  34. Schachter LM, Salome CM, Peat JK, Woolcock AJ. Obesity is a risk for asthma and wheeze but not airway hyperresponsiveness. *Thorax* 2001;56:4-8.
  35. Bibi H, Shoseyov D, Feigenbaum D, Genis M, Friger M, Peled R, Sharff S. The relationship between asthma and obesity in children: is it real or a case of over diagnosis? *J Asthma* 2004; 41 (4): 403-410.
  36. Bibi H, Montgomery M, Pasterkamp H, Chernick V. Relationship between response to inhaled salbutamol and methacholine bronchial provocation in children with suspected asthma. *Pediatr Pulmonol* 1991; 10: 244-248.
  37. Spathopoulos D, Paraskakis E, Trypsianis G, Tsalkidis A, Arvanitidou V, Emporiadou M, Bouros D, Chatzimichael A. The effect of obesity on pulmonary lung function of school aged children in Greece. *Pediatr Pulmonol* 2009; 44: 273-280.
  38. Ülger Z, Demir E, Tanaç R, Gökşen D, Gülen F, Darcan Ş, Can D, Coker M. The effect of childhood obesity on respiratory function tests and airway hyperresponsiveness. *The Turkish Journal of Pediatrics* 2006; 48: 43-50.

- 1  
2  
3 39. van Veen IH, Sterk PJ, Schot R, Gauw SA, Rabe KF, Bel EH. Alveolar nitric oxide versus  
4  
5 measures of peripheral airway dysfunction in severe asthma. *Eur Respir J* 2006; 27 (5): 951-  
6  
7 956.  
8  
9  
10 40. Bacharier LB, Boner A, Carlsen K-H, Eigenmann PA, Frischer T, Götz M, Helms PJ, Hunt  
11  
12 J, Liu A, Papadopoulos N, Platts-Mills T, Pohunek P, Simons FE, Valovirta E, Wahn U,  
13  
14 Wildhaber J; European Pediatric Asthma Group. Diagnosis and treatment of asthma in  
15  
16 childhood: a PRACTALL Consensus Report. *Allergy* 2008; 63: 5-34.  
17  
18  
19 41. Fantuzzi G. Adipose tissue, adipokines, and inflammation. *J Allergy Clin Immunol* 2005;  
20  
21 115: 911-919.  
22  
23  
24 42. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW Jr. Obesity is  
25  
26 associated with macrophage accumulation in adipose tissue. *J Clin Invest* 2003; 112: 1796-  
27  
28 1808.  
29  
30  
31 43. Wellen KE, Hotamisligil GS. Obesity-induced inflammatory changes in adipose tissue. *J*  
32  
33 *Clin Invest* 2003; 112: 1785-1788.  
34  
35  
36 44. Santamaria F, Montella S, De Stefano S, Sperli F, Barbarano F, Valerio G. Relationship  
37  
38 between exhaled nitric oxide and body mass index in children and adolescent. *J Allergy Clin*  
39  
40 *Immunol* 2005; 116: 1163-1164.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1.** Anthropometric measurements and respiratory parameters of the study population.

	ObA	NwA	Ob	Nw	$p^{\dagger}$
<b>Clinical and Anthropometric characteristics</b>					
<b>Gender (M/F)</b>	16/14	16/14	14/16	14/14	NS*
<b>Age (years)</b>	<b>7.03</b> [6.05 – 8.02]	<b>7.06</b> [6.77 – 8.05]	<b>8.04</b> [7.05 – 9.02]	<b>6.1</b> [6.03 – 8.09]	NS
<b>Weight (Kg)</b>	<b>35</b> [31 – 45.25]	<b>25.75</b> [22 – 30.2]	<b>42</b> [36.5 – 48]	<b>24</b> [21 – 31.5]	<b>0.0001</b>
<b>SDS-Height</b>	<b>0.5</b> [0.06 – 1.2]	<b>0.5</b> [0.05 – 1.1]	<b>0.56</b> [0.05 – 1.55]	<b>0.48</b> [0.051 – 1.2]	NS
<b>SDS-BMI</b>	<b>4.6</b> [3.85 – 5.45]	<b>0.4</b> [-0.25 – 1.3]	<b>5</b> [4.55 – 7.1]	<b>0</b> [-0.4 – 0.1]	<b>0.0001</b>
<b>Respiratory inflammation marker</b>					
<b>FeNO (ppb)</b>	<b>5.1</b> [2.8 – 10.3]	<b>7.4</b> [3.5 – 17]	<b>3.5</b> [2.3 – 4.3]	<b>2.3</b> [2.1 – 2.4]	<b>0.0002</b>
<b>Baseline Respiratory Functional Parameters</b>					
<b>FVC (Z-score)</b>	<b>-0.26</b> [-0.78 – 0.53]	<b>-0.11</b> [-0.48 – 0.33]	<b>-0.28</b> [-1.12 – 0.02]	<b>0.25</b> [-0.76 – 1.99]	NS
<b>FEV<sub>1</sub> (Z-score)</b>	<b>-0.14</b> [-0.84 – 0.31]	<b>0.025</b> [-0.35 – 0.44]	<b>0.12</b> [-0.74 – 0.73]	<b>-0.13</b> [-0.66 – 2.01]	NS
<b>FEV<sub>1</sub>/FVC (Z-score)</b>	<b>0.11</b> [-0.57 – 0.69]	<b>0.27</b> [-0.58 – 1]	<b>0.69</b> [-0.37 – 1.45]	<b>0.85</b> [0.62 – 1.10]	<b>0.025</b>
<b>FEF<sub>25-75</sub> (Z-score)</b>	<b>-0.13</b> [-0.49 – 0.25]	<b>0.05</b> [-0.32 – 0.75]	<b>0.35</b> [-0.30 – 0.93]	<b>0.4</b> [0.08 – 0.56]	NS
<b>Respiratory Functional Parameters after “free run” Exercise Challenge Test</b>					
<b>post-FEV<sub>1</sub> (Z-score)</b>	<b>-0.97</b> [-2.12 – 0.33]	<b>-1.11</b> [-1.60 – -0.37]	<b>0.09</b> [-0.69 – 0.85]	<b>-0.03</b> [-1.57 – -0.03]	<b>0.009</b>
<b>post-FVC (Z-score)</b>	<b>-1.01</b> [-1.71 – 0.31]	<b>-1.15</b> [-1.53 – -0.30]	<b>-0.43</b> [-1.15 – 0.36]	<b>0.20</b> [-1.11 – 0.96]	NS
<b>post-FEV<sub>1</sub>/FVC (Z-score)</b>	<b>-0.63</b> [-1.12 – 0.48]	<b>0.18</b> [-0.80 – 0.81]	<b>0.68</b> [-0.14 – 1.95]	<b>1.35</b> [-0.61 – 1.53]	<b>0.01</b>
<b>post-FEF<sub>25-75</sub> (Z-score)</b>	<b>-0.22</b> [-0.65 – 0.13]	<b>-0.06</b> [-0.55 – 0.51]	<b>0.3</b> [-0.21 – 0.49]	<b>0.42</b> [-0.91 – 0.62]	NS
<b>Fall in FEV<sub>1</sub> &gt;12%</b>	<b>5/30</b> 16.7%	<b>7/30</b> 23.3%	<b>2/30</b> 6.7%	<b>0/28</b> 0%	<b>0.008*</b>

Values are Median [Interquartile Range]. NS, no significant difference ( $P > 0.05$ ). M, male; F, female.

$^{\dagger}$  Kruskal Wallis Test

\*Pearson's Chi-square ( $X^2$ ) Test

BMI= Body Mass Index;

FVC= Forced Vital Capacity; FEV<sub>1</sub>= Forced Expiratory Volume in 1 sec; FEV<sub>1</sub>/FVC= Forced Expiratory Volume in 1 sec/ Forced Vital Capacity; FEF<sub>25-75</sub>= Forced Expiratory Flows between 25% and 75% of the Forced Vital Capacity

All the tests are expressed as Z-score.

**Table 2.** Comparison of the respiratory parameters after the 6MWT within each group.

	ObA	NwA	Ob	Nw
<i>FVC (Z-score)</i>	<b>-0.26</b> [-0.78 – 0.53]	<b>-0.11</b> [-0.48 – 0.33]	<b>-0.28</b> [-1.12 – 0.02]	<b>0.25</b> [-0.76 – 1.99]
<i>post-FVC (Z-score)</i>	<b>-1.01</b> [-1.71 – 0.31]	<b>-1.15</b> [-1.53 – -0.30]	<b>-0.43</b> [-1.15 – 0.36]	<b>0.20</b> [-1.11 – 0.96]
<i>p<sup>§</sup></i>	NS	<b>0.0001</b>	NS	NS
<i>FEV<sub>1</sub> (Z-score)</i>	<b>-0.14</b> [-0.84 – 0.31]	<b>0.025</b> [-0.35 – 0.44]	<b>0.12</b> [-0.74 – 0.73]	<b>-0.13</b> [-0.66 – 2.01]
<i>post-FEV<sub>1</sub> (Z-score)</i>	<b>-0.97</b> [-2.12 – 0.33]	<b>-1.11</b> [-1.60 – -0.37]	<b>0.09</b> [-0.69 – 0.85]	<b>-0.03</b> [-0.75 – 2.46]
<i>p<sup>§</sup></i>	NS	<b>0.0001</b>	NS	NS
<i>FEV<sub>1</sub>/FVC (Z-score)</i>	<b>0.11</b> [-0.57 – 0.69]	<b>0.27</b> [-0.58 – 1]	<b>0.69</b> [-0.37 – 1.45]	<b>0.85</b> [0.62 – 1.10]
<i>post-FEV<sub>1</sub>/FVC (Z-score)</i>	<b>-0.63</b> [-1.12 – 0.48]	<b>0.18</b> [-0.80 – 0.81]	<b>0.68</b> [-0.14 – 1.95]	<b>1.35</b> [-0.61 – 1.53]
<i>p<sup>§</sup></i>	<b>0.03</b>	NS	NS	NS
<i>FEF<sub>25-75</sub> (Z-score)</i>	<b>-0.13</b> [-0.49 – 0.25]	<b>0.05</b> [-0.32 – 0.75]	<b>0.35</b> [-0.30 – 0.93]	<b>0.4</b> [0.08 – 0.56]
<i>post-FEF<sub>25-75</sub> (Z-score)</i>	<b>-0.22</b> [-0.65 – 0.13]	<b>-0.06</b> [-0.55 – 0.51]	<b>0.3</b> [-0.21 – 0.49]	<b>0.42</b> [-0.91 – 0.62]
<i>p<sup>§</sup></i>	NS	<b>0.02</b>	NS	NS

Values are Median [Interquartile Range]. NS, no significant difference ( $P > 0.05$ ).

<sup>§</sup> Wilcoxon Signed Ranks Test

FVC= Forced Vital Capacity; FEV<sub>1</sub>= Forced Expiratory Volume in 1 sec; FEV<sub>1</sub>/FVC= Forced Expiratory Volume in 1 sec/ Forced Vital Capacity; FEF<sub>25-75</sub>= Forced Expiratory Flows between 25% and 75% of the Forced Vital Capacity

All the tests are expressed as Z-scores.

**Table 3.** Independent effect of asthma and obesity on lung function and bronchial inflammation.

Source	Dependent Variable	Type III Sum of Squares	F	p <sup>#</sup>
<b>Asthma</b>	<i>FeNO</i>	655,174	3,957	<b>0.048</b>
	<i>MEF<sub>75</sub> (%)</i>	505,481	1,038	NS
	<i>PEF (%)</i>	33,466	0,184	NS
	<i>FEV<sub>1</sub>/FVC (Z-score)</i>	18,871	0,623	NS
<b>Obesity</b>	<i>FeNO</i>	119,116	0,719	NS
	<i>MEF<sub>75</sub> (%)</i>	4156,102	8,534	<b>0.005</b>
	<i>PEF (%)</i>	1131,609	6,232	<b>0.015</b>
	<i>FEV<sub>1</sub>/FVC (Z-score)</i>	5,290	0,175	NS

NS, no significant difference ( $P > 0.05$ ).

<sup>#</sup> Multiple analysis of covariance (MANCOVA)


FeNO= Fractional exhaled Nitric Oxide; MEF<sub>75</sub>= Maximal Expiratory Flow at 75% of the Forced Vital Capacity; PEF= Peak Expiratory Flow; FEV<sub>1</sub>/FVC= Forced Expiratory Volume in 1 sec/ Forced Vital Capacity

FEV<sub>1</sub>/FVC ratio are expressed as Z-score.

MEF<sub>75</sub> and PEF are expressed as percentage of theoretical values for age, gender, and height.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Figure 1. Differences in PEF and MEF<sub>75</sub> obtained at baseline and after 6MWT.**



Values are Median [Interquartile Range].

PEF= Peak Expiratory Flow; MEF<sub>75</sub>= Maximal Expiratory Flow at 75% of the Forced Vital Capacity.

All the tests are expressed as %-predicted.


254x190mm (96 x 96 DPI)

Review



UNIVERSITY OF CHIETI  
- ITALY -  
[www.unich.it/cliped/](http://www.unich.it/cliped/)

**Pediatric Department**  
(Chief: Prof. Francesco Chiarelli)



Chieti, Italy, March 6<sup>th</sup> 2010

Dear Editor,

Thank you for giving us the possibility of revising and resubmitting our manuscript PPUL-09-0317 entitled **“The reciprocal influences of asthma and obesity on lung function testing, AHR and airway inflammation in prepubertal children.”**

We thank you and the referees for the detailed review and for the precious comments, which we have addressed thoroughly thereby improving our manuscript substantially.

We have tried to answer the major comments of referee 1 and 2 and please find enclosed a point-by-point answer sheet for each of them.

In agreement with your suggestion we have now calculated and expressed the results of the lung function tests as Z-score, according to the equations proposed by Stanojevic et al. (AJRCCM 2008). However, we continue to present the values of MEF<sub>75</sub> and PEF as %-expected because for these parameters the equations for Z-score were not available and the main differences detected in our study regarded these indices of lung function.

Furthermore, as requested we added all necessary information concerning our inclusion and exclusion criteria in the method section. We have modified the introduction in order to define in a clear way the main hypothesis of our study. Finally the most important limitation consisting of the cross sectional nature of this study has been discussed.

Please find enclosed the revised version of our manuscript where the changes have been marked-up.

We hope that our findings on this intriguing relationship between obesity and asthma are of interest to your readership and hope that the manuscript is now acceptable for publication in your prestigious journal.

Do not hesitate to contact us should further changes be necessary

Yours Sincerely,

Angelika Mohn

Department of Pediatrics  
University of Chieti  
Ospedale Policlinico  
Via dei Vestini 5  
66100 Chieti, Italy  
Fax: +39 0871-574831  
E-mail: [amohn@unich.it](mailto:amohn@unich.it)