



HAL
open science

On Level One Cuspidal Bianchi Modular Forms

Alexander D. Rahm, Mehmet Haluk Sengun

► **To cite this version:**

Alexander D. Rahm, Mehmet Haluk Sengun. On Level One Cuspidal Bianchi Modular Forms. 2012.
hal-00589184v2

HAL Id: hal-00589184

<https://hal.science/hal-00589184v2>

Preprint submitted on 8 Aug 2012 (v2), last revised 19 Feb 2013 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON LEVEL ONE CUSPIDAL BIANCHI MODULAR FORMS

ALEXANDER D. RAHM AND MEHMET HALUK ŞENGÜN

ABSTRACT. In this paper, we present the outcome of vast computer calculations, locating several of the very rare instances of level one cuspidal Bianchi modular forms that are not lifts of elliptic modular forms.

Bianchi modular forms over an imaginary quadratic field K are automorphic forms of cohomological type associated to the \mathbb{Q} -algebraic group $\text{Res}_{K/\mathbb{Q}}(\text{SL}_2)$. Even though modern studies of Bianchi modular forms go back to the mid 1960's, most of the fundamental problems surrounding their theory are still wide open. In this paper, we report on our remarkably extensive computations that show the paucity of *genuine* level one cuspidal Bianchi modular forms.

Let $S_k(1)$ denote the space of level one weight $k + 2$ cuspidal Bianchi modular forms over $K = \mathbb{Q}(\sqrt{-d})$. In their recent paper [FGT10], Finis, Grunewald and Tiraó computed the dimension of the subspace $L_k(1)$ of $S_k(1)$ which is formed by (twists of) those forms which arise from elliptic cuspidal modular forms via base-change or arise from a quadratic extension of K via automorphic induction (see [FGT10] for these notions). In this paper, we investigate numerically how much of $S_k(1)$ is exhausted by $L_k(1)$. There have been previous reports, however of limited size, in the 2009 paper [CM09] of Calegari and Mazur (the computations in this paper were carried out by Pollack and Stein) and in the 2010 paper [FGT10] of Finis, Grunewald and Tiraó. While the computations in [CM09] were limited to the case $d = 2$, the computations in [FGT10] covered ten imaginary quadratic fields. The precise scope of the computations in [FGT10] is given in Table 1 below.

d	1	2	3	7	11	19	5	6	10	14
$k \leq$	104	141	116	132	153	60	60	60	60	60

TABLE 1. Finis-Grunewald-Tiraó test range

It was observed in [CM09] that for $2k \leq 96$, one has $L_{2k}(1) = S_{2k}(1)$. The computations of [FGT10] extended those of [CM09]. An interesting outcome of the data they collected is that except in two of the 946 spaces they computed, one has $L_k(1) = S_k(1)$. The exceptional cases are $(d, k) = (7, 12)$

Date: 8th August 2012.

The first author is funded by the Irish Research Council.

The second author is funded by a Marie Curie Intra-European Fellowship.

and $(d, k) = (11, 10)$. In both cases, there is a two-dimensional complement to $L_k(1)$ inside $S_k(1)$.

Using a different and more efficient approach, we computed, over more than 800 processor-days, the dimension of 4976 different spaces $S_k(1)$ over 183 different imaginary quadratic fields. The precise scope of our computations is given in Tables 2, 3 and 4, where D and h denote the discriminant and the class number of K respectively. In only 29 of these spaces were we able to observe genuine forms. The precise data about these exceptional cases is provided in Table 5.

$ D $	3	4	7	8	11	15	19	20	23	24	31
h	1	1	1	1	1	2	1	2	3	2	3
$k \leq$	219	216	217	217	217	115	120	100	83	101	74

$ D $	35	39	40	43	47	51	52	55	56	59	67
h	2	4	2	1	5	2	2	4	4	3	1
$k \leq$	86	67	73	83	52	75	65	45	55	60	58

$ D $	68	71	79	83	84	87	88	91	95	103	104
h	4	7	5	3	4	6	2	2	8	5	6
$k \leq$	53	38	33	41	50	36	45	50	30	30	32

$ D $	107	111	115	116	119	120	123	127	131	132	136
h	3	8	2	6	10	4	2	5	5	4	4
$k \leq$	35	28	40	33	25	38	35	25	32	33	32

$ D $	139	143	148	151	152	155	159	163	164	167	168
h	3	10	2	7	6	4	10	1	8	11	4
$k \leq$	29	20	31	21	24	26	19	33	24	18	26

$ D $	179	183	184	187	191	195	199	203	211	212	215
h	5	8	4	2	13	4	9	4	3	6	14
$k \leq$	24	19	25	25	15	27	17	23	21	17	14

TABLE 2. the scope of our computations, part 1(a)

In Section 2, we briefly discuss the conjectural connections between the spaces $S_0(1)$ and Abelian varieties defined over K of GL_2 -type. In Section 3, we make some speculations in light of the data we collect. In particular, we make two conjectures, one about the paucity of genuine cuspidal Bianchi modular forms of level one and the other an analogue of Maeda's conjecture for Bianchi modular forms. Finally in Section 4, we explain in detail how we did our computations. As usual, the starting point of our approach is

$ D $	219	223	227	228	231	232	235	239	244	247	248
h	4	7	5	4	12	2	2	15	6	6	8
$k \leq$	20	14	17	18	13	21	23	12	17	13	16

$ D $	251	255	259	260	263	264	267	271	276	280	283
h	7	12	4	8	13	8	2	11	8	4	3
$k \leq$	15	14	17	14	13	15	21	12	16	16	17

$ D $	287	291	292	295	296	299	303	307	308	311	312
h	14	4	4	8	10	8	10	3	8	19	4
$k \leq$	12	19	16	13	13	13	11	15	13	11	13

$ D $	319	323	327	328	331	335	339	340	344	347	355
h	10	4	12	4	3	18	6	4	10	5	4
$k \leq$	11	12	10	13	14	11	15	14	10	12	13

$ D $	356	359	367	371	372	376	379	383	388	391	395
h	12	19	9	8	4	8	3	17	4	14	8
$k \leq$	11	9	11	10	12	12	13	8	11	9	10

$ D $	399	403	404	407	408	411	415	419	420	424	427
h	16	2	14	16	4	6	10	9	8	6	2
$k \leq$	8	12	9	8	10	12	10	12	11	10	13

TABLE 3. the scope of our computations, part 1(b)

the so called ‘‘Eichler-Shimura-Harder’’ isomorphism which allows us to replace $S_k(1)$ with the cohomology of the relevant Bianchi group with special non-trivial coefficients. Then to compute this cohomology space, we use the program *Bianchi.gp* [Rah10a], which analyzes the structure of the Bianchi group via its action on hyperbolic 3-space (which is isomorphic to the associated symmetric space $\mathrm{SL}_2(\mathbb{C})/\mathrm{SU}_2$). We then feed this group-geometric information into an equivariant spectral sequence that gives us an explicit description of the second cohomology of the Bianchi group, with the relevant coefficients.

Acknowledgments. We wish to thank John Cannon, John Cremona and Stephen S. Gelbart for useful discussions. We are grateful to Dan Yasaki who reworked his Magma program to compute for us the Hecke eigenfield associated to the genuine weight 2 cuspidal Bianchi modular forms that we found for the field $\mathbb{Q}(\sqrt{-455})$. It is a pleasure to acknowledge our debt to Ulrich Görtz and Gabor Wiese for allowing us to use the computer nodes at the Institute for Experimental Mathematics of the University of Duisburg-Essen and the Mathematics Department of the University of Luxembourg respectively. Also, we are grateful to the Weizmann Institute of Science for

$ D $	431	435	436	439	440	443	447	451	452
h	21	4	6	15	12	5	14	6	8
$k \leq$	8	13	11	9	8	12	10	12	11

$ D $	455	456	463	467	471	472	479	483	487
h	20	8	7	7	16	6	25	4	7
$k \leq$	7	10	8	10	7	11	7	11	8

$ D $	488	491	499	520	532	547	555	560	568
h	10	9	3	4	4	3	4	4	4
$k \leq$	9	10	10	9	10	11	11	7	10

$ D $	571	595	627	643	667	696	708	715	723
h	5	4	4	3	4	12	4	4	4
$k \leq$	11	9	11	9	9	4	7	7	9

$ D $	760	763	795	883	907	955	1003	1027	1051
h	4	4	4	3	3	4	4	4	5
$k \leq$	7	7	7	6	7	6	6	5	5

$ D $	1123	1227	1243	1387	1411	1507	1555		
h	5	4	4	4	4	4	4		
$k \leq$	4	5	4	4	4	4	4		

TABLE 4. the scope of our computations, part 2

$ D $	7	11	71	87	91	155	199	223	231	339
k	12	10	1	2	6	4	1	0	4	1
dim	2	2	2	2	2	2	4	2	2	2

$ D $	344	407	408	408	408	415	435	435	435	435
k	1	0	2	5	8	0	2	5	8	11
dim	2	2	2	2	2	2	2	2	2	2

$ D $	455	483	571	571	643	760	1003	1003	1051	
k	0	1	0	1	0	2	0	1	0	
dim	2	2	2	2	2	2	2	2	2	

TABLE 5. the cases where there are *genuine* classes

providing the high performance computing clusters on which the database of cell complexes used for our calculations has been generated. The second author thanks the Algebra and Geometry Group of the Mathematics Department of the University of Barcelona for the post-doctoral fellowship under

which he carried out most of his work that went into this paper. Moreover, he thanks the Mathematical Sciences Research Institute of the University of California and the Max Planck Institute for Mathematics for the wonderful hospitality that he received during his stays.

1. BACKGROUND

Let K be an imaginary quadratic field with ring of integers \mathcal{O} . Let Γ be the Bianchi group $\mathrm{SL}_2(\mathcal{O})$. It is a discrete subgroup of the real Lie group $\mathrm{SL}_2(\mathbb{C})$ and thus acts discontinuously on hyperbolic 3-space. Let Y_Γ be the quotient hyperbolic 3-fold. The Borel-Serre compactification, see [Ser70, appendix], X_Γ of Y_Γ is a compact 3-fold with boundary ∂X_Γ whose interior is homeomorphic to Y_Γ . When the discriminant of K is smaller than -4 , ∂X_Γ consists of h_K disjoint 2-tori where h_K is the class number of K .

Given $n \geq 0$, let $\mathbb{C}[x, y]_n$ denote the space of homogeneous polynomials of degree n on variables x, y with complex coefficients. $\mathrm{SL}_2(\mathbb{C})$ acts on this space in the obvious way permitted by the two variables. Consider the $\mathrm{SL}_2(\mathbb{C})$ -module

$$E_n := \mathbb{C}[x, y]_n \otimes_{\mathbb{C}} \overline{\mathbb{C}[x, y]_n}$$

where the overline on the second factor is to indicate that the action on this factor is twisted with complex conjugation. When considered as a Γ -module, E_n gives rise to a locally constant sheaf \mathcal{E}_n on Y_Γ whose stalks are isomorphic to E_n . Consider the long exact sequence

$$\dots \rightarrow H_c^i(Y_\Gamma; \mathcal{E}_n) \rightarrow H^i(X_\Gamma; \bar{\mathcal{E}}_n) \rightarrow H^i(\partial X_\Gamma; \bar{\mathcal{E}}_n) \rightarrow \dots$$

where H_c^i denotes the compactly supported cohomology and $\bar{\mathcal{E}}_n$ is a certain natural extension of \mathcal{E}_n to X_Γ .

The *cuspidal cohomology* H_{cusp}^i is defined as the image of the compactly supported cohomology. The *Eisenstein cohomology* H_{Eis}^i is the complement of the cuspidal cohomology inside H^i and it is isomorphic to the image of the restriction map inside the cohomology of the boundary. The decomposition $H^i = H_{cusp}^i \oplus H_{Eis}^i$ respects the Hecke action which is defined, as usual, via correspondences on X_Γ .

By construction, the embedding $Y_\Gamma \hookrightarrow X_\Gamma$ is a homotopy invariance. Together with the fact that Y_Γ is a $K(\Gamma, 1)$ -space, we get the isomorphisms

$$H^i(X_\Gamma; \bar{\mathcal{E}}_n) \simeq H^i(Y_\Gamma; \mathcal{E}_n) \simeq H^i(\Gamma; E_n).$$

Via these isomorphisms, we define the cuspidal and Eisenstein parts of $H^i(\Gamma; E_n)$.

Let $S_n(1)$ denote the space of level one cuspidal Bianchi modular forms (over K) of weight $n + 2$. It is well known that

$$S_n(1) \simeq H_{cusp}^1(Y_\Gamma; \mathcal{E}_n) \simeq H_{cusp}^2(Y_\Gamma; \mathcal{E}_n)$$

as Hecke modules. Here the first isomorphism was established by Harder and the second follows from duality, see [AS86].

In [FGT10], a formula for the dimension of the space $L_n(1)$ has been given for all fields K and weights n . We will compare the dimension of $L_n(1)$, which we obtain via their formula, to the dimension of $S_n(1)$, which we will obtain via our computer programs. The following Proposition will allow us to deduce the size of the cuspidal cohomology, and hence of $S_n(1)$, once we have computed the size of the whole cohomology. It is well-known to the specialists, however for the convenience of the reader we include a proof of it.

Proposition 1. *Let K be an imaginary quadratic field of discriminant absolute value greater than 4. Then*

$$\dim H_{Eis}^2(X_\Gamma; \bar{\mathcal{E}}_n) = \begin{cases} h_K - 1, & \text{if } n = 0, \\ h_K, & \text{else.} \end{cases}$$

where h_K is the class number of K .

Proof. It is well-known (see, e.g. [Har75]) that the map

$$H^2(X_\Gamma; \bar{\mathcal{E}}_n) \longrightarrow H^2(\partial X_\Gamma; \bar{\mathcal{E}}_n)$$

is surjective for $n > 0$ and its image has codimension one for $n = 0$. By our assumption on K , the boundary ∂X_Γ is a disjoint union of 2-tori, indexed by the class group of K . Below we prove that the dimension of $H^2(T; \bar{\mathcal{E}}_n)$ is one for every boundary component T of ∂X_Γ , which clearly gives our claim.

Let $c \in K \cup \{\infty\}$ be a cusp and let Γ_c be its stabilizer in Γ (which is a parabolic subgroup). Then Γ_c is the fundamental group of T_c . In fact, T_c is a $K(\Gamma_c, 1)$ -space. Hence we can turn our attention to computing $H^2(\Gamma_c; E_n)$. Composition of the cup product and the well-known perfect pairing $(\cdot, \cdot) : E_n \otimes_{\mathbb{C}} E_n \rightarrow \mathbb{C}$ (see, for example, Section 2.4. [Ber08]) gives us a pairing

$$\begin{array}{ccc} H^0(\Gamma_c; E_n) \times H^2(\Gamma_c; E_n) & \xrightarrow{\cup} & H^2(\Gamma_c; E_n \otimes_{\mathbb{C}} E_n) \\ & & \downarrow (\cdot, \cdot) \\ & & H^2(\Gamma_c; \mathbb{C}) \simeq \mathbb{C}. \end{array}$$

Here the last isomorphism follows from the fact that T_c is a compact 2-fold (see also proof of Prop.3.5. of [Şen11] for a direct algebraic argument). Thus the dimension we are looking for is equal to that of $H^0(\Gamma_c; E_n)$. Clearly, if $n = 0$, the latter dimension is 1 and thus the dimension of $H^2(\partial X_\Gamma; \bar{\mathcal{E}}_n)$ is h_K as desired.

Let us now assume that $n \neq 0$. Conjugation by a matrix in $SL(K)$ which takes c to the cusp at infinity induces an isomorphism $\Gamma_c \simeq \Gamma_\infty = \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \subset SL_2(\mathcal{O}_K)$. Consider the normal subgroup $\Gamma_\infty^+ := \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}$ of Γ_∞ . Then Γ_∞^+ is a free Abelian group on two generators. We are now going to determine the submodule $E_n^{\Gamma_\infty^+}$ of E_n invariant under its action. As the generators are of the form $\begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}$, it is clear that the vector $x^n \otimes x^n$ is fixed by Γ_∞^+ . One

shows, proceeding as in Lemma 2.4. of [Wie07], that there are no other fixed vectors. Hence

$$H^0(\Gamma_\infty^+; E_n) = E_n^{\Gamma_\infty^+} = \langle x^n \otimes x^n \rangle$$

is of complex dimension one. Let $\mu := \Gamma_\infty/\Gamma_\infty^+ = \left\{ \begin{pmatrix} \pm 1 & 0 \\ 0 & \pm 1 \end{pmatrix} \right\}$. As we are considering modules over \mathbb{C} , it follows that

$$H^0(\Gamma_\infty; E_n) \simeq H^0(\Gamma_\infty^+; E_n)^\mu$$

is the invariant submodule under μ . We easily check that the action of μ on E_n is trivial, and so

$$H^0(\Gamma_c; E_n) \simeq H^0(\Gamma_\infty^+; E_n)$$

is again of complex dimension one. This completes the proof. \square

2. ABELIAN VARIETIES OF GL_2 -TYPE

There is a widely believed conjectural connection between Bianchi newforms of weight 2 over K and Abelian varieties of GL_2 -type defined over K (see [EGM82],[Cre92],[Tay95]) which is expressed in terms of the associated L -functions. In particular, an Abelian variety of GL_2 -type over K , that is not definable over \mathbb{Q} nor of CM -type, with everywhere good reduction is expected give rise to newforms in $S_0(1)^+$ that are not in $L_0(1)$. Here $S_0(1)^+$ denotes the *plus-subspace* of $S_0(1)$ in the sense of [EGM82] and [Cre84]. Equivalently, $S_0(1)^+$ can be seen as the space of cuspidal Bianchi modular forms of weight two for $GL_2(\mathcal{O}_K)$.

In the reverse direction, the non-lifted classes in $S_0(1)^+$ are expected to correspond to Abelian varieties of GL_2 -type over K which have everywhere good reduction. As listed in Table 5, we have found eight imaginary quadratic fields for which $S_0(1)$ contained non-lifted classes. For only six of these fields, the non-lifted classes were infact contained in $S_0(1)^+$. In Table 6 below, we list the (necessarily totally real) number field F generated by the Hecke eigenvalues of the non-lifted newforms in these six cases.

$ D $	223	415	455	571	643	1003
F	$\mathbb{Q}(\sqrt{2})$	$\mathbb{Q}(\sqrt{3})$	$\mathbb{Q}(\sqrt{5})$	$\mathbb{Q}(\sqrt{5})$	\mathbb{Q}	$\mathbb{Q}(\sqrt{7})$

TABLE 6. the number field generated by the Hecke eigenvalues of nonlifted newforms in $S_0(1)^+$

We have computed these fields using Dan Yasaki's program, see [Yas], in Magma which computes the Hecke action on $S_0(\Gamma_0(\mathfrak{n}))^+$ for congruence subgroups of type $\Gamma_0(\mathfrak{n})$ of Bianchi groups. Note that as this program only treats GL_2 -cohomology with trivial weight, that is $k = 0$, we could not have used it for our experiment.

Table 6 tells us that there should exist an elliptic curve defined over $\mathbb{Q}(\sqrt{-643})$, and not over \mathbb{Q} , which has everywhere good reduction and it should be *modular*. Indeed we know by Krämer [Krä84] that there is such an

elliptic curve over $\mathbb{Q}(\sqrt{-643})$ and it does seem to be modular, see Scheutzwow [Sch92]. Similarly, there should exist Abelian surfaces defined over $\mathbb{Q}(\sqrt{-d})$ with $d = 223, 415, 455, 571, 1003$ and not over \mathbb{Q} , which have everywhere good reduction and real multiplication by $\sqrt{2}, \sqrt{3}, \sqrt{7}, \sqrt{5}, \sqrt{7}$ respectively and they should be modular. Locating such surfaces is a highly nontrivial task, see [Şen12, Section 8].

3. COMMENTS

The data collected in this paper make it clear that the spaces of cuspidal Bianchi modular forms of level one are generically made of forms which are not genuine. Accordingly, we conjecture the following.

Conjecture 2. *Let K be an imaginary quadratic field. Let $S_k(1)$ denote the space of level one cuspidal Bianchi modular forms over K of weight $k + 2$. Then there are only finitely many k for which the space $S_k(1)$ contains non-lifted forms.*

It is interesting to make a comparison with other types of modular forms. For the case of Hilbert modular forms (this the case of the algebraic group $\text{Res}_{F/\mathbb{Q}}(\text{SL}_2)$ where F is a totally real field) and Siegel modular forms of genus 2 (the case of the algebraic group Sp_4), one has considerable amount of genuine level one cuspidal forms. However, the case of the modular forms for SL_3 is similar to our case. Here in the range of the data collected by Ash and Pollack, see [AP08], the spaces of level one modular forms for SL_3 are completely made of those which are the symmetric square lifts of classical holomorphic modular forms. They in fact conjecture that this is always the case.

It is interesting to note that for the Hilbert and Siegel modular forms of genus 2, where we have plenty of genuine forms, the associated symmetric spaces are Hermitian, while for the Bianchi and SL_3 modular forms, where there is an extreme paucity of genuine forms, the associated symmetric spaces fail to be Hermitian. Is this part of a general phenomenon ?

Next we shall discuss an analogue Maeda's conjecture for non-lifted newforms in $S_k(1)$. The nontrivial automorphism $\sigma \in \text{Gal}(K/\mathbb{Q})$ of K acts on the set of newforms in $S_k(1)$ as an involution, again denoted σ . If f is a newform in $S_k(1)$ then we have $f \mapsto \sigma f$ where the Hecke eigenvalues $c(\cdot, \pi)$ associated to the Hecke operators T_π satisfy the relation

$$c(\sigma f, \pi) = c(f, \sigma(\pi))$$

for every $\pi \in \mathcal{O}$. We will call such forms *twins*. Recall that just as in the case of elliptic modular forms, for a newform f in $S_k(1)$ with Hecke eigenvalue field F , there is a newform f^τ in $S_k(1)$ for every $\tau \in \text{Gal}(F/\mathbb{Q})$ with the property that $c(f^\tau, \pi) = \tau(c(f, \pi))$ for every $\pi \in \mathcal{O}$. We say that f and the f^τ form one *Galois orbit*.

Conjecture 3. Modified Maeda's Conjecture for Bianchi Modular Forms *Let K be an imaginary quadratic field. For every $k \geq 0$, the set of*

non-lifted newforms in $S_k(1)$, modulo the action of $\text{Gal}(K/\mathbb{Q})$, forms one Galois orbit.

In all except one of the cases where we observed non-lifted newforms, the dimension of the non-lifted subspace was only two. In this case, the conjecture above is trivially satisfied as the two non-lifted newforms have to be twins. It is crucial to check the situation $(d, k) = (199, 2)$ where we have a four dimensional subspace of non-lifted forms, however so far we have not implemented Hecke operators in our programs and Yasaki's program only works with $k = 0$. Note that the analogue of this conjecture for Hilbert modular forms over real quadratic fields holds in the range of the computations performed by Doi and Ishii, see [DHI98] p.568.

4. METHOD OF THE COMPUTATIONS

In this section, we will explain how we computed the cohomology of the investigated Bianchi groups.

Let m be a square-free positive integer and $K = \mathbb{Q}(\sqrt{-m})$ be an imaginary quadratic number field with ring of integers \mathcal{O}_{-m} , which we also just denote by \mathcal{O} . Consider the familiar action (we give an explicit formula for it in lemma 4) of the group $\Gamma := \text{SL}_2(\mathcal{O}) \subset \text{GL}_2(\mathbb{C})$ on hyperbolic three-space, for which we will use the upper-half space model \mathcal{H} .

As a set,

$$\mathcal{H} = \{(z, \zeta) \in \mathbb{C} \times \mathbb{R} \mid \zeta > 0\}.$$

Lemma 4 (Poincaré). *If $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{C})$, the action of γ on \mathcal{H} is given by $\gamma \cdot (z, \zeta) = (z', \zeta')$, where*

$$\zeta' = \frac{|\det \gamma| \zeta}{|cz - d|^2 + \zeta^2 |c|^2}, \quad z' = \frac{(\overline{d - cz})(az - b) - \zeta^2 \bar{c}a}{|cz - d|^2 + \zeta^2 |c|^2}.$$

The Bianchi–Humbert theory [Bia92], [Hum15] gives a fundamental domain for the action of Γ on \mathcal{H} , which we shall call the *Bianchi fundamental polyhedron*. It is a polyhedron in hyperbolic space up to the missing vertex ∞ , and up to a missing vertex for each non-trivial ideal class if \mathcal{O}_{-m} is not a principal ideal domain. We observe the following notion of strictness of the fundamental domain: the interior of the Bianchi fundamental polyhedron contains no two points which are identified by Γ . Swan [Swa71] proves a theorem which implies that the boundary of the Bianchi fundamental polyhedron consists of finitely many cells.

4.1. A cell complex for the Bianchi groups. Swan further produces a concept for an algorithm to compute the Bianchi fundamental polyhedron. Such an algorithm has been implemented by Cremona [Cre84] for the five cases where \mathcal{O}_{-m} is Euclidean, and by his students Whitley [Whi90] for the non-Euclidean principal ideal domain cases, Bygott [Byg98] for a case of class number 2 and Lingham ([Lin05], used in [CL07]) for some cases of class number 3; and finally Aranés [Ara10] for arbitrary class numbers.

Another algorithm based on this concept has independently implemented in [Rah10a] for all Bianchi groups; and we make explicit use of the cell complexes it produces. Other results of the employed implementation are described in [Rah10b].

We can check that the computed polyhedron is indeed a fundamental domain for Γ using the following observation of Poincaré [Poi83]: After a cell subdivision which makes the cell stabilizers fix the cells point-wise, the 2-cells (“faces”) of the fundamental polyhedron appear in pairs $(\sigma, \gamma \cdot \sigma)$ with $\gamma \in \Gamma$ — so for every orbit of faces, we have exactly two representatives — such that with the orientation for which the lower side of the face σ lies on the polyhedron, the upper side of $\gamma \cdot \sigma$ lies on the polyhedron. We induce a cell structure on \mathcal{H} by the images under Γ of the faces, edges and vertices of the Bianchi fundamental polyhedron.

4.2. The Flöge cellular complex. In order to obtain a cell complex with compact quotient space, we proceed in the following way due to Flöge [Flö83]. The boundary of \mathcal{H} is the Riemann sphere $\partial\mathcal{H}$, which, as a topological space, is made up of the complex plane \mathbb{C} compactified with the cusp ∞ . The totally geodesic surfaces in \mathcal{H} are the Euclidean vertical planes (we define *vertical* as orthogonal to the complex plane) and the Euclidean hemispheres centered on the complex plane. The action of the Bianchi groups extends continuously to the boundary $\partial\mathcal{H}$. The cellular closure of the refined cell complex in $\mathcal{H} \cup \partial\mathcal{H}$ consists of \mathcal{H} and $(\mathbb{Q}(\sqrt{-m}) \cup \{\infty\}) \subset (\mathbb{C} \cup \{\infty\}) \cong \partial\mathcal{H}$. The $\mathrm{SL}_2(\mathcal{O}_{-m})$ -orbit of a cusp $\frac{\lambda}{\mu}$ in $(\mathbb{Q}(\sqrt{-m}) \cup \{\infty\})$ corresponds to the ideal class $[(\lambda, \mu)]$ of \mathcal{O}_{-m} . It is well-known that this does not depend on the choice of the representative $\frac{\lambda}{\mu}$. We extend the refined cell complex to a cell complex \tilde{X} by joining to it, in the case that \mathcal{O}_{-m} is not a principal ideal domain, the $\mathrm{SL}_2(\mathcal{O}_{-m})$ -orbits of the cusps $\frac{\lambda}{\mu}$ for which the ideal (λ, μ) is not principal. We call the latter cusps the *singular* cusps. At the singular cusps, we equip \tilde{X} with the “horoball topology” described in [Flö83]. This simply means that the set of cusps, which is discrete in $\partial\mathcal{H}$, is located at the hyperbolic extremities of \tilde{X} : No neighborhood of a cusp, except the whole \tilde{X} , contains any other cusp.

We retract \tilde{X} in the following, $\mathrm{SL}_2(\mathcal{O}_{-m})$ -equivariant, way. On the Bianchi fundamental polyhedron, the retraction is given by the vertical projection (away from the cusp ∞) onto its facets which are closed in $\mathcal{H} \cup \partial\mathcal{H}$. The latter are the facets which do not touch the cusp ∞ , and are the bottom facets with respect to our vertical direction. The retraction is continued on \mathcal{H} by the group action. It is proven in [Flö80] that this retraction is continuous. We call the retract of \tilde{X} the *Flöge cellular complex* and denote it by X . So in the principal ideal domain cases, X is a retract of the refined cell complex, obtained by contracting the Bianchi fundamental polyhedron onto its cells which do not touch the boundary of \mathcal{H} . In [RF11], it is checked that the Flöge cellular complex is contractible.

4.3. The spectral sequence. Let X be our Flöge complex constructed as above. Next we will consider the spectral sequence associated to the double complex $\text{Hom}_{\mathbb{Z}\Gamma}(\Theta_*, C_{\mathbb{Z}}^*(X, M))$, where Θ_* is the standard resolution of \mathbb{Z} over $\mathbb{Z}\Gamma$ and $C^*(X, M)$ is the cellular co-chain complex of X with $\mathbb{Z}\Gamma$ -module coefficients M . We can (see [Bro82], p. 164) derive the first-quadrant spectral sequence

$$E_1^{p,q}(M) = \bigoplus_{\sigma \in \Sigma_p} H^q(\Gamma_\sigma; M) \implies H^{p+q}(\Gamma; M)$$

where Σ_p denotes the Γ -conjugacy classes of p -cells of X . Observe that Γ_σ will be a finite group whose order is divisible only by 2 and/or 3 unless σ is the class of a singular cusp, in which case Γ_σ is a free Abelian group on two unipotent generators.

Assume that M admits an additional module structure over a ring where 6 is invertible (in fact we are interested in the case where M is a complex vector space). Then the finite ones among the higher cohomology groups of the Γ_σ vanish. Thus, when there are no singular cusps (equivalently, when the class number of \mathcal{O} is one), the spectral sequence concentrates on the row $q = 0$ and stabilizes on the E^2 -page. Otherwise, the spectral sequence concentrates on the rows $q = 0, 1, 2$ and stabilizes at the E^3 -page.

As we shall see below, the dimension of the module $H^2(\Gamma; M)$, which we want to determine, is the same as the dimension of

$$E_2^{2,0} \simeq E_1^{2,0} / \text{Im}(d_1^{1,0}),$$

where the differential $d_1^{1,0}$ is given as

$$E_1^{1,0} \simeq \bigoplus_{\sigma \in \Sigma_1} M^{\Gamma_\sigma} \longrightarrow \bigoplus_{\sigma \in \Sigma_2} M \simeq E_1^{2,0}.$$

The abutment of the spectral series gives us

$$H^2(\Gamma; M) \simeq E_3^{2,0} \oplus E_3^{0,2}.$$

Here $E_3^{0,2} \simeq \bigoplus_s H^2(\Gamma_s; M)$ where the summation is over Γ -classes of singular cusps s .

Moreover, $E_3^{2,0} = E_2^{2,0} / \text{Im}(d_2^{0,1})$ where the differential $d_2^{0,1}$ is given as

$$\bigoplus_{s \text{ singular}} H^1(\Gamma_s; M) \longrightarrow E_2^{2,0}.$$

We determine the rank of this differential as follows.

Theorem 5 ([Ser70]). *Suppose that the coefficient module M is equipped with a non-degenerate Γ -invariant \mathbb{C} -bilinear form. Then the rank of the map from $H^1(\Gamma; M)$ to the disjoint sum of the $H^1(\Gamma_s; M)$, induced by restriction from $H^1(\Gamma; M)$ to $H^1(\Gamma_s; M)$, equals half of the rank of the disjoint sum of the $H^1(\Gamma_s; M)$.*

Let us assume from now on that $M = E_n$ for some n . As we have seen in the proof of Proposition 1, there is a perfect pairing on M , which is a non-degenerate Γ -invariant \mathbb{C} -bilinear form. So the theorem of Serre applies, and we obtain the following corollary. Note for this purpose that the proof of Proposition 1 shows that

$$\dim H^0(\Gamma_s; M) = \dim H^2(\Gamma_s; M) = 1.$$

As Γ_s is the fundamental group of a torus, we have

$$\dim H^1(\Gamma_s; M) = 2 \cdot \dim H^2(\Gamma_s; M) = 2.$$

Corollary 6. *The rank of the differential*

$$d_2^{0,1} : \bigoplus_{s \text{ singular}} H^1(\Gamma_s; M) \longrightarrow E_2^{2,0}$$

is the number of non-trivial ideal classes.

Remark 7. The above discussion implies that

$$H^2(\Gamma; M) \simeq \left(\bigoplus_{s \text{ singular}} H^2(\Gamma_s; M) \right) \oplus \left(E_2^{2,0} / \text{Im}(d_2^{0,1}) \right),$$

and the dimension of $H^2(\Gamma; M)$ is the same as that of $E_2^{2,0}$.

4.4. The procedure of the computations. We compute the representatives of faces in $E_1^{2,0}$ and the differential $d_1^{2,0}$ of our equivariant spectral sequence with trivial integer coefficients with the program *Bianchi.gp* [Rah10a]. The second author has implemented a MAGMA script that uses the cell stabilizers and identifications obtained with *Bianchi.gp* to compute the action on the coefficient module M that we are interested in. We then deduce the term $E_1^{2,0}$ and the differential $d_1^{2,0}$ with respect to our coefficients. The quotient

$$E_2^{2,0} \cong E_1^{2,0} / \text{image } d_1^{2,0}$$

now admits the dimension of $H^2(\Gamma; M)$ by Remark 7.

As linear algebra over number fields is more expensive compared to working over finite fields, we employ the following shortcut. Recall that by the universal coefficients theorem, the dimension of $H^2(\Gamma; M(\mathbb{F}_p))$ ("the mod p dimension") is greater than or equal to the dimension of $H^2(\Gamma; M(\mathbb{C}))$ ("the complex dimension"). We start computing the mod p -dimensions for primes $p \leq 200$. If we find for a particular p for which the mod p dimension is equal to the lower bound of Finis-Grunewald-Tirao then we infer that the complex dimension is equal to the mod p dimension. Note that by Prop. 3.2 (d) of [Şen11], this implies that $H^2(\Gamma; M(\mathcal{O}))$ has no p -torsion. If this is not the case for the primes in our range, then we compute the complex dimension directly by computing $H^2(\Gamma; M(K))$.

4.5. Execution of the computations. We applied the above described computations to a database of cell complexes for 183 Bianchi groups, which has been established on the computing clusters of the Weizmann Institute of Science, using over fifty processor-months. This database includes all the cases of ideal class numbers 3 and 5, most of the cases of ideal class number 4 and *all* cases with the absolute value of the discriminant less than 500. Almost all of our dimension computations were carried out using the nodes of the computer clusters at the Universities of Duisburg-Essen and Luxembourg.

REFERENCES

- [AP08] A. Ash and D. Pollack, *Everywhere unramified automorphic cohomology for $SL_3(\mathbb{Z})$* , Int. J. Number Theory **4** (2008), no. 4, 663–675.
- [Ara10] M.T. Aranés, *Modular symbols over number fields*, Ph.D. thesis, University of Warwick, 2010.
- [AS86] A. Ash and G. Stevens, *Cohomology of arithmetic groups and congruences between systems of Hecke eigenvalues*, J. Reine Angew. Math. **365** (1986), 192–220.
- [Ber08] T. Berger, *Denominators of Eisenstein cohomology classes for GL_2 over imaginary quadratic fields*, Manuscripta Math. **125** (2008), no. 4, 427–470.
- [Bia92] L. Bianchi, *Sui gruppi di sostituzioni lineari con coefficienti appartenenti a corpi quadratici immaginari*, Math. Ann. **40** (1892), no. 3, 332–412.
- [Bro82] K.S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, vol. **87**, Springer-Verlag, 1982.
- [Byg98] J. Bygott, *Modular forms and modular symbols over imaginary quadratic fields*, Ph.D. thesis, University of Exeter, 1998.
- [CL07] J.E. Cremona and M.P. Lingham, *Finding all elliptic curves with good reduction outside a given set of primes*, Experiment. Math. **16** (2007), no. 3, 303–312.
- [CM09] F. Calegari and B. Mazur, *Nearly ordinary Galois deformations over arbitrary number fields*, J. Inst. Math. Jussieu **8** (2009), no. 1, 99–177.
- [Cre84] J.E. Cremona, *Hyperbolic tessellations, modular symbols, and elliptic curves over complex quadratic fields*, Compositio Math. **51** (1984), no. 3, 275–324.
- [Cre92] ———, *Abelian varieties with extra twist, cusp forms, and elliptic curves over imaginary quadratic fields*, J. London Math. Soc. (2) **45** (1992), no. 3, 404–416.
- [DHI98] K. Doi, H. Hida, and H. Ishii, *Discriminant of Hecke fields and twisted adjoint L -values for $GL(2)$* , Invent. Math. **134** (1998), no. 3, 547–577.
- [EGM82] J. Elstrodt, F. Grunewald, and J. Mennicke, *On the group $PSL_2(\mathbb{Z}[i])$* , Number theory days, 1980 (Exeter), London Math. Soc. Lecture Note Ser., vol. 56, Cambridge Univ. Press, Cambridge, 1982, pp. 255–283.
- [FGT10] T. Finis, F. Grunewald, and P. Tirao, *The cohomology of lattices in $SL(2, \mathbb{C})$* , Experiment. Math. **19** (2010), no. 1, 29–63.
- [Flö80] D. Flöge, *Zur Struktur der PSL_2 über einigen imaginär-quadratischen Zahlringen*, Dissertation, Johann-Wolfgang-Goethe-Universität, Fachbereich Mathematik, 1980.
- [Flö83] ———, *Zur Struktur der PSL_2 über einigen imaginär-quadratischen Zahlringen*, Math. Z. **183** (1983), no. 2, 255–279.
- [Har75] G. Harder, *On the cohomology of discrete arithmetically defined groups*, Discrete subgroups of Lie groups and applications to moduli (Internat. Colloq., Bombay, 1973) (1975), 129–160.
- [Hum15] G. Humbert, *Sur la réduction des formes d’Hermite dans un corps quadratique imaginaire*, C. R. Acad. Sci. Paris **16** (1915), 189–196.

- [Krä84] N. Krämer, *Beiträge zur Arithmetik imaginärquadratischer Zahlkörper*, Ph.D. thesis, Math.-Naturwiss. Fakultät der Rheinischen Friedrich-Wilhelms-Universität Bonn; Bonn. Math. Schr., 1984.
- [Lin05] M. Lingham, *Modular forms and elliptic curves over imaginary quadratic fields*, Ph.D. thesis, University of Nottingham, 2005.
- [Poi83] H. Poincaré, *Mémoire*, Acta Math. **3** (1883), no. 1, 49–92, Les groupes Kleinéens.
- [Rah10a] A. D. Rahm, *Bianchi.gp*, Open source program (GNU general public license), validated by the CNRS: www.projet-plume.org/fiche/bianchigp
Based on Pari/GP, 2010.
- [Rah10b] ———, *(co)homologies and k -theory of Bianchi groups using computational geometric models*, Ph.D. thesis, University of Grenoble and University of Goettingen, 2010.
- [RF11] A.D. Rahm and M. Fuchs, *The integral homology of PSL_2 of imaginary quadratic integers with non-trivial class group*, J. Pure and Applied Algebra **215** (2011), 1443–1472.
- [Sch92] A. Scheutzwow, *Computing rational cohomology and Hecke eigenvalues for Bianchi groups*, J. Number Theory **40** (1992), no. 3, 317–328.
- [Sen11] M.H. Şengün, *On the integral cohomology of Bianchi groups*, Experiment. Math. **20** (2011), no. 4, 487–505.
- [Sen12] ———, *Arithmetic aspects of Bianchi groups*, preprint (2012).
- [Ser70] J.-P. Serre, *Le problème des groupes de congruence pour $SL(2)$* , Ann. of Math. (2) **92** (1970), 489–527.
- [Swa71] R.G. Swan, *Generators and relations for certain special linear groups*, Advances in Math. **6** (1971), 1–77.
- [Tay95] R. Taylor, *Representations of Galois groups associated to modular forms*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994), Birkhäuser, Basel, 1995, pp. 435–442.
- [Whi90] E. Whitley, *Modular symbols and elliptic curves over imaginary quadratic fields*, Ph.D. thesis, University of Exeter, 1990.
- [Wie07] G. Wiese, *On the faithfulness of parabolic cohomology as a Hecke module over a finite field*, J. Reine Angew. Math. **606** (2007), 79–103.
- [Yas] D. Yasaki, *Hyperbolic tessellations associated to Bianchi groups*, 9th International Symposium, Nancy, France, ANTS-IX, July 19-23, 2010, Proceedings.

E-mail address: Alexander.Rahm@nuigalway.ie

URL: <http://www.maths.nuigalway.ie/~rahm/>

NATIONAL UNIVERSITY OF IRELAND AT GALWAY, DEPARTMENT OF MATHEMATICS

E-mail address: M.H.Sengun@warwick.ac.uk

URL: <http://www.uni-due.de/~hm0074/>

MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY, UK