



HAL
open science

Bilinear decompositions and commutators of singular integral operators

Luong Dang Ky

► **To cite this version:**

Luong Dang Ky. Bilinear decompositions and commutators of singular integral operators. 2011.
hal-00589824v1

HAL Id: hal-00589824

<https://hal.science/hal-00589824v1>

Preprint submitted on 2 May 2011 (v1), last revised 13 Jan 2012 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BILINEAR DECOMPOSITIONS AND COMMUTATORS OF SINGULAR INTEGRAL OPERATORS

LUONG DANG KY

ABSTRACT. Let b be a BMO -function. It is well-known that the linear commutator $[b, T]$ of a Calderón-Zygmund operator T does not, in general, map continuously $H^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$. However, Pérez [38] showed that if $H^1(\mathbb{R}^n)$ is replaced by a suitable atomic subspace $\mathcal{H}_b^1(\mathbb{R}^n)$ then the commutator is continuous from $\mathcal{H}_b^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$. In this paper, we find the largest subspace $H_b^1(\mathbb{R}^n)$ such that all commutators of Calderón-Zygmund operators are continuous from $H_b^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$. We also study the commutators $[b, T]$ for T in a class \mathcal{K} of sublinear operators containing almost all important operators in Harmonic analysis. When T is linear, we prove that there exists a bilinear operators \mathfrak{R} map continuously $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$ such that for all $(f, b) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$, we have

$$(1) \quad [b, T](f) = \mathfrak{R}(f, b) + T(\mathfrak{S}(f, b)),$$

where \mathfrak{S} is the bounded bilinear operator from $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$ does not depend on T . In the particular case of T a Calderón-Zygmund operator satisfying $T1 = T^*1 = 0$ and b is in $BMO^{\text{log}}(\mathbb{R}^n)$. We prove that the commutator $[b, T]$ maps continuously $H_b^1(\mathbb{R}^n)$ into $h^1(\mathbb{R}^n)$. Furthermore, if $T^*b = 0$ then the commutator maps continuously $H_b^1(\mathbb{R}^n)$ into $H^1(\mathbb{R}^n)$.

When T is sublinear, we prove that there exists a bounded subbilinear operator $\mathfrak{R} : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ such that for all $(f, b) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$, we have

$$(2) \quad |T(\mathfrak{S}(f, b))| - \mathfrak{R}(f, b) \leq |[b, T](f)| \leq \mathfrak{R}(f, b) + |T(\mathfrak{S}(f, b))|.$$

The bilinear decomposition (1) and the subbilinear decomposition (2) allow us to give a general overview all known weak and strong L^1 -estimates.

1. INTRODUCTION

Given a function b locally integrable on \mathbb{R}^n , and a Calderón-Zygmund operator T , we consider the linear commutator $[b, T]$ defined for smooth, compactly supported functions f by

$$[b, T](f) = bT(f) - T(bf).$$

2010 *Mathematics Subject Classification.* 42B20 (42B30, 42B35, 42B25).

Key words and phrases. Calderón-Zygmund operators, bilinear decompositions, commutators, Hardy spaces, wavelet characterizations, atoms, bilinear operators.

A classical result of R. Coifman, R. Rochberg and G. Weiss (see [10]), states that the commutator $[b, T]$ is continuous on $L^p(\mathbb{R}^n)$ for $1 < p < \infty$, when $b \in BMO(\mathbb{R}^n)$. Unlike the theory of Calderón-Zygmund operators, the proof of this result does not rely on a weak type $(1, 1)$ estimate for $[b, T]$. In fact, it was shown in [38] that, in general, the linear commutator fails to be of weak type $(1, 1)$, when b is in $BMO(\mathbb{R}^n)$. Instead, an endpoint theory was provided for this operator. It is well-known that any singular integral operator maps $H^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$. However, it was observed in [20] that the commutator $[b, H]$ with b in $BMO(\mathbb{R})$, where H is Hilbert transform on \mathbb{R} , does not map, in general, $H^1(\mathbb{R})$ into $L^1(\mathbb{R})$. Instead of this, the weak type estimate (H^1, L^1) for $[b, T]$ is well-known see for example [27, 32, 42]. Remark that intuitively one would like to write

$$[b, T](f) = \sum_{j=1}^{\infty} \lambda_j (b - b_{B_j}) T(a_j) - T\left(\sum_{j=1}^{\infty} \lambda_j (b - b_{B_j}) a_j\right),$$

where $f = \sum_{j=1}^{\infty} \lambda_j a_j$ a atomic decomposition of f and b_{B_j} the average of b on B_j . This is equivalent to ask for a commutation property

$$(1.1) \quad \sum_{j=1}^{\infty} \lambda_j b_{B_j} T(a_j) = T\left(\sum_{j=1}^{\infty} \lambda_j b_{B_j} a_j\right).$$

Even if most authors, for instance in [27, 32, 42, 43, 25, 41, 26], implicitly use (1.1), one must be careful at this point. Indeed, the equality (1.1) is not clear since the two series $\sum_{j=1}^{\infty} \lambda_j b_{B_j} T(a_j)$ and $\sum_{j=1}^{\infty} \lambda_j b_{B_j} a_j$ are not yet well-defined, in general. We refer the reader to [6], Section 3, to see why one must be careful with the equality (1.1).

Although the commutator $[b, T]$ does not map continuously, in general, $H^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$, following Pérez [38] one can find a subspace $\mathcal{H}_b^1(\mathbb{R}^n)$ of $H^1(\mathbb{R}^n)$ such that $[b, T]$ maps continuously $\mathcal{H}_b^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$. Recall that (see [38]) a function a is a b -atom if

- i) $\text{supp } a \subset Q$ for some cube Q ,
- ii) $\|a\|_{L^\infty} \leq |Q|^{-1}$,
- iii) $\int_{\mathbb{R}^n} a(x) dx = \int_{\mathbb{R}^n} a(x) b(x) dx = 0$.

The space $\mathcal{H}_b^1(\mathbb{R}^n)$ consists of the subspace of $L^1(\mathbb{R}^n)$ of functions f which can be written as $f = \sum_{j=1}^{\infty} \lambda_j a_j$ where a_j are b -atoms, and λ_j are complex numbers with $\sum_{j=1}^{\infty} |\lambda_j| < \infty$.

In [38] the author showed that the commutator $[b, T]$ is bounded from $\mathcal{H}_b^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$ by establishing that

$$(1.2) \quad \sup\{\|[b, T](a)\|_{L^1} : a \text{ is a } b\text{-atom}\} < \infty.$$

This leaves a gap in the proof which we fill here. Indeed, as it is pointed out in [6] that there exists a linear operator U defined on the space of all finite

linear combination of $(1, \infty)$ -atoms satisfying

$$\sup\{\|U(a)\|_{L^1} : a \text{ is a } (1, \infty)\text{-atom}\} < \infty,$$

but which does not admit an extension to bounded operator from $H^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$. By this fact, we see that Inequality (1.2) does not suffice to conclude that $[b, T]$ is bounded from $\mathcal{H}_b^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$. In the setting of $H^1(\mathbb{R}^n)$, it is well-known (see [34] for details) that a linear operator U can be extended to a bounded operator from $H^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$ if for some $1 < q < \infty$, we have

$$\sup\{\|U(a)\|_{L^1} : a \text{ is a } (1, q)\text{-atom}\} < \infty.$$

It follows from the fact that the finite atomic norm on $H_{\text{fin}}^{1,q}(\mathbb{R}^n)$ is equivalent with the standard infinite atomic decomposition norm on $H_{\text{ato}}^{1,q}(\mathbb{R}^n)$ through the grand maximal function characterization of $H^1(\mathbb{R}^n)$. However, one can not use this method in the context of $\mathcal{H}_b^1(\mathbb{R}^n)$.

Also, a natural question arising from the fact of the space $\mathcal{H}_b^1(\mathbb{R}^n)$ is could one find *the largest subspace of $H^1(\mathbb{R}^n)$* (of course, this space contains $\mathcal{H}_b^1(\mathbb{R}^n)$, see also Theorem 5.2) such that all commutators $[b, T]$ of Calderón-Zygmund operators are bounded from this space to $L^1(\mathbb{R}^n)$? In this note we give the positive answer (see Remark 5.1). In particular, we consider the space $H_b^1(\mathbb{R}^n)$ consisting of all $f \in H^1(\mathbb{R}^n)$ such that the (sublinear) commutator $[b, \mathfrak{M}]$ of f belongs to $L^1(\mathbb{R}^n)$ where \mathfrak{M} is the nontangential grand maximal operator (see Section 2). The norm on $H_b^1(\mathbb{R}^n)$ is defined by $\|f\|_{H_b^1} := \|f\|_{H^1} \|b\|_{BMO} + \|[b, \mathfrak{M}](f)\|_{L^1}$. Here we just consider b is a non-constant BMO -function since the commutator $[b, T] = 0$ if b is a constant function. We then prove that $[b, T]$ is bounded from $H_b^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$ for every T a Calderón-Zygmund singular integral operator (in fact it holds for all $T \in \mathcal{K}$, see below). Furthermore, $H_b^1(\mathbb{R}^n)$ is *the largest space having this property*. Besides, we also consider the class \mathcal{K} of all sublinear operators T , bounded from $H^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$, satisfying the condition

$$\|(b - b_Q)Ta\|_{L^1} \leq C\|b\|_{BMO}$$

for all BMO -function b , H^1 -atom a related to the cube Q . Here b_Q denotes the average of b on Q , and $C > 0$ is a constant independent of b, a . This class \mathcal{K} contains almost all important operators in Harmonic analysis: Calderón-Zygmund type operators, strongly singular integral operators, multiplier operators, pseudo-differential operators, maximal type operators, the area integral operator of Lusin, Littlewood-Paley type operators, Marcinkiewicz operators, maximal Bochner-Riesz operators, etc (see Section 4). When T is linear and belongs to \mathcal{K} , we prove that there exists a bounded bilinear operators $\mathfrak{R} : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ such that for all $(f, b) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$, we have the following bilinear decomposition

$$(1.3) \quad [b, T](f) = \mathfrak{R}(f, b) + T(\mathfrak{S}(f, b)),$$

where \mathfrak{S} is the bounded bilinear operator from $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$ does not depend on T (see Section 3). This bilinear decomposition is strongly related to our previous result in [4] on paraproduct and product on $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$.

We then prove that $[b, T]$ is bounded from $H_b^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$ (see Theorem 3.3). The main tools in our proof are the bilinear decomposition (1.3) (see Theorem 3.2) and the weak convergence theorem in $H^1(\mathbb{R}^n)$ of Jones and Journé (see Theorem 5.1). Also, the weak type estimate (H^1, L^1) for the commutator $[b, T]$, where T is a Calderón-Zygmund type operator, a strongly singular integral operator, a multiplier operator or a pseudo-differential operator, can be seen as an immediate corollary of the bilinear decomposition (1.3). We should also point out that weak type estimates and Hardy type estimates for the (linear) commutators of multiplier operators [43, 25, 41] and of strongly singular Calderón-Zygmund operators [26] have been studied recently.

In the case of T a Calderón-Zygmund operator satisfying $T1 = T^*1 = 0$ and b is in $BMO^{\text{log}}(\mathbb{R}^n)$ (see Section 2) we prove that the linear commutator $[b, T]$ maps continuously $H_b^1(\mathbb{R}^n)$ into $h^1(\mathbb{R}^n)$. Moreover, if $T^*b = 0$ then the linear commutator $[b, T]$ maps continuously $H_b^1(\mathbb{R}^n)$ into $H^1(\mathbb{R}^n)$. A difficulty point in the proofs of our results is that we have to directly do with $f \in H_b^1(\mathbb{R}^n)$. It will be easier to do with atomic Hardy spaces (for instance $\mathcal{H}_b^1(\mathbb{R}^n)$), and thus is $H_b^1(\mathbb{R}^n)$ if this space has a atomic characterization. However, we do not know that *does there exists a atomic characterization for the space $H_b^1(\mathbb{R}^n)$?* *This is still an open question.*

Let X be a Banach space. We say that an operator $T : X \rightarrow L^1(\mathbb{R}^n)$ is a sublinear operator if for all $f, g \in X$ and $\alpha, \beta \in \mathbb{C}$, we have

$$|T(\alpha f + \beta g)(x)| \leq |\alpha| |Tf(x)| + |\beta| |Tg(x)|.$$

Obviously, a linear operator $T : X \rightarrow L^1(\mathbb{R}^n)$ is a sublinear operator. We also say that a operator $\mathfrak{T} : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ is a subbilinear operator if for all $(f, g) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ the operators $\mathfrak{T}(f, \cdot) : BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ and $\mathfrak{T}(\cdot, g) : H^1(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ are sublinear operators.

In this paper, we also obtain the subbilinear decomposition for sublinear commutator. More precisely, when $T \in \mathcal{K}$ is a sublinear operator, we prove that there exists a bounded subbilinear operator $\mathfrak{R} : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ so that for all $(f, b) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$, we have

$$(1.4) \quad |T(\mathfrak{S}(f, b))| - \mathfrak{R}(f, b) \leq |[b, T](f)| \leq \mathfrak{R}(f, b) + |T(\mathfrak{S}(f, b))|.$$

Then, the strong type estimate (H_b^1, L^1) and the weak type estimate (H^1, L^1) of the commutators of Littlewood-Paley type operators, of Marcinkiewicz operators, and of maximal Bochner-Riesz operators, can be viewed as an immediate corollary of (1.4). When $H_b^1(\mathbb{R}^n)$ is replaced by $\mathcal{H}_b^1(\mathbb{R}^n)$, these type of estimates have also been considered recently (see for example [28, 7, 33, 30, 31, 29]).

Let us emphasize the three main purposes of this paper. First, we want to give the bilinear (resp., subbilinear) decompositions for the linear (resp., sublinear) commutators. Second, we find the largest subspace of $H^1(\mathbb{R}^n)$ such that all commutators of Calderón-Zygmund operators map continuously this space into $L^1(\mathbb{R}^n)$. Finally, we obtain the (H_b^1, h^1) type estimates and the (H_b^1, H^1) type estimates of commutators of Calderón-Zygmund operators.

Our paper is organized as follows. In Section 2 we present some notations and preliminaries to the Calderón-Zygmund operators, the function spaces which we use, and a short introduction to wavelets, a useful tool in our work. In Section 3 we state our two decomposition theorems (Theorem 3.1 and Theorem 3.2), the (H_b^1, L^1) type estimates for commutators (Theorem 3.3), and some remarks. The bilinear type estimates for commutators of Calderón-Zygmund operators (Theorem 3.4) and the boundedness of commutators of Calderón-Zygmund operators on Hardy spaces are also given in this section. In Section 4 we give some examples of operators in the class \mathcal{K} and recall our result from [4] which decomposes a product of f in $H^1(\mathbb{R}^n)$ and g in $BMO(\mathbb{R}^n)$ as a sum of images by four bilinear operators defined through wavelets. These operators are fundamental for the two decomposition theorems. In Section 5 we study the space $H_b^1(\mathbb{R}^n)$. Section 6 and 7 are devoted to the proofs of the two decomposition theorems, the (H_b^1, L^1) type estimates of commutators $[b, T]$ with $T \in \mathcal{K}$, and the boundedness results of commutators of Calderón-Zygmund operators. Finally, in Section 8 we present without proofs some results of commutators of fractional integrals.

Throughout the whole paper, C denotes a positive geometric constant which is independent of the main parameters, but may change from line to line. In \mathbb{R}^n , we denote by $Q = Q[x, r] := \{y = (y_1, \dots, y_n) \in \mathbb{R}^n : \sup_{1 \leq i \leq n} |y_i - x_i| \leq r\}$ a cube with center $x = (x_1, \dots, x_n)$ and radius $r > 0$. For any measurable set E , we denote by χ_E its characteristic function, by $|E|$ its Lebesgue measure, and by E^c the set $\mathbb{R}^n \setminus E$. For a cube Q and f a locally integrable function, we denote by f_Q the average of f on Q .

Acknowledgements. The author would like to thank Prof. Aline Bonami for many very valuable suggestions, discussions and advices to improve this paper. He would also like to thank Prof. Sandrine Grellier for many helpful suggestions, her carefully reading and revision of the manuscript. The author is deeply indebted to them.

2. SOME PRELIMINARIES AND NOTATIONS

As usual, $\mathcal{S}(\mathbb{R}^n)$ denote the Schwartz class of test functions on \mathbb{R}^n , $\mathcal{S}'(\mathbb{R}^n)$ the space of tempered distributions, and $C_0^\infty(\mathbb{R}^n)$ the space of C^∞ -functions with compact support.

2.1. Calderón-Zygmund operators. Let $\delta \in (0, 1]$. A continuous function $K : \mathbb{R}^n \times \mathbb{R}^n \setminus \{(x, x) : x \in \mathbb{R}^n\} \rightarrow \mathbb{C}$ is said to be a δ -Calderón-Zygmund

singular integral kernel if there exists a constant $C > 0$ such that

$$|K(x, y)| \leq \frac{C}{|x - y|^n}$$

for all $x \neq y$, and

$$|K(x, y) - K(x', y)| + |K(y, x) - K(y, x')| \leq C \frac{|x - x'|^\delta}{|x - y|^{n+\delta}}$$

for all $2|x - x'| \leq |x - y|$.

A linear operator $T : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ is said to be a δ -Calderón-Zygmund operator if T can be extended to a bounded operator on $L^2(\mathbb{R}^n)$ and there exists a δ -Calderón-Zygmund singular integral kernel K such that for all $f \in C_0^\infty(\mathbb{R}^n)$ and all $x \notin \text{supp } f$, we have

$$Tf(x) = \int_{\mathbb{R}^n} K(x, y)f(y)dy.$$

We say that T is a Calderón-Zygmund operator if it is a δ -Calderón-Zygmund operator for some $\delta \in (0, 1]$.

We say that the Calderón-Zygmund operator T satisfies the condition $T^*1 = 0$ (resp., $T1 = 0$) if $\int_{\mathbb{R}^n} Ta(x)dx = 0$ (resp., $\int_{\mathbb{R}^n} T^*a(x)dx = 0$) holds for all classical H^1 atoms a . Let b be a locally integrable function on \mathbb{R}^n . We say that the Calderón-Zygmund operator T satisfies the condition $T^*b = 0$ if $\int_{\mathbb{R}^n} b(x)Ta(x)dx = 0$ holds for all classical H^1 atoms a .

2.2. Function spaces. We first consider the subspace \mathcal{A} of $\mathcal{S}(\mathbb{R}^n)$ defined by

$$\mathcal{A} = \left\{ \phi \in \mathcal{S}(\mathbb{R}^n) : |\phi(x)| + |\nabla\phi(x)| \leq (1 + |x|^2)^{-(n+1)} \right\},$$

where $\nabla = (\partial/\partial x_1, \dots, \partial/\partial x_n)$ the gradient. We then define

$$\mathfrak{M}f(x) := \sup_{\phi \in \mathcal{A}} \sup_{|y-x|<t} |f * \phi_t(y)| \quad \text{and} \quad \mathfrak{m}f(x) := \sup_{\phi \in \mathcal{A}} \sup_{|y-x|<t<1} |f * \phi_t(y)|,$$

where $\phi_t(\cdot) = t^{-n}\phi(t^{-1}\cdot)$. The space $H^1(\mathbb{R}^n)$ denotes the space of all tempered distributions f such that $\mathfrak{M}f \in L^1(\mathbb{R}^n)$ equipped with the norm $\|f\|_{H^1} = \|\mathfrak{M}f\|_{L^1}$. The space $h^1(\mathbb{R}^n)$ denotes the space of all tempered distributions f such that $\mathfrak{m}f \in L^1(\mathbb{R}^n)$ equipped with the norm $\|f\|_{h^1} = \|\mathfrak{m}f\|_{L^1}$. The space $H^{\log}(\mathbb{R}^n)$ (see [23, 4]) denotes the space of all tempered distributions f such that $\mathfrak{M}f \in L^{\log}(\mathbb{R}^n)$ equipped with the norm $\|f\|_{H^{\log}} = \|\mathfrak{M}f\|_{L^{\log}}$. Here $L^{\log}(\mathbb{R}^n)$ is the space of all measurable functions f such that $\int_{\mathbb{R}^n} \frac{|f(x)|}{\log(e+|x|) + \log(e+|f(x)|)} dx < \infty$ with the (quasi-)norm

$$\|f\|_{L^{\log}} := \inf \left\{ \lambda > 0 : \int_{\mathbb{R}^n} \frac{\frac{|f(x)|}{\lambda}}{\log(e+|x|) + \log(e + \frac{|f(x)|}{\lambda})} dx \leq 1 \right\}.$$

Clearly, for any $f \in H^1(\mathbb{R}^n)$, we have

$$\|f\|_{h^1} \leq \|f\|_{H^1} \quad \text{and} \quad \|f\|_{H^{\log}} \leq \|f\|_{H^1}.$$

We remark that the local real Hardy space $h^1(\mathbb{R}^n)$, first introduced by Goldberg [18], is larger than $H^1(\mathbb{R}^n)$ and allows more flexibility, since global cancellation conditions are not necessary. For example, the Schwartz space is contained in $h^1(\mathbb{R}^n)$ but not in $H^1(\mathbb{R}^n)$, and multiplication by cutoff functions preserves $h^1(\mathbb{R}^n)$ but not $H^1(\mathbb{R}^n)$. Thus it makes $h^1(\mathbb{R}^n)$ more suitable for working in domains and on manifolds.

It is well-known (see [15] or [40]) that the dual of $H^1(\mathbb{R}^n)$ is $BMO(\mathbb{R}^n)$ the space of all locally integrable functions f with

$$\|f\|_{BMO} := \sup_B \frac{1}{|B|} \int_B |f(x) - f_B| dx < \infty,$$

where the supremum is taken over all balls B . We note $\mathbb{Q} := [0, 1)^n$ and, for f a function in $BMO(\mathbb{R}^n)$,

$$\|f\|_{BMO^+} := \|f\|_{BMO} + |f_{\mathbb{Q}}|.$$

We should also point out that the space $H^{\log}(\mathbb{R}^n)$ arises naturally in the study of pointwise product of functions in $H^1(\mathbb{R}^n)$ with functions in $BMO(\mathbb{R}^n)$, and in the endpoint estimates for the div-curl lemma (see for example [3, 4, 23]).

In [18] it was shown that the dual of $h^1(\mathbb{R}^n)$ can be identified with the space $bmo(\mathbb{R}^n)$, consisting of locally integrable functions f with

$$\|f\|_{bmo} := \sup_{|B| \leq 1} \frac{1}{|B|} \int_B |f(x) - f_B| dx + \sup_{|B| \geq 1} \frac{1}{|B|} \int_B |f(x)| dx < \infty,$$

where the supremums are taken over all balls B .

Clearly, for any $f \in bmo(\mathbb{R}^n)$, we have

$$\|f\|_{BMO} \leq \|f\|_{BMO^+} \leq C \|f\|_{bmo}.$$

We next recall that the space $VMO(\mathbb{R}^n)$ (resp., $vmo(\mathbb{R}^n)$) is the closure of $C_0^\infty(\mathbb{R}^n)$ in $(BMO(\mathbb{R}^n), \|\cdot\|_{BMO})$ (resp., $(bmo(\mathbb{R}^n), \|\cdot\|_{bmo})$). It is well-known that (see [9] and [11]) the dual of $VMO(\mathbb{R}^n)$ (resp., $vmo(\mathbb{R}^n)$) is the Hardy space $H^1(\mathbb{R}^n)$ (resp., $h^1(\mathbb{R}^n)$). We should note that the space $VMO(\mathbb{R}^n)$ (resp., $vmo(\mathbb{R}^n)$) considered by Coifman and Weiss (resp., Dafni [11]) is different from that considered by Sarason, and following Bourdaud [5] it coincides with the space $VMO(\mathbb{R}^n)$ (resp., $vmo(\mathbb{R}^n)$) considered above.

In the study of the pointwise multipliers for $BMO(\mathbb{R}^n)$, Nakai and Yabuta [37] introduced the space $BMO^{\log}(\mathbb{R}^n)$, consisting of locally integrable functions f with

$$\|f\|_{BMO^{\log}} := \sup_{B(a,r)} \frac{|\log r| + \log(e + |a|)}{|B(a,r)|} \int_{B(a,r)} |f(x) - f_{B(a,r)}| dx < \infty.$$

There, the authors proved that a function g is pointwise multiplier for $BMO(\mathbb{R}^n)$ if and only if g belong to $L^\infty(\mathbb{R}^n) \cap BMO^{\log}(\mathbb{R}^n)$. Furthermore, it was also shown in [23] that the space $BMO^{\log}(\mathbb{R}^n)$ is the dual of the space $H^{\log}(\mathbb{R}^n)$.

Definition 2.1. Let b be a locally integrable function and $1 < q \leq \infty$. A function a is called a (q, b) -atom related to the cube Q if

- i) $\text{supp } a \subset Q$,
- ii) $\|a\|_{L^q} \leq |Q|^{1/q-1}$,
- iii) $\int_{\mathbb{R}^n} a(x) dx = \int_{\mathbb{R}^n} a(x)b(x) dx = 0$.

The space $\mathcal{H}_b^{1,q}(\mathbb{R}^n)$ consists of the subspace of $L^1(\mathbb{R}^n)$ of functions f which can be written as $f = \sum_{j=1}^{\infty} \lambda_j a_j$, where a_j 's are (q, b) -atoms, $\lambda_j \in \mathbb{C}$, and $\sum_{j=1}^{\infty} |\lambda_j| < \infty$. As usual, we define on $\mathcal{H}_b^{1,q}(\mathbb{R}^n)$ the norm

$$\|f\|_{\mathcal{H}_b^{1,q}} := \inf \left\{ \sum_{j=1}^{\infty} |\lambda_j| : f = \sum_{j=1}^{\infty} \lambda_j a_j \right\}.$$

Observe that when $q = \infty$, then the space $\mathcal{H}_b^{1,\infty}(\mathbb{R}^n)$ is just the space $\mathcal{H}_b^1(\mathbb{R}^n)$ considered in [38]. Furthermore, $\mathcal{H}_b^{1,\infty}(\mathbb{R}^n) \subset \mathcal{H}_b^{1,q}(\mathbb{R}^n) \subset H^1(\mathbb{R}^n)$ and the inclusions are continuous.

We next introduce the space $H_b^1(\mathbb{R}^n)$ as follows.

Definition 2.2. Let b be a non-constant BMO -function. The space $H_b^1(\mathbb{R}^n)$ consists of all f in $H^1(\mathbb{R}^n)$ such that $[b, \mathfrak{M}](f)(x) = \mathfrak{M}(b(x)f(\cdot) - b(\cdot)f(\cdot))(x)$ belongs to $L^1(\mathbb{R}^n)$. We equipped $H_b^1(\mathbb{R}^n)$ with the norm $\|f\|_{H_b^1} := \|f\|_{H^1} \|b\|_{BMO} + \|[b, \mathfrak{M}](f)\|_{L^1}$.

2.3. Prerequisites on Wavelets. Let us consider a wavelet basis of \mathbb{R} with compact support. More explicitly, we are first given a $\mathcal{C}^1(\mathbb{R})$ wavelet in Dimension one, called ψ , such that $\{2^{j/2}\psi(2^j x - k)\}_{j,k \in \mathbb{Z}}$ form an $L^2(\mathbb{R})$ basis. We assume that this wavelet basis comes for a multiresolution analysis (MRA) on \mathbb{R} , as defined below (see [35]).

Definition 2.3. A multiresolution analysis (MRA) on \mathbb{R} is defined as an increasing sequence $\{V_j\}_{j \in \mathbb{Z}}$ of closed subspaces of $L^2(\mathbb{R})$ with the following four properties

- i) $\bigcap_{j \in \mathbb{Z}} V_j = \{0\}$ and $\overline{\bigcup_{j \in \mathbb{Z}} V_j} = L^2(\mathbb{R})$,
- ii) for every $f \in L^2(\mathbb{R})$ and every $j \in \mathbb{Z}$, $f(x) \in V_j$ if and only if $f(2x) \in V_{j+1}$,

iii) for every $f \in L^2(\mathbb{R})$ and every $k \in \mathbb{Z}$, $f(x) \in V_0$ if and only if $f(x-k) \in V_0$,

iv) there exists a function $\phi \in L^2(\mathbb{R})$, called the scaling function, such that the family $\{\phi_k(x) = \phi(x-k) : k \in \mathbb{Z}\}$ is an orthonormal basis for V_0 .

It is classical that, given an (MRA) on \mathbb{R} , one can find a wavelet ψ such that $\{2^{j/2}\psi(2^jx-k)\}_{k \in \mathbb{Z}}$ is an orthonormal basis of W_j , the orthogonal complement of V_j in V_{j+1} . Moreover, by Daubechies Theorem (see [12]), it is possible to find a suitable (MRA) so that ϕ and ψ are $C^1(\mathbb{R})$ and compactly supported, ψ has mean 0 and $\int x\psi(x)dx = 0$, which is known as the moment condition. We could content ourselves, in the following theorems, to have ϕ and ψ decreasing sufficiently rapidly at ∞ , but proofs are simpler with compactly supported wavelets. More precisely we can choose $m > 1$ such that ϕ and ψ are supported in the interval $1/2+m(-1/2, +1/2)$, which is obtained from $(0, 1)$ by a dilation by m centered at $1/2$.

Going back to \mathbb{R}^n , we recall that a wavelet basis of \mathbb{R}^n is found as follows. We call E the set $E = \{0, 1\}^n \setminus \{(0, \dots, 0)\}$ and, for $\sigma \in E$, put $\psi^\sigma(x) = \phi^{\sigma_1}(x_1) \cdots \phi^{\sigma_n}(x_n)$, with $\phi^{\sigma_j}(x_j) = \phi(x_j)$ for $\sigma_j = 0$ while $\phi^{\sigma_j}(x_j) = \psi(x_j)$ for $\sigma_j = 1$. Then the set $\{2^{nj/2}\psi^\sigma(2^jx-k)\}_{j \in \mathbb{Z}, k \in \mathbb{Z}^n, \sigma \in E}$ is an orthonormal basis of $L^2(\mathbb{R}^n)$. As it is classical, for I a dyadic cube of \mathbb{R}^n , which may be written as the set of x such that $2^jx-k \in (0, 1)^n$, we note

$$\psi_I^\sigma(x) = 2^{nj/2}\psi^\sigma(2^jx-k).$$

We also note $\phi_I = 2^{nj/2}\phi_{(0,1)^n}(2^jx-k)$, with $\phi_{(0,1)^n}$ the scaling function in n variables, given by $\phi_{(0,1)^n}(x) = \phi(x_1) \cdots \phi(x_n)$. In the sequel, the letter I always refers to dyadic cubes. Moreover, we note kI the cube of same center dilated by the coefficient k . Because of the assumption on the supports of ϕ and ψ , the functions ψ_I^σ and ϕ_I are supported in the cube mI .

The wavelet basis $\{\psi_I^\sigma\}$, obtained by letting I vary among dyadic cubes and σ in E , comes from an (MRA) in \mathbb{R}^n , which we still note $\{V_j\}_{j \in \mathbb{Z}}$, obtained by taking tensor products of the one-dimensional ones.

The following theorem gives the wavelet characterization of $H^1(\mathbb{R}^n)$ (cf. [35, 21]).

Theorem 2.1. *There exists a constant $C > 0$ such that $f \in H^1(\mathbb{R}^n)$ if and only if $\mathcal{W}_\psi f := \left(\sum_I \sum_{\sigma \in E} |\langle f, \psi_I^\sigma \rangle|^2 |I|^{-1} \chi_I \right)^{1/2} \in L^1(\mathbb{R}^n)$, moreover,*

$$C^{-1} \|f\|_{H^1} \leq \|\mathcal{W}_\psi f\|_{L^1} \leq C \|f\|_{H^1}.$$

A function $a \in L^2(\mathbb{R}^n)$ is called a ψ -atom related to the (not necessarily dyadic) cube R if it may be written as

$$a = \sum_{I \subset R} \sum_{\sigma \in E} a_{I,\sigma} \psi_I^\sigma$$

with $\|a\|_{L^2} \leq |R|^{-1/2}$. Remark that a is compactly supported in mR and has mean 0, so that it is a classical atom related to mR , up to the multiplicative constant $m^{n/2}$. It is standard that an atom is in $H^1(\mathbb{R}^n)$ with norm bounded by a uniform constant. The atomic decomposition gives the converse.

Theorem 2.2 (Atomic decomposition). *There exists a constant $C > 0$ such that all functions $f \in H^1(\mathbb{R}^n)$ can be written as the limit in the distribution sense and in $H^1(\mathbb{R}^n)$ of an infinite sum*

$$(2.1) \quad f = \sum_{j=1}^{\infty} \lambda_j a_j$$

with a_j ψ -atoms related to some dyadic cubes R_j and λ_j constants such that

$$C^{-1} \|f\|_{H^1} \leq \sum_{j=1}^{\infty} |\lambda_j| \leq C \|f\|_{H^1}.$$

This theorem is a small variation of a standard statement which can be found in [21], Section 6.5. Remark that the interest of dealing with finite atomic decompositions has been underlined recently, for instance in [34, 23].

Now, we denote by $H_{\text{fin}}^1(\mathbb{R}^n)$ the vector space of all finite linear combinations of ψ -atoms, that is,

$$f = \sum_{j=1}^k \lambda_j a_j,$$

where a_j 's are ψ -atoms. Then, the norm of f in $H_{\text{fin}}^1(\mathbb{R}^n)$ is defined by

$$\|f\|_{H_{\text{fin}}^1} = \inf \left\{ \sum_{j=1}^k : f = \sum_{j=1}^k \lambda_j a_j \right\}.$$

By the atomic decomposition theorem, the set $H_{\text{fin}}^1(\mathbb{R}^n)$ is dense in $H^1(\mathbb{R}^n)$ for the norm $\|\cdot\|_{H^1}$.

The following two wavelet characterizations of $L^p(\mathbb{R}^n)$, $1 < p < \infty$, and $BMO(\mathbb{R}^n)$ are well-known (see [35]).

Theorem 2.3. *Let $1 < p < \infty$. Then the norms $\|f\|_{L^p}$, $\|(\sum_I \sum_{\sigma \in E} |\langle f, \psi_I^\sigma \rangle|^2 |I|^{-1} \chi_I)^{1/2}\|_{L^p}$ and $\|(\sum_I \sum_{\sigma \in E} |\langle f, \psi_I^\sigma \rangle|^2 (\psi_I^\sigma)^2)^{1/2}\|_{L^p}$ are equivalent on $L^p(\mathbb{R}^n)$.*

Theorem 2.4. *A function $g \in BMO(\mathbb{R}^n)$ if and only if $\frac{1}{|R|} \sum_{I \subset R} \sum_{\sigma \in E} |\langle g, \psi_I^\sigma \rangle|^2 < \infty$ for all (not necessarily dyadic) cubes R . Moreover, there exists a constant $C > 0$ such that for all $g \in BMO(\mathbb{R}^n)$,*

$$C^{-1} \|g\|_{BMO} \leq \sup_R \left(\frac{1}{|R|} \sum_{I \subset R} \sum_{\sigma \in E} |\langle g, \psi_I^\sigma \rangle|^2 \right)^{1/2} \leq C \|g\|_{BMO},$$

where the supremum is taken over all cubes R .

By Theorem 2.3, Theorem 2.4 and John-Nirenberg inequality, we obtain the following lemma. The proof is easy and will be omitted.

Lemma 2.1. *Let f be a ψ -atom related to the cube R and $b \in BMO(\mathbb{R}^n)$. Then, $\sum_{I \subset R} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \langle b, \psi_I^\sigma \rangle (\psi_I^\sigma)^2 \in L^q(\mathbb{R}^n)$ for any $q \in (1, 2)$.*

3. BILINEAR, SUBBILINEAR DECOMPOSITIONS AND COMMUTATORS

Recall that \mathcal{K} is the set of all sublinear operators T bounded from $H^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$ satisfying

$$\|(b - b_Q)Ta\|_{L^1} \leq C\|b\|_{BMO},$$

for all $b \in BMO(\mathbb{R}^n)$, a a H^1 -atom supported in the cube Q , where $C > 0$ a constant independent of b, a . This class \mathcal{K} contains almost all important operators in Harmonic analysis: Calderón-Zygmund type operators, strongly singular integral operators, multiplier operators, pseudo-differential operators, maximal type operators, the area integral operator of Lusin, Littlewood-Paley type operators, Marcinkiewicz operators, maximal Bochner-Riesz operators, etc (see Section 4).

Here and what in follows the bilinear operator \mathfrak{S} is defined by

$$\mathfrak{S}(f, g) = - \sum_I \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \langle g, \psi_I^\sigma \rangle (\psi_I^\sigma)^2.$$

In [4], the authors show that \mathfrak{S} is a bounded bilinear operator from $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$.

3.1. Two decomposition theorems and (H_b^1, L^1) type estimates. Let b be a locally integrable function and $T \in \mathcal{K}$. As usual, the (sublinear) commutator $[b, T]$ of the operator T is defined by $[b, T](f)(x) := T\left((b(x) - b(\cdot))f(\cdot)\right)(x)$.

Theorem 3.1 (Subbilinear decomposition). *Let $T \in \mathcal{K}$. There exists a bounded subbilinear operator $\mathfrak{R} : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ such that for all $(f, b) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$, we have*

$$|T(\mathfrak{S}(f, b))| - \mathfrak{R}(f, b) \leq |[b, T](f)| \leq \mathfrak{R}(f, b) + |T(\mathfrak{S}(f, b))|.$$

Corollary 3.1. *Let $T \in \mathcal{K}$ such that T is of weak type $(1, 1)$. Then, the bilinear operator $\mathfrak{P}(f, g) = [g, T](f)$ maps continuously $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into weak- $L^1(\mathbb{R}^n)$. In particular, the commutator $[b, T]$ is of weak type (H^1, L^1) if $b \in BMO(\mathbb{R}^n)$.*

We remark that the class of operators $T \in \mathcal{K}$ of weak type $(1, 1)$ contains Calderón-Zygmund operators, strongly singular integral operators, multiplier operators, pseudo-differential operators whose symbols in the Hörmander class $S_{\varrho, \delta}^m$ with $0 < \varrho \leq 1, 0 \leq \delta < 1, m \leq -n((1 - \varrho)/2 + \max\{0, (\delta - \varrho)/2\})$, maximal type operators, the area integral operator of Lusin, Littlewood-Paley

type operators, Marcinkiewicz operators, maximal Bochner-Riesz operators T_*^δ with $\delta > (n-1)/2$, etc.

When T is linear and belongs to \mathcal{K} . We obtain the bilinear decomposition for the linear commutator $[b, T]$ of f , $[b, T](f) = bT(f) - T(bf)$, instead of the subbilinear decomposition as stated in Theorem 3.1.

Theorem 3.2 (Bilinear decomposition). *Let T be a linear operator in \mathcal{K} . Then, there exists a bounded bilinear operator $\mathfrak{R} : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ such that for all $(f, b) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$, we have*

$$[b, T](f) = \mathfrak{R}(f, b) + T(\mathfrak{S}(f, b)).$$

The following give (H_b^1, L^1) type estimates for commutators $[b, T]$ with T belongs to the class \mathcal{K} .

Theorem 3.3. *Let b be a non-constant BMO-function and $T \in \mathcal{K}$. Then, the commutator $[b, T]$ maps continuously $H_b^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$.*

Remark that in the particular case of T a 1-Calderón-Zygmund operator and $H_b^1(\mathbb{R}^n)$ is replaced by $\mathcal{H}_b^1(\mathbb{R}^n)$, the author in [38] proved

$$(3.1) \quad \sup\{\|[b, T](a)\|_{L^1} : a \text{ is a } (\infty, b)\text{-atom}\} < \infty.$$

Then he conclude that the (linear) commutator $[b, T]$ maps continuously $\mathcal{H}_b^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$. Notice that $\mathcal{H}_b^1(\mathbb{R}^n) \subset \mathcal{H}_b^{1,q}(\mathbb{R}^n) \subset H_b^1(\mathbb{R}^n)$, $1 < q \leq \infty$, and the inclusions are continuous (see Section 5). However, as mentioned in the introduction, Inequality (3.1) does not suffice to conclude that the (linear) commutator $[b, T]$ is bounded from $\mathcal{H}_b^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$. We should also point out that the (H^1, L^1) weak type estimates and the (\mathcal{H}_b^1, L^1) type estimates for the (linear) commutators of multiplier operators (see [43, 25, 41]), strongly singular Calderón-Zygmund operators (see [26]) and for the (sublinear) commutators of Littlewood-Paley type operators (see [28]), Marcinkiewicz operators (see [33]), maximal Bochner-Riesz operators (see [30, 31, 29]) have been studied recently. However, they just prove Inequality (1.2) (that is Inequality (3.1)) and use Equality (1.1) which leaves a gap as pointed out in the introduction.

3.2. Boundedness of linear commutators on Hardy spaces. In this subsection, we consider \mathcal{K}_0 the set of all Calderón-Zygmund operators T satisfying the condition $T1 = T^*1 = 0$.

Analogously as Hardy estimates for bilinear operators of Coifman and Grafakos [8] (see also [14]), we also obtain the following bilinear estimates which improve Corollary 3.1.

Theorem 3.4. *Let T be a linear operator in \mathcal{K} . Assume that A_i, B_i , $i = 1, \dots, K$, be Calderón-Zygmund operators satisfying $A_i1 = A_i^*1 = B_i1 = B_i^*1 =$*

0, and for every f and g in $L^2(\mathbb{R}^n)$,

$$\int_{\mathbb{R}^n} \left(\sum_{i=1}^K A_i f \cdot B_i g \right) dx = 0.$$

Then, the bilinear operator \mathfrak{T} , defined by

$$\mathfrak{T}(f, g) = \sum_{i=1}^K [B_i g, T](A_i f),$$

maps continuously $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$.

Two next theorems give the boundedness of commutators of Calderón-Zygmund operators on Hardy spaces.

Theorem 3.5. *Let b be a non-constant BMO^{\log} -function and $T \in \mathcal{K}_0$. Then, the linear commutator $[b, T]$ maps continuously $H_b^1(\mathbb{R}^n)$ into $h^1(\mathbb{R}^n)$.*

Theorem 3.6. *Let b be a non-constant BMO^{\log} -function and $T \in \mathcal{K}_0$. Assume that the operator T satisfies $T^*(b) = 0$. Then, the linear commutator $[b, T]$ maps continuously $H_b^1(\mathbb{R}^n)$ into $H^1(\mathbb{R}^n)$.*

Observe that the condition $T^*(b) = 0$ is also "necessary" in the sense if the linear commutator $[b, T]$ maps continuously $H_b^1(\mathbb{R}^n)$ into $H^1(\mathbb{R}^n)$, then $\int_{\mathbb{R}^n} b(x)Ta(x)dx = 0$ hold for all (q, b) -atoms a , $1 < q \leq \infty$.

4. THE CLASS \mathcal{K} AND FOUR BILINEAR OPERATORS ON $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$

4.1. The class \mathcal{K} . The purpose of this subsection is to give some examples of operators in the class \mathcal{K} . More precisely, the class \mathcal{K} contains almost all important operators in Harmonic analysis: Calderón-Zygmund type operators, strongly singular integral operators, multiplier operators, pseudo-differential operators with symbols in the Hörmander class $S_{\varrho, \delta}^m$ with $0 < \varrho \leq 1, 0 \leq \delta < 1, m \leq -n((1 - \varrho)/2 + \max\{0, (\delta - \varrho)/2\})$ (see [2, 1]), maximal type operators, the area integral operator of Lusin, Littlewood-Paley type operators, Marcinkiewicz operators, maximal Bochner-Riesz operators T_*^δ with $\delta > (n - 1)/2$ (cf. [24]), etc. It is well-known that these operators T are bounded from $H^1(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$. So, in order to establish that these ones are in the class \mathcal{K} , we just need to show that

$$(4.1) \quad \|(b - b_Q)Ta\|_{L^1} \leq C\|b\|_{BMO}$$

for all BMO -function b , H^1 -atom a related to a cube $Q = Q[x_0, r]$ with constant $C > 0$ independent of b, a .

Observe that the nontangential grand maximal operator \mathfrak{M} belongs to \mathcal{K} since it satisfies the inequality (4.1) (cf. [40]). See also [20] for the (sublinear) commutators $[b, M_{\varphi, \alpha}]$ of the maximal operators $M_{\varphi, \alpha}$ with a note that $M_{\varphi, 0}$ lies in \mathcal{K} .

Here we just give the proofs for Calderón-Zygmund operators (linear operators) and the area integral operator of Lusin (sublinear operator). For the other operators, we leave the proofs to the interested reader.

One first recall that $P(x) = \frac{1}{(1+|x|^2)^{(n+1)/2}}$ is the Poisson kernel and $u_f(x, t) := f * P_t(x)$ the Poisson integral of f . Then the area integral operator S of Lusin is defined by

$$S(f)(x) = \left(\int_{\Gamma(x)} |\nabla u_f(y, t)|^2 t^{1-n} dy dt \right)^{1/2},$$

where $\Gamma(x)$ is the cone $\{(y, t) \in \mathbb{R}_+^{n+1} : |y - x| < t\}$ with vertex at x , while $\nabla u_f = (\partial u_f / \partial x_1, \dots, \partial u_f / \partial x_n, \partial u_f / \partial t)$ the gradient of u_f on $\mathbb{R}_+^{n+1} = \mathbb{R}^n \times (0, \infty)$.

Proposition 4.1. *Let $\delta \in (0, 1]$ and T be a δ -Calderón-Zygmund operator. Then T satisfies the inequality (4.1), and thus $T \in \mathcal{K}$.*

Proof. We cut the integral of $|(b - b_Q)Ta|$ into two parts. By Schwarz inequality and the boundedness of T on $L^2(\mathbb{R}^n)$, we have

$$\begin{aligned} \int_{2Q} |b(x) - b_Q| |Ta(x)| dx &\leq C \left(\int_{2Q} |b(x) - b_Q|^2 dx \right)^{1/2} \|a\|_{L^2} \\ &\leq C \|b\|_{BMO} \end{aligned}$$

here one used the fact $|b_{2Q} - b_Q| \leq C \|b\|_{BMO}$. Next, for $x \notin 2Q$,

$$\begin{aligned} |Ta(x)| &= \left| \int_Q (K(x, y) - K(x, x_0)) a(y) dy \right| \\ &\leq C \int_Q \frac{|y - x_0|^\delta}{|x - x_0|^{n+\delta}} |a(y)| dy \\ &\leq C \frac{r^\delta}{|x - x_0|^{n+\delta}}. \end{aligned}$$

Therefore,

$$\int_{(2Q)^c} |b(x) - b_Q| |Ta(x)| dx \leq C \int_{Q^c} |b(x) - b_Q| \frac{r^\delta}{|x - x_0|^{n+\delta}} dx \leq C \|b\|_{BMO},$$

since the last inequality is classical (cf. [40]). This finishes the proof. \square

Corollary 4.1. *Let $\mathcal{R}_j, j = 1, \dots, n$, are classical Riesz transforms. Then, $\mathcal{R}_j \in \mathcal{K}$ for all $j = 1, \dots, n$.*

Proposition 4.2. *The area integral operator S satisfies the inequality (4.1), and thus $S \in \mathcal{K}$.*

Proof. We also cut the integral of $|(b - b_Q)S(a)|$ into two parts. By Schwarz inequality and the boundedness of S on $L^2(\mathbb{R}^n)$, we have

$$\begin{aligned} \int_{2Q} |b(x) - b_Q| |S(a)(x)| dx &\leq C \left(\int_{2Q} |b(x) - b_Q|^2 dx \right)^{1/2} \|a\|_{L^2} \\ &\leq C \|b\|_{BMO}. \end{aligned}$$

Next, for $x \notin 2Q$, by using the equality

$$u_a(y, t) = \int_{\mathbb{R}^n} \frac{1}{t^n} \left(P\left(\frac{y-z}{t}\right) - P\left(\frac{y-x_0}{t}\right) \right) a(z) dz,$$

since $\int_{\mathbb{R}^n} a(z) dz = 0$, it is easy to establish that

$$S(a)(x) = \left(\int_{\Gamma(x)} |\nabla u_a(y, t)|^2 t^{1-n} dy dt \right)^{1/2} \leq C \frac{r}{|x - x_0|^{n+1}}.$$

Therefore,

$$\int_{(2Q)^c} |b(x) - b_Q| |S(a)(x)| dx \leq C \int_{Q^c} |b(x) - b_Q| \frac{r}{|x - x_0|^{n+1}} dx \leq C \|b\|_{BMO},$$

which ends the proof. \square

We should point out that the Littlewood-Paley type operators also can be viewed as vector-valued Calderón-Zygmund operators (see [39]). See also [20] in the context of vector-valued commutators.

4.2. Four bilinear operators on $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$. Let us consider four bilinear operators on $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ which are fundamental for our bilinear decomposition theorem.

Now, we state some lemmas whose proofs can be found in [4].

Lemma 4.1. *The bilinear operator Π_3 , defined on $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ by*

$$\Pi_3(f, g) = \sum_I \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \langle g, \psi_I^\sigma \rangle (\psi_I^\sigma)^2,$$

is a bounded bilinear operator from $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$.

Observe that $\mathfrak{S}(f, g) = -\Pi_3(f, g)$ for all $(f, g) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$.

Lemma 4.2. *The bilinear operator Π_4 , defined on $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ by*

$$\Pi_4(f, g) = \sum_{I, I'} \sum_{\sigma, \sigma' \in E} \langle f, \psi_I^\sigma \rangle \langle g, \psi_{I'}^{\sigma'} \rangle \psi_I^\sigma \psi_{I'}^{\sigma'},$$

the sums being taken over all dyadic cubes I, I' and $\sigma, \sigma' \in E$ such that $(I, \sigma) \neq (I', \sigma')$, is a bounded bilinear operator from $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $H^1(\mathbb{R}^n)$.

Lemma 4.3. *The bilinear operator Π_1 defined by*

$$\Pi_1(a, g) = \sum_{|I|=|I'|} \sum_{\sigma \in E} \langle a, \phi_I \rangle \langle g, \psi_{I'}^\sigma \rangle \phi_I \psi_{I'}^\sigma,$$

for a ψ -atom and $g \in BMO(\mathbb{R}^n)$, can be extended into a bounded bilinear operator from $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $H^1(\mathbb{R}^n)$.

Lemma 4.4. *The bilinear operator Π_2 defined by*

$$\Pi_2(a, g) = \sum_{|I|=|I'|} \sum_{\sigma \in E} \langle a, \psi_I^\sigma \rangle \langle g, \phi_{I'} \rangle \psi_I^\sigma \phi_{I'},$$

for a ψ -atom related to the cube R and $g \in BMO(\mathbb{R}^n)$, can be extended into a bounded bilinear operator from $H^1(\mathbb{R}^n) \times BMO^+(\mathbb{R}^n)$ into $H^{\log}(\mathbb{R}^n)$. Furthermore, we can write

$$(4.2) \quad \Pi_2(a, g) = h^{(1)} + \kappa g_R h^{(2)}$$

where $\|h^{(1)}\|_{H^1} \leq C\|g\|_{BMO}$, $h^{(2)}$ is an atom related to mR , and κ a uniform constant, independent of a and g .

The following remarks are useful in our proofs in Section 6 and Section 7.

Remark 4.1. *i) If $g \in BMO(\mathbb{R}^n)$ and $f \in H^1(\mathbb{R}^n)$ such that $fg \in L^1(\mathbb{R}^n)$, then*

$$\int_{\mathbb{R}^n} fg dx = \int_{\mathbb{R}^n} \mathfrak{S}(f, g) dx = \sum_I \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \langle g, \psi_I^\sigma \rangle.$$

ii) For any $(f, g) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ and c a constant, we have

$$\Pi_i(f, g) = \Pi_i(f, g + c), \quad i = 1, 3, 4.$$

iii) As a consequence of Lemma 4.4, if $g_R = 0$ then Equality (4.2) gives that $\Pi_2(a, g) \in H^1(\mathbb{R}^n)$. Moreover, $\|\Pi_2(a, g)\|_{H^1} \leq C\|g\|_{BMO}$.

In [4], the authors have shown the following decomposition theorem for the product space $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$.

Theorem 4.1 (Decomposition theorem). *Let $f \in H^1(\mathbb{R}^n)$ and $g \in BMO(\mathbb{R}^n)$. Then, we have the following decomposition*

$$fg = \Pi_1(f, g) + \Pi_2(f, g) + \Pi_3(f, g) + \Pi_4(f, g),$$

that is

$$fg = \Pi_1(f, g) + \Pi_2(f, g) + \Pi_4(f, g) - \mathfrak{S}(f, g).$$

5. THE SPACE $H_b^1(\mathbb{R}^n)$

Let b be a non-constant BMO -function. In this section, we study the space $H_b^1(\mathbb{R}^n)$. In particular, we give some characterizations of the space $H_b^1(\mathbb{R}^n)$ (see Theorem 5.1), and the comparison with the space $\mathcal{H}_b^1(\mathbb{R}^n)$ of Pérez (see Theorem 5.2).

First, let us consider the class $\tilde{\mathcal{K}}$ of all $T \in \mathcal{K}$ such that T can characterize the space $H^1(\mathbb{R}^n)$, that means $f \in H^1(\mathbb{R}^n)$ if and only if $Tf \in L^1(\mathbb{R}^n)$. Clearly, the class $\tilde{\mathcal{K}}$ contain the maximal operator \mathfrak{M} , the area integral operator S of Lusin, the Littlewood-Paley g -operator (see [15]), the Littlewood-Paley g_λ^* -operator with $\lambda > 3n$ (see [19]), etc. We now can give the characterization of $H_b^1(\mathbb{R}^n)$ as follows.

Theorem 5.1. *Let b be a non-constant BMO -function and $T \in \tilde{\mathcal{K}}$. For $f \in H^1(\mathbb{R}^n)$, then the following conditions are equivalent:*

- i) $f \in H_b^1(\mathbb{R}^n)$.*
- ii) $\mathfrak{S}(f, b) \in H^1(\mathbb{R}^n)$.*
- iii) $[b, \mathcal{R}_j](f) \in L^1(\mathbb{R}^n)$ for all $j = 1, \dots, n$.*
- iv) $[b, T](f) \in L^1(\mathbb{R}^n)$.*

Remark 5.1. *Theorem 3.3 and Theorem 5.1 give that $[b, T]$ is bounded from $H_b^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$ for every T a Calderón-Zygmund singular integral operator. Furthermore, $H_b^1(\mathbb{R}^n)$ is the largest space having this property.*

Here and what in follows, the symbol $f \approx g$ mean that $C^{-1}f \leq g \leq Cf$.

Proof of Theorem 5.1. (i) \Leftrightarrow (ii) By Theorem 3.1, there exists a bounded subbilinear operator $\mathfrak{R} : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ such that

$$\mathfrak{M}(\mathfrak{S}(f, b)) - \mathfrak{R}(f, b) \leq |[b, \mathfrak{M}](f)| \leq \mathfrak{R}(f, b) + \mathfrak{M}(\mathfrak{S}(f, b)).$$

Consequently, $\mathfrak{S}(f, b) \in H^1(\mathbb{R}^n)$ if and only if $[b, \mathfrak{M}](f) \in L^1(\mathbb{R}^n)$. Moreover,

$$\|f\|_{H_b^1} \approx \|f\|_{H^1} \|b\|_{BMO} + \|\mathfrak{S}(f, b)\|_{H^1}.$$

(ii) \Leftrightarrow (iii). By Theorem 3.2, there exist n bounded bilinear operators $\mathfrak{R}_j : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$, $j = 1, \dots, n$, such that

$$[b, \mathcal{R}_j](f) = \mathfrak{R}_j(f, b) + \mathcal{R}_j(\mathfrak{S}(f, b)).$$

Consequently, $\mathfrak{S}(f, b) \in H^1(\mathbb{R}^n)$ if and only if $[b, \mathcal{R}_j](f) \in L^1(\mathbb{R}^n)$ for all $j = 1, \dots, n$. Moreover,

$$\|f\|_{H^1} \|b\|_{BMO} + \|\mathfrak{S}(f, b)\|_{H^1} \approx \|f\|_{H^1} \|b\|_{BMO} + \sum_{j=1}^n \|[b, \mathcal{R}_j](f)\|_{L^1}.$$

(iii) \Leftrightarrow (iv). By Theorem 3.1, there exists a bounded subbilinear operator $\mathfrak{R} : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ such that

$$|T(\mathfrak{S}(f, b))| - \mathfrak{R}(f, b) \leq |[b, T](f)| \leq \mathfrak{R}(f, b) + |T(\mathfrak{S}(f, b))|.$$

Consequently, $\mathfrak{S}(f, b) \in H^1(\mathbb{R}^n)$ if and only if $[b, T](f) \in L^1(\mathbb{R}^n)$ since $T \in \widetilde{\mathcal{K}}$. Moreover,

$$\|f\|_{H^1} \|b\|_{BMO} + \|\mathfrak{S}(f, b)\|_{H^1} \approx \|f\|_{H^1} \|b\|_{BMO} + \|[b, T](f)\|_{L^1}.$$

□

Observe that the constant $C > 0$ in the last equivalence depends on T , though it is independent of f, b .

The following lemma is an immediate corollary of the weak convergence theorem in $H^1(\mathbb{R}^n)$ of Jones and Journé. See also [11] in the setting of $h^1(\mathbb{R}^n)$.

Lemma 5.1. *Let $\{f_k\}_{k \geq 1}$ be a bounded sequence in $H^1(\mathbb{R}^n)$ (resp., in $h^1(\mathbb{R}^n)$) such that f_k tends to f in $L^1(\mathbb{R}^n)$. Then f in $H^1(\mathbb{R}^n)$ (resp., in $h^1(\mathbb{R}^n)$), and*

$$\|f\|_{H^1} \leq \varliminf_{k \rightarrow \infty} \|f_k\|_{H^1} \quad (\text{resp., } \|f\|_{h^1} \leq \varliminf_{k \rightarrow \infty} \|f_k\|_{h^1}).$$

Theorem 5.2. *Let b be a non-constant BMO-function and $1 < q \leq \infty$. Then, $\mathcal{H}_b^{1,q}(\mathbb{R}^n) \subset H_b^1(\mathbb{R}^n)$ and the inclusion is continuous.*

Proof. Let a be a (q, b) -atom related to the cube Q . Then, $\text{supp } (b - b_Q)a \subset \text{supp } a \subset Q$ and $\int_{\mathbb{R}^n} (b(x) - b_Q)a(x)dx = \int_{\mathbb{R}^n} b(x)a(x)dx - b_Q \int_{\mathbb{R}^n} a(x)dx = 0$. Moreover, by Hölder inequality and John-Nirenberg inequality, we get

$$\|(b - b_Q)a\|_{L^{(\tilde{q}+1)/2}} \leq \|(b - b_Q)\chi_Q\|_{L^{\tilde{q}(\tilde{q}+1)/(\tilde{q}-1)}} \|a\|_{L^{\tilde{q}}} \leq C \|b\|_{BMO} |Q|^{(-\tilde{q}+1)/(\tilde{q}+1)},$$

where $\tilde{q} = q$ if $1 < q < \infty$, $\tilde{q} = 2$ if $q = \infty$, and $C > 0$ is independent of b, a . This proves that $(b - b_Q)a$ is $C \|b\|_{BMO}$ times a classical $(\tilde{q} + 1)/2$ -atom, and thus $\|(b - b_Q)a\|_{H^1} \leq C \|b\|_{BMO}$.

By Theorem 3.2, there exist n bounded bilinear operators $\mathfrak{R}_j : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$, $j = 1, \dots, n$, such that

$$[b, \mathcal{R}_j](a) = \mathfrak{R}_j(a, b) + \mathcal{R}_j(\mathfrak{S}(a, b)),$$

since \mathcal{R}_j is linear and belongs to \mathcal{K} (see Corollary 4.1). Consequently, for all $j = 1, \dots, n$, as $\mathcal{R}_j \in \mathcal{K}$,

$$\begin{aligned} \|\mathcal{R}_j(\mathfrak{S}(a, b))\|_{L^1} &= \|(b - b_Q)\mathcal{R}_j(a) - \mathcal{R}_j((b - b_Q)a) - \mathfrak{R}_j(a, b)\|_{L^1} \\ &\leq \|(b - b_Q)\mathcal{R}_j(a)\|_{L^1} + \|\mathcal{R}_j\|_{H^1 \rightarrow L^1} \|((b - b_Q)a)\|_{H^1} + \|\mathfrak{R}_j(a, b)\|_{L^1} \\ &\leq C \|b\|_{BMO}. \end{aligned}$$

This proves that $\mathfrak{S}(a, b) \in H^1(\mathbb{R}^n)$ since $\|\mathfrak{S}(a, b)\|_{L^1} \leq C \|b\|_{BMO}$, and moreover that

$$(5.1) \quad \|\mathfrak{S}(a, b)\|_{H^1} \leq C \|b\|_{BMO}.$$

Now, for any $f \in \mathcal{H}_b^{1,q}(\mathbb{R}^n)$, there exists an expansion $f = \sum_{j=1}^{\infty} \lambda_j a_j$ where the a_j 's are (q, b) -atoms and $\sum_{j=1}^{\infty} |\lambda_j| \leq 2 \|f\|_{\mathcal{H}_b^{1,q}}$. Then the sequence $\{\sum_{j=1}^k \lambda_j a_j\}_{k \geq 1}$ converges to f in $\mathcal{H}_b^{1,q}(\mathbb{R}^n)$ and thus in $H^1(\mathbb{R}^n)$. Hence, Lemma

4.1 implies that the sequence $\left\{ \mathfrak{S}\left(\sum_{j=1}^k \lambda_j a_j, b\right) \right\}_{k \geq 1}$ converges to $\mathfrak{S}(f, b)$ in $L^1(\mathbb{R}^n)$. In addition, by (5.1),

$$\left\| \mathfrak{S}\left(\sum_{j=1}^k \lambda_j a_j, b\right) \right\|_{H^1} \leq \sum_{j=1}^k |\lambda_j| \|\mathfrak{S}(a_j, b)\|_{H^1} \leq C \|f\|_{\mathcal{H}_b^{1,q}} \|b\|_{BMO}.$$

We then use Lemma 5.1 to conclude that $\mathfrak{S}(f, b) \in H^1(\mathbb{R}^n)$, and thus $f \in H_b^1(\mathbb{R}^n)$ (see Theorem 5.1). Moreover,

$$\begin{aligned} \|f\|_{H_b^1} &\leq C(\|f\|_{H^1} \|b\|_{BMO} + \|\mathfrak{S}(f, b)\|_{H^1}) \\ &\leq C\left(\|f\|_{\mathcal{H}_b^{1,q}} \|b\|_{BMO} + \lim_{k \rightarrow \infty} \left\| \mathfrak{S}\left(\sum_{j=1}^k \lambda_j a_j, b\right) \right\|_{H^1}\right) \\ &\leq C\|f\|_{\mathcal{H}_b^{1,q}} \|b\|_{BMO}, \end{aligned}$$

which ends the proof. \square

From Theorem 3.3 and Theorem 5.1, we get the following corollary.

Corollary 5.1. *Let b be a BMO-function, $T \in \mathcal{K}$ and $1 < q \leq \infty$. Then the linear commutator $[b, T]$ maps continuously $\mathcal{H}_b^{1,q}(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$.*

6. PROOF OF THEOREM 3.1, THEOREM 3.2, THEOREM 3.3

In order to prove the decomposition theorems (Theorem 3.2 and Theorem 3.1), we need the following two lemmas.

Lemma 6.1. *Let $T \in \mathcal{K}$ and f be a classical atom related to the cube mQ . Then, there exists a positive constant $C = C(m)$ such that*

$$\|(g - g_Q)Tf\|_{L^1} \leq C\|g\|_{BMO}, \text{ for all } g \in BMO(\mathbb{R}^n).$$

Proof. Since $T \in \mathcal{K}$ and since $|g_Q - g_{mQ}| \leq C(m)\|g\|_{BMO}$, we have

$$\|(g - g_Q)Tf\|_{L^1} \leq C(m)\|g\|_{BMO}\|Tf\|_{L^1} + \|(g - g_{mQ})Tf\|_{L^1} \leq C\|g\|_{BMO}.$$

\square

Lemma 6.2. *The norms $\|\cdot\|_{H^1}$ and $\|\cdot\|_{H_{\text{fin}}^1}$ are equivalent on $H_{\text{fin}}^1(\mathbb{R}^n)$.*

We should point out that in the proof below we use some results in [21], Theorem 5.12. Even though the proofs in [21] are worked out in the one-dimension case, they can be easily carried out in higher dimension as well.

The proof of Lemma 6.2. Obviously, $H_{\text{fin}}^1(\mathbb{R}^n) \subset H^1(\mathbb{R}^n)$ and for all $f \in H_{\text{fin}}^1(\mathbb{R}^n)$, we have $\|f\|_{H^1} \leq C\|f\|_{H_{\text{fin}}^1}$. We now have to show that there exists a constant $C > 0$ such that for all $f \in H_{\text{fin}}^1(\mathbb{R}^n)$,

$$\|f\|_{H_{\text{fin}}^1} \leq C\|f\|_{H^1}.$$

By homogeneity, we can assume that $f = \sum_{j=1}^{N_0} \lambda_j a_j$, a_j 's are ψ -atoms related to the cubes R_j 's, with $\|f\|_{H^1} = 1$. Since $f \in L^2(\mathbb{R}^n) \cap H^1(\mathbb{R}^n)$, there exists a ψ -atomic decomposition (see [21], Theorem 5.12)

$$f = \sum_I \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma = \sum_{k \in \mathbb{Z}} \sum_{i \in \Lambda_k} \left(\sum_{I \subset \tilde{I}_k^i, I \in \mathcal{B}_k} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma \right)$$

where $\sum_{I \subset \tilde{I}_k^i, I \in \mathcal{B}_k} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma = \lambda(k, i) a_{k,i}$ with $a_{k,i}$ ψ -atoms related to the cubes $m\tilde{I}_k^i$ and

$$(6.1) \quad \sum_{k \in \mathbb{Z}} \sum_{i \in \Lambda_k} |\lambda(k, i)| \leq C \|f\|_{H^1} = C.$$

We note that $\text{supp } a_{k,i} \subset \bigcup_{j=1}^{N_0} mR_j$ for all $k \in \mathbb{Z}, i \in \Lambda_k$. Recall that

$$\begin{aligned} \mathcal{W}_\psi f &= \left(\sum_I \sum_{\sigma \in E} |\langle f, \psi_I^\sigma \rangle|^2 |I|^{-1} \chi_I \right)^{1/2} \\ &= \left(\sum_{j=1}^{N_0} \sum_{I \subset R_j} \sum_{\sigma \in E} |\langle f, \psi_I^\sigma \rangle|^2 |I|^{-1} \chi_I \right)^{1/2} \end{aligned}$$

and $\Omega_k = \{x \in \mathbb{R}^n : \mathcal{W}_\psi f(x) > 2^k\}$ for any $k \in \mathbb{Z}$. Clearly, $\text{supp } \mathcal{W}_\psi f \subset \bigcup_{j=1}^{N_0} mR_j$. So, there exists a cube Q such that $\Omega_k \subset \text{supp } \mathcal{W}_\psi f \subset \bigcup_{j=1}^{N_0} mR_j \subset Q$ for all $k \in \mathbb{Z}$. We now denote by k' the largest integer k such that $2^k \leq |Q|^{-1}$. Then, we define the functions g and ℓ by

$$g = \sum_{k \leq k'} \sum_{i \in \Lambda_k} \left(\sum_{I \subset \tilde{I}_k^i, I \in \mathcal{B}_k} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma \right) \text{ and } \ell = \sum_{k > k'} \sum_{i \in \Lambda_k} \left(\sum_{I \subset \tilde{I}_k^i, I \in \mathcal{B}_k} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma \right).$$

Obviously, $f = g + \ell$, moreover, $\text{supp } g \subset Q$ and $\text{supp } \ell \subset Q$. On the other hand, it follows from [21] (Theorem 5.12) that $\sum_{I \subset \tilde{I}_k^i, I \in \mathcal{B}_k} \sum_{\sigma \in E} |\langle f, \psi_I^\sigma \rangle|^2 \leq C 2^{2k} |\tilde{I}_k^i \cap \Omega_k|$. Hence, as the dyadic cubes \tilde{I}_k^i are disjoint (see also [21]), we get

$$\begin{aligned} \|g\|_{L^2}^2 &\leq C \sum_{k \leq k'} \sum_{i \in \Lambda_k} \sum_{I \subset \tilde{I}_k^i, I \in \mathcal{B}_k} \sum_{\sigma \in E} |\langle f, \psi_I^\sigma \rangle|^2 \\ &\leq C \sum_{k \leq k'} \sum_{i \in \Lambda_k} 2^{2k} |\tilde{I}_k^i \cap \Omega_k| \leq C \sum_{k \leq k'} 2^{2k} |\Omega_k| \\ &\leq C 2^{2k'} |Q| \leq C |Q|^{-1}. \end{aligned}$$

This prove that $C^{-1/2}g$ is a ψ -atom related to the cube Q .

Now, for any positive integer K , set $F_K = \{(k, i) : k > k', |k| + |i| \leq K\}$ and $\ell_K = \sum_{(k,i) \in F_K} \left(\sum_{I \subset \tilde{I}_k^i, I \in \mathcal{B}_k} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma \right)$. Observe that since $f \in L^2(\mathbb{R}^n)$,

the series $\sum_{k>k'} \sum_{i \in \Lambda_k} \left(\sum_{I \subset \tilde{I}_k^i, I \in \mathcal{B}_k} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma \right)$ converges to ℓ in $L^2(\mathbb{R}^n)$. So, for any $\varepsilon > 0$, if K is large enough, $\varepsilon^{-1}(\ell - \ell_K)$ is a ψ -atom related to the cube Q . Therefore, $f = g + \ell_K + (\ell - \ell_K)$ is a finite linear combination of atoms for f , and thus

$$\begin{aligned} \|f\|_{H_{\text{fin}}^1} &\leq C(\|g\|_{H_{\text{fin}}^1} + \|\ell_K\|_{H_{\text{fin}}^1} + \|\ell - \ell_K\|_{H_{\text{fin}}^1}) \\ &\leq C\left(C + \sum_{k \in \mathbb{Z}} \sum_{i \in \Lambda_k} |\lambda(k, i)| + \varepsilon\right) \leq C \end{aligned}$$

since (6.1), which ends the proof. \square

Proof of Theorem 3.1. We define the subbilinear operator \mathfrak{R} by

$$\mathfrak{R}(f, b)(x) := \left| T\left(b(x)f(\cdot) - \Pi_2(f, b)(\cdot)\right)(x) \right| + |T(\Pi_1(f, b))(x)| + |T(\Pi_4(f, b))(x)|$$

for all $(f, b) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$. Then, by Theorem 4.1, we obtain that

$$|T(\mathfrak{S}(f, b))| - \mathfrak{R}(f, b) \leq |[b, T](f)| \leq \mathfrak{R}(f, b) + |T(\mathfrak{S}(f, b))|.$$

By Lemma 4.1, Lemma 4.2 and Lemma 4.3, it is sufficient to show that the subbilinear operator

$$\mathfrak{U}(f, b)(x) := \left| T\left(b(x)f(\cdot) - \Pi_2(f, b)(\cdot)\right)(x) \right|$$

is bounded from $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$.

One first consider $b \in BMO(\mathbb{R}^n)$ and f a ψ -atom related to the cube Q . Then, by Remark 4.1, we have

$$\mathfrak{U}(f, b)(x) = \mathfrak{U}(f, b - b_Q)(x) \leq |(b(x) - b_Q)Tf(x)| + |T(\Pi_2(f, b - b_Q))(x)|.$$

Consequently, by Remark 4.1, Lemma 6.1 and the fact f is C times a classical atom related to the cube mQ , we obtain that

$$(6.2) \quad \|\mathfrak{U}(f, b)\|_{L^1} \leq \|(b - b_Q)Tf\|_{L^1} + \|T\|_{H^1 \rightarrow L^1} \|\Pi_2(f, b - b_Q)\|_{H^1} \leq C\|b\|_{BMO},$$

where $C > 0$ independent of f, b .

Now, for any $b \in BMO(\mathbb{R}^n)$ and $f \in H_{\text{fin}}^1(\mathbb{R}^n)$. By Lemma 6.2, there exists a finite decomposition $f = \sum_{j=1}^k \lambda_j a_j$ such that $\sum_{j=1}^k |\lambda_j| \leq C\|f\|_{H^1}$. Consequently, by (6.2), we obtain that

$$\|\mathfrak{U}(f, b)\|_{L^1} \leq \sum_{j=1}^k |\lambda_j| \|\mathfrak{U}(a_j, b)\|_{L^1} \leq C\|f\|_{H^1} \|b\|_{BMO},$$

which ends the proof as $H_{\text{fin}}^1(\mathbb{R}^n)$ is dense in $H^1(\mathbb{R}^n)$ for the norm $\|\cdot\|_{H^1}$. \square

Proof of Theorem 3.2. We define the bilinear operators \mathfrak{R} by

$$\mathfrak{R}(f, b) = bTf - T(\Pi_1(f, b) + \Pi_2(f, b) + \Pi_4(f, b)),$$

for all $(f, b) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$. By Theorem 4.1, we have

$$(6.3) \quad [b, T](f) = \mathfrak{R}(f, b) + T(\mathfrak{S}(f, b)).$$

One now needs to prove that \mathfrak{R} is bounded from $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $L^1(\mathbb{R}^n)$. Indeed, we first consider $b \in BMO(\mathbb{R}^n)$ and f a ψ -atom related to the cube Q . Then, the equality $[b, T](f) = [b - b_Q, T](f)$ together with Equality (6.3) and Remark 4.1 imply that

$$\mathfrak{R}(f, b) = (b - b_Q)Tf - T(\Pi_1(f, b - b_Q) + \Pi_2(f, b - b_Q) + \Pi_4(f, b - b_Q)).$$

Combining this equality, Lemma 4.2, Lemma 4.3, Remark 4.1, Lemma 6.1 and the fact f is C time a classical atom related to the cube mQ , we obtain that

$$(6.4) \quad \|\mathfrak{R}(f, b)\|_{L^1} \leq C\|b\|_{BMO},$$

where $C > 0$ independent of f, b .

Now, for any $b \in BMO(\mathbb{R}^n)$ and $f \in H_{\text{fin}}^1(\mathbb{R}^n)$. By Lemma 6.2, there exists a finite decomposition $f = \sum_{j=1}^k \lambda_j a_j$ such that $\sum_{j=1}^k |\lambda_j| \leq C\|f\|_{H^1}$. Consequently, by (6.4), we obtain that

$$\|\mathfrak{R}(f, b)\|_{L^1} \leq \sum_{j=1}^k |\lambda_j| \|\mathfrak{R}(a_j, b)\|_{L^1} \leq C\|f\|_{H^1} \|b\|_{BMO},$$

which ends the proof as $H_{\text{fin}}^1(\mathbb{R}^n)$ is dense in $H^1(\mathbb{R}^n)$ for the norm $\|\cdot\|_{H^1}$. \square

Proof of Theorem 3.3. Theorem 3.3 is an immediate corollary of Theorem 3.1 and Theorem 5.1. \square

7. PROOF OF THEOREM 3.4, THEOREM 3.5 AND THEOREM 3.6

First we recall the following well-known result. It can be found in [8] or [14].

Theorem 7.1. *Let T be a Calderón-Zygmund operator satisfying $T1 = T^*1 = 0$, $1 < q < \infty$ and $1/p + 1/q = 1$. Then, $fTg - gT^*f \in H^1(\mathbb{R}^n)$ for all $f \in L^p(\mathbb{R}^n)$, $g \in L^q(\mathbb{R}^n)$.*

Now, in order to prove the bilinear type estimates and the Hardy type theorems for the commutators of Calderón-Zygmund operators, we need the following three technical lemmas.

Lemma 7.1. *Let $\delta \in (0, 1]$, and A, B be two δ -Calderón-Zygmund operators such that $A1 = A^*1 = B1 = B^*1 = 0$. Then, there exists a constant $C = C(n, \delta)$ such that*

$$\sum_{I, I', I''} \sum_{\sigma, \sigma', \sigma'' \in E} |\langle f, \psi_I^\sigma \rangle \langle g, \psi_{I'}^{\sigma'} \rangle \langle A\psi_I^\sigma, \psi_{I''}^{\sigma''} \rangle \langle B\psi_{I'}^{\sigma'}, \psi_{I''}^{\sigma''} \rangle| \leq C\|f\|_{H^1} \|g\|_{BMO}$$

for all $f \in H^1(\mathbb{R}^n), g \in BMO(\mathbb{R}^n)$.

Lemma 7.2. *Let $\delta \in (0, 1]$, and $A_i, B_i, i = 1, \dots, K$, be δ -Calderón-Zygmund operators satisfying $A_i 1 = A_i^* 1 = B_i 1 = B_i^* 1 = 0$, and for every f and g in $L^2(\mathbb{R}^n)$,*

$$\int_{\mathbb{R}^n} \left(\sum_{i=1}^K A_i f \cdot B_i g \right) dx = 0.$$

Then, the bilinear operator \mathfrak{P} , defined by $\mathfrak{P}(f, g) = \sum_{i=1}^K \mathfrak{S}(A_i f, B_i g)$, maps continuously $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $H^1(\mathbb{R}^n)$.

Corollary 7.1. *Let T be a Calderón-Zygmund operator satisfying $T1 = T^*1 = 0$. Then the bilinear operator \mathfrak{P} , defined by $\mathfrak{P}(f, g) = \mathfrak{S}(Tf, g) - \mathfrak{S}(f, T^*g)$, maps continuously $H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$ into $H^1(\mathbb{R}^n)$.*

Lemma 7.3. *Let b be a non-constant BMO-function and $T \in \mathcal{K}_0$. Assume that $f \in H_b^1(\mathbb{R}^n)$ with the wavelet decomposition $f = \sum_{j=1}^{\infty} \sum_{I \subset R_j} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma$ where R_j 's are dyadic cubes and $\sum_{I \subset R_j} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma$ are multiples of ψ -atoms related to the cubes R_j . Set $f_k = \sum_{j=1}^k \sum_{I \subset R_j} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma$, $k = 1, 2, \dots$. Then, the sequence $\{[b, T](f_k)\}_{k \geq 1}$ tends to $[b, T](f)$ in the sense of distributions $\mathcal{S}'(\mathbb{R}^n)$.*

Proof of Lemma 7.1. We first remark (see [36], Proposition 1) that there exists a constant $C > 0$ such that for all dyadic cubes I, I' and $\sigma, \sigma' \in E$, we have

$$(7.1) \quad \max\{|\langle A\psi_I^\sigma, \psi_{I'}^{\sigma'} \rangle|, |\langle B\psi_I^\sigma, \psi_{I'}^{\sigma'} \rangle|\} \leq C 2^{-|j-j'|(\delta+n/2)} \left(\frac{2^{-j} + 2^{-j'}}{2^{-j} + 2^{-j'} + |x_I - x_{I'}|} \right)^{n+\delta}.$$

Consequently,

$$(7.2) \quad \max\{|\langle A\psi_I^\sigma, \psi_{I'}^{\sigma'} \rangle|, |\langle B\psi_I^\sigma, \psi_{I'}^{\sigma'} \rangle|\} \leq C p_\delta(I, I')$$

with

$$p_\delta(I, I') = \frac{2^{-|j-j'|(\delta/2+n/2)}}{1 + |j - j'|^2} \left(\frac{2^{-j} + 2^{-j'}}{2^{-j} + 2^{-j'} + |x_I - x_{I'}|} \right)^{n+\delta/2}.$$

Here $|I| = 2^{-jn}$ and $|I'| = 2^{-j'n}$, while x_I and $x_{I'}$ denote the centers of the two cubes. On the other hand, it follows from [[14], Lemma 1.3] that there exists a constant $C = C(n, \delta) > 0$ such that

$$(7.3) \quad \sum_{I''} p_\delta(I, I'') p_\delta(I', I'') \leq C p_\delta(I, I').$$

Combining (7.2) and (7.3), we obtain

$$\sum_{I, I', I''} \sum_{\sigma, \sigma', \sigma'' \in E} |\langle f, \psi_I^\sigma \rangle \langle g, \psi_{I'}^{\sigma'} \rangle \langle A\psi_I^\sigma, \psi_{I''}^{\sigma''} \rangle \langle B\psi_{I'}^{\sigma'}, \psi_{I''}^{\sigma''} \rangle| \leq C \sum_{I, I'} \sum_{\sigma, \sigma' \in E} p_\delta(I, I') |\langle f, \psi_I^\sigma \rangle| |\langle g, \psi_{I'}^{\sigma'} \rangle|.$$

It is easy to establish that the matrix $\{p_\delta(I, I')\}_{I, I'}$ is almost diagonal (by taking $\varepsilon = \delta/4$ in the definition (3.1) of Frazier and Jawerth [16]) and thus is bounded on $f_1^{0,2}$ the space of all sequences $(a_I)_I$ such that $\left(\sum_I |a_I|^2 |I|^{-1} \chi_I\right)^{1/2}$ is in $L^1(\mathbb{R}^n)$. We then use the wavelet characterization of $H^1(\mathbb{R}^n)$ (see Theorem 2.1) and the fact that (cf. [16])

$$\sum_{I'} \sum_{\sigma' \in E} |\langle h, \psi_{I'}^{\sigma'} \rangle| |\langle g, \psi_{I'}^{\sigma'} \rangle| \leq C \|h\|_{H^1} \|g\|_{BMO},$$

for all $h \in H^1(\mathbb{R}^n)$, to conclude that

$$\sum_{I, I', I''} \sum_{\sigma, \sigma', \sigma'' \in E} |\langle f, \psi_I^\sigma \rangle \langle g, \psi_{I'}^{\sigma'} \rangle \langle A \psi_I^\sigma, \psi_{I''}^{\sigma''} \rangle \langle B \psi_{I'}^{\sigma'}, \psi_{I''}^{\sigma''} \rangle| \leq C \|f\|_{H^1} \|g\|_{BMO}.$$

□

Proof of Lemma 7.2. By Lemma 7.1, we have

$$\begin{aligned} \mathfrak{P}(f, g) &= \sum_{i=1}^K \mathfrak{S}(A_i f, B_i g) \\ &= \sum_{i=1}^K \sum_{I, I', I''} \sum_{\sigma, \sigma', \sigma'' \in E} \langle f, \psi_I^\sigma \rangle \langle g, \psi_{I'}^{\sigma'} \rangle \langle A_i \psi_I^\sigma, \psi_{I''}^{\sigma''} \rangle \langle B_i \psi_{I'}^{\sigma'}, \psi_{I''}^{\sigma''} \rangle (\psi_{I''}^{\sigma''})^2 \end{aligned}$$

where all the series converge in $L^1(\mathbb{R}^n)$. For any dyadic cubes $I, I', \sigma, \sigma' \in E$, we have

$$\begin{aligned} &\sum_{i=1}^K \sum_{I''} \sum_{\sigma'' \in E} \langle f, \psi_I^\sigma \rangle \langle g, \psi_{I'}^{\sigma'} \rangle \langle A_i \psi_I^\sigma, \psi_{I''}^{\sigma''} \rangle \langle B_i \psi_{I'}^{\sigma'}, \psi_{I''}^{\sigma''} \rangle (\psi_{I''}^{\sigma''})^2 \\ &= \sum_{i=1}^K \sum_{I''} \sum_{\sigma'' \in E} \langle f, \psi_I^\sigma \rangle \langle g, \psi_{I'}^{\sigma'} \rangle \langle A_i \psi_I^\sigma, \psi_{I''}^{\sigma''} \rangle \langle B_i \psi_{I'}^{\sigma'}, \psi_{I''}^{\sigma''} \rangle \left((\psi_{I''}^{\sigma''})^2 - (\psi_I^\sigma)^2 \right) \end{aligned}$$

since (see Remark 4.1)

$$\sum_{i=1}^K \sum_{I''} \sum_{\sigma'' \in E} \langle A_i \psi_I^\sigma, \psi_{I''}^{\sigma''} \rangle \langle B_i \psi_{I'}^{\sigma'}, \psi_{I''}^{\sigma''} \rangle = \int_{\mathbb{R}^n} \left(\sum_{i=1}^K A_i \psi_I^\sigma \cdot B_i \psi_{I'}^{\sigma'} \right) dx = 0.$$

An explicit computation gives that $|\psi_{I''}^{\sigma''}|^2 - |\psi_I^\sigma|^2$ is in $H^1(\mathbb{R}^n)$, with

$$\| |\psi_{I''}^{\sigma''}|^2 - |\psi_I^\sigma|^2 \|_{H^1} \leq C \left(\log(2^{-j} + 2^{-j''})^{-1} + \log(|x_I - x_{I''}| + 2^{-j} + 2^{-j''}) \right).$$

Here $|I| = 2^{-jn}$ and $|I''| = 2^{-j''n}$, while x_I and $x_{I''}$ denote the centers of the two cubes. Consequently, by (7.1) and (7.3), we get

$$\begin{aligned}
 & \left\| \sum_{i=1}^K \sum_{I''} \sum_{\sigma'' \in E} \langle f, \psi_I^\sigma \rangle \langle g, \psi_{I'}^{\sigma'} \rangle \langle A_i \psi_I^\sigma, \psi_{I''}^{\sigma''} \rangle \langle B_i \psi_{I'}^{\sigma'}, \psi_{I''}^{\sigma''} \rangle (\psi_{I''}^{\sigma''})^2 \right\|_{H^1} \\
 & \leq \sum_{i=1}^K \sum_{I''} \sum_{\sigma'' \in E} |\langle f, \psi_I^\sigma \rangle \langle g, \psi_{I'}^{\sigma'} \rangle \langle A_i \psi_I^\sigma, \psi_{I''}^{\sigma''} \rangle \langle B_i \psi_{I'}^{\sigma'}, \psi_{I''}^{\sigma''} \rangle| \left\| (\psi_{I''}^{\sigma''})^2 - (\psi_I^\sigma)^2 \right\|_{H^1} \\
 & \leq C \sum_{i=1}^K \sum_{I''} \sum_{\sigma'' \in E} |\langle f, \psi_I^\sigma \rangle \langle g, \psi_{I'}^{\sigma'} \rangle| p_\delta(I, I'') p_\delta(I', I'') \\
 & \leq C p_\delta(I, I') |\langle f, \psi_I^\sigma \rangle| |\langle g, \psi_{I'}^{\sigma'} \rangle|,
 \end{aligned}$$

here we used the fact that

$$(1 + |j - j''|^2) \log \left(\frac{|x_I - x_{I''}| + 2^{-j} + 2^{-j''}}{2^{-j} + 2^{-j''}} \right) \leq C(\delta) 2^{|j - j''|\delta/2} \left(\frac{|x_I - x_{I''}| + 2^{-j} + 2^{-j''}}{2^{-j} + 2^{-j''}} \right)^{\delta/2}.$$

Thus, the same argument as in the proof of Lemma 7.1 allows to conclude that

$$\begin{aligned}
 \|\mathfrak{B}(f, g)\|_{H^1} & \leq C \sum_{I, I'} \sum_{\sigma, \sigma' \in E} p_\delta(I, I') |\langle f, \psi_I^\sigma \rangle| |\langle g, \psi_{I'}^{\sigma'} \rangle| \\
 & \leq C \|f\|_{H^1} \|g\|_{BMO},
 \end{aligned}$$

which ends the proof. \square

Before giving the proof of Lemma 7.3, let us recall the following lemma. It can be found in [17] (Lemma 2.3).

Lemma 7.4. *Let T be a Calderón-Zygmund operator satisfying $T1 = 0$. Then T maps $\mathcal{S}(\mathbb{R}^n)$ into $L^\infty(\mathbb{R}^n)$. Moreover, there exists a constant $C > 0$, depending only on T , such that for any $\phi \in \mathcal{S}(\mathbb{R}^n)$ with $\text{supp } \phi \subset B(x_0, r)$, we have*

$$\|T\phi\|_{L^\infty} \leq C(\|\phi\|_{L^\infty} + r\|\nabla\phi\|_{L^\infty})$$

Proof of Lemma 7.3. By Theorem 3.2, it is sufficient to prove that

$$\lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} T(\mathfrak{S}(f_k, b)) h dx = \int_{\mathbb{R}^n} T(\mathfrak{S}(f, b)) h dx,$$

for all $h \in \mathcal{S}(\mathbb{R}^n)$. Because of the hypothesis, we observe that $\mathfrak{S}(f, b) \in H^1(\mathbb{R}^n)$ and $\mathfrak{S}(f_k, b) \in L^q(\mathbb{R}^n)$, $k = 1, 2, \dots$, for some $q \in (1, 2)$ (see Lemma 2.1).

For every $h \in \mathcal{S}(\mathbb{R}^n)$. Let $\mathfrak{S}(f, b) = \sum_{j=1}^\infty \lambda_j a_j$ be a classical L^q -atomic decomposition of $\mathfrak{S}(f, b)$. Then, $T(\sum_{j=1}^k \lambda_j a_j)$ tends to $T(\mathfrak{S}(f, b))$ in $L^1(\mathbb{R}^n)$ (in fact, it also holds in $H^1(\mathbb{R}^n)$ since $T^*1 = 0$). Hence, as $h \in \mathcal{S}(\mathbb{R}^n) \subset$

$L^\infty(\mathbb{R}^n) \cap L^{q'}(\mathbb{R}^n)$ where $1/q + 1/q' = 1$, $\mathfrak{S}(f_k, b), a_j \in L^q(\mathbb{R}^n)$ and $T^*h \in L^\infty(\mathbb{R}^n)$ since $T^*1 = 0$ (see Lemma 7.4), by Theorem 7.1 we get

$$\begin{aligned} \int_{\mathbb{R}^n} T(\mathfrak{S}(f, b))h dx &= \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} T\left(\sum_{j=1}^k \lambda_j a_j\right)h dx = \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} \left(\sum_{j=1}^k \lambda_j a_j\right)T^*h dx \\ &= \int_{\mathbb{R}^n} \mathfrak{S}(f, b)T^*h dx = \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} \mathfrak{S}(f_k, b)T^*h dx \\ &= \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} T(\mathfrak{S}(f_k, b))h dx, \end{aligned}$$

since $\mathfrak{S}(f_k, b)$ tends to $\mathfrak{S}(f, b)$ in $L^1(\mathbb{R}^n)$ by f_k tends to f in $H^1(\mathbb{R}^n)$ (see Theorem 3.2). This finishes the proof. \square

Proof of Theorem 3.4. Let $(f, g) \in H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n)$. By Theorem 3.2 and Lemma 7.2, we obtain $\mathfrak{T}(f, g) = \sum_{i=1}^K [B_i g, T](A_i f) \in L^1(\mathbb{R}^n)$, moreover,

$$\begin{aligned} \|\mathfrak{T}(f, g)\|_{L^1} &\leq \sum_{i=1}^K \|\mathfrak{R}(A_i f, B_i g)\|_{L^1} + \left\| T\left(\sum_{i=1}^K \mathfrak{S}(A_i f, B_i g)\right) \right\|_{L^1} \\ &\leq C \sum_{i=1}^K \|A_i f\|_{H^1} \|B_i g\|_{BMO} + \|T\|_{H^1 \rightarrow L^1} \left\| \sum_{i=1}^K \mathfrak{S}(A_i f, B_i g) \right\|_{H^1} \\ &\leq C \|f\|_{H^1} \|g\|_{BMO}. \end{aligned}$$

This completes the proof. \square

Proof of Theorem 3.5. Let $f \in H_b^1(\mathbb{R}^n)$, we prove $[b, T](f) \in h^1(\mathbb{R}^n)$ via the fact that $BMO^{\log}(\mathbb{R}^n)$ is the dual of $H^{\log}(\mathbb{R}^n)$ (see [23]). Indeed, by Theorem 2.2, there exists a decomposition $f = \sum_{j=1}^\infty \sum_{I \subset R_j} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma$ where $\sum_{I \subset R_j} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma$ are multiples of ψ -atoms related to the dyadic cubes R_j . Set $f_k = \sum_{j=1}^k \sum_{I \subset R_j} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \psi_I^\sigma$, $k = 1, 2, \dots$. Then, the sequence $[b, T](f_k)$ tends to $[b, T](f)$ in the sense of distributions $\mathcal{S}'(\mathbb{R}^n)$ (see Lemma 7.3), and thus

$$(7.4) \quad \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} [b, T](f_k)h dx = \int_{\mathbb{R}^n} [b, T](f)h dx,$$

for all $h \in C_0^\infty(\mathbb{R}^n)$. Notice that $[b, T](f_k) \in L^2(\mathbb{R}^n)$ and $[b, T](f) \in L^1(\mathbb{R}^n)$.

For every $h \in C_0^\infty(\mathbb{R}^n)$. By Lemma 4.2, Lemma 4.3, Lemma 4.4, Remark 4.1 and Corollary 7.1, we have $hT(f_k) - f_k \left(T^*h - (T^*h)_\mathbb{Q} \right) \in H^{\log}(\mathbb{R}^n)$. More

precisely,

$$\begin{aligned}
 & \left\| hT(f_k) - f_k \left(T^*h - (T^*h)_\mathbb{Q} \right) \right\|_{H^{\log}} \\
 \leq & C \left\{ \left\| \mathfrak{S}(T(f_k), h) - \mathfrak{S}(f_k, T^*h - (T^*h)_\mathbb{Q}) \right\|_{H^1} + \right. \\
 & \left. + \sum_{j=1,4} \left(\left\| \Pi_j(T(f_k), h) \right\|_{H^1} + \left\| \Pi_j(f_k, T^*h - (T^*h)_\mathbb{Q}) \right\|_{H^1} \right) + \right. \\
 & \left. + \left\| \Pi_2(T(f_k), h) \right\|_{H^{\log}} + \left\| \Pi_2(f_k, T^*h - (T^*h)_\mathbb{Q}) \right\|_{H^{\log}} \right\} \\
 \leq & C \left\{ \left\| f_k \right\|_{H^1} \left\| h \right\|_{BMO} + \left\| T(f_k) \right\|_{H^1} \left\| h \right\|_{BMO} + \left\| f_k \right\|_{H^1} \left\| T^*h - (T^*h)_\mathbb{Q} \right\|_{BMO} + \right. \\
 & \left. + \left\| T(f_k) \right\|_{H^1} \left\| h \right\|_{BMO^+} + \left\| f_k \right\|_{H^1} \left\| T^*h - (T^*h)_\mathbb{Q} \right\|_{BMO^+} \right\} \\
 \leq & C \left(\left\| f_k \right\|_{H^1} \left\| h \right\|_{bmo} + \left\| f_k \right\|_{H^1} \left\| T^*h \right\|_{BMO} \right) \leq C \left\| f \right\|_{H^1} \left\| h \right\|_{bmo},
 \end{aligned}$$

here one used $\mathfrak{S}(f, T^*h - (T^*h)_\mathbb{Q}) = \mathfrak{S}(f, T^*h)$, $\|T^*h - (T^*h)_\mathbb{Q}\|_{BMO^+} = \|T^*h\|_{BMO}$ and $\|f_k\|_{H^1} \leq C\|f\|_{H^1}$. As the L^2 - functions f_k have compact support, $b \in BMO^{\log}(\mathbb{R}^n) \subset BMO(\mathbb{R}^n)$, we deduce that $bhT(f_k)$, $hT(bf_k)$, $bf_kT^*h \in L^1(\mathbb{R}^n)$. Moreover, $\int_{\mathbb{R}^n} hT(bf_k)dx = \int_{\mathbb{R}^n} bf_kT^*hdx$ since $hT(bf_k) - bf_kT^*h \in H^1(\mathbb{R}^n)$ (see Theorem 7.1). Therefore, as $BMO^{\log}(\mathbb{R}^n)$ is the dual of $H^{\log}(\mathbb{R}^n)$ (see [23]), we get

$$\begin{aligned}
 \left| \int_{\mathbb{R}^n} [b, T](f_k)hdx \right| &= \left| \int_{\mathbb{R}^n} b(hT(f_k) - f_kT^*h)dx \right| \\
 &\leq \left| \int_{\mathbb{R}^n} b \left(hT(f_k) - f_k \left(T^*h - (T^*h)_\mathbb{Q} \right) \right) dx \right| + |(T^*h)_\mathbb{Q}| \left| \int_{\mathbb{R}^n} bf_kdx \right| \\
 &\leq C\|b\|_{BMO^{\log}} \left\| hT(f_k) - f_k \left(T^*h - (T^*h)_\mathbb{Q} \right) \right\|_{H^{\log}} + |(T^*h)_\mathbb{Q}| \left| \int_{\mathbb{R}^n} bf_kdx \right| \\
 &\leq C\|b\|_{BMO^{\log}} \|f\|_{H^1} \|h\|_{bmo} + |(T^*h)_\mathbb{Q}| \left| \sum_{j=1}^k \sum_{ICR_j} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \langle b, \psi_I^\sigma \rangle \right|.
 \end{aligned}$$

The above inequality and (7.4) imply that for all $h \in C_0^\infty(\mathbb{R}^n)$, we obtain

$$\left| \int_{\mathbb{R}^n} [b, T](f) h dx \right| \leq C \|b\|_{BMO^{\log}} \|f\|_{H^1} \|h\|_{bmo}$$

since $\mathfrak{S}(f, b) \in H^1(\mathbb{R}^n)$ (see Theorem 5.1) and thus (see Remark 4.1)

$$\lim_{k \rightarrow \infty} \sum_{j=1}^k \sum_{I \subset R_j} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \langle b, \psi_I^\sigma \rangle = \int_{\mathbb{R}^n} \mathfrak{S}(f, b) dx = 0.$$

This proves that $[b, T](f) \in h^1(\mathbb{R}^n)$ since $h^1(\mathbb{R}^n)$ is the dual of $vmo(\mathbb{R}^n)$ (see Section 2). Furthermore,

$$\|[b, T](f)\|_{h^1} \leq C \|b\|_{BMO^{\log}} \|f\|_{H^1} \leq C \|b\|_{BMO^{\log}} \|b\|_{BMO}^{-1} \|f\|_{H_b^1},$$

which ends the proof of Theorem 3.5. \square

Proof of Theorem 3.6. The idea of the proof is similar to the one of Theorem 3.5. Let $f \in H_b^1(\mathbb{R}^n)$ and the sequence $\{f_k\}_{k \geq 1}$ as in Theorem 3.5. For every $h \in C_0^\infty(\mathbb{R}^n)$. Similarly to the argument used in the proof of Theorem 3.5, we obtain that $(h - h_{\mathbb{Q}})T(f_k) - f_k(T^*h - (T^*h)_{\mathbb{Q}}) \in H^{\log}(\mathbb{R}^n)$, moreover,

$$\left\| (h - h_{\mathbb{Q}})T(f_k) - f_k(T^*h - (T^*h)_{\mathbb{Q}}) \right\|_{H^{\log}} \leq C \|f\|_{H^1} \|h\|_{BMO}.$$

Here one used $\|h - h_{\mathbb{Q}}\|_{BMO} = \|h\|_{BMO}$ and $\|h - h_{\mathbb{Q}}\|_{BMO^+} = \|h\|_{BMO}$. Hence,

$$\begin{aligned} \left| \int_{\mathbb{R}^n} [b, T](f_k) \cdot h dx \right| &= \left| \int_{\mathbb{R}^n} b(hT(f_k) - f_k T^*h) dx \right| \\ &\leq \left| \int_{\mathbb{R}^n} b \left((h - h_{\mathbb{Q}})T(f_k) - f_k(T^*h - (T^*h)_{\mathbb{Q}}) \right) dx \right| + |(T^*h)_{\mathbb{Q}}| \left| \int_{\mathbb{R}^n} b f_k dx \right| \\ &\leq C \|b\|_{BMO^{\log}} \|f\|_{H^1} \|h\|_{BMO} + |(T^*h)_{\mathbb{Q}}| \left| \sum_{j=1}^k \sum_{I \subset R_j} \sum_{\sigma \in E} \langle f, \psi_I^\sigma \rangle \langle b, \psi_I^\sigma \rangle \right|, \end{aligned}$$

since $\int_{\mathbb{R}^n} bT(f_k) dx = 0$ by $T^*b = 0$ and f_k is a finite linear combination of classical atoms. Similarly to the argument used in the proof of Theorem 3.5, we also get

$$\left| \int_{\mathbb{R}^n} [b, T](f) \cdot h dx \right| \leq C \|b\|_{BMO^{\log}} \|f\|_{H^1} \|h\|_{BMO}$$

This proves that $[b, T](f) \in H^1(\mathbb{R}^n)$ since $H^1(\mathbb{R}^n)$ is the dual of $VMO(\mathbb{R}^n)$ (see Section 2). Furthermore,

$$\|[b, T](f)\|_{H^1} \leq \|b\|_{BMO^{\log}} \|b\|_{BMO}^{-1} \|f\|_{H_b^1},$$

which ends the proof of Theorem 3.6. □

8. COMMUTATORS OF FRACTIONAL INTEGRALS

Given $0 < \alpha < n$, the fractional integral operator I_α is defined by

$$I_\alpha f(x) = \int_{\mathbb{R}^n} \frac{f(y)}{|x - y|^{n-\alpha}} dy.$$

Let b be a locally integrable function. We consider the linear commutator $[b, I_\alpha]$ is defined by

$$[b, I_\alpha](f) = bI_\alpha f - I_\alpha(bf).$$

We end this article by presenting some results related to commutators of fractional integrals as follows.

Theorem 8.1. *Let $0 < \alpha < n$. There exist a bounded bilinear operator $\mathfrak{R} : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^{n/(n-\alpha)}(\mathbb{R}^n)$ and a bounded bilinear operator $\mathfrak{S} : H^1(\mathbb{R}^n) \times BMO(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ such that*

$$[b, I_\alpha](f) = \mathfrak{R}(f, b) + I_\alpha(\mathfrak{S}(f, b)).$$

Corollary 8.1. *Let $0 < \alpha < n$ and $b \in BMO(\mathbb{R}^n)$. Then, the linear commutator $[b, I_\alpha]$ maps continuously $H^1(\mathbb{R}^n)$ into weak- $L^{n/(n-\alpha)}(\mathbb{R}^n)$.*

Theorem 8.2. *Let $0 < \alpha < n$, $b \in BMO(\mathbb{R}^n)$, and $1 < q \leq \infty$. Then, the linear commutator $[b, I_\alpha]$ maps continuously $H_b^1(\mathbb{R}^n)$ into $L^{n/(n-\alpha)}(\mathbb{R}^n)$.*

The results above can be proved similarly to Theorem 3.2 and Theorem 3.3. We leave the proofs to the interested readers. In the case $H_b^1(\mathbb{R}^n)$ is replaced by $\mathcal{H}_b^1(\mathbb{R}^n)$, Theorem 8.2 was considered by the authors in [13]. There, they proved that

$$\sup\{\|[b, I_\alpha](a)\|_{L^{n/(n-\alpha)}} : a \text{ is a } (\infty, b)\text{-atom}\} < \infty.$$

However, it is now well-known that this argument does not suffice to conclude that $[b, I_\alpha]$ is bounded from $\mathcal{H}_b^1(\mathbb{R}^n)$ to $L^{n/(n-\alpha)}(\mathbb{R}^n)$.

REFERENCES

- [1] J. Alvarez, R.J. Bagby, D.S. Kurtz and C. Pérez, Weighted estimates for commutators of linear operators. *Studia Math.* 104 (1993), no. 2, 195–209.
- [2] J. Alvarez and J. Hounie, Estimates for the kernel and continuity properties of pseudo-differential operators. *Ark. Mat.* 28 (1990), no. 1, 1–22.
- [3] A. Bonami, J. Feuto and S. Grellier, Endpoint for the div-curl lemma in Hardy spaces. *Publ. Mat.* 54, No. 2 (2010), 341–358.

- [4] A. Bonami, S. Grellier and L. D. Ky, Paraproducts and products of functions in $BMO(\mathbb{R}^n)$ and $H^1(\mathbb{R}^n)$ through wavelets, accepted to J. Math. Pure Appl.
- [5] G. Bourdaud, Remarques sur certains sous-espaces de $BMO(\mathbb{R}^n)$ et de $bmo(\mathbb{R}^n)$. (French) [Remarks on some subspaces of $BMO(\mathbb{R}^n)$ and $bmo(\mathbb{R}^n)$] Ann. Inst. Fourier (Grenoble) 52 (2002), no. 4, 1187–1218.
- [6] M. Bownik, Boundedness of operators on Hardy spaces via atomic decompositions. Proc. Amer. Math. Soc. 133 (2005), 3535–3542.
- [7] Y. Chen and Y. Ding, Commutators of Littlewood-Paley operators. Sci. China Ser. A 52 (2009), no. 11, 2493–2505.
- [8] R. Coifman and L. Grafakos, Hardy space estimates for multilinear operators. I. Rev. Mat. Iberoamericana 8 (1992), no. 1, 45–67.
- [9] R. R. Coifman and G. Weiss, Extensions of Hardy spaces and their use in analysis. Bull. Amer. Math. Soc. 83 (1977), 569–645.
- [10] R.R. Coifman, R. Rochberg and G. Weiss, Factorization theorems for Hardy spaces in several variables. Ann. of Math., 103, 611–635 (1976)
- [11] G. Dafni, Local VMO and weak convergence in H^1 , Canad. Math. Bull. 45 (2002), p. 46–59.
- [12] I. Daubechies, Orthonormal basis of compactly supported wavelets, Comm. Pure Appl. Math. 41 (1988), 909–996.
- [13] Y. Ding, S. Z. Lu and P. Zhang, Continuity of higher order commutators on certain Hardy spaces. Acta Math. Sinica. 18 (2002), 391–404.
- [14] S. Dobyinsky, La "version ondelettes" du théorème du Jacobien. (French) [The "wavelet version" of the theorem of the Jacobian] Rev. Mat. Iberoamericana 11 (1995), no. 2, 309–333.
- [15] C. Fefferman, E.M. Stein, H^p spaces of several variables. Acta Math. 129 (1972), 137–193.
- [16] M. Frazier and B. Jawerth, A discrete transform and decompositions of distribution spaces. J. Funct. Anal. 93 (1990), no. 1, 34–170.
- [17] M. Frazier, R. Torres and G. Weiss, The boundedness of Calderón-Zygmund operators on the spaces $\dot{F}_p^{\alpha,q}$. Rev. Mat. Iberoamericana, 4 (1988), no. 1, 41–72.
- [18] D. Golberg, A local version of Hardy spaces, Duke J. Math. 46 (1979), 27–42.
- [19] Y. Han, D. Müller and D. Yang, Littlewood-Paley characterizations for Hardy spaces on spaces of homogeneous type. Math. Nachr. 279 (2006), No. 13-14, 1505–537.
- [20] E. Harboure, C. Segovia and J. L. Torrea, Boundedness of commutators of fractional and singular integrals for the extreme values of p . Illinois J. Math. 41 (1997), 676–700.
- [21] E. Hernández and G. Weiss, A first course on wavelets, with a foreword by Yves Meyer, Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1996.
- [22] P. W. Jones and J-L. Journé, On weak convergence in $H^1(\mathbb{R}^n)$. Proc. Amer. Math. Soc. 120 (1994), no. 1, 137–138.
- [23] L. D. Ky, New Hardy spaces of Musielak-Orlicz type and boundedness of sublinear operators, submitted.
- [24] M-Y. Lee, Weighted norm inequalities of Bochner-Riesz means. J. Math. Anal. Appl. 324 (2006), no. 2, 1274–1281.
- [25] W. Li, S. Lu and H. Mo, The boundedness for commutators of multipliers. Acta Math. Appl. Sin. Engl. Ser. 23 (2007), no. 1, 113–122.
- [26] Y. Lin and S. Lu, Boundedness of commutators on Hardy-type spaces. Integr. equ. oper. theory 57 (2007), 381–396.

- [27] Z. Liu and S. Z. Lu, Endpoint estimates for commutators of Calderón-Zygmund type operators. *Kodai. Math. J.* 25 (2002), 79–88.
- [28] L.Z. Liu, Continuity for commutators of Littlewood-Paley operator on certain Hardy spaces. *J. Korean Math. Soc.* 40 (2003), No. 1, 41–60.
- [29] L. Z. Liu and S. Z. Lu, Continuity for maximal multilinear Bochner-Riesz operators on Hardy and Herz-Hardy spaces. *Acta Math. Sin. (Engl. Ser.)* 22 (2006), no. 1, 69–76.
- [30] Z. Liu, G. Lu and S. Lu, Continuity properties for the maximal operator associated with the commutator of the Bochner-Riesz operator. *Publ. Mat.* 47 (2003), no. 1, 45–69.
- [31] L. Liu and Q. Tong, Continuity for maximal commutator of Bochner-Riesz operators on some weighted Hardy spaces. *Int. J. Math. Math. Sci.* 2005, no. 2, 195–201.
- [32] S. Z. Lu, Q. Wu and D. C. Yang, Boundedness of commutators on Hardy type spaces. *Sci. China Ser. A.* 45 (2002), 984–997.
- [33] S. Z. Lu and L. F. Xu, Boundedness of some Marcinkiewicz integral operators related to higher order commutators on Hardy spaces. *Acta Math. Sinica.* 22 (2006), 105–114.
- [34] S. Meda, P. Sjögren and M. Vallarino, On the H^1-L^1 boundedness of operators. *Proc. Amer. Math. Soc.* 136 (2008), 2921–2931.
- [35] Y. Meyer, *Wavelets and operators*. Advanced mathematics. Cambridge University Press, 1992.
- [36] Y. Meyer and R. Coifman, *Wavelets, Calderón-Zygmund and multilinear operators*. Advanced mathematics. Cambridge University Press, 1997.
- [37] E. Nakai and K. Yabuta, Pointwise multipliers for functions of bounded mean oscillation. *J. Math. Soc. Japan.* 37 (1985), 207–218.
- [38] C. Pérez, Endpoint estimates for commutators of singular integral operators. *J. Func. Anal.* 128 (1995), 163–185.
- [39] J.L. Rubio de Francia, F.J. Ruiz and J.L. Torrea, Calderón-Zygmund theory for operator-valued kernels. *Adv. in Math.* 62 (1986), no. 1, 7–48.
- [40] E.M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*. Princeton Univ. Press, Princeton, NJ, 1993.
- [41] K. Wang and L. Liu, Boundedness for multilinear commutator of multiplier operator on Hardy spaces. *Sci. Ser. A Math. Sci. (N.S.)* 17 (2009), 19–26.
- [42] D. Y. Yan, G. E. Hu and J. C. Lan, Weak-type endpoint estimates for multilinear singular integral operators. *Acta Math. Sinica.* 21 (2005), 209–214.
- [43] P. Zhang and J. Hua, Commutators of multipliers on Hardy spaces. *Anal. Theory Appl.* 21 (2005), no. 3, 226–234.

MAPMO-UMR 6628, DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ D'ORLÉANS,
45067 ORLÉANS CEDEX 2, FRANCE

E-mail address: dangky@math.cnrs.fr