

Accepted Manuscript

Title: A historical declination curve for Munich from different data sources

Authors: Monika Korte, Mioara Manda, Jürgen Matzka


PII: S0031-9201(09)00179-4
DOI: doi:10.1016/j.pepi.2009.08.005
Reference: PEPI 5200

To appear in: *Physics of the Earth and Planetary Interiors*

Received date: 27-5-2009
Revised date: 14-8-2009
Accepted date: 19-8-2009

Please cite this article as: Korte, M., Manda, M., Matzka, J., A historical declination curve for Munich from different data sources, *Physics of the Earth and Planetary Interiors* (2008), doi:10.1016/j.pepi.2009.08.005

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



A historical declination curve for Munich from different data sources

Monika Korte, Mioara Mandea^a, Jürgen Matzka^b

^a*Helmholtz Zentrum Potsdam, Deutsches GeoForschungsZentrum GFZ, Telegrafenberg, 14473 Potsdam, Germany*

^b*Danish Meteorological Institute, Lyngbyvej 100, 2100 Copenhagen, Denmark*

Abstract

Long series of geomagnetic field changes are important for studying processes in the Earth's core. We have compiled 635 historical declination values for southern Germany and surrounding areas. Indirect estimations, including the oldest ones from back in the 15th century, come from 185 sundials and compasses with declination information, and 15 historical maps. Measurements carried out by monks in the time interval 1668 to 1854 amount to 122 annual mean values and data related to the orientation of mine shafts contributes by 313 annual values for several locations. All these data can be useful to improve historical geomagnetic field models. Previously compiled German church orientations, however, are shown to be no reliable sources of the past declination. The compiled new declination curve for Munich shows general agreement with previously published curves for London and Paris and allows to detect geomagnetic jerks with a temporal uncertainty of ± 10 yrs. More or less regular impulses, on a decadal time-scale ranging from 30 to 60 years, are identified for most of the time interval 1400 to 2000, but the century from about 1760 to 1860 seems to be devoid of sudden secular variation changes.

Key words: Geomagnetism, declination, historical data, geomagnetic jerks

1. Introduction

The Earth's magnetic field generated by dynamo processes in the core varies on a broad range of time-scales, from years to millions of years, known as secular variation. A highly

4 detailed picture of the current secular variation is obtained from satellite magnetic vec-
5 tor measurements (Ørsted and CHAMP satellite), available since 1999. Systematic vector
6 measurements at locations all over the world, leading to the present network of geomag-
7 netic observatories, started in the early 19th century with the initiative of Alexander von
8 Humboldt and Carl Friedrich Gauss. Moreover, a major improvement was the develop-
9 ment of a method to determine the absolute field intensity by Gauss (Gauss, 1833). Time
10 series longer than 200 years are, however, desirable to study decadal to centennial secu-
11 lar variation processes. Archaeo- and paleomagnetic data provide information on longer
12 intervals, but mostly with limited time resolution and lower accuracy than direct measure-
13 ments. Magnetic data from historical sources are then an important complement to study
14 the field evolution of the past centuries.

15 The first measured geomagnetic field element was the declination, linked to the early
16 use of compasses. The magnetic compass has been known in Europe since the 12th century
17 (e.g. Merrill et al., 1996), but it is not clear when its deviation from geographic north, i.e.
18 declination, became known there. In China, both the compass and declination had been
19 known some centuries before. The earliest known European declination measurements
20 start with an observation by Hartmann in Rome in 1510 (Hellmann, 1899). The discovery
21 of declination in the European area has often been ascribed to Christopher Columbus
22 in 1492, but Chapman and Bartels (1962) describe evidence from ancient sundials and
23 compasses indicating that the deviation of a compass needle from geographic north was
24 known in Europe at least since the early 15th century. This knowledge, however, might not
25 have been widely distributed. Moreover, some of the first observed deviations might have
26 been interpreted as resulting either from the construction of specific instruments (see e.g.
27 Wolkenhauer, 1904) or from different magnetisation directions of the loadstone, used to
28 magnetise the needle, depending on its location of origin (see e.g. Balmer, 1965; Körber,
29 1965). The oldest declination value given by a magnetic compass known to us dates from
30 1451 (Zinner, 1939). This instrument was made by Peurbach in Vienna. However, it is
31 not clear if Peurbach understood the deviation from geographic north as a property of the
32 magnetic field or as one of the specific instrument. Three more compasses made by him at

33 the same location between 1451 and 1456 indicate different declination values, although the
34 discovery of change of declination with time (i.e. secular variation) is generally assumed
35 to lie only in the early 17th century. It has first been described by Henry Gellibrand
36 in 1634 and probably been noticed before by Edmund Gunter in 1622 (Chapman and
37 Bartels, 1962). It is generally assumed that inclination was discovered in 1544 by Georg
38 Hartmann at Nuremberg but probably first measured correctly by Robert Norman in 1576
39 (e.g. Chapman and Bartels, 1962).

40 Measurements of the two field directions, declination and inclination, date back further
41 than full vector observations. Frequent measurements were taken particularly from ship-
42 board for navigational purposes from the late 16th century on, and Jonkers et al. (2003)
43 compiled a large global set of such data. Their database also contains a few land measure-
44 ments prior to the start of systematic observations. Individual time series of declination
45 and sometimes inclination have been compiled by Malin and Bullard (1981); Cafarella et al.
46 (1992); Barraclough (1995); Alexandrescu et al. (1996) and Soare et al. (1998) for London,
47 Rome, Edinburgh, Paris and Romania, respectively.

48 When we noticed a historical declination curve for southern Germany based on declina-
49 tion information from sundials (Wagner, 1997) and a number of historical measurements
50 carried out at monasteries and not included in the Jonkers et al. (2003) database, we
51 started new efforts of finding historical data over Germany and surrounding areas. In the
52 course of this work we noticed even more largely unexploited sources of declination data:
53 compass roses printed on old maps (Mandea and Korte, 2007), declination measurements
54 used for mining activities and probably even orientations of churches. These data sources
55 have also been mentioned by Knothe (1987, 1988), who actually compiled declination data
56 from mining activities in Europe, but only published and preserved resulting curves and
57 not the values.

58 Here, we first give an overview of the previously published data compilations for the
59 German region from 1300 on. Then, we discuss the different data sources explored in this
60 study and present the new data compilation for (southern) Germany. We compare the
61 data to predictions from the *gufm1* global geomagnetic field model (Jackson et al., 2000),

62 which covers the time interval from 1590 to 1990, and to the archeomagnetic data from
63 earlier times. A smoothed declination curve for Munich is obtained. Its similarities and
64 differences to the declination curves for London and Paris are shown and the occurrence of
65 geomagnetic jerks from 1400 to present is investigated.

66 **2. Existing German declination data since 1400**

67 *2.1. Geomagnetic observatories*

68 Started in the early 19th century and initiated by Alexander von Humboldt, regular
69 measurements of the geomagnetic field were carried out on a daily basis at fixed times at
70 a growing number of geomagnetic observatories . At the Sternwarte Berlin such measure-
71 ments were carried out from 1836 to 1865 (Encke, 1840, 1844, 1848, 1857, 1884). The
72 annual mean values are included in the supplementary data file. In the late 19th and
73 early 20th century the magnetic field was recorded at several locations in and nearby Ger-
74 many, and annual mean values are archived at the World Data Center (WDC) Edinburgh
75 (<http://www.wdc.bgs.ac.uk>). The time series of three active geomagnetic observatories go
76 back to the 19th century when they are combined with data from one or two predecesing
77 stations located not far away from the present observatories. Table 1 indicates them with
78 their location changes. The data are published in the yearbooks of the observatories and
79 are also available from the WDC Edinburgh. We only consider the MNH-MAS-FUR time
80 series in the following, as most of our newly compiled data come from southern Germany
81 and our aim is to construct a regional declination curve from all values adjusted to Munich.

83 *2.2. Gufm1 and its data basis*

84 A global compilation of historical geomagnetic data has been published by Jonkers
85 et al. (2003). It spans the times from 1510 to 1930 and consists mainly of measurements
86 made on ships for navigational purposes during the voyages over the oceans, but also
87 some measurements on land. From that database, we have extracted declination values for
88 the region 47-55°N, 6-15°E, covering the present day German territory and parts of some




Figure 1: Locations of declination data sources. Stars mark locations of found declination values from sundials, compasses and historical maps. Direct measurements at mines and monasteries are marked by black dots. Crosses mark the locations of historical data available from the Jonkers et al. (2003) database (region 47-55°N,6-15°E) and gray diamonds of archeomagnetic data compiled for Germany by Schnepp et al. (2004). Labels (letters) are given for some locations referred to in the text.

Table 1: German geomagnetic observatories with long data series

	location 1	location 2	location 3
Name	Wilhemshaven	Wingst	
IAGA Code	WLH	WNG	
Time interval	1884-1932	since 1939	
Name	Potsdam	Seddin	Niemegk
IAGA Code	POT	SED	NGK
Time interval	1890-1907	1908-1931	since 1932
Name	Munich	Maisach	Fürstenfeldbruck
IAGA Code	MNH	MAS	FUR
Time interval	1841-1926	1927-1935	since 1939

89 surrounding countries. The locations of the resulting 326 declination values from 1523 to
 90 1895 are shown in Fig. 1.

91 Jackson et al. (2000) constructed a global, time-varying magnetic field model spanning
 92 the time 1590 to 1990, named *gufm1*. Predictions of declination for the area of Germany
 93 from the *gufm1* model, in 100 year intervals, are shown in Fig. 2a, complemented by the
 94 declination prediction of the International Geomagnetic Reference Field (IGRF) for 2000
 95 (Macmillan et al., 2003). In Fig. 2b the declination curves from the *gufm1* model for the
 96 four corners of the considered region and its center (51.0°N, 10.5°E) are shown. A strong
 97 change of the declination gradient with time in east-west direction is very clear. Around
 98 1700, declination values all over Germany are on the order of -8°, while around 1900 they
 99 range from -13° (western part) to -9° (eastern part). The epoch without strong spatial
 100 declination gradient around 1700 also marks a change in the isogonic lines over Germany,
 101 from more easterly declination values in the west than the east to more westerly declination
 102 values in the west than the east (see Fig. 2b).

103 2.3. Archeomagnetic data

104 A catalogue of German archaeomagnetic data has been compiled by Schnepf et al.
 105 (2004). Among others, this catalogue contains a consecutive time series of data from a




Figure 2: a) Declination distribution over Germany from 1600 to 2000 according to global models *gufm1* and IGRF. b) Declination predictions from *gufm1* for the center and four corners of the region shown in a).

106 bread oven-floor sequence in Lübeck (Schnepp et al., 2003), spanning approximately the
107 time interval 1300 to 1800. The locations of 51 declination values available since 1300 from
108 that catalogue are also shown in Fig. 1. The mean age uncertainty of these archeomagnetic
109 data is 75 yrs, and the mean declination data uncertainty we computed based on the
110 given α_{95} uncertainties is 3.6° . Both the previous historical and archaeomagnetic data are
111 discussed together with the newly compiled data in section 4.

112 **3. Newly compiled declination data**

113 Reports of direct declination observations are the most reliable sources of information
114 about this magnetic element. However, additional historical sources of declination exist.
115 Here, we describe the different sources we have explored during this study, and discuss
116 their accuracy and reliability. The new data are supplied in the supplemental data file.
117 The locations of the newly compiled declination data are shown in Fig. 1.

118 *3.1. Sundials and compasses*

119 Portable sundials have to be oriented in the right direction in order to show the correct
120 time. The easiest way to achieve this is by means of a magnetic compass. The manu-
121 facturers of sundials obviously were aware of the deviation of the compass needle from
122 the true north since the 15th century (Hellmann, 1899; Zinner, 1939). Ancient sundials
123 or compasses often have a mark of declination angle (Körber, 1965), and this information
124 can generally be found in books and catalogues describing collections of historical sundials
125 (e.g. Körber, 1964; Gouk, 1988; Wagner, 1997). The southern German cities Nuremberg
126 and Augsburg (N and A in Fig. 1) were centers of sundial manufacturing from the 15th to
127 18th century. For many of the ancient sundials an indication about the place and the epoch
128 when they were made is included in the descriptions. Wagner (1997) compiled a list of
129 declination values given by well-dated historical sundials mainly produced at Nuremberg.

130 Wagner (1997) only included data from instruments of which the exact years of man-
131 ufacturing were known. Here, we also include values from instruments that were only
132 approximately dated. The assigned age error estimates depend on the accuracy of the
133 dating. If a time interval was given we assigned the mean time \pm the difference between

134 mean and interval borders. A dating of, e.g., 16th century became 1550 ± 50 , of , e.g., first
135 half of 16th century became 1525 ± 25 , and datings like “about 1670” became 1670 ± 5 . Note
136 that these are not statistical error estimates and probably not even completely consistent
137 within the dataset due to the many different formulations for the age estimates by different
138 authors. In general, they should be close to the maximum age uncertainties and like that
139 useful when error bars are used to decide on data consistency and reliability.

140 We do not have a clear idea what uncertainty to expect as we do not know by which
141 method exactly and under which conditions the values were determined. In general the
142 values are given as full degrees. Körber (1965) notes that the values might have been given
143 according to a 32- or 36-parts wind rose, where approximately 6° or 5° is half a mark on
144 the rose. We assume that 1° to 3° is a reasonable estimate for the general uncertainties of
145 these data.

146 Appendix A lists the declination values obtained from sundials and compasses, together
147 with the references, and data are plotted in Fig. 3a for comparison with other data. Fig. 4
148 is an enlargement of that panel with error bars when the year of manufacturing is unknown.
149 Interestingly, the ages of the older instruments, in general, are better known than of those
150 from 18th and 19th century. The older declination data with known manufacturing dates are
151 much more consistent than the most recent ones. Unless the declination has not been taken
152 into account properly for those instruments, either several of the estimated ages are not
153 correct, or those sundials have been built for significantly different locations. The number
154 of data from sundials amounts to 133 values from instruments with known locations, plus
155 52 values from unknown locations.

156 3.2. *Historical maps*

157 Printed historical maps occasionally show compass roses with an indication of declina-
158 tion. Assuming that geographic north is at the top of the map, the declination can be read
159 from the orientation of the printed compass needle. The accuracy to which the declina-
160 tion can be estimated is in the order of 1° to 5° and 15 declination values from historical
161 maps have been compiled here. Further values, including earlier ones (Wolkenhauer, 1904,




Figure 3: Declination data for the indicated locations provided by several sources: a) from sundials and compasses b) from historical maps; c) from measurements in monasteries and d) from measurements in mines (due to the close agreement several of the mine data are hardly distinguishable). In c) and d) predictions from the *gufm1* model for the same locations are shown. Abbreviation letters refer to those shown in Fig. 1.




Figure 4: Declination data from sundials and compasses from Fig. 3a with estimated error bars when the year of manufacturing is unknown.

162 1907), certainly exist, but as it will be discussed in the following this kind of data seems
 163 less reliable than the sundial data. Here, we limited our work to the easily verifiable values
 164 published by Körber (1965) and Kleinschmidt (1989). Indeed we found some discrepancies
 165 between the values provided by these two sources and confirmed all values from the atlas
 166 of historical maps (Bachmann, 1941, 1942, 1961). All these values, including differing pub-
 167 lished ones, are listed in Appendix B and plotted in Fig. 3b. All declination values compiled
 168 here come from within one century. The figure clearly shows a large scatter among these
 169 values, indicating a much lower reliability of these data compared to those obtained from
 170 sundials. Apparently the printed compass roses do not represent very exact declination
 171 values or the maps themselves are not oriented accurately to geographic north. The three
 172 strongly westward declination values even suggest that declination has been applied with
 173 the wrong sign to these maps.

174 3.3. Monasteries and meteorological stations

175 During the 18th and 19th centuries, measurements of declination were carried out at
 176 several places in southern Germany and Austria by monks.




Figure 5: Declination data measured at monasteries from Fig. 3 enlarged to show the differences in detail.

177 At Hohenpeissenberg, magnetic declination measurements were initiated together with
 178 meteorological measurements in 1781 by the Societas Meteorologica Palatina, the Academy
 179 of the Palatinate. Declination measurements were conducted three times a day by monks
 180 and local priests until 1839. Annual means from these measurements, published in year-
 181 books (e.g. Hemmer, 1783) and a later compilation by Lamont (1851) have been digitized.

182 Results from magnetic observations carried out at monasteries at Augsburg and Kremsmünster
 183 (Austria) are published in a series of yearbooks, like Stark (1814) and Reslhuber (1857),
 184 respectively. Monthly mean values based on at least three measurements a day exist for
 185 Augsburg from 1813 to 1837, and we computed annual means based on them. The time
 186 series for Kremsmünster consists of single data points between 1740 and 1790. From 1815
 187 on the digitized values are means of several measurements per year and from 1832 several
 188 measurements per month with a declinatorium manufactured by Brander in Augsburg.
 189 This is the same instrument as was used in Hohenpeissenberg and Augsburg. Around 1840
 190 a gradual shift to classic geomagnetic observatory instruments and routines with regular
 191 daily observations started.

192 The locations and time series of these observations are included in Figs. 1 and 3c. Fig. 5
 193 shows the same data on an enlarged scale to visualize the detailed differences between these
 194 data mainly coming from the time interval 1750 to 1850. The data show a good internal
 195 consistency and generally good agreement with the *gufm1* model. The Augsburg data,
 196 however, present somewhat suspect values as they are nearly constant over the whole time

197 span they cover.

198 The overall number of data from these sources amounts to 122 values spanning the time
199 1668 to 1854.

200 3.4. Mining activities

201 Since the 13th century, compasses were also used in mining to determine the direction
202 of the mine legs (Ludwig and Schmidtchen, 1997). Christian Doppler first realized that
203 declination information can be gained either by comparing old mining maps with newer ones
204 or that declination values are given in mining publications (Doppler, 1849). Knothe (1987,
205 1988) compiled several data from different European locations, but only plots and no values
206 are published or have been preserved by the author (Knothe, pers. comm.). Schreyer (1886)
207 compiled a large number of declination measurements carried out for mining purposes in
208 Saxony (at that time Kingdom of Saxony) from 1575 to 1885. A number of measurements
209 in Berlin and Regensburg between 1717 and 1788 are included.

210 For two main mining areas in Freiberg and Clausthal 10 year interval data series from
211 1545 to 1885 are given by Schreyer (1886). All found data within each decade have been
212 reduced to the central epoch by a simple assumption of secular variation and averaged
213 “taking into account their reliability” (Schreyer, 1886), but details of this averaging are
214 not given. A few measurements reported by Schreyer (1886) from Freiberg between 1773
215 and 1790 come from different, insufficiently documented sources and are averages of 9 to
216 24 measurements over the year or individual measurements.

217 Annual data for several locations for the 19th century are best documented. Those
218 values had mostly been published annually in a series of mining calendars, e.g. Königl.
219 Bergakademie zu Freiberg (1850), which have been scanned by the Technical University
220 Bergakademie Freiberg and today are available on a website ([http://www.tu-freiberg.de/ub/el-](http://www.tu-freiberg.de/ub/el-bibl/jb_sachsen/jb_sachsen.html)
221 [bibl/jb_sachsen/jb_sachsen.html](http://www.tu-freiberg.de/ub/el-bibl/jb_sachsen/jb_sachsen.html)). Schreyer (1886) reports that if several observations ex-
222 isted within a year, they have been reduced to the middle of the year by an estimate of
223 secular variation for that time and averaged. Moreover, if the time of observation was
224 given the values had first been reduced to the daily mean by rules determined by Johann

225 von Lamont for Munich. The values are rounded to 0.1'. The individual measurements
226 on which these averages are based were done by means of mine surveyors' compasses and
227 their accuracy is estimated as 10' for the earlier and 3-5' for the later years by Schreyer
228 (1886). Note, that the signs of all these values in our digital supplement are opposite to
229 the ones in the Schreyer (1886) publication, in order to agree with today's convention of
230 positive sign for eastern declination and negative for western.

231 Time series of all these 313 declination measurements are shown in Fig. 3d. A good
232 accuracy of these data with very good agreement to *gufm1* from 1700 onward can be seen.
233 The long data series from Clausthal (Ct in Fig. 1) further west and Freiberg (Fb) further
234 east clearly reflect the change in declination gradient around 1725 again. Assuming that
235 the accuracy of these averaged declination data is nearly equally high throughout the 17th
236 century, these data have the potential to improve details of models like *gufm1*, which fits
237 the earlier data less well.

238 3.5. Church orientations

239 A few publications describe church orientations as a possible source of magnetic decli-
240 nation values. Motivated by historical documents proving the use of a compass for church
241 orientation in 1516, Wehner (1905) studied the deviation from the geographic east of the
242 axes of some 300 churches. He concluded that several churches were oriented by magnetic
243 compass and provides a list of 45 German churches with orientations. However, his claim
244 that most of them were oriented by magnetic compass is based on the assumption that
245 the declination varies strictly periodically. Figure 6 shows the declination values resulting
246 from the deviation of church orientations from the geographic east. The associated ages
247 are the years of the church foundation and error bars of 50 years are shown when the year
248 is not exactly known. Nippold (1916) supports Wehner's conclusion (Wehner, 1905) by
249 direct comparison of church axes deviations and geomagnetic field data, however, for lack
250 of available data, again under the assumption of periodicity. Today we know from archeo-
251 and paleomagnetism that we cannot expect a strictly periodic declination variation. Abra-
252 hamsen (1985) picked up the topic again by studying Romanesque churches in Denmark




Figure 6: Declinations for 45 locations in Germany, as determined from church axes deviations from the geographic east under the assumption that orientation was done by magnetic compass. Black dots are the values presumed to represent magnetic declination by Wehner (1905), stars are the ones he considered not in agreement with orientation by magnetic compass. Archeomagnetic declination data (gray dots, see sec. 2.3) and the *gufm1* model prediction for the center of Germany (gray line) are also shown for comparison.

253 in the 12th century. From a statistical analysis of orientation of more than 500 churches he
 254 concluded that about 25% of all Danish Romanesque churches were oriented by means of
 255 a magnetic compass (Abrahamsen, 1990).

256 The declination values obtained from church orientations published by Wehner (1905)
 257 are based on an a priori assumption about the geomagnetic field. Apart from the fact that
 258 the assumption is most likely wrong, as e.g. a comparison with the archaeomagnetic data
 259 described in section 2.3 (Fig. 6) suggests, the values are not independent and cannot be
 260 used to study past declination. A rigorous statistical analysis of all German churches from
 261 a given epoch might provide true information on past declination, but a complete dataset
 262 is not readily available and such a study is outside the scope of this work. We did not
 263 consider the values from church orientations any further in this work.

264 4. A declination curve for southern Germany

All new and previous historical and archeomagnetic data from the German region, spanning the time interval 1300 to 1950 are combined in Fig. 7a on their original locations. Then the values D have been adjusted to the location $M = (11.57E, 48.13N)$ of Munich (M in Fig. 1) from their original locations $x = (longitude, latitude)$ (Fig. 7b) by using

adjustment values ΔD determined from the *gufm1* model for the respective epoch t :

$$D(M, t) = D(x, t) + \Delta D(t) \quad (1)$$

with

$$\Delta D(t) = Dg(M, t) - Dg(x, t), \quad (2)$$

265 where $Dg(M, t)$ and $Dg(x, t)$ are *gufm1* model predictions at time t for locations M and
 266 x , respectively. For data prior to the validity of the model, i.e. $t < 1590$, the adjustment
 267 values were computed with $t = 1590$. Data from the sundials without known location have
 268 not been considered and are omitted in Fig. 7b and in the following determination of a
 269 smoothed curve. From the historical map data, only the values determined/confirmed by
 270 us have been used. Most of the adjusted data are consistent within 10° , but some outliers
 271 exist among the sundials and maps data.

272 A smooth declination curve for Munich for the time span 1400 to 2000 has been created
 273 by fitting a smoothing spline (Constable and Parker, 1988) to 1105 reduced data. In
 274 order to avoid any artificial end effects of the spline function the archaeomagnetic data
 275 have been considered from 1300 on and the combined annual declination means from the
 276 geomagnetic observatories at Munich-Maisach-Fürstfeldbruck from 1840 to 2000 and
 277 reduced to Munich have been included (147 values). We do not have good uncertainty
 278 estimates for the data and for this reason no weighting has been applied. The knot-point
 279 spacing of the spline function has been set to 25 years. The minimum root mean square
 280 (rms) misfit of this spline curve is 2.89° . The residuals between data and curve are not
 281 normally distributed, but show a symmetric distribution with good fit to the majority of the
 282 data and some significant outliers. In order to avoid an influence from obviously erroneous
 283 data we rejected several outliers. As the residuals are not truly normally distributed and we
 284 have no information on the uncertainty distribution of the data, we somewhat arbitrarily
 285 rejected all data lying further than three standard deviations away from this initial curve
 286 (26 rejected values). We fit a new smoothing spline to the remaining declination values,
 287 which can achieve a new minimum rms misfit of 1.45° . In order to keep the variability
 288 of the whole curve comparable with the recent end, the fit to the observatory values, we




Figure 7: All new (black symbols) data presented in Fig. 3a to d together with previously published historical and archeomagnetic data (gray symbols) on a) their original locations and b) adjusted to Munich (11.57E 48.13N). c) Historical declination curves for Munich (thick black line, see text), London (long dashed line, Malin and Bullard (1981)) and Paris (short dashed line, Alexandrescu et al. (1996)). The data used and rejected for the Munich curve are shown here as gray dots and stars, respectively. The time series from the observatories in Munich, Maisach and Fürstenfeldbruck (adjusted to Munich) is given as thin black line.

289 applied a slight smoothing. Our preferred curve has an rms misfit of 1.48° and is shown as
290 thick solid line in Fig. 7c. The curve is also provided in the supplemental data file.

291 5. Comparison to other declination curves

292 The declination curves by Malin and Bullard (1981) and Alexandrescu et al. (1996) for
293 London and Paris, respectively, are included in Fig. 7c. The agreement in general shape
294 of European declination variation is obvious. The intersection between the Munich and
295 Paris curves confirms again the change from more easterly declination further west (Paris)
296 to more westerly declination there than further east (Munich) in the early 18th century.
297 The new Munich curve shows a somewhat different variability in the 16th and 17th century
298 than the previous curves. Geomagnetic jerks, characterised by a sharp extrema of secular
299 variation like the 1969 event, can be identified as rather broad minimum or maxima in
300 the curve. The well-known 1978 and 1990 geomagnetic jerks, however, are not detected as
301 they occur near the very end of the investigated period. Moreover, a comparison of secular
302 variation in form of the first derivative of the spline function with first differences of the
303 MNH-MAS-FUR data series (Fig. 8a), slightly smoothed by an 11-yr running average, to
304 minimise the solar cycle related variations, shows that for the recent decades the spline
305 function is not able to represent the very short time-scale changes, recently defined as
306 “rapid secular variation fluctuations” by Mandeia and Olsen (2009).

307 The occurrence time of maxima and minima of the curve, representing geomagnetic
308 jerks, can be determined more easily by looking for changes of sign in the second derivative,
309 secular acceleration, which is also shown in Fig. 8a. Note that the sharp angles in this
310 secular acceleration curve are a consequence of the 25 year knot-point spacing of the cubic
311 spline basis functions, but the zero crossings accurately represent the times where maxima
312 and minima occur in the secular variation curve. Indeed, the sign changes of the Munich
313 secular acceleration around 1932, 1889 and 1861 roughly agree with the known geomagnetic
314 jerks in 1925, 1901 and 1870 described by Alexandrescu et al. (1995, 1997). The comparison
315 between the observatory data and the smoothed historical curve around 1925 suggests
316 that the uncertainty in dating these jerks lies in the smoothed representation and we

317 consequently should adopt an uncertainty of about ± 10 yrs for all earlier events shown by
318 our curve. The new Munich curve suggests events at 1763, 1741, 1708, 1693, 1661, 1558,
319 1508, 1448 and 1410, although the two earliest have to be regarded with caution because
320 they are constrained only by few data. Two more geomagnetic jerks might have occurred
321 around 1598 and 1603, but are less clearly resolved.

322 Alexandrescu et al. (1997) describe five possible jerks between 1680 and 1870, based
323 on their study of the Paris data series. They are dated 1700, 1730, 1750, 1770 and 1785.
324 Considering the data noise and some baseline problems for the first two centuries of the
325 Paris curve, Alexandrescu et al. (1997) conclude that none of these jerks is deeply sup-
326 ported by their data and the presence of any geomagnetic jerk between 1680 and 1870
327 could be doubted. Comparing their and our geomagnetic jerk occurrence times, the sign
328 change of secular acceleration (maximum or minimum of secular variation), and taking into
329 account the significant temporal uncertainties, the events dated 1700/1708, 1730/1741 and
330 1750/1763 could represent the same events. On the other hand, our compilation identifies
331 no events in 1770 and 1785.

332 The identification of geomagnetic jerks by Alexandrescu et al. (1997) was carried out on
333 a significantly less smoothed data series. We applied a similar spline fit to the Paris data
334 series for a more direct comparison to our analysis. We used the annual data presented by
335 Alexandrescu et al. (1996), which go back to 1541, and added 26 archeomagnetic declination
336 values from France for the time span 1300 to 1540, mainly compiled by Bucur (1994) and
337 digitally available from the global compilation by Korte et al. (2005) or the GEOMAGIA
338 V.2 database (<http://www.geomagia.ucsd.edu>). The declination series by Alexandrescu
339 et al. (1996) is adjusted to the location of Chambon-la-Forêt (2.27E, 48.02N), the location
340 of the present observatory near Paris. We adjusted the archeomagnetic data to the same
341 location using the *gufm1* model as described for the German data in section 4. The data
342 were fit by smoothing splines for the same time interval (1300 to 2000) and with the same
343 knot-point spacing (25 yrs) as the Munich curve. A comparable variability of the curve was
344 obtained by applying weak smoothing and fitting the data to an rms misfit of 0.69° . Secular
345 variation and acceleration at Paris are represented by the first and second derivative of this

346 curve in Fig. 8b.

347 The comparison of the smoothed Munich and Paris secular variation is best described
348 in three time intervals: 1400 – 1580, 1580 – 1770 and 1770 – 2000. During the recent time
349 interval, from about 1770 onwards, secular variation and acceleration at both locations
350 broadly agree, with the jerk seen in 1901 in the original French data also represented slightly
351 earlier by the smooth Paris curve, but agreeing within our estimated dating uncertainty of
352 ± 10 yrs due to the smoothing. The earliest time interval from 1400 to about 1580, least well
353 supported by data particularly in the Paris curve, also shows surprisingly consistent secular
354 variation and acceleration between the smoothed curves from the two locations. Larger
355 time lags between similar patterns in this case might be influenced by relatively large
356 dating uncertainties associated with the archeomagnetic data, but artificial oscillations in
357 the spline fit resulting from the sparse data until 1550 can also not be ruled out.

358 Significant differences between the two locations are seen in the time interval between
359 1580 and 1770. The geomagnetic jerks suggested between 1580 and 1680 by the Munich
360 curve are not confirmed by the Paris curve. During this interval, however, the Paris
361 curve might lack some information as it is based only on 25 annual values derived from
362 35 individual measurements, while 128 data points support the Munich curve within this
363 interval. After 1680, when a reasonable amount of data exists for both locations, the
364 smoothed Paris curve only shows one of the five jerks suggested by Alexandrescu et al.
365 (1997) based on the original data (at 1700). However, no deceleration (i. e. secular
366 acceleration $\dot{j} < 0$) appears in the smoothed curve for nearly two centuries after that event.
367 The existence and tentative link between suggested jerks about 1700/1708, 1730/1741 and
368 1750/1763 thus neither can be confirmed nor excluded by this comparison. Both analyses
369 agree, however, that the century from about 1765 to about 1865 is devoid of strong rapid
370 secular variation changes. This is the time when the declination in Europe has reached its
371 most western values and changes its trend, i.e. the minimum seen in the declination curve
372 (Fig. 7c).




Figure 8: Secular variation and acceleration for Munich (a) and Paris (b). Secular variation is given by the first temporal derivative of the smoothed curves fit to the declination data (black) and by first differences of the data series smoothed by 11-yr running averages (gray, MNH-MAS-FUR (a) and of the declination data from Alexandrescu et al. (1996) (b), respectively). Secular acceleration (dashed lines) of the declination curves is shown with right-side label axes.

373 6. Conclusions

374 We have compiled 635 historical declination values from southern Germany and sur-
375 rounding areas from different sources. The accuracy of declination values determined from
376 sundials and old compasses from the 15th to 19th century are in the same order as that of
377 the available archeomagnetic data. Measurements in mines and those made by monks from
378 the 17th to 19th century show an accuracy better than 1°. All these data can be useful to
379 improve historical geomagnetic field models and to link archeomagnetic and historical field
380 reconstructions. Declination values obtained from church orientations, however, have to be
381 taken with caution. They require more comprehensive statistical investigations than cur-
382 rently available for Germany in order to be considered as a source of historical declination
383 information.

384 The compiled data have been adjusted to the location of Munich together with available
385 archeomagnetic and previously published historical data. A smooth declination curve has
386 been fit to the data, extending the existing observatory record from MNH-MAS-FUR
387 backward to AD 1400. The comparison to declination curves for Paris and London shows
388 a broadly uniform European declination variation, but with a significant spatial gradient
389 change in the early 18th century.

390 The smooth secular variation description provided by the Munich declination curve
391 indicates several geomagnetic jerks with an uncertainty of about ± 10 years, as can be
392 estimated for the well-known jerks of the 20th century. The geomagnetic jerk shown by
393 the Munich curve around 1861 is presumably the one proposed around 1870 based on data
394 from Paris and four other European locations by Alexandrescu et al. (1997). Three jerks
395 suggested by Alexandrescu et al. (1997) from the Paris data over the 18th century might
396 be confirmed by this new compilation: 1750/1763, 1730/1741 and 1700/1708 (first date is
397 based on the Paris original data, the second one is based on the smoothed Munich curve).
398 However, the two more recent events are not seen in the Paris curve smoothed in the same
399 way as the Munich one. Going back in time, several earlier events are suggested by the
400 Munich curve around the following epochs: 1693, 1661, 1558, 1508, and perhaps, but less

401 clearly resolved or supported by data, in 1603, 1598, 1448 and 1410. An analysis of the
402 similarly smoothed, sparse Paris data over the 17th and 16th century, however, suggests
403 significantly different secular variation and acceleration during these times and does not
404 support the suggested jerks.

405 More data are necessary and regional or global modelling might help to resolve whether
406 true small-scale field structure or insufficient data cause the observed differences. Note also
407 that jerks in relatively quick succession, like e.g. the 1979 and 1990 events, can in general
408 not be resolved by a smoothed declination reconstruction. In summary, our study suggests
409 that geomagnetic jerks, as defined by Manda and Olsen (2009), occurred more or less
410 regularly on a decadal time scale (from some three to six decades) during most of the
411 studied six centuries. However, the time span from about 1760 to 1860 seems to have been
412 devoid of sudden secular variation changes.

413 We expect that this new data compilation will be useful to improve historical geomag-
414 netic field models and to better track the different temporal variations revealed by the
415 Earth's magnetic field. We also hope that our work will encourage the search for unknown
416 ancient geomagnetic field data from all around the world.

417 **Acknowledgements**

418 We thank Klaus Eichholz for information on sundials and historical compass scales, and
419 Nils Olsen for drawing our attention to the work of Nils Abrahamsen. We thank P. Amand
420 Kraml (Kremsmünster) and P. Gregor Helms (Augsburg) for support in their monasteries
421 and Peter Winkler for support at Hohenpeissenberg and helpful discussions. We particu-
422 larly acknowledge efforts by Hannelore Podewski, Francesca Overbeck and Martin Feller for
423 documenting and digitizing all the data from the literature. We also thank two anonymous
424 reviewers for constructive comments which helped us to improve the original manuscript.

425 A. Declination from sundials and compasses

426

[h]

Year	ΔT	D (deg.)	Location	Manufacturer	Reference	Ref. No.	Given Age
1451	0	10.0	Vienna	G. Peuerbach	Wagner (1997), Zinner (1979)		
1451	0	11.0	Vienna	G. Peuerbach	Wagner (1997), Zinner (1979)		
1453	0	14.0	Vienna	G. Peuerbach	Wagner (1997), Zinner (1979)		
1455	0	10.0	Vienna	G. Peuerbach	Zinner (1979)	4525	
1456	0	12.5	Vienna	G. Peuerbach	Wagner (1997), Zinner (1979)		
1463	0	13.0	Budapest	J. Regiomontian	Zinner (1979)	122	
1465	0	6.0	Nuremberg	J. Regiomontian	Wagner (1997), Zinner (1979)		
1466	2	6.0	Nuremberg	J. Regiomontian	Zinner (1979)	WI 7	1464-1467
1466	2	10.0	Nuremberg	J. Regiomontian	Zinner (1979)	F 1361	1464-1467
1476	0	10.0	(Budapest)	C. Dorn	Zinner (1979)	12364	
1479	0	14.0	(Budapest)	C. Dorn	Zinner (1979)	288	
1480	0	10.0	(Budapest)	C. Dorn	Zinner (1979)		
1481	0	14.0	(Budapest)	C. Dorn	Zinner (1979)		
1483	3	10.0	(Budapest)	C. Dorn	Zinner (1979)	G 425	
1484	2	14.0	unknown	W. Faber	Zinner (1979)		1480-1486
1485	5	7.0	(Budapest)	C. Dorn	Zinner (1979)	1893	presum. 1484
1486	0	12.0	(Budapest)	C. Dorn	Zinner (1979)		1480-1490
1491	0	11.0	Vienna	C. Dorn	Zinner (1979)	94	
1501	0	10.0	Nuremberg	E. Eitzlaub	Wagner (1997), Gouk (1988)		
1511	0	15.0	Nuremberg	E. Eitzlaub	Zinner (1979)	WI 28	
1514	0	10.0	unknown	unknown	Körber (1964)	DI 65	
1525	25	7.0	Kassel	B. Emck	Hamel (2000), Mackensen	Mat U 24	1 st half 16. th cent.
1527	0	10.5	Nuremberg	G. Hartmann	Zinner (1979)		
1535	0	10.5	Nuremberg	G. Hartmann	Wagner (1997), Gouk (1988)		
1538	0	11.5	Nuremberg	G. Hartmann	Wagner (1997), Gouk (1988)		
1539	0	10.25	Nuremberg	G. Hartmann	Zinner (1979)		
1539	0	10.25	Nuremberg	G. Hartmann	Zinner (1979)	Rar 434	
1542	0	9.5	Nuremberg	G. Hartmann	Wagner (1997), Gouk (1988)	WI 182	
1555	0	12.0	Nuremberg	G. Reimann	Zinner (1979)	WI 267	
1556	0	7.0	Nuremberg	J. Gebhard	Wagner (1997), Bryden (1988)	1681	
1557	0	10.0	Augsburg	C. Schissler	Wagner (1997)		
1558	0	9.5	Nuremberg	H. Reimann	Wagner (1997), Syndram (1989)	N. 12	
1559	0	9.0	Augsburg	C. Schissler	Wagner (1997), Zinner (1979)		
1560	10	5.0	Dresden	Göbe	Kleinschmidt (1989), Körber (1964)	DI 63	about 1560
1561	0	8.0	unknown	unknown	Hamel (2000), Zinner (1979)	MATB75	
1561	0	12.0	unknown	unknown	Wagner (1997), Körber (1964)	DI 12	
1562	0	10.0	Augsburg	C. Schissler	Wagner (1997), Körber (1964)	DI 37	
1565	0	5.0	Augsburg	(C. Schissler)	Wagner (1997), Körber (1964)	DI 22	
1565	0	25.0	(Nuremberg)	C. Heiden	Hamel (2000), Zinner (1979)	MAT B 29	
1567	0	8.5	Augsburg	C. Schissler	Wagner (1997), Zinner (1979)		
1569	0	6.5	Augsburg	C. Heiden	Wagner (1997), Zinner (1979)		
1574	0	8.0	Nuremberg	H. Tucher	Wagner (1997), Zinner (1979)		

Year	ΔT	D (deg.)	Location	Manufacturer	Reference	Ref. No.	Given Age
1579	0	8.0	Nuremberg	H. Tucher	Wagner (1997), Körber (1964)	1	
1580	0	9.5	Nuremberg	H. Ducher	Wagner (1997), Syndram (1989)	N. 14	
1581	0	9.13	Augsburg	T. Klieber	Zinner (1979)		
1582	0	15.0	(Kassel)	J. Bürgi	Zinner (1979)		
1583	0	8.0	Nuremberg	P. Reimann	Wagner (1997), Zinner (1979)		
1584	0	11.5	Nuremberg	P. Reimann	Wagner (1997), Zinner (1979)		
1595	0	7.0	Nuremberg	H. Tucher	Wagner (1997), Lloyd (1992)		
1598	0	6.0	(Antwerpen)	unknown	Wagner (1997), Syndram (1989)	Nr. 17	
1600	0	8.0	Bamberg	J. Bonus	Glasmann (1999)	Nr. 53	
1600	0	8.0	Nuremberg	H. Troschel	Wagner (1997), Lloyd (1992)		
1600	5	8.0	unknown	"R"	Hamel (2000), Zinner (1979)	MAT B 72	about 1600
1601	0	6.0	Nuremberg	Troschel	Wagner (1997), Körber (1964)	2	
1602	0	6.5	Nuremberg	P. Reimann	Wagner (1997), Zinner (1979)		
1604	2	6.0	Nuremberg	unknown	Kleinschmidt (1989), Körber (1964)		
1605	2	9.0	Dresden	unknown	Kleinschmidt (1989), Körber (1964)	AI 44	after 1602 after 1603
1607	0	12.0	Nuremberg	P. Reimann	Wagner (1997), Zinner (1979)		
1610	0	4.0	Nuremberg	Lösel	Wagner (1997), Körber (1964)	4	
1610	0	-20.0	(Augsburg)	L. Miller	Zinner (1979)		
1611	0	5.0	Nuremberg	H. Troschel	Wagner (1997), Lloyd (1992)	7534	
1611	0	7.0	Nuremberg	H. Tucher	Wagner (1997), Lloyd (1992)	7577	
1613	0	-6.0	Kassel	J. Bürgi	Zinner (1979)		
1613	0	2.0	Nuremberg	L. Miller	Wagner (1997), Gouk (1988)	1684	
1613	0	5.0	Nuremberg	L. Miller	Wagner (1997), Lloyd (1992)	7459	
1614	0	6.5	Nuremberg	H. Tucher	Wagner (1997)	H 5820	
1616	0	7.0	Nuremberg	L. Miller	Wagner (1997), Lloyd (1992)	7565	
1619	0	0.0	Nuremberg	G. Karner	Wagner (1997), Zinner (1979)		
1620	0	4.0	Nuremberg	C. Karner	Wagner (1997), Lloyd (1992)	7542	
1620	0	5.5	Nuremberg	C. Trechsler	Wagner (1997), Syndram (1989)	N. 37	
1622	0	5.0	Nuremberg	C. Karner	Wagner (1997), Lloyd (1992)	7552	
1624	0	7.0	Nuremberg	H. Tucher	Wagner (1997), Zinner (1979)		
1625	0	6.0	Nuremberg	L. Miller	Wagner (1997)		
1625	0	6.0	Nuremberg	L. Miller	Wagner (1997), Zinner (1979)		
1626	0	5.0	Nuremberg	H. Troschel	Wagner (1997), Lloyd (1992)	7458	
1626	0	7.0	Nuremberg	C. Karner	Hamel (2000), Zinner (1979), Gouk (1988)		
1626	0	10.0	Nuremberg	H. Troschel	Hamel (2000), Zinner (1979), Gouk (1988)		
1626	0	22.0	changed	C. Karner	Hamel (2000), Zinner (1979), Gouk (1988)		
1629	0	4.0	Nuremberg	L. Miller	Wagner (1997), Lloyd (1992)	7560	
1630	0	3.0	Nuremberg	C. Karner	Wagner (1997), Lloyd (1992)	7553	
1630	0	4.0	Nuremberg	C. Karner	Wagner (1997), Lloyd (1992)	7554	
1630	0	7.0	Nuremberg	L. Miller	Wagner (1997), Lloyd (1992)	7567	
1634	0	6.0	Augsburg	L. Miller	Wagner (1997), Körber (1964)	5	
1636	0	4.0	Nuremberg	J. Karner	Wagner (1997), Lloyd (1992)	7550	
1636	0	4.0	Nuremberg	L. Miller	Wagner (1997), Syndram (1989)	N. 19	
1636	0	5.0	Nuremberg	L. Miller	Wagner (1997), Lloyd (1992)	7568	
1637	0	8.0	Nuremberg	L. Miller	Hamel (2000), Gouk (1988), Zinner (1979)		
1637	0	12.0	changed	L. Miller	Hamel (2000), Gouk (1988), Zinner (1979)		

Year	ΔT	D (deg.)	Location	Manufacturer	Reference	Ref. No.	Given Age
1639	0	2.0	Nuremberg	J. Karner	Wagner (1997), Gouk (1988)	634	
1640	0	0.0	Nuremberg	L. Miller	Wagner (1997), Gouk (1988)	1684	
1640	0	4.0	Nuremberg	L. Miller	Wagner (1997), Lloyd (1992)	7562	
1640	0	5.0	Nuremberg	L. Miller	Wagner (1997), Gouk (1988)	1683	
1642	0	2.0	South Ger.	L. Hartmann	Wagner (1997)		
1644	0	1.0	Nuremberg	L. Miller	Wagner (1997), Lloyd (1992)	7563	
1646	0	0.0	Nuremberg	L. Miller	Wagner (1997), Gouk (1988)	184	
1649	0	3.0	Nuremberg	N. Miller	Wagner (1997), Lloyd (1992)	1770	
1649	0	3.5	Nuremberg	N. Miller	Wagner (1997), Syndram (1989)	N. 21	
1649	0	-5.0	Nuremberg	N. Miller	Wagner (1997), Gouk (1988)	1686	
1650	5	-28.0	unknown	unknown	Hamel (2000)	MAT B 81	about 1650
1650	50	-10.0	unknown	unknown	Körber (1964)	7	ca. 17 th cent.
1650	50	-11.0	unknown	unknown	Körber (1964)	compass	ca. 17 th cent.
1652	0	0.0	Nuremberg	L. Miller	Wagner (1997), D.M.Mnchen	69503	
1652	0	5.0	Nuremberg	A. Karner	Wagner (1997), Lloyd (1992)	7540	
1661	0	0.0	Nuremberg	N. Miller	Wagner (1997), Zinner (1979)		
1666	13	-17.0	unknown	J. Koch	Zinner (1979)	1880-36	
1673	0	-7.0	Nuremberg	unknown	Wagner (1997), Zinner (1979)		
1680	10	-22.0	Augsburg	J. Martin	Körber (1964)	11	approx. after 1670
1689	0	-17.0	(Kassel)	J.W. Schulze	Hamel (2000), Zinner (1979)	MATB23	
1694	0	-5.5	Nuremberg	M. Karner	Wagner (1997)		
1695	5	-17.0	Cologne	S. Krigner	Hamel (2000), Zinner (1979)	MAT 1997-6	1690-1700
1699	0	-11.0	unknown	Richardus	Wagner (1997), Körber (1964)	26	
1700	0	-5.0	Nuremberg	G. Karner	Wagner (1997), Lloyd (1992)	7525	
1700	5	-7.0	Augsburg	J. Martin	Hamel (2000), Zinner (1979)	MAT B 25	about 1700
1700	5	-7.0	Augsburg	J. Martin	Hamel (2000), Zinner (1979)	MAT	about 1700
1700	5	-7.5	Augsburg	J. Martin	Hamel (2000), Zinner (1979)	MAT B 62	about 1700
1700	5	-15.0	changed	J. Martin	Hamel (2000), Zinner (1979)	MAT B 62	about 1700
1700	5	-21.0	unknown	(Kriger)	Körber (1964)	DI 80	
1750	50	-11.0	Augsburg	Höldrich	Körber (1964)	DI 64	
1750	50	-13.0	unknown	unknown	Körber (1964)	DI 88	
1750	50	-13.0	Augsburg	A. Braunmüller	Zinner (1979)	6625	
1710	10	-10.0	Augsburg	J. Willebrand	Glasemann (1999)		beg. 18 th cent.
1710	10	-15.0	German	unknown	Glasemann (1999)	Nr. 38	early 18 th cent.
1710	10	-15.0	unknown	H.G. Welligen	Hamel (2000), Hausmann	MAT A 27	beg. 18 th cent.
1710	10	-17.0	German	unknown	Glasemann (1999)	Nr. 37	early 18 th cent.
1720	5	-5.0	Augsburg	J. Willebrand	Hamel (2000), Zinner (1979)	MAT B 49	about 1720
1720	10	-9.0	Augsburg	J. Willebrand	Körber (1964)	DI 94	about 1720
1725	25	-9.0	unknown	unknown	Hamel (2000)	MAT B 21	1st half 18 th cent.
1725	25	-12.0	(Hessen)	unknown	??	MAT B 26	1st half 18 th cent.
1725	25	-20.0	Nuremberg	L.A. Karner	Hamel (2000), Gouk (1988), Zinner (1979)	MAT B 78	1st half 18 th cent.
1737	12	-17.0	unknown	unknown	Glasemann (1999)	Nr. 25	2 nd quarter 18 th cent.
1750	0	-14.0	Augsburg	L.T. Müller	Hamel (2000), Zinner (1979)	MATB59	
1750	0	-17.0	Augsburg	L.T. Müller	Hamel (2000), Zinner (1979)	MATB84	
1750	5	-15.0	unknown	unknown	Hamel (2000)	MAT B 61	about 1750
1750	10	-5.0	unknown	unknown	Hamel (2000)	MAT B 82	middle 18 th cent.

Year	ΔT	D (deg.)	Location	Manufacturer	Reference	Ref. No.	Given Age
1750	10	-12.0	Augsburg	J.P. Bihler	Syndram (1989)	H-W 70	middle 18 th cent.
1750	50	-5.0	unknown	"B"	Körber (1964)	19	about 18 th cent.
1750	50	-10.0	unknown	unknown	Hamel (2000)	MAT B 32	18 th cent.
1750	50	-15.0	unknown	unknown	Körber (1964)	49 compass	about 18 th cent.
1750	50	-17.0	Augsburg	A. Vogler	Hamel (2000), Zimmer (1979)	MAT B 87	18 th cent.
1750	50	-19.0	unknown	Pfersich	Körber (1964)	12	about 18 th cent.
1750	50	-20.0	unknown	unknown	Körber (1964)	17	about 18 th cent.
1750	50	-22.0	unknown	"K"	Körber (1964)	23	about 18 th cent.
1751	0	5.0	Lüttich	Vinon	Körber (1964)	compass	about 18 th cent.
1752	0	-22.0	Nuremberg	L.A. Karner	Wagner (1997)	22	about 1760
1758	0	-22.0	unknown	A.F.	Wagner (1997), Körber (1964)		ca. 1760
1760	5	-15.0	Augsburg	L.T. Miller	Glasemann (1999) Körber (1964)	Nr. 32	about 1760
1760	5	-15.0	Augsburg	L.T. Miller	Glasemann (1999) Körber (1964)	DI 94	about 1760
1760	5	-15.0	Nuremberg	D. Beringer	Kleinschmidt (1989), Körber (1964)	DI 7	about 1760
1760	5	-15.0	Reinharz	J.G. Zimmer	Kleinschmidt (1989), Körber (1964)	DI 96	about 1760
1760	5	-20.0	Augsburg	J.G. Vogler	Kleinschmidt (1989), Körber (1964)	80/239	
1769	0	-17.0	unknown	unknown	Wagner (1997), D.M.Mnchen	DI 89	about 1775
1775	5	-16.0	Augsburg	L. Grassl	Kleinschmidt (1989), Körber (1964)		2 nd half 18 th cent.
1775	25	-13.0	Augsburg	A. Vogler	Glasemann (1999)	Nr. 30	2 nd half 18 th cent.
1775	25	-15.0	Augsburg	L. Grassl	Glasemann (1999)	Nr. 33	2 nd half 18 th cent.
1775	25	-15.0	German	unknown	Glasemann (1999)	Nr. 41	2 nd half 18 th cent.
1775	25	-20.0	German	unknown	Glasemann (1999)	Nr. 40	2 nd half 18 th cent.
1777	2	-17.0	Nuremberg	D. Beringer	Hamel (2000), Zimmer (1979)	MAT B 93	1775-1780
1780	15	-18.0	German	unknown	Syndram (1989)	H-W	last third 18 th cent.
1787	12	-17.0	Ansbach	K.C. Keller	Glasemann (1999)	Nr. 35	last quarter 18 th cent.
1790	10	-17.0	Augsburg	J.N. Hölderich	Hamel (2000), Zimmer (1979)	MAT B 63	end 18 th cent.
1790	10	-20.0	Nuremberg	D. Beringer	Hamel (2000), Zimmer (1979)	MAT B 110	end 18 th cent.
1790	10	-20.0	Nuremberg	D. Beringer	Hamel (2000), Zimmer (1979)	MAT B 67	end 18 th cent.
1790	10	-20.0	Nuremberg	J.B. Bauer	Glasemann (1999)	Nr. 81	late 18 th cent.
1790	10	-20.0	Nuremberg	P.P. Beringer	Hamel (2000), Zimmer (1979)	MAT B 113	end 18 th cent.
1790	10	-20.0	Nuremberg	P.P. Beringer	Hamel (2000), Zimmer (1979)	MAT B 68	end 18 th cent.
1795	5	-20.0	Augsburg	J. Schretteger	Kleinschmidt (1989), Körber (1964)	DI 91	ca. after 1790
1795	5	-22.0	Augsburg	J. Schretteger	Kleinschmidt (1989), Körber (1964)	DI 95	ca. after 1790
1798	0	-15.0	unknown	I.C.R.	Wagner (1997), Körber (1964)		
1800	5	-7.0	unknown	unknown	Hamel (2000)	MAT B 3	about 1800
1800	5	-7.0	unknown	unknown	Hamel (2000)	MAT B 6	about 1800
1800	5	-17.0	unknown	unknown	Hamel (2000)	MAT B 65	about 1800
1800	5	-20.0	unknown	unknown	Hamel (2000)	MAT B 71	about 1800
1800	5	-22.0	unknown	unknown	Hamel (2000)	MAT B 69	about 1800
1800	5	-22.0	unknown	unknown	Hamel (2000), Zimmer (1979)	MAT B 70	about 1800
1800	10	-7.0	Cologne	E. Schmaldt	Hamel (2000), Zimmer (1979)	MAT B 88	about 1800
1800	10	-20.0	Fürth	Stockkert	Hamel (2000), Zimmer (1979)	MAT B 66	about 1800
1800	50	-11.0	unknown	unknown	Körber (1964)	24	about 18/19 th cent.
1800	50	-12.0	unknown	unknown	Körber (1964)	DI 85	18/19 th cent.
1800	50	-13.0	unknown	unknown	Körber (1964)	DI 86	18/19 th cent.

Year	ΔT	D (deg.)	Location	Manufacturer	Reference	Ref. No.	Given Age
1810	10	-14.0	unknown	unknown	Hamel (2000)	MAT B 14	begin. 19 th cent.
1820	5	-18.0	unknown	unknown	Kleinschmidt (1989), Körber (1964)	DI 82	about 1820
1850	10	-20.0	German	unknown	Glasemann (1999)	Nr. 35	middle 19 th cent.
1850	50	-7.0	unknown	unknown	Hamel (2000)	MAT B 16	19 th cent.
1850	50	-22.0	unknown	unknown	Körber (1964)	25	about 19 th cent.
1850	50	-10.0	unknown	unknown	Körber (1965)	DI 77	19 th cent.

427 **B. Declination from maps**

428 Differing declination estimates by different authors are given in one line of the table.
 429 All values with references Bachmann (1941, 1942, 1961) are the values taken from these
 430 reproductions of the original maps by us.

Year	D (deg.)	Location	Reference
1566	11	Zürich	Körber (1965)
1599	6 / 11 / 8	Nuremberg	Körber (1965) / Kleinschmidt (1989) / Bachmann (1942)
1599	6	Munich	Wagner (1997)
1602	11 / 10 / 12	Bamberg	Körber (1965) / Kleinschmidt (1989) / Bachmann (1942)
1603	16	Konstanz	Kleinschmidt (1989) (Bodenseekarte by J.G. Tibian)
1613	8 / 11 / 11	Munich	Körber (1965) / Kleinschmidt (1989) / Bachmann (1942)
1614	6 / 9	Landshut	Körber (1965) / Bachmann (1942)
1619	25	Thierhaupten	Kleinschmidt (1989), Bachmann (1942)
1619	-25	Donauwörth	Kleinschmidt (1989), Bachmann (1942)
1628	13.5	Bunde	Kleinschmidt (1989) (map by J. Sems)
1633	22 / 29 / 22	Bamberg	Körber (1965) / Kleinschmidt (1989) / Bachmann (1942)
1643	-29 / -21	Wolfegg	Kleinschmidt (1989) / Bachmann (1961)
1643	10	Leutkirchen	Kleinschmidt (1989), Bachmann (1961)
1643	26	Giengen	Kleinschmidt (1989), Bachmann (1961)
1657	11	Minden	Körber (1965), Bachmann (1941)

431 **References**

- 432 Abrahamsen, N., 1985. Romanske kirkers orientering og den magnetiske misvisning i 11-
 433 tallet i Danmark, Orientation of Romanesque churches and magnetic declination in the
 434 12th century in denmark. GeoSkrifter 23, Geologisk Institut, Aarhus Universitet.
- 435 Abrahamsen, N., May 1990. Orientation of Romanesque churches in Denmark suggest
 436 common use of magnetic compass in the 12th century, unpublished manuscript.
- 437 Alexandrescu, M., Courtillot, V., LeMouël, J.-L., 1996. Geomagnetic field direction in
 438 Paris since the mid-sixteenth century. Phys. Earth. Planet. Inter. 98, 321–360.
- 439 Alexandrescu, M., Courtillot, V., LeMouël, J.-L., 1997. High resolution secular variation
 440 of the geomagnetic field in western Europe over the last 4 centuries: Comparison and
 441 integration of historical data from paris and london. J. Geophys. Res. 102, 20,245–20,258.

- 442 Alexandrescu, M., Gibert, D., Hulot, G., Mouël, J.-L. L., Saracco, G., 1995. Detection of
443 geomagnetic jerks using wavelet analysis. *J. Geophys. Res.* 100, 12557–12572.
- 444 Bachmann, F. (Ed.), 1941. *Die alte deutsche Stadt. Ein Bilderatlas der Städteansichten*
445 *bis zum Ende des 30jährigen Kriegs. Bd. I: Der Nordwesten.* Hiersemann, Leipzig.
- 446 Bachmann, F. (Ed.), 1942. *Die alte deutsche Stadt. Ein Bilderatlas der Städteansichten*
447 *bis zum Ende des 30jährigen Kriegs. Bd. II: Der Südosten. Teil 1: Bayern.* Hiersemann,
448 Leipzig.
- 449 Bachmann, F., 1961. *Die alte deutsche Stadt. Bd. IV: Baden-Württemberg.* Hiersemann,
450 Stuttgart.
- 451 Balmer, H., 1965. Beiträge zur Geschichte der Erkenntnis des Erdmagnetismus. Vol. 20 of
452 *Veröffentl. Schw. Ges. f. Gesch. d. Medizin u. Naturwissensch.* Aarau.
- 453 Barraclough, D., 1995. Observations of the Earth's magnetic field in Edinburgh, from 1670
454 to the present day. *Tran. R. Soc. Edinburgh Earth Sci.* 85, 239–252.
- 455 Bryden, D., 1988. *Sundials and Related Instruments.* Whipple Museum Publications, Cam-
456 bridge.
- 457 Bucur, I., 1994. The direction of the terrestrial magnetic field in France during the last 21
458 centuries. *Recent progress. Phys. Earth. Planet. Interiors* 67, 95–109.
- 459 Cafarella, L., DeSantis, A., Meloni, A., 1992. Secular variation in Italy from historical
460 geomagnetic field measurements. *Phys. Earth. Planet. Inter.* 73, 206–221.
- 461 Chapman, S., Bartels, J., 1962. *Geomagnetism.* Oxford University Press, London.
- 462 Constable, C. G., Parker, R. L., 1988. Smoothing, splines and smoothing splines: Their
463 application in geomagnetism. *J. Comput. Phys.* 78, 493–508.

- 464 Doppler, C., 1849. über eine bisher unbenützte Quelle magnetischer Declinations-
465 Beobachtungen. Sitzungsberichte math.-nat. Classe, k. Akademie d. Wissenschaften,
466 Wien 2, 249–261.
- 467 Encke, J., 1840, 1844, 1848, 1857, 1884. Astronomische Beobachtungen auf der Königlichen
468 Sternwarte zu Berlin. vol. 1 to 5, Berlin.
- 469 Gauss, C., 1833. Intensitas vis magneticae terrestris ad mensuram absolutam revocata.
470 Göttingische Gelehrte Anzeigen 205, 2041–2058.
- 471 Glasemann, R., 1999. Erde, Sonne, Mond & Sterne. Globen, Sonnenuhren und astronomis-
472 che Instrumente im Historischen Museum Frankfurt am Main. Schriften des Historischen
473 Museums, Frankfurt/M.
- 474 Gouk, P., 1988. The Ivory Sundials of Nuremberg 1500-1700. Cambridge.
- 475 Hamel, J., 2000. Die Sonnenuhren des Museums für Astronomie und Technikgeschichte
476 Kassel - Bestandskatalog. Thun, Frankfurt am Main.
- 477 Hellmann, G., 1899. The beginning of magnetic observations. Terr. Magn. Atmos. Elect.
478 IV, 73–86.
- 479 Hemmer, J., 1783. 1783 Ephemerides Societatis Meteorologicae Palatinae. Historia et Ob-
480 servationes 1781. Societatis Meteorologicae Palatinae.
- 481 Jackson, A., Jonkers, A. R. T., Walker, M. R., 2000. Four centuries of geomagnetic secular
482 variation from historical records. Phil. Trans. R. Soc. Lond. A 358, 957–990.
- 483 Jonkers, A., Jackson, A., Murray, A., 2003. Four centuries of geomagnetic data from his-
484 torical records. Rev. Geophys. 41,2, doi:10.1029/2002RG000115.
- 485 Kleinschmidt, R., 1989. Eine Methode zur Gewinnung von Werten der Säkularvariation
486 im Mittelalter - An einem Glasschmelzofen im Niematal und einem Kamin auf der Burg
487 Plesse dargestellt. Ph.D. thesis, Diploma thesis, Institut für Geophysik der Gerog August
488 Universität zu Göttingen.

- 489 Knothe, C., 1987. Secular variation of the magnetic declination in middle Europe during
490 the last 500 years, derived mostly from mine surveying. HHI-Report 21, 90–98.
- 491 Knothe, C., 1988. Herleitung und Bedeutung der säkularen magnetischen deklinationskur-
492 ven, speziell der des Freiburger Bergreviers. *Z. geol. Wiss.* 16, 37–42.
- 493 Körber, H.-G., 1964. Katalog der Hellmanschen Sammlung von Sonnenuhren und Kom-
494 passen des 16. bis 19. Jahrhunderts im Geomagnetische Institut Potsdam. In: *Jahrbuch*
495 *1962 des Adolf-Schmidt-Observatoriums für Erdmagnetismus in Niemegek.* Akademie-
496 Verlag, Berlin, pp. 149–172.
- 497 Körber, H.-G., 1965. Zur Geschichte der Konstruktion von Sonnenuhren und Kompassen
498 des 16. bis 18. Jahrhunderts. Vol. 3 of *Veröffentlichungen des Staatlichen Mathematisch-*
499 *Physikalischen Salons.* VEB Deutscher Verlag der Wissenschaften, Berlin.
- 500 Korte, M., Genevey, A., Constable, C. G., Frank, U., Schnepf, E., 2005. Continuous
501 geomagnetic field models for the past 7 millennia: 1. A new global data compilation.
502 *Geochem., Geophys., Geosys.* 6, Q02H15, doi:10.1029/2004GC000800.
- 503 Lamont, J., 1851. Beobachtungen des Meteorologischen Observatoriums auf dem Hohen-
504 peissenberg von 1792 - 1850. *Annalen der Münchner Sternwarte, I. Supplementband.*
505 Hübschmann, München.
- 506 Lloyd, S., 1992. *Ivory Diptych Sundials 1570-1750.* Harvard.
- 507 Ludwig, K.-H., Schmidtchen, V., 1997. *Metalle und Macht 1000-1600.* Propyläen Tech-
508 nikgeschichte, Berlin.
- 509 Macmillan, S., Maus, S., Bondar, T., Chambodut, A., Golovkov, V., Holme, R., Langlais,
510 B., Lesur, V., Lowes, F., Lühr, H., Mai, W., Mandea, M., Olsen, N., Rother, M., Sabaka,
511 T., Thomson, A., Wardinski, I., 2003. The 9th-generation International Geomagnetic
512 Reference Field. *Geophys. J. Int.* 155, 1051–1056.

- 513 Malin, S., Bullard, E., 1981. The direction of the Earth's magnetic field at London, 1570-
514 1975. *Phil. Trans. R. Soc. Lond.* 299, 357–423.
- 515 Mandeá, M., Korte, M., 2007. Ancient sundials and maps reveal historical declination
516 values. *EOS, Trans.* 88(31), 310–311.
- 517 Mandeá, M., Olsen, N., 2009. Geomagnetic and archeomagnetic jerks: where do we stand?
518 *EOS, Trans.* 90(24), 208.
- 519 Merrill, R., McElhinny, M., McFadden, P., 1996. *The Magnetic Field of the Earth*. Aca-
520 demic Press, San Diego.
- 521 Nippold, A., 1916. Ein Beitrag zur Frage der Ausrichtung der Kirchenachsen mit dem
522 Magneten. *Archiv der Naturwissenschaften und der Technik* 7, 109–114, 236–244.
- 523 Reslhuber, A., 1857. *Magnetische Beobachtungen zu Kremsmünster im Jahre 1854*.
524 Kaiserlich-Königliche Hof- und Staatsdruckerei.
- 525 Schnepf, E., Pucher, R., Geodicke, C., Manzano, A., Müller, U., Lanos, P., 2003. Paleo-
526 magnetic directions and thermoluminescence dating from a bread oven-floor sequence in
527 Lübeck (Germany): A record of 450 years of geomagnetic secular variation. *J. Geophys.*
528 *Res.* 108.
- 529 Schnepf, E., Pucher, R., Reinders, J., Hambach, U., Soffel, H., Hedley, I., 2004. A German
530 catalogue of archaeomagnetic data. *Geophys. J. Int.* 157, 64–78.
- 531 Schreyer, O., 1886. *Erdmagnetische Beobachtungen im Königreich Sachsen*. Gerlachsche
532 Buchdruckerei, Freiberg.
- 533 Soare, A., Cucu, G., Alexandrescu, M. M., 1998. Historical geomagnetic measurements in
534 Romania. *Annali di Geofisica* 41, 539–554.
- 535 Stark, A., 1814. *Meteorologisches Jahrbuch von 1813*. Augsburg.
- 536 Syndram, D., 1989. *Wissenschaftliche Instrumente und Sonnenuhren*. Callwey, Munich.

- 537 Wagner, G., 1997. Die magnetische Deklination und was die Sonnenuhrmacher davon
538 wussten. In: Sonnenuhren und wissenschaftliche Instrumente - Aus den Sammlungen des
539 Mainfränkischen Museums Würzburg. Katalog des Mainfränkischen Museums Würzburg,
540 Band 9, pp. 36–54.
- 541 Wehner, H., 1905. Ueber die Kenntnis der magnetischen Nordweisung im frühen Mittelal-
542 ter, I - III. Das Weltall 18-20, 319–324, 340–347, 351–356.
- 543 Wolkenhauer, A., 1904. Beiträge zur Geschichte der Kartographie und Nautik des 15. bis
544 17. Jahrhunderts. Mitteil. d. Geogr. Ges. München 1, 161–260.
- 545 Wolkenhauer, A., 1907. Der Nürnberger Kartograph Erhard Etzlaub. Dt. geogr. Blätter
546 30, 1–23.
- 547 Zinner, E., 1939. Die ältesten Räderuhren und modernen sonnenuhren. In: 28. Bericht der
548 naturforschenden Gesellschaft. Bamberg, pp. 1–148.
- 549 Zinner, E., 1979. Deutsche und niederländische astronomische Instrumente des 11.-18.
550 Jahrhunderts. C.H. Beck'sche Verlagsbuchhandlung, München.
- 551 zu Freiberg, K. B. (Ed.), 1850. Kalender für den Sächsischen Berg- und Hüttenmann auf
552 das Jahr 1850. Gerlach'sche Buchdruckerei Freiberg.