



On estimation of the shape parameter of the gamma distribution

A. Zaigraev, A. Podraza-Karakulska

► To cite this version:

A. Zaigraev, A. Podraza-Karakulska. On estimation of the shape parameter of the gamma distribution. Statistics and Probability Letters, 2009, 78 (3), pp.286. 10.1016/j.spl.2007.07.003 . hal-00594453

HAL Id: hal-00594453

<https://hal.science/hal-00594453>

Submitted on 20 May 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Author's Accepted Manuscript

On estimation of the shape parameter of the gamma distribution

A. Zaigraev, A. Podraza-Karakulska


PII: S0167-7152(07)00240-4
DOI: doi:10.1016/j.spl.2007.07.003
Reference: STAPRO 4706

To appear in: *Statistics & Probability Letters*

Received date: 8 December 2006
Revised date: 17 May 2007
Accepted date: 24 July 2007

Cite this article as: A. Zaigraev and A. Podraza-Karakulska, On estimation of the shape parameter of the gamma distribution, *Statistics & Probability Letters* (2007), doi:[10.1016/j.spl.2007.07.003](https://doi.org/10.1016/j.spl.2007.07.003)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



www.elsevier.com/locate/stapro

On estimation of the shape parameter of the gamma distribution

A. Zaigraev* and A. Podraza-Karakulska
Nicolaus Copernicus University of Toruń

Abstract

The problem of estimation of an unknown shape parameter under the sample drawn from the gamma distribution, where the scale parameter is also unknown, is considered. A new estimator, called the maximum likelihood scale invariant estimator, is proposed. It is established that both the bias and the variance of this estimator are less than that of the usual maximum likelihood estimator. A property of the psi function is also obtained.

2000 Subject Classifications: Primary: 62F10
Secondary: 33B15

Keywords: gamma distribution, maximum likelihood scale invariant estimator, maximum likelihood estimator, psi function

*Postal address: Faculty of Mathematics and Computer Sciences, N.Copernicus University, Chopin str. 12/18, 87-100 Toruń, POLAND. Email address: alzaig@mat.uni.torun.pl

1 Introduction

Let a sample $x = (x_1, x_2, \dots, x_n)$ be drawn from the gamma distribution $\Gamma(\alpha, \sigma)$ with an unknown shape parameter $\alpha > 0$ and an unknown scale parameter $\sigma > 0$, whose density function has the form

$$p(u; \alpha, \sigma) = \frac{u^{\alpha-1} e^{-u/\sigma}}{\sigma^\alpha \Gamma(\alpha)}, \quad u > 0.$$

Consider the problem of estimation of α . One of the most popular estimators is the well-known maximum likelihood estimator (ML-estimator) (e.g. [3, Sections 9.3, 9.4], [4], [5], [6]). Let

$$\mathbf{p}(x; \alpha, \sigma) = \sigma^{-n\alpha} (\Gamma(\alpha))^{-n} \left(\prod_{j=1}^n x_j \right)^{\alpha-1} \exp \left(-\frac{1}{\sigma} \sum_{k=1}^n x_k \right)$$

be the corresponding likelihood function. The ML-estimators of α and σ are determined by the equations:

$$\begin{cases} \ln \sigma + \Psi(\alpha) = \sum_{j=1}^n \ln x_j / n, \\ \alpha - \sum_{k=1}^n x_k / (n\sigma) = 0, \end{cases}$$

where $\Psi(\alpha) := \frac{d}{d\alpha} \ln \Gamma(\alpha) := (\ln \Gamma(\alpha))'$ is the so-called Euler psi (digamma) function.

From those equations one can obtain the ML-estimators α^* and σ^* of α and σ , respectively. Namely, α^* is the root of the equation

$$g(\alpha) := \ln \alpha - \Psi(\alpha) = \ln \bar{x} - \frac{1}{n} \sum_{j=1}^n \ln x_j$$

while

$$\sigma^* = \frac{\bar{x}}{\alpha^*}.$$

Here, \bar{x} is the sample mean, i.e.

$$\bar{x} = \frac{1}{n} \sum_{k=1}^n x_k.$$

Observe that the function g is strictly decreasing and takes values in $(0, \infty)$ (e.g. Theorem 1 of [2]). Therefore, the estimator α^* is well-defined and unique.

Observe also that the estimator α^* is scale invariant. Furthermore, one can easily see that

$$E_{(\alpha, \sigma)} \left(\ln \bar{x} - \frac{1}{n} \sum_{j=1}^n \ln x_j \right) = \Psi(n\alpha) - \Psi(\alpha) - \ln n = g(\alpha) - g(n\alpha) := g_n(\alpha).$$

The question arises: *why wouldn't one take the root of the equation*

$$g_n(\alpha) = \ln \bar{x} - \frac{1}{n} \sum_{j=1}^n \ln x_j \quad (1)$$

as an estimator of an unknown shape parameter α ? Such an estimator would coincide with that based on the method of moments.

It turns out that such a choice has quite a deep reasoning. Since in our scheme σ becomes a nuisance parameter, it is natural to apply the maximum likelihood principle to the measure defined on the σ -algebra of the scale invariant sets generated by the underlying gamma distribution. As it is known (e.g. [7, Subsection 3.2.2], [8, Section 8.3]), the density corresponding to this measure, with respect to that generated by the standard normal distribution, is given as follows:

$$\mathbf{q}(x; \alpha) = \frac{\int_0^\infty t^{n-1} \mathbf{p}(tx; \alpha, \sigma) dt}{\int_0^\infty t^{n-1} \mathbf{s}(tx) dt} = \frac{2\pi^{n/2} \Gamma(n\alpha) (\sum_{i=1}^n x_i^2)^{n/2} (\prod_{i=1}^n x_i)^{\alpha-1}}{\Gamma(n/2) (\Gamma(\alpha))^n (\sum_{i=1}^n x_i)^{n\alpha}},$$

where

$$\mathbf{s}(x) = (2\pi)^{-n/2} \exp\left(-\frac{1}{2} \sum_{k=1}^n x_k^2\right).$$

Then by direct calculations one can obtain that the maximum likelihood scale invariant estimator (IML-estimator) $\alpha^{**} \in \arg \max_{\alpha > 0} \mathbf{q}(x; \alpha)$ is the root of the equation (1).




Figure 1: Graphs of the functions g, g_n, g', g'_n for $n = 2$.

The estimator α^{**} is also scale invariant, well-defined and unique since the function g_n is strictly decreasing and takes values in $(0, \infty)$ (see Lemma 1 in Section 3).

It is worth noting that the scale invariance of the maximum likelihood estimator of a shape parameter, is a quite common property in the case when the scale is also unknown. Indeed, consider the likelihood function

$$\mathbf{p}(x; \alpha, \sigma) = \sigma^{-n} \prod_{j=1}^n p(\sigma^{-1}x_j; \alpha, 1),$$

where α is a shape parameter taking values in (α_-, α_+) . Assume that

$$\max_{\alpha \in (\alpha_-, \alpha_+), \sigma > 0} \mathbf{p}(x; \alpha, \sigma) = \max_{\alpha \in (\alpha_-, \alpha_+)} \max_{\sigma > 0} \mathbf{p}(x; \alpha, \sigma).$$

Let

$$\hat{\sigma}(x; \alpha) \in \arg \max_{\sigma > 0} \mathbf{p}(x; \alpha, \sigma).$$

Observe that for any $\lambda > 0$

$$\mathbf{p}(\lambda x; \alpha, \sigma) = \lambda^{-n} \mathbf{p}(x; \alpha, \sigma/\lambda),$$

whence

$$\hat{\sigma}(\lambda x; \alpha) = \lambda \hat{\sigma}(x; \alpha).$$

Thus,

$$\begin{aligned} (\alpha^*, \sigma^*) &\in \arg \max_{\alpha \in (\alpha_-, \alpha_+), \sigma > 0} \mathbf{p}(x; \alpha, \sigma) = \arg \max_{\alpha \in (\alpha_-, \alpha_+)} \mathbf{p}(x; \alpha, \hat{\sigma}(x; \alpha)) \\ &= \arg \max_{\alpha \in (\alpha_-, \alpha_+)} \left((\hat{\sigma}(x; \alpha))^{-n} \prod_{j=1}^n p((\hat{\sigma}(x; \alpha))^{-1}x_j; \alpha, 1) \right). \end{aligned}$$

It is evident that $\alpha^*(\lambda x) = \alpha^*(x)$, i.e. the estimator α^* is scale invariant. Therefore, it is reasonable to apply the method presented here also for other distributions.

The goal of this paper is to compare two estimators of α : the ML-estimator and the IML-estimator. We show that both the bias and the variance of the IML-estimator α^{**} are less than that of the ML-estimator α^* .

The paper is organized as follows. In Section 2 we establish the main result while all the auxiliary results are proved in Section 3.

2 Main result

Theorem. *If a sample $x = (x_1, x_2, \dots, x_n)$ is drawn from $\Gamma(\alpha, \sigma)$ distribution, then*

$$\mathbb{E}\alpha^* > \mathbb{E}\alpha^{**} > \alpha, \quad \text{Var } \alpha^* > \text{Var } \alpha^{**}. \quad (2)$$

Proof. Since the functions g and g_n are strictly decreasing, the inverse functions g^{-1} and g_n^{-1} are well-defined. According to the definitions,

$$\alpha^* = g^{-1}(T(x)), \quad \alpha^{**} = g_n^{-1}(T(x)),$$

where

$$T(x) := \ln \bar{x} - \frac{1}{n} \sum_{j=1}^n \ln x_j.$$

Denote

$$f(u) := g^{-1}(u) - g_n^{-1}(u), \quad u > 0.$$

The function f is positive. Indeed, since

$$g_n(g^{-1}(u)) = g(g^{-1}(u)) - g(ng^{-1}(u)) = u - g(ng^{-1}(u)) < u$$

and the function g_n is strictly decreasing, we obtain

$$g^{-1}(u) > g_n^{-1}(u), \quad u > 0.$$

Therefore, $E\alpha^* > E\alpha^{**}$.

Now let us show that for any $\alpha > 0$

$$E\alpha^{**} = E g_n^{-1}(T(x)) > \alpha.$$

Observe that the function g_n^{-1} is strictly convex since from Lemma 1 in Section 3 it follows that

$$\frac{\partial^2 g_n^{-1}(u)}{\partial u^2} = -\frac{g_n''(g_n^{-1}(u))}{(g_n'(g_n^{-1}(u)))^3} > 0, \quad u > 0.$$

Since $ET(x) = g_n(\alpha)$, by the Jensen inequality we get

$$E g_n^{-1}(T(x)) > g_n^{-1}(ET(x)) = g_n^{-1}(g_n(\alpha)) = \alpha.$$

It remains to prove the inequality for variances. From the evident equalities

$$\alpha^* - E\alpha^{**} = \alpha^* - \alpha^{**} + \alpha^{**} - E\alpha^{**}, \quad \alpha^* - E\alpha^{**} = \alpha^* - E\alpha^* + E\alpha^* - E\alpha^{**}$$

one can obtain

$$\text{Var } \alpha^* + (E\alpha^* - E\alpha^{**})^2 = \text{Var } \alpha^{**} + E(\alpha^* - \alpha^{**})^2 + 2E(\alpha^* - \alpha^{**})(\alpha^{**} - E\alpha^{**}).$$

Therefore,

$$\text{Var } \alpha^* - \text{Var } \alpha^{**} = \text{Var } (\alpha^* - \alpha^{**}) + 2E(\alpha^* - \alpha^{**})(\alpha^{**} - E\alpha^{**}).$$

In order to show the second part of (2), it is enough to establish that

$$\mathbb{E}(\alpha^* - \alpha^{**})(\alpha^{**} - \mathbb{E}\alpha^{**}) > 0.$$

Denote $c := \mathbb{E}\alpha^{**}$ and observe that

$$\begin{aligned} \mathbb{E}(\alpha^* - \alpha^{**})(\alpha^{**} - \mathbb{E}\alpha^{**}) &= \mathbb{E}\left(g^{-1}(T(x)) - g_n^{-1}(T(x))\right)\left(g_n^{-1}(T(x)) - c\right) \\ &= \int_0^\infty (g^{-1}(u) - g_n^{-1}(u))(g_n^{-1}(u) - c)p_{T(x)}(u)du, \end{aligned}$$

where $p_{T(x)}$ is the density of $T(x)$.

Let

$$\Delta(u) := g_n^{-1}(u) - c, \quad u > 0.$$

Observe that

$$\Delta(u) > 0, \quad 0 < u < g_n(c),$$

$$\Delta(u) = 0, \quad u = g_n(c),$$

$$\Delta(u) < 0, \quad u > g_n(c).$$

Since, as it is proved in Lemma 3 of Section 3, the function f is strictly decreasing, we obtain

$$\int_0^{g_n(c)} f(u)\Delta(u)p_{T(x)}(u)du > f(g_n(c)) \int_0^{g_n(c)} \Delta(u)p_{T(x)}(u)du$$

and

$$\int_{g_n(c)}^\infty f(u)\Delta(u)p_{T(x)}(u)du > f(g_n(c)) \int_{g_n(c)}^\infty \Delta(u)p_{T(x)}(u)du.$$

Therefore,

$$\begin{aligned} \mathbb{E}(\alpha^* - \alpha^{**})(\alpha^{**} - \mathbb{E}\alpha^{**}) &= \int_0^\infty f(u)\Delta(u)p_{T(x)}(u)du > f(g_n(c)) \int_0^\infty \Delta(u)p_{T(x)}(u)du \\ &= f(g_n(c))\mathbb{E}(\alpha^{**} - \mathbb{E}\alpha^{**}) = 0. \end{aligned} \quad \triangle$$

Remark 1. Since

$$\mathbb{E}(\alpha^* - \alpha)^2 = \text{Var } \alpha^* + (E\alpha^* - \alpha)^2, \quad \mathbb{E}(\alpha^{**} - \alpha)^2 = \text{Var } \alpha^{**} + (E\alpha^{**} - \alpha)^2,$$

from the theorem it follows that

$$\mathbb{E}(\alpha^* - \alpha)^2 > \mathbb{E}(\alpha^{**} - \alpha)^2.$$

Let us apply the Monte-Carlo simulation to confirm the results of the theorem. Given α and σ , we generate 10000 samples drawn from $\Gamma(\alpha, \sigma)$ distribution for $n = 10, 20, 30, 50, 100$. Next, we solve numerically two equations: $g(\alpha) = T(x)$ and $g_n(\alpha) = T(x)$ with respect to α , and obtain 10000 values of α^* and α^{**} . Taking their means, we estimate $E\alpha^*$ and $E\alpha^{**}$ and then calculate $(E\alpha^* - \alpha)^2$, $(E\alpha^{**} - \alpha)^2$, $\text{Var } \alpha^*$, $\text{Var } \alpha^{**}$. The corresponding programs are written in C++. The results are presented in Table 1.

$\alpha = 0.125$					$\alpha = 0.5$			
n	v^*	v^{**}	$\text{Var } \alpha^*$	$\text{Var } \alpha^{**}$	v^*	v^{**}	$\text{Var } \alpha^*$	$\text{Var } \alpha^{**}$
10	0.00159	0.00126	0.00119	0.00118	0.00410	0.00140	0.02770	0.02760
20	0.00120	0.00097	0.00100	0.00100	0.00258	0.00114	0.02008	0.01941
30	0.00100	0.00084	0.00080	0.00079	0.00120	0.00052	0.01392	0.01347
50	0.00082	0.00072	0.00048	0.00047	0.00056	0.00027	0.00849	0.00821
100	0.00068	0.00064	0.000228	0.000225	0.00011	0.00005	0.00367	0.00361
$\alpha = 1$					$\alpha = 2$			
n	v^*	v^{**}	$\text{Var } \alpha^*$	$\text{Var } \alpha^{**}$	v^*	v^{**}	$\text{Var } \alpha^*$	$\text{Var } \alpha^{**}$
10	0.0145	0.0042	0.1171	0.1165	0.0536	0.0136	0.4885	0.4876
20	0.0112	0.0048	0.0865	0.0841	0.0463	0.0197	0.3744	0.3705
30	0.0078	0.0038	0.0647	0.0625	0.0320	0.0148	0.2785	0.2706
50	0.0026	0.0012	0.0371	0.0358	0.0128	0.0059	0.1672	0.1615
100	0.0006	0.0003	0.0174	0.0171	0.0029	0.0013	0.0771	0.0756
$\alpha = 4$					$\alpha = 16$			
n	v^*	v^{**}	$\text{Var } \alpha^*$	$\text{Var } \alpha^{**}$	v^*	v^{**}	$\text{Var } \alpha^*$	$\text{Var } \alpha^{**}$
10	0.216	0.051	2.027	2.026	42.83	12.07	198.44	127.58
20	0.186	0.077	1.541	1.535	3.09	1.21	25.69	25.44
30	0.142	0.062	1.190	1.137	2.04	0.86	19.02	18.62
50	0.062	0.029	0.71	0.69	1.03	0.47	12.50	12.05
100	0.014	0.007	0.335	0.328	0.24	0.11	5.556	5.446
$\alpha = 32$					$\alpha = 128$			
n	v^*	v^{**}	$\text{Var } \alpha^*$	$\text{Var } \alpha^{**}$	v^*	v^{**}	$\text{Var } \alpha^*$	$\text{Var } \alpha^{**}$
10	174.38	2.43	774.92	560.11	191.55	32.38	2102.56	2099.77
20	11.82	4.38	103.39	102.07	173.42	62.90	1619.45	1615.24
30	8.62	3.59	78.95	76.43	143.47	60.76	1288.96	1247.04
50	3.81	1.69	47.84	46.48	55.77	23.48	781.47	759.32
100	0.89	0.38	21.96	21.53	14.14	5.982	357.58	350.46

Table 1: Numerical calculations of $v^* = (E\alpha^* - \alpha)^2$, $v^{**} = (E\alpha^{**} - \alpha)^2$, $\text{Var } \alpha^*$, $\text{Var } \alpha^{**}$.

3 Proofs of auxiliary results

Lemma 1. *For any $n > 1$ the function $g_n(u) = \Psi(nu) - \Psi(u) - \ln n$, $u > 0$, is strictly decreasing and strictly convex.*

Proof. The integral representation

$$\Psi'(u) = \int_0^{\infty} \frac{te^{-ut}}{1 - e^{-t}} dt, \quad u > 0 \quad (3)$$

(e.g. [1, formula 6.4.1]) yields

$$g'_n(u) = n\Psi'(nu) - \Psi'(u) = \int_0^{\infty} \frac{t}{1 - e^{-t}} (ne^{-nut} - e^{-ut}) dt, \quad u > 0. \quad (4)$$

Denote

$$h(t) := \frac{t}{1 - e^{-t}}, \quad \Delta_{n,u}(t) := ne^{-nut} - e^{-ut}, \quad t > 0.$$

The function h is strictly increasing since

$$h'(t) = \frac{1 - (1+t)e^{-t}}{(1 - e^{-t})^2} > 0, \quad t > 0$$

due to the inequality $e^t > 1 + t$, $t > 0$. The function $\Delta_{n,u}$ is such that

$$\Delta_{n,u}(t) > 0, \quad 0 < t < t_n(u) = \frac{\ln n}{(n-1)u},$$

$$\Delta_{n,u}(t) = 0, \quad t = t_n(u),$$

$$\Delta_{n,u}(t) < 0, \quad t > t_n(u).$$

This implies that

$$\int_0^{t_n(u)} h(t) \Delta_{n,u}(t) dt < h(t_n(u)) \int_0^{t_n(u)} \Delta_{n,u}(t) dt$$

and

$$\int_{t_n(u)}^{\infty} h(t) \Delta_{n,u}(t) dt < h(t_n(u)) \int_{t_n(u)}^{\infty} \Delta_{n,u}(t) dt.$$

Therefore,

$$g'_n(u) < h(t_n(u)) \int_0^{\infty} \Delta_{n,u}(t) dt = 0$$

since

$$\int_0^{\infty} \Delta_{n,u}(t) dt = 0.$$

Further, by differentiation of both sides of (4) we obtain

$$g_n''(u) = \int_0^{\infty} \frac{t}{1 - e^{-t}} (te^{-ut} - n^2 te^{-nut}) dt, \quad u > 0.$$

Denote

$$\bar{\Delta}_{n,u}(t) := te^{-ut} - n^2 te^{-nut}, \quad t > 0$$

and observe that

$$\bar{\Delta}_{n,u}(t) > 0, \quad t > \bar{t}_n(u) = \frac{2 \ln n}{(n-1)u},$$

$$\bar{\Delta}_{n,u}(t) = 0, \quad t = \bar{t}_n(u),$$

$$\bar{\Delta}_{n,u}(t) < 0, \quad 0 < t < \bar{t}_n(u).$$

This implies that

$$\int_0^{\bar{t}_n(u)} h(t) \bar{\Delta}_{n,u}(t) dt > h(\bar{t}_n(u)) \int_0^{\bar{t}_n(u)} \bar{\Delta}_{n,u}(t) dt$$

and

$$\int_{\bar{t}_n(u)}^{\infty} h(t) \bar{\Delta}_{n,u}(t) dt > h(\bar{t}_n(u)) \int_{\bar{t}_n(u)}^{\infty} \bar{\Delta}_{n,u}(t) dt.$$

Therefore,

$$g_n''(u) > h(\bar{t}_n(u)) \int_0^{\infty} \bar{\Delta}_{n,u}(t) dt = 0$$

since

$$\int_0^{\infty} \bar{\Delta}_{n,u}(t) dt = 0. \quad \triangle$$

In the next lemma a property of the psi function is established. This result will be helpful in proving the last lemma.

Lemma 2. *Let the function θ be defined by the equality*

$$g'(\theta(u)) = g_n'(u), \quad u > 0.$$

Then

$$g(\theta(u)) > g_n(u), \quad u > 0.$$

Proof. First of all, observe that the function θ is well-defined since both the functions g' and g'_n take values in $(-\infty, 0)$, are negative and strictly increasing (see Theorem 1 of [2] and Lemma 1 above).

Now we prove that

$$g' \left(\sqrt{n/(n-1)}u \right) > g'_n(u), \quad u > 0. \quad (5)$$

Consider the function

$$\rho(u) := g'_n(u) - g' \left(\sqrt{n/(n-1)}u \right) = g'(u) - ng'(nu) - g' \left(\sqrt{n/(n-1)}u \right), \quad u > 0.$$

Making use of (3) one can write the representation (Binet's formula):

$$g(u) = \int_0^\infty \varphi(t) e^{-ut} dt, \quad u > 0,$$

where

$$\varphi(t) := \frac{1}{1-e^{-t}} - \frac{1}{t}, \quad t > 0.$$

Therefore,

$$\rho(u) = \int_0^\infty \varphi(t) \left(nte^{-nut} + te^{-\sqrt{n/(n-1)}ut} - te^{-ut} \right) dt, \quad u > 0.$$

The function φ is strictly increasing and

$$\lim_{t \rightarrow 0} \varphi(t) = \frac{1}{2}, \quad \lim_{t \rightarrow \infty} \varphi(t) = 1$$

(see the proof of Theorem 1 in [2]). Denote

$$\begin{aligned} \Delta_n(z) &:= nze^{-nz} + ze^{-\sqrt{n/(n-1)}z} - ze^{-z} \\ &= ze^{-z} \left(ne^{-(n-1)z} + e^{-(\sqrt{n/(n-1)}-1)z} - 1 \right), \quad z > 0. \end{aligned}$$

Then

$$\rho(u) = \frac{1}{u^2} \int_0^\infty \varphi \left(\frac{z}{u} \right) \Delta_n(z) dz, \quad u > 0.$$

Observe that there exists $z_0 > 0$ such that

$$\Delta_n(z) > 0, \quad 0 < z < z_0,$$

$$\Delta_n(z) = 0, \quad z = z_0,$$

$$\Delta_n(z) < 0, \quad z > z_0.$$

The standard reasoning leads to the inequalities

$$\begin{aligned} \int_0^{z_0} \varphi\left(\frac{z}{u}\right) \Delta_n(z) dz &< \varphi\left(\frac{z_0}{u}\right) \int_0^{z_0} \Delta_n(z) dz, \\ \int_{z_0}^{\infty} \varphi\left(\frac{z}{u}\right) \Delta_n(z) dz &< \varphi\left(\frac{z_0}{u}\right) \int_{z_0}^{\infty} \Delta_n(z) dz. \end{aligned}$$

Therefore,

$$u^2 \rho(u) = \int_0^{\infty} \varphi\left(\frac{z}{u}\right) \Delta_n(z) dz < \varphi\left(\frac{z_0}{u}\right) \int_0^{\infty} \Delta_n(z) dz = 0$$

since

$$\int_0^{\infty} \Delta(z) dz = 0,$$

and (5) holds.

Since the function g' is strictly increasing, from (5) it follows that

$$\theta(u) < \sqrt{n/(n-1)}u, \quad u > 0.$$

On the other hand,

$$\theta(u) > u, \quad u > 0$$

since

$$g'_n(u) - g'(u) = -ng'(nu) > 0, \quad u > 0.$$

Denote

$$\lambda(u) := g(\theta(u)) - g_n(u), \quad u > 0.$$

Since $\theta'(u) > 1$ and $g'(u) < 0$, we obtain

$$\lambda'(u) = g'(\theta(u))\theta'(u) - g'_n(u) = g'(\theta(u))(\theta'(u) - 1) < 0, \quad u > 0.$$

Hence, the function λ is strictly decreasing. Now we establish that $\lambda(u) > 0$ for all sufficiently large u . Due to the representations

$$g(u) = \frac{1}{2u}(1 + o(1)), \quad g_n(u) = \frac{1}{2u}\left(1 - \frac{1}{n}\right)(1 + o(1)), \quad u \rightarrow \infty$$

(e.g. [1, formula 6.3.18]), we get

$$\lambda(u) = \frac{1}{2} \left(\frac{1}{\theta(u)} - \frac{1}{u} \left(1 - \frac{1}{n} \right) \right) (1 + o(1)), \quad u \rightarrow \infty.$$

Simple calculations leads to the conclusion that $\lambda(u) > 0$, as $u \rightarrow \infty$, for $\theta(u) < \sqrt{n/(n-1)}u$. Thus, $\lambda(u) > 0$ for all $u > 0$. \triangle

In the proof of Lemma 2 it is established that $u < \theta(u) < \sqrt{n/(n-1)}u$. The numerical calculations, made with the help of Maple for $n = 2, 10, 50$, and presented in Table 2, confirm those inequalities.

	$n = 2$		$n = 10$		$n = 50$	
u	$\theta(u)$	$\sqrt{n/(n-1)}u$	$\theta(u)$	$\sqrt{n/(n-1)}u$	$\theta(u)$	$\sqrt{n/(n-1)}u$
0.1	0.134913918	0.141421356	0.103577225	0.105409255	0.100565420	0.101015254
0.5	0.651452797	0.707106781	0.517553390	0.527046277	0.503191582	0.505076272
1	1.316748502	1.414213562	1.039115016	1.054092553	1.007252736	1.010152544
3	4.090593081	4.242640686	3.140261683	3.162277660	3.026250755	3.030457633
5	6.905201515	7.071067810	5.246643827	5.270462768	5.046219652	5.050762721
10	13.96570362	14.14213562	10.51571989	10.54092553	10.09672350	10.10152544
50	70.52586082	70.71067810	52.67831719	52.70462768	50.50261897	50.50762721
100	141.2355025	141.4213562	105.3828081	105.4092553	101.0102207	101.0152544

Table 2: Numerical calculations of bounds for $\theta(u)$.

Lemma 3. For any $n > 1$ the function $f(u) = g^{-1}(u) - g_n^{-1}(u)$, $u > 0$, is strictly decreasing.

Proof. Clearly,

$$f'(u) = \frac{1}{g'(g^{-1}(u))} - \frac{1}{g'_n(g_n^{-1}(u))}, \quad u > 0.$$

For any $u > 0$ we have

$$g(g^{-1}(u)) = g_n(g_n^{-1}(u)) < g(\theta(g_n^{-1}(u))), \quad (6)$$

where the function θ is defined as in Lemma 2. Since the function g is strictly decreasing, from (6) it follows that

$$g^{-1}(u) > \theta(g_n^{-1}(u)), \quad u > 0$$

while since the function g' is strictly increasing, we get

$$g'_n(g_n^{-1}(u)) = g'(\theta(g_n^{-1}(u))) < g'(g^{-1}(u)), \quad u > 0.$$

Thus, $f'(u) < 0$ for all $u > 0$. \triangle

Acknowledgement

The authors are grateful to the referee for very helpful comments and useful suggestions.

References

- [1] Abramovitz, M. and Stegun, I. A., eds. (1965), *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables* (Dover, New York).
- [2] Alzer, H. (1997), *On some inequalities for the gamma and psi functions*, Math. Comp. **66**, 373–389.
- [3] Barndorff-Nielsen, O. (1978), *Information and Exponential families in Statistical Theory* (Wiley, New York).
- [4] Bowman, K. O. and Shenton, L. R. (1988), *Properties of Estimators for the Gamma Distribution* (Marcel Dekker, New York).
- [5] Dang, H. and Weerakkody, G. (2000), *Bounds for the maximum likelihood estimates in two-parameter gamma distribution*, J. Math. Anal. Appl. **245**, 1–6.
- [6] Crain, B. R. (1976), *Exponential models, maximum likelihood estimation, and the Haar condition*, J. Amer. Statist. Assoc. **71**, 737–740.
- [7] Hájek, J., Šidák, Z. and Sen, Pranab K. (1999), *Theory of Rank Tests*. Second Edition (Academic Press, San Diego).
- [8] Nagaev, A. V. (1996), *Limit Theorems under Testing Hypotheses* (UMK, Torun).