

# The almost sure central limit theorems in the joint version for the maxima and sums of certain stationary Gaussian sequences

Marcin Dudziński

### ► To cite this version:

Marcin Dudziński. The almost sure central limit theorems in the joint version for the maxima and sums of certain stationary Gaussian sequences. Statistics and Probability Letters, 2009, 78 (4), pp.347. 10.1016/j.spl.2007.07.007. hal-00594454

## HAL Id: hal-00594454 https://hal.science/hal-00594454

Submitted on 20 May 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

### Author's Accepted Manuscript

The almost sure central limit theorems in the joint version for the maxima and sums of certain stationary Gaussian sequences

Marcin Dudziński

PII: DOI: Reference:

To appear in:

S0167-7152(07)00247-7 doi:10.1016/j.spl.2007.07.007 e: STAPRO 4713

Statistics & Probability Letters



www.elsevier.com/locate/stapro

Received date:3 November 2004Revised date:30 March 2007Accepted date:25 July 2007

Cite this article as: Marcin Dudziński, The almost sure central limit theorems in the joint version for the maxima and sums of certain stationary Gaussian sequences, *Statistics & Probability Letters* (2007), doi:10.1016/j.spl.2007.07.007

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The almost sure central limit theorems in the joint version for the maxima and sums of certain stationary Gaussian sequences

Marcin Dudziński<sup>1</sup>

Department of Econometrics and Statistics,

Warsaw Agricultural University, ul. Nowoursynowska 166, 02-787 Warszawa, Poland

#### Abstract

Suppose that  $X_1, X_2, ...$  is a standardized stationary Gaussian sequence. Let:  $M_n := \max(X_1, ..., X_n), S_n := \sum_{i=1}^n X_i, \sigma_n := \sqrt{Var(S_n)}$ , and  $a_n > 0, b_n$  denote suitable normalizing constants. Our goal is to prove the almost sure central limit theorem for the sequence  $\{a_n (M_n - b_n), S_n / \sigma_n\}$ , under certain additional assumptions on the covariance function  $r(t) := Cov(X_1, X_{1+t})$ . MSC: primary 60F15, secondary 60F05.

Key words and phrases: Almost sure central limit theorem; Extreme values; Partial sums; Stationary Gaussian sequences; Associated random variables; Slowly varying functions at infinity; Karamata's theorem.

#### 1. Introduction

The almost sure central limit theorem (ASCLT) has become an intensively studied subject in recent research. In our paper, we are concerned with the ASCLT in its two-dimensional version. A general outline of the investigations concerning the ASCLT in such a form may be described as follows. Suppose that  $X_1, X_2, \ldots$  is a sequence of r.v.'s,  $f_1, f_2, \ldots, f_k, \ldots$  and  $g_1, g_2, \ldots, g_k, \ldots$  are some real-valued measurable functions, defined on  $\mathbb{R}, \mathbb{R}^2, \ldots, \mathbb{R}^k, \ldots$ , respectively. We seek conditions under which, for some nondegenerate d.f. H, defined on  $\mathbb{R}^2$ ,

$$\frac{1}{D_N} \sum_{n=1}^N d_n I\left\{f_n\left(X_1, X_2, ..., X_n\right) \le x, g_n\left(X_1, X_2, ..., X_n\right) \le y\right\} \xrightarrow{\text{a.s.}} H\left(x, y\right) \quad (1)$$

for all  $(x, y) \in C_H$ , where:  $\{d_n\}$  is some sequence of weights,  $D_N = \sum_{n=1}^N d_n$ , I stands for the indicator function,  $C_H$  denotes the set of continuity points of H.

In our investigations, we will restrict ourselves to the situation, when (1)

is satisfied with:  $d_n = 1/n$ ,  $D_N \sim \log N$ ,  $f_n(X_1, ..., X_n) := a_n(M_n - b_n)$ ,  $g_n(X_1, ..., X_n) := S_n/\sigma_n$ , where:  $M_n := \max(X_1, ..., X_n)$ ,  $S_n := X_1 + ... + X_n$ ,  $\sigma_n := \sqrt{Var(S_n)}$ , and  $a_n > 0$ ,  $b_n$  are certain normalizing constants. The purpose of our paper is to prove that if:  $X_1, X_2, ...$  is a standardized stationary Gaussian sequence, the covariance function  $r(t) := Cov(X_1, X_{1+t})$  satisfies

$$r(t) = L(t) / t^{\alpha}, t = 1, 2, ...,$$
 (2)

where  $\alpha > 0$  and  $L(\cdot)$  is a positive, slowly varying function at infinity, and if moreover, the numerical sequence  $\{u_n\}$  fulfills the relation

$$\lim_{n \to \infty} n \left( 1 - \Phi \left( u_n \right) \right) = \tau \text{ for some } \tau, \ 0 \le \tau < \infty, \tag{3}$$

 $<sup>{}^1\</sup>textit{E-mail addresses: mdudzinski@poczta.onet.pl, mdudzinski@mors.sggw.waw.pl}$ 

where  $\Phi$  is the standard normal d.f., then

$$\lim_{N \to \infty} \frac{1}{\log N} \sum_{n=1}^{N} \frac{1}{n} I\left\{ M_n \le u_n, S_n / \sigma_n \le y \right\} = e^{-\tau} \Phi\left(y\right) \text{ a.s. for any } y \in \mathbb{R}.$$
(4)

As a direct conclusion, we will also show that if, for  $n \ge 3$ :

$$a_n := (2\log n)^{1/2}, \ b_n := (2\log n)^{1/2} - \frac{1}{2} (2\log n)^{-1/2} (\log\log n + \log 4\pi), \ (5)$$

then for all  $x, y \in \mathbb{R}$ 

$$\lim_{N \to \infty} \frac{1}{\log N} \sum_{n=1}^{N} \frac{1}{n} I\{a_n (M_n - b_n) \le x, S_n / \sigma_n \le y\} = \exp(-e^{-x}) \Phi(y) \text{ a.s.}$$
(6)

It should be mentioned that, although  $r(t) = L(t)/t^{\alpha}$  in (2) is defined only for positive integers, the slowly varying functions  $L(\cdot)$  are in general defined on the real line or on the  $\mathbb{R}_+$  semi-line.

The earlier given notations -  $M_n$ ,  $S_n$ ,  $\sigma_n$ , r(t),  $\Phi$  - will be intensively used throughout the paper. For our convenience, we will also introduce the following one  $M_{m,n} := \max(X_{m+1}, ..., X_n)$ . Furthermore,  $f(n) \ll g(n)$  and  $f(n) \sim g(n)$ will stand for  $f(n) = \mathcal{O}(g(n))$  and  $f(n)/g(n) \to 1$ , as  $n \to \infty$ , respectively.

#### 2. Main result

Our main result is the following almost sure central limit theorem in the joint version for the maxima and sums of some standardized stationary Gaussian sequences.

**Theorem 1** . Let  $X_1, X_2, ...$  be a standardized stationary Gaussian sequence. Assume that the covariance function  $r(t) := Cov(X_1, X_{1+t})$  satisfies (2) for some  $\alpha > 0$ . Then:

(i) If the numerical sequence {u<sub>n</sub>} fulfills (3), then (4) holds,
(ii) If {a<sub>n</sub>}, {b<sub>n</sub>} are such as in (5), then (6) holds.

#### 3. Auxiliary results

In this section, we state and prove some lemmas, which will be needed for the proof of our main result. Here, we present the first one.

**Lemma 1**. Let  $X_1, X_2, ...$  be a standardized stationary Gaussian sequence. Assume that the covariance function  $r(t) := Cov(X_1, X_{1+t})$  satisfies assumption (2) with some  $\alpha > 0$ . Suppose moreover that the sequence  $\{u_n\}$  fulfills (3), and that m < n. Then for any  $y \in \mathbb{R}$  and some  $\gamma > 0$ 

$$E |I\{M_n \le u_n, S_n / \sigma_n \le y\} - I\{M_{m,n} \le u_n, S_n / \sigma_n \le y\}| \ll 1/n^{\gamma} + m/n.$$
(7)

**Proof.** Assume first that (2) holds with some  $0 < \alpha < 1$ . Let  $1 \le i \le n$ . We have

$$0 < Cov(X_i, S_n / \sigma_n) < \frac{2}{\sigma_n} \sum_{t=0}^{n-1} r(t) = \frac{2}{\sigma_n} + \frac{2}{\sigma_n} \sum_{t=1}^{n-1} \frac{L(t)}{t^{\alpha}} \text{ for some } 0 < \alpha < 1.$$
(8)

As  $0 < \alpha < 1$ , then, by the two times application of Karamata's theorem, we get (see also Mielniczuk (2002), p. 394)

$$\sigma_n \sim C(\alpha) L(n)^{1/2} n^{1-\alpha/2}$$
, where  $C(\alpha) = \{2/(1-\alpha)(2-\alpha)\}^{1/2}$ . (9)

Due to (8), (9), we have for some  $0 < \alpha < 1$  that

$$Cov\left(X_{i}, S_{n}/\sigma_{n}\right) \ll \frac{1}{L\left(n\right)^{1/2} n^{1-\alpha/2}} \sum_{t=1}^{n-1} \frac{L\left(t\right)}{t^{\alpha}} \ll \frac{L\left(n\right) n^{1-\alpha}}{L\left(n\right)^{1/2} n^{1-\alpha/2}} = \frac{L\left(n\right)^{1/2}}{n^{\alpha/2}}$$

Since L(n) is a slowly varying function at infinity, then  $L(n)^{1/2} \ll n^{\epsilon}$  for arbitrary  $\epsilon > 0$ . Consequently

$$Cov(X_i, S_n/\sigma_n) \ll \frac{n^{\epsilon}}{n^{\alpha/2}}$$
 for any  $\epsilon > 0$  and some  $0 < \alpha < 1$ .

Therefore

$$0 < \sup_{1 \le i \le n} Cov\left(X_i, S_n/\sigma_n\right) \ll 1/n^{\beta} \text{ for some } 0 < \beta < 1/2.$$
(10)

Since  $\lim_{n\to\infty} 1/n^{\beta} = 0$ , there exist numbers  $\lambda$ ,  $n_0$ , such that

$$0 < \sup_{1 \le i \le n} Cov\left(X_i, S_n/\sigma_n\right) < \lambda < \beta/(1-\beta) < 1 \text{ for any } n > n_0.$$
(11)

In turn, by assumption (2), we have  $\lim_{t\to\infty} r(t) = \lim_{t\to\infty} L(t)/t^{\alpha} = 0$ . Thus, there exist numbers  $\delta$ ,  $n_1$ , such that

$$0 < \sup_{t > n_1} r(t) = \delta < \alpha/(2 - \alpha) < 1.$$
(12)

Relations (10)-(12) will be intensively used in the next stages of our proof.

Let y be an arbitrary real number and m < n. We have

$$E | I \{ M_n \le u_n, S_n / \sigma_n \le y \} - I \{ M_{m,n} \le u_n, S_n / \sigma_n \le y \}$$
  
=  $P \{ M_{m,n} \le u_n, S_n / \sigma_n \le y \} - P \{ M_n \le u_n, S_n / \sigma_n \le y \}.$ 

Suppose moreover that  $Y_n$  is a random variable, which has the same distribution as  $S_n/\sigma_n$ , but is independent of  $(X_1, ..., X_n)$ . We can write that

$$E |I \{ M_n \le u_n, S_n / \sigma_n \le y \} - I \{ M_{m,n} \le u_n, S_n / \sigma_n \le y \} |$$

$$\le |P \{ M_n \le u_n, S_n / \sigma_n \le y \} - P \{ M_n \le u_n \} P \{ Y_n \le y \} |$$

$$+ |P \{ M_{m,n} \le u_n, S_n / \sigma_n \le y \} - P \{ M_{m,n} \le u_n \} P \{ Y_n \le y \} |$$

$$+ (P \{ M_{m,n} \le u_n \} - P \{ M_n \le u_n \}) =: A_1 + A_2 + A_3.$$
(13)

Since  $(X_1, ..., X_n, S_n/\sigma_n)$ ,  $(X_1, ..., X_n, Y_n)$  are standard normal random vectors and (11), (12) hold, then, by Theorem 4.2.1 in Leadbetter et al. (1983), the fact that  $Y_n$  is independent of  $(X_1, ..., X_n)$  and relation (10), we obtain

$$A_1 + A_2 \ll \sum_{i=1}^n Cov\left(X_i, S_n/\sigma_n\right) \exp\left(-\frac{u_n^2}{2\left(1+\lambda\right)}\right) \ll n\frac{1}{n^\beta} \exp\left(-\frac{u_n^2}{2\left(1+\lambda\right)}\right) \tag{14}$$

for some  $0 < \beta < 1/2$ , where  $\lambda$  is such as in (11). As  $\{u_n\}$  satisfies (3), then, by (4.3.4(i)), (4.3.4(ii)) in Leadbetter et al. (1983),

$$\exp\left(-\frac{u_n^2}{2(1+\lambda)}\right) \sim L\frac{\left(\log n\right)^{\frac{1}{2(1+\lambda)}}}{n^{\frac{1}{1+\lambda}}} \text{ for some } L, \text{ not depending on } n.$$
(15)

From (14) and (15), we have

$$A_{1} + A_{2} \ll n \frac{1}{n^{\beta}} \frac{(\log n)^{\frac{1}{2(1+\lambda)}}}{n^{\frac{1}{1+\lambda}}} = \frac{(\log n)^{\frac{1}{2(1+\lambda)}}}{n^{\beta+\frac{1}{1+\lambda}-1}}.$$
(16)

As, by (11),  $\lambda < \beta / (1 - \beta)$ , we have  $\beta + \frac{1}{1+\lambda} - 1 > 0$ . Thus, by (16),

$$A_1 + A_2 \ll 1/n^{\gamma} \text{ for some } \gamma > 0.$$
(17)

It remains only to estimate the term  $A_3$  in (13). By going through the first lines in the proof of Lemma 2.4 from Csaki and Gonchigdanzan (2002), we see that

$$A_{3} \leq |P\{M_{n} \leq u_{n}\} - \Phi^{n}(u_{n})| + |P\{M_{m,n} \leq u_{n}\} - \Phi^{n-m}(u_{n})| + \{\Phi^{n-m}(u_{n}) - \Phi^{n}(u_{n})\} =: B_{1} + B_{2} + B_{3}.$$
(18)

Our goal now is to give the bounds for  $B_1$ - $B_3$  in (18). Let  $\delta$  be such as in (12). By Theorem 4.2.1 in Leadbetter et al. (1983), relation (15) and assumption (2),  $B_1 + B_2 \ll n \exp\left(-\frac{u_n^2}{1+\delta}\right) \sum_{t=1}^{n-1} r(t) \ll \frac{(\log n)^{\frac{1}{1+\delta}}}{n^{\frac{2}{1+\delta}-1}} \sum_{t=1}^{n-1} \frac{L(t)}{t^{\alpha}} \ll$  $\frac{(\log n)^{\frac{1}{1+\delta}}L(n)n^{1-\alpha}}{n^{\frac{2}{1+\delta}-1}} = \frac{(\log n)^{\frac{1}{1+\delta}}L(n)}{n^{\frac{2}{1+\delta}+\alpha-2}}.$  As, by (12),  $\delta < \alpha/(2-\alpha)$ , we have  $\frac{2}{1+\delta} + \alpha - 2 > 0. \text{ Since } (\log n)^{\frac{1}{1+\delta}} L(n) \ll n^{\epsilon} \text{ for arbitrary } \epsilon > 0, \text{ we obtain}$  $B_1 + B_2 \ll 1/n^{\gamma} \text{ for some } \gamma > 0. \tag{1}$ 

(19)

Moreover, the bound

$$B_3 \le m/n \tag{20}$$

follows from the fact that  $x^{n-m} - x^n \le m/n$ , if  $0 \le x \le 1$ . By (18)-(20)

$$A_3 \ll 1/n^{\gamma} + m/n \text{ for some } \gamma > 0.$$
(21)

Relations (13), (17), (21) yield (7), provided (2) holds with some  $0 < \alpha < 1$ .

Assume now that (2) holds with some  $\alpha \geq 1$ . Since  $r(\cdot)$  is positive, we get  $\sigma_n = \sqrt{n + 2\sum_{t=1}^{n-1} (n-t)r(t)} \ge n^{1/2}$ . This and the fact that  $\alpha \ge 1$  imply  $2 - 2 - \frac{n-1}{L}L(t) = L(n)\log n \quad \text{ind} \quad 0 \le 1$ 0

$$0 < Cov \left(X_i, S_n / \sigma_n\right) < \frac{2}{n^{1/2}} + \frac{2}{n^{1/2}} \sum_{t=1}^{L(t)} \frac{L(t)}{t} \ll \frac{L(t)\log n}{n^{1/2}}. \text{ As } L(n)\log n \ll n^{t}$$

for any  $\epsilon > 0$ , we obtain that (10), (11) hold also in the case, when (2) is satisfied for some  $\alpha \ge 1$ . Let  $\mathbb{A}_1$ - $\mathbb{A}_3$  be defined as  $A_1$ - $A_3$  in (13), but for the case, when (2) holds with some  $\alpha \ge 1$ . By using Theorem 4.2.1 in Leadbetter et al. (1983) and reasoning as in (14)-(17), we get  $\mathbb{A}_1 + \mathbb{A}_2 \ll 1/n^{\gamma}$  for some  $\gamma > 0$ . Let moreover,  $\mathbb{A}_3$ ,  $\mathbb{B}_1$ - $\mathbb{B}_3$  be defined as  $A_3$ ,  $B_1$ - $B_3$  in (13), (18), but for the case, when (2) holds with some  $\alpha \ge 1$ . Since  $\lim_{t\to\infty} r(t) = 0$ , then (see the remark before Lemma 4.3.2 in Leadbetter et al. (1983))  $0 < \sup_{t\ge 1} r(t) = \delta < 1$ . By Theorem

4.2.1 in Leadbetter et al. (1983), we have  $\mathbb{B}_1 + \mathbb{B}_2 \ll n \exp\left(-\frac{u_n^2}{1+\delta}\right) \sum_{t=1}^{n-1} r(t) \ll \frac{(\log n)^{\frac{1}{1+\delta}+1} L(n)}{n^{\frac{2}{1+\delta}-1}} \sum_{t=1}^{n-1} \frac{L(t)}{t} \ll \frac{(\log n)^{\frac{1}{1+\delta}+1} L(n)}{n^{\frac{2}{1+\delta}-1}}$ . As  $(\log n)^{\frac{1}{1+\delta}+1} L(n) \ll n^{\epsilon}$  for any

 $\epsilon > 0$ , we obtain that  $\mathbb{B}_1 + \mathbb{B}_2 \ll 1/n^{\gamma}$  for some  $\gamma > 0$ . As moreover,  $\mathbb{B}_3 \leq m/n$ , we get  $\mathbb{A}_3 \leq \mathbb{B}_1 + \mathbb{B}_2 + \mathbb{B}_3 \ll 1/n^{\gamma} + m/n$  for some  $\gamma > 0$ . Consequently  $\mathbb{A}_1 + \mathbb{A}_2 + \mathbb{A}_3 \ll 1/n^{\gamma} + m/n$  for some  $\gamma > 0$ , which yields (7) in the case, when (2) holds with some  $\alpha \geq 1$ . This completes the whole proof of Lemma 1.

We also need to prove the following statement.

**Lemma 2**. Suppose that  $X_1, X_2, ...$  is a standardized stationary Gaussian sequence, the sequence  $\{u_n\}$  fulfills condition (3), and m < n. We have: (a) If the covariance function  $r(t) := Cov(X_1, X_{1+t})$  satisfies assumption (2) with some  $0 < \alpha < 1$ , then for any  $y \in \mathbb{R}$  and some  $\gamma > 0$ 

$$|Cov \left( I \left\{ M_m \le u_m, S_m / \sigma_m \le y \right\}, I \left\{ M_{m,n} \le u_n, S_n / \sigma_n \le y \right\} \right)| \\ \ll \quad \frac{1}{n^{\gamma}} + \frac{m \left( n + 2m \right)^{1-\alpha} L \left( n + 2m \right)}{L \left( m \right)^{1/2} m^{1-\alpha/2} L \left( n \right)^{1/2} n^{1-\alpha/2}} + \frac{L \left( m \right)^{1/2} m^{1-\alpha/2}}{L \left( n \right)^{1/2} n^{1-\alpha/2}}, \quad (22)$$

(b) If the covariance function  $r(t) := Cov(X_1, X_{1+t})$  satisfies assumption (2) with some  $\alpha > 1$ , then for any  $y \in \mathbb{R}$  and some  $\gamma > 0$ 

$$|Cov(I\{M_m \le u_m, S_m / \sigma_m \le y\}, I\{M_{m,n} \le u_n, S_n / \sigma_n \le y\})| \ll \frac{1}{n^{\gamma}} + \frac{m^{1/2}}{n^{1/2}},$$
(23)

(c) If the covariance function  $r(t) := Cov(X_1, X_{1+t})$  satisfies assumption (2) with  $\alpha = 1$ , then for any  $y \in \mathbb{R}$  and some  $\gamma > 0$ 

$$\|Cov \left(I \left\{ M_m \le u_m, S_m / \sigma_m \le y \right\}, I \left\{ M_{m,n} \le u_n, S_n / \sigma_n \le y \right\} \right)\|$$

$$\ll \frac{1}{n^{\gamma}} + \frac{m^{1/2} \left( \widetilde{L} \left( n + 2m \right) - \widetilde{L} \left( m + 1 \right) \right)}{n^{1/2} \widetilde{L} \left( m \right)^{1/2}} + \frac{m^{1/2} \widetilde{L} \left( m \right)^{1/2}}{n^{1/2} \widetilde{L} \left( n \right)^{1/2}}, \quad (24)$$

where  $\tilde{L}(n) := 1 + 2\sum_{t=1}^{n-1} r(t).$ 

**Proof of Lemma 2(a).** We assume that (2) holds with some  $0 < \alpha < 1$ . Let  $i \ge m + 1$ . By assumption (2) and relation (9), we have

$$0 < Cov (X_i, S_m / \sigma_m) \le \frac{1}{\sigma_m} \sum_{t=i-m}^{i-1} \frac{L(t)}{t^{\alpha}} \sim \frac{1}{C(\alpha) L(m)^{1/2} m^{1-\alpha/2}} \sum_{t=i-m}^{i-1} \frac{L(t)}{t^{\alpha}}.$$
  
Since  $L(t) \ll t^{\epsilon}$  for arbitrary  $\epsilon > 0$ , we can write that  
 $Cov (X_i, S_m / \sigma_m) \ll \frac{1}{L(m)^{1/2} m^{1-\alpha/2}} \sum_{t=i-m}^{i-1} \frac{t^{\alpha/4}}{t^{\alpha}} \le \frac{1}{L(m)^{1/2} m^{1-\alpha/2}} \sum_{t=1}^m \frac{1}{t^{3\alpha/4}}$   
 $= \frac{1}{L(m)^{1/2} m^{\alpha/4} m^{1-3\alpha/4}} \sum_{t=1}^m \frac{1}{t^{3\alpha/4}} \le \frac{1}{L(m)^{1/2} m^{\alpha/4}} \sum_{t=1}^m \frac{1}{t^{1-3\alpha/4} t^{3\alpha/4}}$   
 $\ll \frac{\log m}{L(m)^{1/2} m^{\alpha/4}} \ll \frac{m^{\epsilon}}{m^{\alpha/4}} \text{ for any } \epsilon > 0.$ 

Hence, we get that  $0 < \sup_{i \ge m+1} Cov(X_i, S_m/\sigma_m) \ll 1/m^{\eta}$  for some  $0 < \eta < 1/4$ . Consequently, there exist numbers  $\mu$ ,  $m_0$ , such that

$$0 < \sup_{i \ge m+1} Cov \left( X_i, S_m / \sigma_m \right) < \mu < \alpha / (2 - \alpha) < 1 \text{ for all } m > m_0.$$
(25)

We will use (25) later in our proof. Let: y be an arbitrary real number, m < n. We know that:  $(-X_1, ..., -X_m, -S_m/\sigma_m, -X_{m+1}, ..., -X_n, -S_n/\sigma_n)$  is a standard Gaussian vector, and (since  $r(\cdot)$  is positive) that:  $Cov(-X_i, -X_j) > 0$ ,  $Cov(-X_i, -S_j/\sigma_j) > 0$ ,  $Cov(-S_i/\sigma_i, -S_j/\sigma_j) > 0$  for all i, j. Therefore, by Theorem in Pitt (1991),  $-X_1, ..., -X_m, -S_m/\sigma_m, -X_{m+1}, ..., -X_n, -S_n/\sigma_n$  are the associated r.v.'s. Hence, due to Theorem 2 in Newman (1984),

$$0 \leq Cov \left( I \left\{ M_m \leq u_m, S_m / \sigma_m \leq y \right\}, I \left\{ M_{m,n} \leq u_n, S_n / \sigma_n \leq y \right\} \right)$$
  
$$\leq \sum_{i=1}^{m} \sum_{j=m+1}^{n} \left( P \left\{ -X_i > -u_m, -X_j > -u_n \right\} - P \left\{ -X_i > -u_m \right\} P \left\{ -X_j > -u_n \right\} \right)$$
  
$$+ \sum_{i=1}^{m} \left( P \left\{ -X_i > -u_m, -S_n / \sigma_n > -y \right\} - P \left\{ -X_i > -u_m \right\} P \left\{ -S_n / \sigma_n > -y \right\} \right)$$
  
$$+ \left( P \left\{ -X_i > -u_n, -S_m / \sigma_m > -y \right\} - P \left\{ -X_i > -u_n \right\} P \left\{ -S_m / \sigma_m > -y \right\} \right)$$
  
$$+ \left( P \left\{ -S_m / \sigma_m > -y, -S_n / \sigma_n > -y \right\} - P \left\{ -S_m / \sigma_m > -y \right\} P \left\{ -S_n / \sigma_n > -y \right\} \right)$$
  
$$=: D_1 + D_2 + D_3 + D_4. \tag{26}$$

Our task now is to estimate all the components  $D_1$ - $D_4$  in (26).

$$m \exp\left(-\frac{u_m^2 + u_n^2}{2(1+\delta)}\right) \sum_{t=1}^{n-1} r(t) \ll \frac{m(\log n)^{1+\delta}}{m^{\frac{1}{1+\delta}} n^{\frac{1}{1+\delta}}} \sum_{t=1}^{n-1} \frac{L(t)}{t^{\alpha}} \ll \frac{m(\log n)^{1+\delta} L(n) n^{1-\delta}}{m^{\frac{1}{1+\delta}} n^{\frac{1}{1+\delta}}}$$

$$=\frac{m^{1-\frac{1}{1+\delta}}\left(\log n\right)^{\frac{1}{1+\delta}}L\left(n\right)}{n^{1-\frac{1}{1+\delta}}n^{\frac{2}{1+\delta}+\alpha-2}}\ll\frac{\left(\log n\right)^{\frac{1}{1+\delta}}L\left(n\right)}{n^{\frac{2}{1+\delta}+\alpha-2}}\ll\frac{n^{\epsilon}}{n^{\frac{2}{1+\delta}+\alpha-2}} \text{ for any } \epsilon>0.$$
  
Since, by (12),  $\delta<\alpha/\left(2-\alpha\right)$ , then  $\frac{2}{1+\delta}+\alpha-2>0$  and

$$D_1 \ll 1/n^{\gamma} \text{ for some } \gamma > 0.$$
 (27)

In order to estimate the component  $D_2$  in (26), let us denote by  $Y_n$  a standard normal random variable, which is independent of  $S_n/\sigma_n$ . We can write that  $D_2 = \sum_{i=1}^m (P\{X_i \leq u_m, S_n/\sigma_n \leq y\} - P\{Y_n \leq u_m, S_n/\sigma_n \leq y\})$ . Let  $\lambda$  satisfy (11). As the vectors  $(X_i, S_n/\sigma_n)$ ,  $(Y_n, S_n/\sigma_n)$  are standard normal and (11) holds, then, due to Theorem 4.2.1 in Leadbetter et al. (1983), and (10), (15),

$$D_2 \ll \sum_{i=1}^m Cov\left(X_i, S_n/\sigma_n\right) \exp\left(-\frac{u_m^2}{2\left(1+\lambda\right)}\right) \ll \frac{m\left(\log m\right)^{\frac{1}{2\left(1+\lambda\right)}}}{n^\beta m^{\frac{1}{1+\lambda}}} \ll 1/n^\gamma$$
(28)

for some  $\gamma > 0$ , where the last relation follows from the fact that  $\beta > 1 - \frac{1}{1+\lambda}$ . We also wish to estimate  $D_3$  in (26). Let  $\mu$  satisfy (25). By using similar reasoning to that, which led us to the first derivation in (28), we obtain

$$D_3 \ll \exp\left(-\frac{u_n^2}{2\left(1+\mu\right)}\right) \sum_{i=m+1}^n Cov\left(X_i, S_m/\sigma_m\right).$$
<sup>(29)</sup>

Notice that  $\sum_{i=m+1}^{n} Cov(X_i, S_m/\sigma_m) = \frac{1}{\sigma_m} \left( \sum_{t=m}^{n-1} r(t) + \dots + \sum_{t=1}^{n-m} r(t) \right) < \frac{m}{\sigma_m} \sum_{t=1}^{n-1} r(t).$ Thus, by (29), (15), assumption (2) and relation (9),

$$D_{3} \ll \frac{(\log n)^{\frac{1}{2(1+\mu)}}}{n^{\frac{1}{1+\mu}}} \frac{m}{L(m)^{1/2} m^{1-\alpha/2}} \sum_{t=1}^{n-1} \frac{L(t)}{t^{\alpha}} \ll \frac{(\log n)^{\frac{1}{2(1+\mu)}} m^{\alpha/2} L(n) n^{1-\alpha}}{n^{\frac{1}{1+\mu}} L(m)^{1/2}} \\ \ll \frac{(\log n)^{\frac{1}{2(1+\mu)}} n^{\alpha/2} L(n)}{n^{\frac{1}{1+\mu}+\alpha-1}} = \frac{(\log n)^{\frac{1}{2(1+\mu)}} L(n)}{n^{\frac{1}{1+\mu}+\alpha/2-1}} \ll \frac{n^{\epsilon}}{n^{\frac{1}{1+\mu}+\alpha/2-1}} \text{ for any } \epsilon > 0.$$
  
Since, by (25),  $\mu < \alpha/(2-\alpha)$ , then  $\frac{1}{1+\mu} + \alpha/2 - 1 > 0$  and

$$D_3 \ll 1/n^{\gamma}$$
 for some  $\gamma > 0.$  (30)

It remains to estimate  $D_4$  in (26). Notice that, if  $Cov(S_m/\sigma_m, S_n/\sigma_n) \leq 1/2$ , then, by Theorem 4.2.1 in Leadbetter et al. (1983),

$$D_4 \le \frac{Cov\left(S_m/\sigma_m, S_n/\sigma_n\right)}{2\pi\sqrt{1 - \left\{Cov\left(S_m/\sigma_m, S_n/\sigma_n\right)\right\}^2}} \le \frac{Cov\left(S_m/\sigma_m, S_n/\sigma_n\right)}{\pi\sqrt{3}}.$$
 (31)

In turn, provided  $Cov(S_m/\sigma_m, S_n/\sigma_n) > 1/2$ , we can estimate  $D_4$  in (26) as follows. Observe that, since  $S_m/\sigma_m$ ,  $S_n/\sigma_n$  are associated and have bounded,

continuus density, then, by Lemma 2.2 in Bagai and Prakasa Rao (1991), there exists a positive constant  $C_1$ , neither depending on m, n nor on y, such that  $D_4 \leq C_1 \{Cov (S_m/\sigma_m, S_n/\sigma_n)\}^{1/3}$ . Thus, if  $Cov (S_m/\sigma_m, S_n/\sigma_n) > 1/2$ , then

$$D_{4} \leq C_{1} \frac{Cov \left(S_{m}/\sigma_{m}, S_{n}/\sigma_{n}\right)}{\left\{Cov \left(S_{m}/\sigma_{m}, S_{n}/\sigma_{n}\right)\right\}^{2/3}} < C_{1} 2^{2/3} Cov \left(S_{m}/\sigma_{m}, S_{n}/\sigma_{n}\right).$$
(32)

Put  $C := \max(1/\pi\sqrt{3}, C_1 2^{2/3})$ . By (31), (32),  $D_4 \leq C \cdot Cov(S_m/\sigma_m, S_n/\sigma_n)$ . Consequently

$$D_4 \ll Cov \left( S_m / \sigma_m, S_n / \sigma_n \right). \tag{33}$$

Let us estimate the covariance on the r.h.s. of (33). We have

$$Cov\left(\frac{S_m}{\sigma_m}, \frac{S_n}{\sigma_n}\right) = Cov\left(\frac{S_m}{\sigma_m}, \frac{S_{n+2m} - S_{2m}}{\sigma_n}\right) + E\left(\frac{S_m}{\sigma_m}\right)\left(\frac{S_n}{\sigma_n} - \frac{S_{n+2m} - S_{2m}}{\sigma_n}\right)$$
$$-E\left(\frac{S_m}{\sigma_m}\right)E\left(\frac{S_n}{\sigma_n} - \frac{S_{n+2m} - S_{2m}}{\sigma_n}\right) =: F_1 + F_2 + F_3. \tag{34}$$

We now estimate  $F_1$ - $F_3$  in (34). By (2), (9) and Karamata's theorem, we obtain

$$F_{1} \ll \frac{1}{L(m)^{1/2} m^{1-\alpha/2} L(n)^{1/2} n^{1-\alpha/2}} m \sum_{t=m+1}^{n+2m-1} \frac{L(t)}{t^{\alpha}} \\ \ll \frac{1}{L(m)^{1/2} m^{1-\alpha/2} L(n)^{1/2} n^{1-\alpha/2}} m(n+2m)^{1-\alpha} L(n+2m).$$
(35)

In order to bound  $F_2$  in (34), notice that  $F_2 \leq E \left| \frac{S_m}{\sigma_m} \right| \left| \frac{(S_{n+2m}-S_n)-S_{2m}}{\sigma_n} \right|$ . By the Cauchy-Schwarz inequality, the stationarity of the sequence  $\{X_i\}$  and the fact that  $EX_iX_j > 0$  for all i, j, we deduce that

$$F_2 \le \sqrt{E\left(\frac{S_m}{\sigma_m}\right)^2 E\left(\frac{(S_{n+2m} - S_n) - S_{2m}}{\sigma_n}\right)^2} \le \sqrt{2\frac{\sigma_{2m}^2}{\sigma_n^2}} = \sqrt{2\frac{\sigma_{2m}}{\sigma_n}}.$$
 (36)

This, relation (9) and definition of the slowly varying function at infinity imply

$$F_2 \ll \frac{\sigma_{2m}}{\sigma_n} \sim \frac{L \left(2m\right)^{1/2} \left(2m\right)^{1-\alpha/2}}{L \left(n\right)^{1/2} n^{1-\alpha/2}} \ll \frac{L \left(m\right)^{1/2} m^{1-\alpha/2}}{L \left(n\right)^{1/2} n^{1-\alpha/2}}.$$
 (37)

Since, in view of the stationarity of the sequence  $\{X_i\}$ , the last component  $F_3$  in (34) is equal to zero, then, by (33)-(35) and (37),

$$D_4 \ll \frac{m \left(n+2m\right)^{1-\alpha} L \left(n+2m\right)}{L \left(m\right)^{1/2} m^{1-\alpha/2} L \left(n\right)^{1/2} n^{1-\alpha/2}} + \frac{L \left(m\right)^{1/2} m^{1-\alpha/2}}{L \left(n\right)^{1/2} n^{1-\alpha/2}}.$$
 (38)

Finally, due to (26)-(28), (30) and (38), relation (22) is fulfilled in the case, when assumption (2) is satisfied with some  $0 < \alpha < 1$ .

**Proof of Lemma 2(b).** Suppose that (2) holds for some  $\alpha > 1$ . We have

$$\begin{aligned} |Cov \left( I \left\{ M_m \le u_m, S_m / \sigma_m \le y \right\}, I \left\{ M_{m,n} \le u_n, S_n / \sigma_n \le y \right\} \right)| \\ =: \mathbb{D}_1 + \mathbb{D}_2 + \mathbb{D}_3 + \mathbb{D}_4, \end{aligned}$$
(39)

where  $\mathbb{D}_1$ - $\mathbb{D}_4$  are defined as  $D_1$ - $D_4$  in (26), but for the case (2) holds for some  $\alpha > 1$ . By applying Theorem 4.2.1 in Leadbetter et al. (1983), it is easy to check that

$$\mathbb{D}_1 + \mathbb{D}_2 + \mathbb{D}_3 \ll 1/n^{\gamma} \text{ for some } \gamma > 0.$$
(40)

Thus, it remains to estimate the component  $\mathbb{D}_4$ . By (33)

$$\mathbb{D}_4 \ll Cov \left( S_m / \sigma_m, S_n / \sigma_n \right) =: \mathbb{F}_1 + \mathbb{F}_2 + \mathbb{F}_3, \tag{41}$$

with  $\mathbb{F}_1$ - $\mathbb{F}_3$ , defined as  $F_1$ - $F_3$  in (34), but for the case (2) holds with some  $\alpha > 1$ . As  $\mathbb{F}_1 = Cov\left(\frac{S_m}{\sigma_m}, \frac{S_{n+2m} - S_{2m}}{\sigma_n}\right) \le \frac{1}{\sigma_m \sigma_n} m\left(\sum_{t=m+1}^{\infty} \frac{L(t)}{t^{\alpha}} - \sum_{t=n+2m}^{\infty} \frac{L(t)}{t^{\alpha}}\right)$ , it follows from the facts that:  $\alpha > 1$ ,  $\sigma_i \ge i^{1/2}$ , and from Karamata's theorem

It follows from the facts that:  $\alpha > 1$ ,  $\sigma_i \ge i^{2/2}$ , and from Karamata's theorem that

$$\mathbb{F}_1 \ll (m^{1/2}/n^{1/2})(L(m)/m^{\alpha-1}) \ll m^{1/2}/n^{1/2}.$$
(42)

In order to estimate  $\mathbb{F}_2 = E\left(\frac{S_m}{\sigma_m}\right)\left(\frac{S_n}{\sigma_n} - \frac{S_{n+2m} - S_{2m}}{\sigma_n}\right)$ , notice that, as (2) holds for some  $\alpha > 1$ , then  $\sigma_{2m} \ll m^{1/2}$ . Since in addition,  $\sigma_n \ge n^{1/2}$ , we have, by (36),

$$\mathbb{F}_2 \ll \sigma_{2m} / \sigma_n \ll m^{1/2} / n^{1/2}.$$
(43)

As moreover,  $\mathbb{F}_3 = -E\left(\frac{S_m}{\sigma_m}\right) E\left(\frac{S_n}{\sigma_n} - \frac{S_{n+2m} - S_{2m}}{\sigma_n}\right) = 0$ , then, due to (41)-(43),  $\mathbb{D} = m^{1/2} / n^{1/2}$  (44)

$$\mathbb{D}_4 \ll m^{1/2}/n^{1/2}.$$
 (44)

Relations (39), (40), (44) imply (23) in the case (2) holds for some  $\alpha > 1$ . **Proof of Lemma 2(c).** Assume that (2) is satisfied with  $\alpha = 1$ . We have

$$|Cov \left(I \left\{ M_m \le u_m, S_m / \sigma_m \le y \right\}, I \left\{ M_{m,n} \le u_n, S_n / \sigma_n \le y \right\} \right)|$$
  
=:  $\mathcal{D}_1 + \mathcal{D}_2 + \mathcal{D}_3 + \mathcal{D}_4,$  (45)

where  $\mathcal{D}_1$ - $\mathcal{D}_4$  are defined as  $D_1$ - $D_4$  in (26), but for the case (2) holds with  $\alpha = 1$ . By using Theorem 4.2.1 in Leadbetter et al. (1983), it is easy to verify that

$$\mathcal{D}_1 + \mathcal{D}_2 + \mathcal{D}_3 \ll 1/n^{\gamma} \text{ for some } \gamma > 0.$$
(46)

Thus, it remains to estimate the component  $\mathcal{D}_4$ . By (33)

$$\mathcal{D}_4 \ll Cov \left( S_m / \sigma_m, S_n / \sigma_n \right) =: \mathcal{F}_1 + \mathcal{F}_2 + \mathcal{F}_3, \tag{47}$$

with  $\mathcal{F}_1$ - $\mathcal{F}_3$  defined as  $F_1$ - $F_3$  in (34), but for the case, when (2) holds with  $\alpha = 1$ . We have

$$\mathcal{F}_{1} = \frac{1}{\sigma_{m}\sigma_{n}}Cov\left(S_{m}, S_{n+2m} - S_{2m}\right) \le \frac{1}{\sigma_{m}\sigma_{n}}m\sum_{t=m+1}^{n+2m-1}r\left(t\right).$$
 (48)

By the same reasoning as in the excerpt of the proof of Theorem 1 in Mielniczuk (2002) (see p. 396, lines 7-15), we obtain

$$\sigma_n^2 = n + 2\sum_{t=1}^{n-1} (n-t) r(t) \sim n\left(1 + 2\sum_{t=1}^{n-1} r(t)\right) = n\widetilde{L}(n).$$
(49)

By definition of  $\overset{\sim}{L}(n)$ , we have  $m \sum_{t=m+1}^{n+2m-1} r(t) = m \left( \overset{\sim}{L}(n+2m) - \overset{\sim}{L}(m+1) \right) / 2$ . This and relations (48), (49) yield

$$\mathcal{F}_{1} \ll \frac{1}{m^{1/2}\widetilde{L}(m)^{1/2} n^{1/2} \widetilde{L}(n)^{1/2}} m\left(\widetilde{L}(n+2m) - \widetilde{L}(m+1)\right).$$
(50)

In turn, by (36),  $\mathcal{F}_2 = E\left(\frac{S_m}{\sigma_m}\right)\left(\frac{S_n}{\sigma_n} - \frac{S_{n+2m} - S_{2m}}{\sigma_n}\right) \leq E\left|\frac{S_m}{\sigma_m}\right| \left|\frac{(S_{n+2m} - S_n) - S_{2m}}{\sigma_n}\right| \ll \sigma_{2m}/\sigma_n$ . Hence, from (49) and the fact that  $\stackrel{\sim}{L}(\cdot)$  is slowly varying, we get

$$\mathcal{F}_2 \ll \left( m^{1/2} \widetilde{L}\left(m\right)^{1/2} \right) / \left( n^{1/2} \widetilde{L}\left(n\right)^{1/2} \right).$$
(51)

As moreover,  $F_3 = 0$ , then, by (47), (50) and (51),

$$\mathcal{D}_{4} \ll \frac{m^{1/2} \left(\widetilde{L} \left(n+2m\right) - \widetilde{L} \left(m+1\right)\right)}{n^{1/2} \widetilde{L} \left(m\right)^{1/2} \widetilde{L} \left(n\right)^{1/2}} + \frac{m^{1/2} \widetilde{L} \left(m\right)^{1/2}}{n^{1/2} \widetilde{L} \left(n\right)^{1/2}}.$$
 (52)

Relations (45), (46), (52) imply (24) in the case, when (2) is satisfied with  $\alpha = 1$ . This completes the whole proof of Lemma 2.

The following lemma will be also used in our further considerations.

**Lemma 3** (for comparison - see also Ho and Hsing (1996)). Under the assumptions of Theorem 1 on  $X_1, X_2, ..., and r(t), \{u_n\}$ , we have

$$\lim_{n \to \infty} P\left\{M_n \le u_n, S_n / \sigma_n \le y\right\} = e^{-\tau} \Phi\left(y\right) \text{ for all } y \in \mathbb{R} \text{ and } \tau \text{ defined in (3).}$$

**Proof.** Notice that, as  $X_1, X_2, ...$  is a standard normal sequence, r(t),  $\{u_n\}$  satisfy (2), (3), respectively, then, by Theorem 4.3.3 in Leadbetter et al. (1983),

$$\lim_{n \to \infty} P\left\{M_n \le u_n\right\} = e^{-\tau} \text{ for } \tau \text{ defined in (3).}$$
(53)

Let y be an arbitrary real number and let, for each  $n \geq 1$ ,  $Y_n$  denote a random variable, which has the same distribution as  $S_n/\sigma_n$ , but is independent of  $(X_1, ..., X_n)$ . From the estimations of  $A_1$ ,  $A_1$  in the proof of Lemma 1,  $|P\{M_n \leq u_n, S_n/\sigma_n \leq y\} - P\{M_n \leq u_n\} P\{Y_n \leq y\}| \ll 1/n^{\gamma}$  for some  $\gamma > 0$ . As: (53) holds,  $1/n^{\gamma} \to 0$ , as  $n \to \infty$ , and, for each n,  $Y_n$  is standard normal, we get  $\lim_{n\to\infty} P\{M_n \leq u_n, S_n/\sigma_n \leq y\} = \lim_{n\to\infty} P\{M_n \leq u_n\} P\{Y_n \leq y\} = e^{-\tau}\Phi(y)$ , which yields the desired result.

#### 4. Proof of main result

In this section, we present the proof of Theorem 1. In our proof, we make an extensive use of the earlier proved Lemmas 1-3.

**Proof of Theorem 1(i).** First, we will show that for any real y

$$\frac{1}{\log N} \sum_{n=1}^{N} \frac{1}{n} \left( I \left\{ M_n \le u_n, S_n / \sigma_n \le y \right\} - P \left\{ M_n \le u_n, S_n / \sigma_n \le y \right\} \right) \xrightarrow{\text{a.s.}} 0, \quad (54)$$

as  $N \to \infty$ . By Lemma 3.1 in Csaki and Gonchigdanzan (2002), in order to prove (54), it suffices to show that the following property occurs for some  $\varepsilon > 0$ 

$$Var\left(\sum_{n=1}^{N} \frac{1}{n} I\left\{M_n \le u_n, S_n/\sigma_n \le y\right\}\right) \ll \left(\log N\right)^2 \left(\log\log N\right)^{-(1+\varepsilon)}.$$
 (55)

Put  $\xi_n := I \{ M_n \leq u_n, S_n / \sigma_n \leq y \}$ . We have

$$Var\left(\sum_{n=1}^{N} \frac{1}{n} I\left\{M_{n} \le u_{n}, S_{n}/\sigma_{n} \le y\right\}\right) = Var\left(\sum_{n=1}^{N} \frac{1}{n}\xi_{n}\right)$$
$$\le \sum_{n=1}^{N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{1 \le m < n \le N} \frac{1}{mn} |Cov\left(\xi_{m}, \xi_{n}\right)| =: \sum_{n=1}^{N} \sum_{n=1}^{N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le n \le N} \frac{1}{mn} |Cov\left(\xi_{n}, \xi_{n}\right)| =: \sum_{n \le N} \sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} |Cov\left(\xi_{n}, \xi_{n}\right)| =: \sum_{n \le N} \sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} |Cov\left(\xi_{n}, \xi_{n}\right)| =: \sum_{n \le N} \frac{1}{n^{2}} \left(\sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} \left(\sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} \left(\sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} \left(\sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} \left(\sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} \left(\sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} \left(\sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} \left(\sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} \left(\sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} \left(\sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \le N} \frac{1}{n^{2}} \left(\sum_{n \le N} \frac{1}{n^{2}} Var\left(\xi_{n}\right) + \sum_{n \ge$$

Since  $\xi_n^2 \leq 1$ , we obtain that

$$\sum_{1} \le \sum_{n=1}^{\infty} \frac{1}{n^2} < \infty.$$
(57)

Thus, we only need to estimate  $\sum_{2}$  in (56). Let m < n. Observe that

$$\begin{aligned} |Cov(\xi_m,\xi_n)| &\leq 2E \left| I \left\{ M_n \leq u_n, S_n/\sigma_n \leq y \right\} - I \left\{ M_{m,n} \leq u_n, S_n/\sigma_n \leq y \right\} | \\ &+ \left| Cov(I \left\{ M_m \leq u_m, S_m/\sigma_m \leq y \right\}, I \left\{ M_{m,n} \leq u_n, S_n/\sigma_n \leq y \right\} ) \right|. \end{aligned}$$

This and Lemmas 1, 2 imply that: (a) if condition (2) is satisfied with some  $0 < \alpha < 1$ , then for some  $\gamma > 0$ 

$$|Cov(\xi_m,\xi_n)| \ll \frac{1}{n^{\gamma}} + \frac{m}{n} + \frac{m(n+2m)^{1-\alpha}L(n+2m)}{L(m)^{1/2}m^{1-\alpha/2}L(n)^{1/2}n^{1-\alpha/2}} + \frac{L(m)^{1/2}m^{1-\alpha/2}}{L(n)^{1/2}n^{1-\alpha/2}},$$
(58)

(b) if condition (2) is satisfied with some  $\alpha > 1$ , then for some  $\gamma > 0$ 

$$Cov(\xi_m,\xi_n)| \ll 1/n^{\gamma} + m^{1/2}/n^{1/2},$$
(59)

(c) if condition (2) is satisfied with  $\alpha = 1$ , then for some  $\gamma > 0$ 

$$|Cov(\xi_m,\xi_n)| \ll \frac{1}{n^{\gamma}} + \frac{m}{n} + \frac{m^{1/2} \left(\widetilde{L}(n+2m) - \widetilde{L}(m+1)\right)}{n^{1/2} \widetilde{L}(m)^{1/2} \widetilde{L}(n)^{1/2}} + \frac{m^{1/2} \widetilde{L}(m)^{1/2}}{n^{1/2} \widetilde{L}(n)^{1/2}}.$$
(60)

Let us consider the case (a) first. By (56) and (58), we get

$$\sum_{2} \ll \sum_{1 \le m < n \le N} \frac{1}{mn} \frac{1}{n^{\gamma}} + \sum_{1 \le m < n \le N} \frac{1}{mn} \frac{m}{n}$$
$$+ \sum_{1 \le m < n \le N} \frac{1}{mn} \frac{m \left(n + 2m\right)^{1-\alpha} L\left(n + 2m\right)}{L\left(m\right)^{1/2} m^{1-\alpha/2} L\left(n\right)^{1/2} n^{1-\alpha/2}} + \sum_{1 \le m < n \le N} \frac{1}{mn} \frac{L\left(m\right)^{1/2} m^{1-\alpha/2}}{L\left(n\right)^{1/2} n^{1-\alpha/2}}$$
$$=: \sum_{3} + \sum_{4} + \sum_{5} + \sum_{6} \ll \log N + \sum_{5} + \sum_{6}, \qquad (61)$$

where the last relation follows from the obvious fact that  $\sum_3 + \sum_4 \ll \log N$ . Thus, it remains to estimate  $\sum_5$ ,  $\sum_6$  in (61). It is easy to check that

$$\sum_{5} \ll \sum_{m=1}^{N-1} \frac{1}{L(m)^{1/2} m^{1-\alpha/2}} \sum_{n=m+1}^{N} \left(1 + \frac{2m}{n}\right)^{1-\alpha} \frac{L(n+2m)}{L(n)^{1/2}} \frac{1}{n^{1+\alpha/2}}.$$
 (62)

For a fixed m, put  $\overline{L}(n) := L(n+2m)/L(n)^{1/2}$ . By using (62) and the same reasoning as in the excerpt of the proof of Theorem 1 in Mielniczuk (2002) (see p. 395, lines 7-14), we obtain

$$\sum_{5} \ll \sum_{m=1}^{N-1} \frac{\bar{L}(m) m^{-\alpha/2}}{L(m)^{1/2} m^{1-\alpha/2}} = \sum_{m=1}^{N-1} \frac{\bar{L}(m)}{L(m)^{1/2}} \frac{1}{m} = \sum_{m=1}^{N-1} \frac{L(3m)}{L(m)} \frac{1}{m} \ll \log N.$$
(63)

In addition, we can estimate  $\sum_{6}$  in (61) as follows. Observe that, by the facts that  $L(\cdot)$  is slowly varying and  $0 < \alpha < 1$ , we have

$$\sum_{6} \ll \sum_{m=1}^{N-1} \frac{L(m)^{1/2}}{m^{\alpha/2}} \sum_{n=m+1}^{\infty} \frac{1}{L(n)^{1/2} n^{2-\alpha/2}} \ll \sum_{m=1}^{N-1} \frac{L(m)^{1/2}}{m^{\alpha/2}} \frac{1}{L(m)^{1/2} m^{1-\alpha/2}}$$

Hence  $\sum_{6} \ll \log N$ . This and (61), (63) yield

$$\sum_{2} \ll \log N. \tag{64}$$

Due to (56), (57) and (64), we conclude that (55) holds for arbitrary real y and any positive  $\varepsilon$ . Consequently, by the already mentioned Lemma 3.1 in Csaki and Gonchigdanzan (2002), condition (54) is satisfied as well. This, Lemma 3 and the regularity property of logarithmic means imply the statement (i) of Theorem 1 for the case, when (2) is satisfied with some  $0 < \alpha < 1$ .

We now consider the case (b), i.e. the case, when (2) holds with some  $\alpha > 1$ . By (56) and (59), we get  $\sum_{2} \ll \sum_{1 \le m < n \le N} \frac{1}{mn} \frac{1}{n^{\gamma}} + \sum_{1 \le m < n \le N} \frac{1}{mn} \frac{m^{1/2}}{n^{1/2}} \ll \log N$ . This, (56) and (57) imply that condition (55) holds for arbitrary real y and any positive  $\varepsilon$ . Therefore (54) occurs. Hence, also in the case, when (2) is satisfied

with some  $\alpha > 1$ , the statement (i) of Theorem 1 is true. Finally, let us consider the case (c), assuming that (2) holds with  $\alpha = 1$ . It follows from (56) and (60) that

$$\sum_{2} \ll \sum_{1 \le m < n \le N} \frac{1}{mn} \frac{1}{n^{\gamma}} + \sum_{1 \le m < n \le N} \frac{1}{mn} \frac{m}{n}$$

$$+ \sum_{1 \le m < n \le N} \frac{1}{mn} \frac{m^{1/2} \left(\widetilde{L} \left(n + 2m\right) - \widetilde{L} \left(m + 1\right)\right)}{n^{1/2} \widetilde{L} \left(m\right)^{1/2}} + \sum_{1 \le m < n \le N} \frac{1}{mn} \frac{m^{1/2} \widetilde{L} \left(m\right)^{1/2}}{n^{1/2} \widetilde{L} \left(n\right)^{1/2}}$$

$$=: \sum_{7} + \sum_{8} + \sum_{9} + \sum_{10} \ll \log N + \sum_{9} + \sum_{10} \ldots$$
(65)

Let us estimate the term  $\sum_{9}$  in (65). By reasoning identically as in Mielniczuk Let us estimate the term  $\sum_{9} m$  (05). By reasoning identically as in Memiczuk (2002), p. 397, we get  $\sum_{9} = \sum_{m=1}^{N-1} \frac{1}{m^{1/2}\widetilde{L}(m)^{1/2}} \sum_{n=m+1}^{N} \frac{\widetilde{L}(n+2m) - \widetilde{L}(m+1)}{n^{3/2}\widetilde{L}(n)^{1/2}} \ll \sum_{m=1}^{N-1} \frac{1}{m^{1/2}\widetilde{L}(m)^{1/2}} \frac{\widetilde{L}(3m) - \widetilde{L}(m+1)}{m^{1/2}\widetilde{L}(m)^{1/2}} = \sum_{m=1}^{N-1} \frac{\widetilde{L}(3m) - \widetilde{L}(m+1)}{m\widetilde{L}(m)} \ll \log N.$ In order to estimate  $\sum_{10}$  in (65), observe that, as  $\widetilde{L}(\cdot)$  is slowly varying,

$$\sum_{10} \ll \sum_{m=1}^{N-1} \frac{L(m)^{1/2}}{m^{1/2}} \sum_{n=m+1}^{N} \frac{1}{n^{3/2} \widetilde{L}(n)^{1/2}} \ll \sum_{m=1}^{N-1} \frac{L(m)^{1/2}}{m^{1/2}} \frac{1}{m^{1/2} \widetilde{L}(m)^{1/2}} \ll \log N.$$
The estimates for  $\sum_{n=m+1}^{N} \sum_{m=1}^{N-1} \frac{1}{n^{3/2} \widetilde{L}(n)^{1/2}} \ll \log N.$ 

The estimates for  $\sum_{9}$ ,  $\sum_{10}$ , together with (65), yield  $\sum_{2} \ll \log N$ , which, by (56), (57), implies that (55) holds for arbitrary real y and any  $\varepsilon > 0$ . Therefore (54) occurs. Thus, (4) holds also in the case (2) is satisfied with  $\alpha = 1$ .

**Proof of Theorem 1(ii).** It is easily seen that Theorem 1(ii) is a special case of the earlier proved Theorem 1(i), with:  $u_n := x/a_n + b_n, \tau := e^{-x}$ . Acknowledgement. I thank an anonymous Referee for valuable remarks. References

Bagai, I., Prakasa Rao, B.L.S., 1991. Estimation of the survival function for stationary associated processes, Statist. Probab. Lett. 12, 385-391.

Csaki, E., Gonchigdanzan, K., 2002. Almost sure limit theorems for the maximum of stationary Gaussian sequences, Statist. Probab. Lett. 58, 195-203.

Dudziński, M., 2003. An almost sure limit theorem for the maxima and sums of stationary Gaussian sequences, Probab. Math. Statist. 23, 139-152.

Ho, H.C., Hsing, T., 1996. On the asymptotic joint distribution of the sum and maximum of stationary normal random variables, J. Appl. Probab. 33, 138-145. Leadbetter, M.R., Lindgren, G., Rootzen, H., 1983. Extremes and related properties of random sequences and processes, Springer, New York.

Mielniczuk, J., 2002. Some remarks on the almost sure central limit theorem for dependent sequences, in: Limit theorems in Probability and Statistics, Vol. 2, Janos Bolyai Math. Soc., Budapest, 391-403.

Newman, C.M., 1984. Asymptotic independence and limit theorems for positively and negatively dependent random variables, in: Inequalities in Statistics and Probability, Vol. 5, IMS Lecture Notes-Monograph Series, 127-140.

Pitt, L.D., 1982. Positively correlated normal variables are associated. Ann. Probab. 10, 496-499.

Resnick, S., 1987. Extreme values, regular variation and point processes, Springer, New York.