



**HAL**  
open science

## The role of population PK-PD modelling in paediatric clinical research

Roosmarijn F. W. Cock, Chiara Piana, Elke H. J. Krekels, Meindert Danhof,  
Karel Allegaert, Catherijne A. J. Knibbe

► **To cite this version:**

Roosmarijn F. W. Cock, Chiara Piana, Elke H. J. Krekels, Meindert Danhof, Karel Allegaert, et al..  
The role of population PK-PD modelling in paediatric clinical research. *European Journal of Clinical Pharmacology*, 2010, 67 (S1), pp.5-16. 10.1007/s00228-009-0782-9 . hal-00594491

**HAL Id: hal-00594491**

**<https://hal.science/hal-00594491>**

Submitted on 20 May 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**The role of population PK-PD modeling in paediatric clinical research**

Journal:	<i>European Journal of Clinical Pharmacology</i>
Manuscript ID:	EJCP-2009-0223.R2
Type of submission:	Review Article
Date Submitted by the Author:	21-Dec-2009
Complete List of Authors:	De Cock, Roosmarijn; Leiden University (Leiden/Amsterdam Centre for Drug Research), Pharmacology Piana, Chiara; Leiden University (Leiden/Amsterdam Centre for Drug Research), Pharmacology Krekels, Elke; Leiden University (Leiden/Amsterdam Centre for Drug Research), Pharmacology; Erasmus MC - Sophia Children's Hospital, Pediatric Surgery Danhof, Meindert; Leiden University (Leiden/Amsterdam Centre for Drug Research), Pharmacology Allegaert, Karel; University Hospital Leuven, Neonatal Intensive Care Unit Knibbe, Catherijne; St Antonius Hospital, Clinical Pharmacy; Leiden University (Leiden/Amsterdam Centre for Drug Research), Pharmacology; Erasmus MC - Sophia Children's Hospital, Pediatric Surgery



**The role of population PK-PD modeling in paediatric clinical research**

*Roosmarijn F. W. De Cock<sup>1</sup>, Chiara Piana<sup>1</sup>, Elke H.J. Krekels<sup>1,2</sup>, Meindert Danhof<sup>1</sup>,  
Karel Allegaert<sup>4</sup>, Catherijne A.J. Knibbe<sup>1,2,3</sup>*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
Division of Pharmacology, Leiden/Amsterdam Centre for Drug Research,  
Leiden University,  
Leiden, the Netherlands

2  
Department of Pediatric Intensive Care  
Erasmus MC - Sophia Children's Hospital  
Rotterdam, the Netherlands

3  
Department of Clinical Pharmacy  
St. Antonius Hospital  
P.O. Box 2500  
3430 EM Nieuwegein, the Netherlands  
Email: [c.knibbe@antoniuziekenhuis.nl](mailto:c.knibbe@antoniuziekenhuis.nl)

4  
Neonatal Intensive Care Unit  
University Hospital Leuven, Belgium

RDC and CP equally contributed to this paper

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Children differ from adults in their response to drugs. While this may be the result of changes in dose-exposure (pharmacokinetics (PK)) and/or exposure-response (pharmacodynamics (PD)) relationships, the magnitude of these changes may not be solely reflected by differences in bodyweight. As a consequence, dosing recommendations empirically derived from adults dosing regimens using linear extrapolations based on bodyweight, can result in therapeutic failure, occurrence of adverse effect or even fatalities. In order to define rational, patient tailored dosing schemes, population PK-PD studies in children are needed. For the analysis of the data, population modeling using non-linear mixed effect modeling is the preferred tool since this approach allows for the analysis of sparse and unbalanced datasets. Additionally it permits the exploration of the influence of different covariates such as bodyweight, age and other covariates, to explain the variability in drug response. Finally, using this approach, these PK-PD studies can be designed in the most efficient manner in order to obtain the maximum information on the PK-PD parameters with the highest precision. Once a population PK-PD model is developed, internal and external validations should be performed. If the model performs well in these validation procedures, model simulations can be used to define a dosing regimen which in turn needs to be tested and challenged in a prospective clinical trial. This methodology will improve the efficacy/safety balance of dosing guidelines which will be of benefit to the individual child.

Keywords: Pharmacokinetics – Pharmacodynamics – Population modeling – Paediatric population – Validation

1  
2  
3 Introduction:

4 Children differ from adults in their response to drugs. These differences may be caused  
5 by changes in the pharmacokinetics (PK) and/or pharmacodynamics (PD) between  
6 children and adults and may also vary between children of different ages. The PK of a  
7 drug includes processes of absorption, distribution, metabolism and elimination of a drug  
8 whereas the PD comprises the physiological and biological response to the administered  
9 drug and therefore may represent both efficacy and safety measures. While a child grows,  
10 enzyme pathways (involved in the PK), function and expression of receptors and proteins  
11 (involved in the PD) mature, which can be referred to as 'developmental changes' in  
12 childhood. The maturation rates of these developmental changes vary however between  
13 the pathways and receptors and often do not correlate solely with the increase in  
14 bodyweight of the child. The question is therefore how to obtain data in children that  
15 allow for the study of these developmental changes ultimately resulting in evidence based  
16 dosing regimens for drugs in children.  
17  
18  
19  
20  
21  
22  
23

24  
25 To date, only a small number of drugs used in children is licensed for use in this specific  
26 group. Up to 70% of the drugs in paediatric intensive care, and 90% of the drugs in  
27 neonatal intensive care, are prescribed in an off-label or unlicensed manner [1-4].  
28 Paediatric dosing regimens are usually empirically derived from adult regimens using  
29 linear extrapolations based on bodyweight. Since these developmental changes are non-  
30 linear dynamic processes, this dosing paradigm may result in under or over-dosing  
31 particularly in specific age groups. This may cause therapeutic failure, occurrence of  
32 severe adverse effects or even fatalities such as fatalities occurring after long-term  
33 sedation with high doses of propofol [5, 6] and occurrence of the grey baby syndrome in  
34 neonates after treatment with chloramphenicol [7, 8]. As a result, dose adjustments in the  
35 younger age groups are often proposed. For vancomycin for example lower doses are  
36 administered in neonates younger than 1 week (20 mg/kg/day) compared to 1-4 week-old  
37 neonates (30 mg/kg/day) and children between 1 month and 18 years (40 mg/kg/day) [9].  
38  
39  
40  
41  
42  
43  
44  
45

46 Instead of the *a priori* use of bodyweight for dosing guidelines in children, detailed  
47 information on PK and potentially also the PD needs to be considered in order to define  
48  
49  
50

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

effective and safe dosing regimens throughout the paediatric age range. The lack of PK and PD information on drugs in children has led to the European Regulation which entered into force in 2007. This law imposes pharmaceutical companies to perform research in the whole paediatric age-range for all drugs that are developed for the European market, by requiring the submission of a paediatric investigational plan (PIP) in the early stages of the development of a new drug. In this PIP, a full description has to be given of the studies and of drug formulation in the paediatric population. In case little information is available about efficacy and safety of a drug, studies in children are only performed after more information is obtained in the adult population to increase the safety of the paediatric study [10-12]. The main targets of introducing the paediatric Regulation were to facilitate development and availability of medicines in children between 0 and 17 years, to improve the availability of information about medicines used in children, to ensure that the medicines are of high quality, can be administered in a safe and effective way and that paediatric studies are performed in an ethically correct way [10]. The reward for this effort is a six month supplementary production certificate for the pharmaceutical company.

Both for industry and for academic researchers, performing (PK-PD) studies in children in order to develop rational dosing schemes is very challenging because of ethical and practical issues. Unlike studies in healthy adults, research in healthy children is considered to be unethical, so all paediatric studies are performed in the vulnerable group of children suffering from a disease. In all clinical trials, an informed consent has to be signed by the patient before he or she can be enrolled into a trial. In paediatric trials, this informed consent can not be obtained by the patient that participates in the trial, and is therefore replaced by the consent of the parents or guardians. In older age groups, in addition to this consent, an assent is used in which the aim of the study is explained in an age-appropriate language so that children can understand. [1, 4, 13].

Apart from ethical issues, practical challenges also occur when performing studies in children. There are limitations to the number and volume of samples that can be obtained, resulting in infrequent sampling possibilities and the need for advanced drug assay techniques with improved sensitivity. Another complicating factor is the limited available

1  
2 number of subjects that suffers from the same disease. Finally, pharmacodynamic  
3 endpoints that measure the efficacy of the drug, and which are validated for children may  
4 be lacking. All these factors call for highly advanced study designs and analysis  
5 techniques so that the burden for each child can be kept to a minimum while still  
6 addressing all the study objectives.  
7  
8  
9

10  
11 This paper aims to inform clinical pharmacologists, paediatricians and pharmacists about  
12 population PK-PD modeling in paediatric drug research. Advanced statistical tools are  
13 discussed that can be used to develop rational dosing schemes based on the PK and PD of  
14 a drug in children, despite practical and ethical restrictions. Using these tools, covariates  
15 can be identified in order to define appropriate doses and dosing intervals based on  
16 individual characteristics of each child with minimal burden to each patient. The paper  
17 also describes how to evaluate the predictive performance of the models by different  
18 validation methods including a prospective clinical trial. Ultimately, the efforts result in  
19 an individualized dosing regimen based on the PK-PD relation through the paediatric  
20 age-range.  
21  
22  
23  
24  
25  
26

#### 27 28 PK-PD in children

29  
30 Developmental changes in childhood can affect all PK processes from absorption until  
31 elimination as well as the pharmacodynamic effects. For example, in neonates intra-  
32 gastric pH is elevated (>4) which may increase the bioavailability of acid-labile  
33 compounds (penicillin G) and decrease the bioavailability of weak acids (phenobarbital)  
34 when given orally [14]. Additionally, gastric emptying in neonates is delayed, which  
35 means that also the absorption of drugs e.g. paracetamol is slower in neonates [15, 16].  
36 Other examples are changes in metabolizing enzyme capacity in children. Although most  
37 uridine 5'-diphosphate (UDP)-glucuronosyltransferases (UGTs) and P-450 cytochromes  
38 (CYPs) are expressed during the first week of life, the activity at birth in comparison with  
39 adults is often low, e.g. UGT2B7 activity at birth is around 10% of the adult level and  
40 maturation rates of different enzyme systems are known to mature at different rates [14,  
41 17-20].  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

In addition, renal function and liver flow are influenced by physiological changes depending on age, e.g. the glomerular filtration rate in mL/min/70kg in full term neonates is 35% of the adult value, while mL/min/70kg adult values are reached at approximately 1 year old [21]). When using units of mL/min/70kg however, it should be realized that actual values of GFR in children are still very low compared to adult values because of correction for differences in total body weight between adults and infants.

Furthermore the body composition of children changes continuously resulting in an age-dependent proportion of body water and fat, which influences the distribution of drugs. For example, the total amount of body water (80-90 % of the bodyweight) is higher in neonates compared to adults (55-60%). Hydrophilic drugs like aminoglycosides have a larger volume of distribution in neonates which can be explained by larger extra-cellular fluid (45% of the bodyweight) compared to adults (20%) [14, 22].

In order to characterize the specific influence of developmental changes in childhood on the PK of a drug, concentration-time profiles are necessary, which require measurements of drug concentrations. For ethical reasons, in paediatric studies, discomfort, like pain and anxiety associated with venipuncture, must be restricted and practical issues limit the volume and amount of blood samples that can be obtained. Therefore, sensitive analysis techniques requiring only small blood samples should be used. While HPLC methods have reported to require only 50  $\mu$ L of blood [23], more recently LC-MS methods can measure up to ten different drugs in volumes as low as 50-100  $\mu$ L [24]. Additionally, also alternative matrices such as saliva should be explored as a non-invasive, more child-friendly alternative to measure a drug concentration. An example in this respect is a LC-MS/MS method which was developed and validated for the measurement of busulphan in saliva [24]. Also the use of a dried blood spot method e.g. for tacrolimus can facilitate the measurement of drugs in children [25]. Another method is capillary electrophoresis which requires only a low sample volume for the quantification of drugs in biological fluids [26].

Changes between children and adults may also result from differences in the pharmacodynamics of a drug in children, e.g. by changes in the relative number and function of receptors. These age-related PD differences are until present rarely reported in literature, but one of the few examples is the increased sensitivity to d-tubocurarine, an



1  
2 antagonist of nicotinic neuromuscular acetylcholine receptors, in neonates and infants  
3 compared to children and adults [27]. Other examples are the observed lower minimum  
4 alveolar concentration (MAC) of isoflurane in preterm neonates compared to full-term  
5 neonates and older children [28, 29] and the different sensitivity to bronchodilators  
6 because of the lack of smooth muscles in the airways in neonates [30].  
7  
8  
9

10 To study the PD of a drug in children, the use of a PD endpoint which is validated for use  
11 in children is a prerequisite. An illustrative example is the measurement of pain in young  
12 children. Since they are not able to report their pain using a visual analogue scale, an  
13 observational scale has been developed. This comfort behavioral (COMFORT-B) scale  
14 was developed and validated for use in children under the age of three years [31]. The  
15 scale assesses six behavioral items: alertness, calmness, muscle tone, body movement,  
16 facial tension, and crying (non-ventilated children) or respiratory response (ventilated  
17 children). All items range from 1 (no distress) to 5 (severe distress), resulting in a total  
18 score of varying from 6 to 30. This validated scale can then be used as a PD endpoint for  
19 the development of PD models for pain and/or sedation in children of different ages [32-  
20 34].  
21  
22  
23  
24  
25  
26

27 The influence of covariates such as the developmental changes, disease status and  
28 genetics on the PK and PD of drugs in children is depicted in Figure 1.  
29  
30  
31  
32

33 When both the PK and PD of a drug in children are characterized, the developed models  
34 can be used to derive rational dosing regimens with predictable efficacy and  
35 concentration profiles. An example of such a PK-PD model with a derived dosing  
36 regimen is an article published by Peeters *et al.* In this paper both the PK and the PD  
37 were characterized in children, the latter with the use of the COMFORT-B scale as  
38 pharmacodynamic endpoint [33]. Based on the model it was found that propofol  
39 clearance is two times higher in non-ventilated children compared to ventilated children  
40 and adults. For the PD, a model was derived in which an effect of propofol was  
41 characterized within a naturally occurring sleep pattern of children in the ICU. Both  
42 models (PK as well as PD) were used to simulate concentrations as well as the effects  
43 that could be expected using different dosing schemes (Figure 2). As a result, based on  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 this PK-PD model, a propofol dose of 30mg/h was recommended for a child of 10 kg  
4 which will result in adequate COMFORT-B scales in the night following craniofacial  
5 surgery.  
6  
7

#### 8 9 Methods to analyse data: standard two-stage or population approach

10  
11 When concentration-time and concentration-effect datasets obtained in children are  
12 considered for analysis, two different methods can be applied: the standard two-stage  
13 approach and the population approach using non-linear mixed effect models [35-38].  
14 When using the standard two-stage approach or classical approach, in a first step  
15 parameters are estimated in each individual based on individual concentration-time  
16 profiles (figure 3A). In a second step, these parameters are summarized by calculating the  
17 mean or median of the parameters and the variability between subjects (SE or IQR). A  
18 major drawback of this methodology is that this approach requires a relatively high  
19 number of samples in each individual patient (Figure 3A) while each patient has to  
20 contribute roughly the same number of samples. Moreover it is very difficult to  
21 distinguish between inter-individual (variability between subjects), intra-individual and  
22 residual variability (variability within one subject, measurement error, and model  
23 misspecification) and as a result inter-individual variability is often overestimated [39].  
24 Since usually only a limited number of observations can be obtained in paediatric  
25 subjects, the population approach using non-linear mixed effect modeling to obtain PK  
26 and PD parameters, is the preferred approach [37]. The population approach differs from  
27 the standard two stage-approach in the fact that the analysis is based on simultaneous  
28 analysis of all data of the entire population while still taking into account that different  
29 observations come from different patients (Figure 3B). Additionally the population  
30 approach allows not only for the analysis of dense data but also for sparse (limited  
31 number of observations per individual) and unbalanced data (unequal distribution of  
32 observations in various parts of the concentration-time profile in the individuals) or a  
33 combination of both. Finally both the interindividual and intra-individual variability are  
34 separately estimated in the dataset using this approach.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

47 As a result of this methodology, when designing a paediatric study of which the data will  
48 be analyzed using the population approach, it is advisable to collect samples at different  
49  
50

times (or time-windows) or to set alternating sampling schemes in subgroups of patients. This also means that (part of the) samples can be collected during routine clinical sampling. Consequently, the burden for the child that participates in the trial is reduced and the statistical power to develop a model describing the concentration-time or concentration-effect profile is not affected or improved.

The term 'mixed' in non-linear mixed effects modeling stands for a mixture of fixed and random effects. For the fixed effects, a structural model describing the PK or PD is chosen (e.g. a two-compartment model for PK or an  $E_{\max}$  model for PD). The random effects quantify the variability that is not explained by the fixed effects. These random effects include inter-subject and intra-subject and random variability (Figure 4), which are both simultaneously and separately estimated. It is often assumed that the variability between subjects follows a normal distribution with a mean of zero and variance  $\omega^2$ . Equation 1 is used to describe the relationship between individual and population parameter estimates.

$$\theta_i = \theta_{mean} \cdot e^{\eta_i} \quad (\text{equation 1})$$

where  $\theta_i$  represents the parameter of the  $i$ th subject,  $\theta_{mean}$  the population mean, and  $\eta_i$  the variability between subjects. The residual error is in generally described using a proportional error (error is dependent on the concentration, which means a higher absolute error at higher concentrations (equation 2)) or additive error (constant for all observations (equation 3)) or a combination of both. This means for the  $j$ th observed concentration of the  $i$ th individual the relation ( $Y_{ij}$ ):

$$Y_{ij} = c_{pred,ij} \cdot (1 + \varepsilon_{ij}) \quad (\text{equation 2})$$

$$Y_{ij} = c_{pred,ij} + \varepsilon_{ij} \quad (\text{equation 3})$$

where  $c_{pred}$  is predicted concentration and  $\varepsilon_{ij}$  is a random variable with mean zero and variance  $\sigma^2$ .

1  
2  
3  
4 In general, model building requires three different steps. First a structural model (fixed  
5 effects) has to be designed, then a statistical sub-model (random effects) has to be  
6 developed and in the final step a covariate sub-model is identified.

7  
8 The structural model describes the overall trend in the data. The choice of structural  
9 model (e.g. one, two or three-compartment model for PK and an  $E_{\max}$  model for PD) is to  
10 be based upon the best *a priori* information about the drug to be studied [40]. The  
11 structural model uses fixed effects parameters such as clearance and volume of  
12 distribution for PK or  $E_{\max}$  and  $EC_{50}$  for PD. The population values for these parameters  
13 are called typical values (TV).

14  
15 After selecting the structural model, the statistical sub-model which accounts for the  
16 inter-individual as well as the residual variability is chosen and tested. Information on  
17 both inter- and intra- and residual variability is of clinical value, because it describes  
18 differences in clinical response between and within patients and may therefore provide  
19 guidance to rational dose adjustments. With the population approach, both these random  
20 effects are obtained, apart from estimates of both the population values (TV) and the  
21 individual values of PK and PD parameters (so called *post hoc* parameter estimates).

22  
23 In the final step the covariate sub-model is determined which expresses relationships  
24 between covariates and parameters of the structural model (e.g. influence of bodyweight  
25 on volume of distribution or clearance). Covariates can be individual-specific (age,  
26 bodyweight, genetic profile, etc) or time-varying (renal function, hemodynamic  
27 parameters, body temperature etc). The covariate analysis will be explained more in  
28 details in the following section.

29  
30 As these three models are interrelated, the choice of the structural (and statistical) model  
31 may affect the choice of the covariate model and vice versa. The process of finding a  
32 model that adequately describes the data is thus an elaborate task, where model  
33 checking/refining is performed in several steps. To assess model fit in relation to the  
34 observed concentrations or effect measures, scatter plots or the so called goodness-of-fit  
35 plots are created (see Validation of the PK-PD models). Free software packages (Xpose,  
36 PSN etc.) are available to generate these plots.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 The most commonly used software package for model building, which is also supported  
3 by the European Medicines Agency (EMA) is the nonlinear mixed-effect modeling  
4 program NONMEM (GloboMax/ ICON, Ellicott City, MD) [4, 41-43]. NONMEM  
5 estimates parameters (e.g. clearance, volume of distribution or  $EC_{50}$ ) via a maximum  
6 likelihood approach. This means that with the given data, the estimations of the  
7 parameters are the estimations which occur with the highest probability. Alternative  
8 software packages that can be used are for example Monolix, WinNonMix, USC\*PAC  
9 which uses nonparametric maximum likelihood methods [44] or ADAPT using maximum  
10 *a posteriori* (MAP) methods [45].  
11  
12  
13  
14  
15  
16

### 17 Covariate analysis

18 To determine the optimal dose based on the individual characteristics of a patient, a  
19 covariate analysis has to be performed [40, 46, 47]. The aim of the covariate analysis is to  
20 identify specific predictors (covariates) of PK and PD variability and can typically be  
21 studied in population models. Covariate analysis involves the modeling of the distribution  
22 of the individual parameter estimates as a function of covariates which can be of  
23 demographic (e.g. age, bodyweight, gender), patho-physiological (e.g. renal or hepatic  
24 function), and genetic/environmental origin and/or be the result of the concomitant use of  
25 other drugs, which may influence the PK and/or PD. The identification of predictive  
26 covariates for variability provides the scientific basis for rational and individualized,  
27 patient tailored dosing schemes.  
28  
29  
30  
31  
32  
33

34 The influence of developmental changes in childhood can be explored primarily by using  
35 size and/or age as covariates. Size (bodyweight) can be incorporated into the model using  
36 two different approaches. The first approach or 'allometric size approach' includes size *a*  
37 *priori* by using a bodyweight based exponential equation with a fixed exponent of 0.75  
38 for clearance and 1 for volume of distribution [48-52]. Once size is incorporated in the  
39 model using this fixed manner, the influence of age is investigated, being the difference  
40 between actual value of the PK parameter and the 0.75 allometric equation. When  
41 incorporating age as a covariate, different age descriptors may be used like postmenstrual  
42 age (PMA), gestational age (GA) or postnatal age (PNA) [53]. The choice for any of  
43 these age descriptors is based on the results of the systematic covariate analysis as  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 described below [50, 54]. In the second approach or ‘systematic covariate analysis’,  
3 bodyweight is regarded as a covariate as any other which means that the descriptive  
4 properties on the PK parameters are evaluated in a systematic covariate analysis as  
5 described below [55-57].  
6  
7  
8  
9

10 In a systematic analysis, when studying the influence of covariates, scatter plots and  
11 summary plots of individual parameter estimates and/or weighted residuals *versus*  
12 covariates are used to screen for appropriate covariates to include in the covariate sub-  
13 model. Additionally these plots are used to explore the nature of the influence of the  
14 covariate (linear, exponential, allometric, subpopulations etc). Likely candidate  
15 covariates are then added to the model (forward inclusion). The influence of each  
16 covariate on the parameters is examined separately and compared to the simple model  
17 (no covariates). To assess whether the model with covariate statistically improved the fit  
18 to the data, the difference between their objective function value, referred to as log-  
19 likelihood ratio, is calculated. This ratio is assumed to be Chi-square distributed, which  
20 means that a reduction in objective function of 3.84 is considered to be significant  
21 ( $P < 0.05$ ) [43, 58]. Beside the reduction in objective function, goodness-of-fit plots of the  
22 simple model and covariate model are explored for diagnostic purposes. Furthermore, the  
23 confidence interval of the parameter estimates, the correlation matrix (indicates the  
24 relationship between two structural parameters) and visual improvement of the individual  
25 plots are used to evaluate the model. Finally, a superior model is expected to reduce the  
26 inter-subject variance and/or the residual error terms. This procedure of covariate  
27 modeling implies that each covariate is only implemented if this can be fully justified by  
28 the data and the results of the statistic evaluations.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 When two or more covariates are found to significantly improve the model the covariate  
40 that reduced the objective function most is included in the model after which the other  
41 covariates are tested again for their significance. After all covariates that significantly  
42 improved the objective function are added to the simple model, a backward deletion is  
43 performed, which means that each covariate is removed from the full model, one at a  
44 time (the one which causes the smallest increase in objective function first). Retaining or  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 removing the covariate is statistically tested by the use of the objective function (Chi-  
3 square test) until each covariate has been tested.  
4  
5

6  
7 In datasets containing sparse data, there may not be enough information to accurately  
8 estimate inter- and intra-individual variability. This causes the values of these parameters  
9 to shrink to 0, resulting in individual parameter estimates that are closer to the population  
10 parameter estimates than they really are. This phenomenon is called shrinkage [59].  
11 Shrinkage may cause individual predictions, individual parameter estimates and  
12 diagnostics based on them to be less reliable. It can also hide, falsely introduce or distort  
13 the shape of covariate relationships.  
14  
15

16 Shrinkage is the result of properties of the data and is therefore difficult to avoid. One can  
17 only be aware of the presence of shrinkage, realize the influence it may have on the  
18 covariate analysis and use diagnostics other than those based on individual predictions or  
19 individual weighted residuals in the model building and model evaluation procedures.  
20  
21  
22  
23  
24

#### 25 Validation of PK-PD models

26 The objective of a PK or PK-PD modeling exercise is usually not just to describe the  
27 dataset of the sample of individuals that were studied. Generally, models are used to  
28 simulate which concentrations and/or effects and their variability can be expected when  
29 different doses are given to future patients. These simulations may therefore lead to  
30 optimized dosing recommendations or to optimization of new studies for the entire  
31 population where the sample of individuals belongs to. It is often said that 'all models are  
32 wrong, but some are useful' [60]. In order to define whether a model is useful and valid  
33 for clinical and trial simulations, thorough evaluation and validation of the model is  
34 necessary. Although validations of PK models are only performed in 17% of the  
35 published paediatric studies [4] and in 28% of the adults studies [61], proper model  
36 validations are an essential step in model building. For this purpose, different evaluation  
37 and validation methods are available. As described before [62], a proper validation and  
38 evaluation procedure includes an internal model evaluation followed by an external  
39 evaluation and a prospective clinical study.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The first evaluation method is the basic internal model validation used to assess whether the model is able to describe the learning dataset (dataset used to develop the model) accurately and without bias. This evaluation should actually be considered the final stage of the model building procedure. Subsequently, in the external evaluation it is assessed whether the model is able to describe one or more external datasets (datasets other than the one used to develop the model) adequately. Alternatively if a dataset is sufficiently large the original dataset may be split in two so that the model is developed using one part (about two thirds) of the dataset and evaluated externally using the other part (one third) of the dataset. In paediatric studies, it is then especially important to stratify the data correctly and ascertain that all age groups are represented in equal proportions in both datasets.

Various techniques are available for the validation and evaluation of population PK and PK-PD models (both for internal and external validation procedures).

- Basic goodness-of-fit plots ((1) individual predicted versus observed concentrations, 2) population predicted versus observed, 3) (conditional) weighted residuals versus time and 4) (conditional) weighted residuals versus dependent variable plots).

WRES and CWRES are calculated as the following:

$$WRES = \frac{\bar{y}_i - E_{FO}(\bar{y}_i)}{\sqrt{COV_{FO}(\bar{y}_i)}} \quad (\text{equation 4})$$

$$CWRES = \frac{\bar{y}_i - E_{FOCE}(\bar{y}_i)}{\sqrt{COV_{FOCE}(\bar{y}_i)}} \quad (\text{equation 5})$$

Where  $\bar{y}_i$  is the vector of measurements,  $E(\bar{y}_i)$  is the expectation of the data and  $Cov(\bar{y}_i)$  is the covariance matrix of the data [63].

These plots are used in model building, but can also be used to ascertain that there is no trend or bias in the model predictions of the final model. Furthermore, these plots can also be used for both the internal and external evaluation of the model.

- In a bootstrap analysis new datasets are generated by resampling from the original dataset and is therefore an internal validation of the model. The new datasets are



1  
2 subsequently refitted to the original model, yielding mean values and standard  
3 errors for every model parameter.

4  
5 A bootstrap analysis provides information on the stability of the model and its  
6 dependence on specific individuals in the learning dataset. With the freely available  
7 PSN or Wings for NONMEM software packages an automated bootstrap analysis  
8 can be performed.  
9

- 10  
11 • In a visual predictive check (VPC) [64] a PK or PD profile is simulated a 100 to  
12 1000 times and lines for the median values and their 90% prediction interval are  
13 plotted in a graph. The observed values in the internal or external dataset are  
14 subsequently plotted on top of this. It can then be visually checked whether 90% of  
15 the observations are within the indicated prediction interval and whether there is no  
16 bias in the observations compared to prediction interval. In figure 5, two examples  
17 of a VPC are given, showing when a model does not work and when a model does  
18 work on the same data.

19  
20 The VPC is a simulation-based diagnostic that can be used when the PK or PD  
21 profiles for all individuals in the dataset are similar and it allows for easy  
22 interpretation of the result. For this diagnostic tool, there are not statistical tests and  
23 all evaluations are based on visual evaluations. When the individual profiles are  
24 expected to deviate largely from one another because there is for instance a large  
25 variability in the time and amount of dose administrated, or when there are many  
26 covariates, the use of this diagnostic becomes more difficult.

Deleted: This

- 27  
28 • Another simulation-based diagnostic which can be used for both internal and  
29 external validations is the normalized prediction distribution error (NPDE) [65]. An  
30 example of an NPDE published before is shown in figure 6 [55]. This method yields  
31 information on how accurate the model predicts the median value of the  
32 observations and the variability within them. The interpretation of this diagnostic is  
33 less straightforward than for the VPC, but the advantage of this method is that it can  
34 be used when the variability in dosing regimen (both in time, amounts and rates) is  
35 high or when there is a large number of covariates in the model. This can for  
36 instance be the case for data obtained during routine paediatric clinical practice.  
37 Software (e.g. NPDE add-on package for R) [66] to perform this analysis is freely  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Deleted: 4

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

available. For the NPDE, beside visual evaluation of the plots, statistical tests are available. These statistical tests are however reported to be highly sensitive and powerful, so that decisions for the model should primarily be based on visual assessments. An example is the statistically significant deviation of zero of the mean value because of the large number of data, while the actual deviation is small (e.g. 0.074) and not of clinical relevance.

If the model performed well in both the evaluation procedures, the dosing algorithm that results from the PK-PD model needs to be tested and challenged in a prospective (clinical) trial. If the predictive performance of the model is corroborated by the trial it can be used with confidence in clinical practice.

#### Optimal design of paediatric studies

When new population PK-PD studies are performed, it is important to design these studies in the most efficient manner possible to obtain maximum information about the PK and PD parameters so that they can be determined with the highest precision [51, 67]. When designing PK-PD studies in paediatrics certain factors need to be taken into account e.g. age-range of the paediatric group, therapeutic index, possibility to collect blood samples, availability of validated PD endpoints for children, and the availability of sensitive analytical methods.

When optimizing a PK or PK-PD study design, using literature data from adults or children of different age-ranges or possible *in vitro* or pre-clinical data, a concentration-time or effect-time profile for a study can be simulated. This can help to identify possible shortcomings in the design or to perform a power-analysis. Alternatively software packages are available (WINPOPT [68], PopED [67] and PFIM [69]) that can help to identify the optimal number and time points of observations in a study based on the prior information on a drug [70]. To determine the appropriate sample size certain factors, which are summarized in Table 1, need to be taken into account. Each of these factors can influence the required number of patients and/or samples in a positive or negative way. In a study of Peeters *et al.* [32] only 24 patients (aged between 3 and 24 months) were required to determine both the PK and PD since rich sampling was performed (median of

1  
2 11 samples per child) and no covariates were found in the relatively homogenous  
3 population. This in contrast to a study performed by Knibbe *et al.* [55] in which 250  
4 children were included. This higher number was required because in addition to the large  
5 dispersion in age from (preterm) neonates up to toddlers of 3 years of age, only 1 to 4  
6 samples were available for each subject. Moreover infusion rates and additional bolus  
7 doses varied for each child during the study to obtain the desired analgesic effect. In the  
8 another example [71], only 6 patients (aged between 1 and 5 years) were required in  
9 which 7 samples per patient were collected. This lower number of patients (N=6)  
10 compared to the study of Peeters *et al.* (n=24) can be explained because there often exist  
11 a lower variability in PK than in PD which results in a lower required number of patients  
12 (Table 1).  
13  
14  
15  
16  
17  
18  
19

### 20 Conclusions and perspectives

21 In view of the European Regulation which came into force in 2007, it seems now time to  
22 use the progress that has been made in the field of integrated PK-PD modelling [72] to  
23 develop rational and individualized dosing schemes for children. Because of the  
24 possibility to analyse sparse and unbalanced datasets thereby minimizing the burden for  
25 each child, population PK-PD modelling and simulation using non-linear mixed effect  
26 modelling has become the preferred tool to develop effective and safe dosing regimens  
27 for children. Specifically in paediatrics where the developmental changes have to be  
28 taken into account, which may influence the PK and/or the PD of the drugs, this  
29 advanced statistical tool is of critical value.  
30

31 Before dosing regimens can be tested in clinical practice, proper validations of the  
32 models should be performed, for which recently adequate tools have been developed.  
33 Beside internal and external validations, prospective clinical trials, which allow for the  
34 evaluation of the model based dosing regimens, are needed, not only to adjust the  
35 proposed dosing regimen but also to convince paediatricians to use the information that  
36 has been generated using these modelling exercises.  
37

38 Furthermore, one of the future goals may be to explore possibilities for cross-validation  
39 of the models, in which the reported influences of developmental changes on a certain PK  
40 or PD parameter of one drug are evaluated for use in another drug that go through the  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

1  
2 same metabolic route or share the same mechanism of action. In this respect,  
3 physiologically-based pharmacokinetic (PBPK) models are needed. PBPK models  
4 consider the physiological and biochemical processes by using *in vitro* data to describe  
5 the PK of drugs [73, 74]. The combination of these two approaches may use the  
6 information that is already available in a more optimal way in defining effective and safe  
7 dosing regimens for every individual patient.  
8  
9

10  
11 In conclusion, analyses of paediatric data using population PK-PD modelling and  
12 covariate analysis will result in individualized dosing regimens for children of different  
13 age, bodyweight and genetic background. Thus population PK-PD modelling constitutes  
14 an innovative approach to the study of drug effects in this very special patient population,  
15 which is otherwise difficult to study.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 Acknowledgements:

3  
4 The clinical research of K Allegaert is supported by the Fund for Scientific Research,  
5 Flanders (Belgium) (FWO Vlaanderen) by a Fundamental Clinical Investigatorship  
6 (1800209N)  
7

8 The clinical research of C.A.J. Knibbe is supported by the Innovational Research  
9 Incentives Scheme (Veni grant, July 2006) of the Dutch Organization for Scientific  
10 Research (NWO).  
11

12 The research of E.H.J. Krekels is supported by the Top Institute Pharma project number  
13 D2-104  
14

15 The authors would like to thank Pieter De Cock, Hospital Pharmacist, for critically  
16 reading the manuscript.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References:

- 1 Ernest TB, Elder DP, Martini LG, et al (2007) Developing paediatric medicines: identifying the needs and recognizing the challenges. *J Pharm Pharmacol* 59 (8): 1043-1055
- 2 t Jong GW, Vulto AG, de Hoog M, et al (2001) A survey of the use of off-label and unlicensed drugs in a Dutch children's hospital. *Pediatrics* 108 (5): 1089-1093
- 3 Conroy S, Choonara I, Impicciatore P, et al (2000) Survey of unlicensed and off label drug use in paediatric wards in European countries. *European Network for Drug Investigation in Children. BMJ* 320 (7227): 79-82
- 4 Tod M, Jullien V, Pons G (2008) Facilitation of drug evaluation in children by population methods and modelling. *Clin Pharmacokinet* 47 (4): 231-243
- 5 Bray RJ (1998) Propofol infusion syndrome in children. *Paediatr Anaesth* 8 (6): 491-499
- 6 Parke TJ, Stevens JE, Rice AS, et al (1992) Metabolic acidosis and fatal myocardial failure after propofol infusion in children: five case reports. *BMJ* 305 (6854): 613-616
- 7 Sutherland JM (1959) Fatal cardiovascular collapse of infants receiving large amounts of chloramphenicol. *AMA J Dis Child* 97 (6): 761-767
- 8 McIntyre J, Choonara I (2004) Drug toxicity in the neonate. *Biol Neonate* 86 (4): 218-221
- 9 Nederland Kenniscentrum Farmacotherapie bij Kinderen. (<http://www.kinderformularium.nl> Accessed on 29 June 2009).
- 10 European Medicines Agency (EMA): Medicines for children. (<http://www.emea.europa.eu/htms/human/paediatrics/introduction.htm> Accessed on 17 June 2009).
- 11 Dunne J (2007) The European Regulation on medicines for paediatric use. *Paediatr Respir Rev* 8 (2): 177-183
- 12 Consultant PR (2006) EU paediatric regulation for testing medicines in children. (<http://www.child-medicines-research-info.com/paediatric-research-consultancy/eu-paediatric-regulation.asp> Accessed on 23 June 2009).
- 13 Saint Raymond A, Brasseur D (2005) Development of medicines for children in Europe: ethical implications. *Paediatr Respir Rev* 6 (1): 45-51
- 14 Kearns GL, Abdel-Rahman SM, Alander SW, et al (2003) Developmental pharmacology--drug disposition, action, and therapy in infants and children. *N Engl J Med* 349 (12): 1157-1167
- 15 Grand RJ, Watkins JB, Torti FM (1976) Development of the human gastrointestinal tract. A review. *Gastroenterology* 70 (5 PT.1): 790-810
- 16 Anderson BJ, van Lingen RA, Hansen TG, et al (2002) Acetaminophen developmental pharmacokinetics in premature neonates and infants: a pooled population analysis. *Anesthesiology* 96 (6): 1336-1345
- 17 Hines RN, McCarver DG (2002) The ontogeny of human drug-metabolizing enzymes: phase I oxidative enzymes. *J Pharmacol Exp Ther* 300 (2): 355-360
- 18 de Wildt SN, Kearns GL, Leeder JS, et al (1999) Glucuronidation in humans. Pharmacogenetic and developmental aspects. *Clin Pharmacokinet* 36 (6): 439-452

- 1  
2  
3 19 Leeder JS (2004) Translating pharmacogenetics and pharmacogenomics into drug  
4 development for clinical pediatrics and beyond. *Drug Discov Today* 9 (13): 567-  
5 573  
6 20 Krekels EH, van den Anker JN, Baiardi P, et al (2007) Pharmacogenetics and  
7 paediatric drug development: issues and consequences to labelling and dosing  
8 recommendations. *Expert Opin Pharmacother* 8 (12): 1787-1799  
9 21 Rhodin MM, Anderson BJ, Peters AM, et al (2009) Human renal function  
10 maturation: a quantitative description using weight and postmenstrual age. *Pediatr*  
11 *Nephrol* 24 (1): 67-76  
12 22 Bartelink IH, Rademaker CM, Schobben AF, et al (2006) Guidelines on paediatric  
13 dosing on the basis of developmental physiology and pharmacokinetic  
14 considerations. *Clin Pharmacokinet* 45 (11): 1077-1097  
15 23 Knibbe CA, Koster VS, Deneer VH, et al (1998) Determination of propofol in  
16 low-volume samples by high-performance liquid chromatography with  
17 fluorescence detection. *J Chromatogr B Biomed Sci Appl* 706 (2): 305-310  
18 24 Rauh M, Stachel D, Kuhlén M, et al (2006) Quantification of busulfan in saliva  
19 and plasma in haematopoietic stem cell transplantation in children : validation of  
20 liquid chromatography tandem mass spectrometry method. *Clin Pharmacokinet*  
21 45 (3): 305-316  
22 25 Edelbroek PM, van der Heijden J, Stolk LM (2009) Dried blood spot methods in  
23 therapeutic drug monitoring: methods, assays, and pitfalls. *Ther Drug Monit* 31  
24 (3): 327-336  
25 26 Jabeen R, Payne D, Wiktorowicz J, et al (2006) Capillary electrophoresis and the  
26 clinical laboratory. *Electrophoresis* 27 (12): 2413-2438  
27 27 Fisher DM, O'Keeffe C, Stanski DR, et al (1982) Pharmacokinetics and  
28 pharmacodynamics of d-tubocurarine in infants, children, and adults.  
29 *Anesthesiology* 57 (3): 203-208  
30 28 LeDez KM, Lerman J (1987) The minimum alveolar concentration (MAC) of  
31 isoflurane in preterm neonates. *Anesthesiology* 67 (3): 301-307  
32 29 Blusse van Oud-Alblas HJ PM, Brill MJE, Knibbe CAJ, Klein J, Tibboel D,  
33 Weber F (2009) The pharmacodynamics of isoflurane in children using Bispectral  
34 index and composite auditory evoked potentials. [.\(www.page-  
35 meeting.org/?abstract=1576](http://www.page-meeting.org/?abstract=1576) Accessed on  
36 30 Stephenson T (2005) How children's responses to drugs differ from adults. *Br J*  
37 *Clin Pharmacol* 59 (6): 670-673  
38 31 van Dijk M, de Boer JB, Koot HM, et al (2000) The reliability and validity of the  
39 COMFORT scale as a postoperative pain instrument in 0 to 3-year-old infants.  
40 *Pain* 84 (2-3): 367-377  
41 32 Peeters MY, Prins SA, Knibbe CA, et al (2006) Pharmacokinetics and  
42 pharmacodynamics of midazolam and metabolites in nonventilated infants after  
43 craniofacial surgery. *Anesthesiology* 105 (6): 1135-1146  
44 33 Peeters MY, Prins SA, Knibbe CA, et al (2006) Propofol pharmacokinetics and  
45 pharmacodynamics for depth of sedation in nonventilated infants after major  
46 craniofacial surgery. *Anesthesiology* 104 (3): 466-474  
47 34 van Dijk M, Peters JW, van Deventer P, et al (2005) The COMFORT Behavior  
48 Scale: a tool for assessing pain and sedation in infants. *Am J Nurs* 105 (1): 33-36  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3 35 Jonsson EN, Karlsson MO (1999) Xpose--an S-PLUS based population  
4 pharmacokinetic/pharmacodynamic model building aid for NONMEM. Comput  
5 Methods Programs Biomed 58 (1): 51-64  
6 36 FDA C (1999) Guidance for industry. Population  
7 pharmacokinetics. (<http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm072137.pdf> Accessed on  
8  
9 37 FDA C (1998) Guidance for industry. General considerations for pediatric  
10 pharmacokinetic studies for drugs and biological products (Draft  
11 Guidance). (<http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm072114.pdf> Accessed on 22 June 2009).  
12  
13 38 Beal SL, Sheiner LB (1982) Estimating population kinetics. Crit Rev Biomed Eng  
14 8 (3): 195-222  
15 39 Ette EI, Williams PJ, Lane JR (2004) Population pharmacokinetics III: design,  
16 analysis, and application of population pharmacokinetic Studies. Ann  
17 Pharmacother 38 (12): 2136-2144  
18 40 Wade JR, Beal SL, Sambol NC (1994) Interaction between structural, statistical,  
19 and covariate models in population pharmacokinetic analysis. J Pharmacokinet  
20 Biopharm 22 (2): 165-177  
21 41 Aarons L (1999) Software for population pharmacokinetics and  
22 pharmacodynamics. Clin Pharmacokinet 36 (4): 255-264  
23 42 Beal SL SL, Broeckman AJ (2006) NONMEM Users Guides. Icon Development  
24 Solutions, Ellicott City, Maryland, USA.  
25 43 Broeckman AJ SL, Beal SL (1994) NONMEM Users Guide - Part V. University  
26 of California at San Francisco, San Francisco.  
27 44 Jelliffe R SA, Van Guilder M, et al. (1995) User manual for version 10.7 of the  
28 USC\*PAC Collection of PC Programs, USC Laboratory of Applied  
29 Pharmacokinetics. USC School of Medicine, Los Angeles.  
30 45 Bustad A, Terziivanov D, Leary R, et al (2006) Parametric and nonparametric  
31 population methods: their comparative performance in analysing a clinical dataset  
32 and two Monte Carlo simulation studies. Clin Pharmacokinet 45 (4): 365-383  
33 46 Mentre F, Mallet A (1994) Handling covariates in population pharmacokinetics.  
34 Int J Biomed Comput 36 (1-2): 25-33  
35 47 Mandema JW, Verotta D, Sheiner LB (1992) Building population  
36 pharmacokinetic--pharmacodynamic models. I. Models for covariate effects. J  
37 Pharmacokinet Biopharm 20 (5): 511-528  
38 48 Anderson BJ, Holford NH (2008) Mechanism-based concepts of size and maturity  
39 in pharmacokinetics. Annu Rev Pharmacol Toxicol 48: 303-332  
40 49 Anderson BJ, Allegaert K, Holford NH (2006) Population clinical pharmacology  
41 of children: general principles. Eur J Pediatr 165 (11): 741-746  
42 50 Anderson BJ, Allegaert K, Holford NH (2006) Population clinical pharmacology  
43 of children: modelling covariate effects. Eur J Pediatr 165 (12): 819-829  
44 51 Meibohm B, Laer S, Panetta JC, et al (2005) Population pharmacokinetic studies  
45 in pediatrics: issues in design and analysis. AAPS J 7 (2): E475-487  
46 52 Anderson BJ, Holford NH (2009) Mechanistic basis of using body size and  
47 maturation to predict clearance in humans. Drug Metab Pharmacokinet 24 (1): 25-  
48 36  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3 53 Engle WA (2004) Age terminology during the perinatal period. *Pediatrics* 114 (5):  
4 1362-1364  
5 54 Allegaert K, Anderson BJ, Cossey V, et al (2006) Limited predictability of  
6 amikacin clearance in extreme premature neonates at birth. *Br J Clin Pharmacol*  
7 61 (1): 39-48  
8 55 Knibbe CA, Krekels EH, van den Anker JN, et al (2009) Morphine  
9 glucuronidation in preterm neonates, infants and children younger than 3 years.  
10 *Clin Pharmacokinet* 48 (6): 371-385  
11 56 Mahmood I (2007) Prediction of drug clearance in children: impact of allometric  
12 exponents, body weight, and age. *Ther Drug Monit* 29 (3): 271-278  
13 57 Mahmood I (2009) Application of fixed exponent 0.75 to the prediction of human  
14 drug clearance: an inaccurate and misleading concept. *Drug Metabol Drug*  
15 *Interact* 24 (1): 57-81  
16 58 Sheiner LB, Ludden TM (1992) Population pharmacokinetics/dynamics. *Annu*  
17 *Rev Pharmacol Toxicol* 32: 185-209  
18 59 Karlsson MO, Savic RM (2007) Diagnosing model diagnostics. *Clin Pharmacol*  
19 *Ther* 82 (1): 17-20  
20 60 Box G (1979) Robustness in the strategy of scientific model building. In: Launer  
21 RL Wilkinson GN, editors. *Robustness in Statistics*. New York: Academic Press,  
22 pp202.  
23 61 Brendel K, Dartois C, Comets E, et al (2007) Are population pharmacokinetic  
24 and/or pharmacodynamic models adequately evaluated? A survey of the literature  
25 from 2002 to 2004. *Clin Pharmacokinet* 46 (3): 221-234  
26 62 Ince I, de Wildt SN, Tibboel D, et al (2009) Tailor-made drug treatment for  
27 children: creation of an infrastructure for data-sharing and population PK-PD  
28 modeling. *Drug Discov Today* 14 (5-6): 316-320  
29 63 Hooker AC, Staats CE, Karlsson MO (2007) Conditional weighted residuals  
30 (CWRES): a model diagnostic for the FOCE method. *Pharm Res* 24 (12): 2187-  
31 2197  
32 64 Fang L, Holford NH, Hinkle G, et al (2007) Population pharmacokinetics of  
33 humanized monoclonal antibody HuCC49deltaCH2 and murine antibody CC49 in  
34 colorectal cancer patients. *J Clin Pharmacol* 47 (2): 227-237  
35 65 Brendel K, Comets E, Laffont C, et al (2006) Metrics for external model  
36 evaluation with an application to the population pharmacokinetics of gliclazide.  
37 *Pharm Res* 23 (9): 2036-2049  
38 66 Comets E, Brendel K, Mentre F (2008) Computing normalised prediction  
39 distribution errors to evaluate nonlinear mixed-effect models: the npde add-on  
40 package for R. *Comput Methods Programs Biomed* 90 (2): 154-166  
41 67 Foracchia M, Hooker A, Vicini P, et al (2004) POPED, a software for optimal  
42 experiment design in population kinetics. *Comput Methods Programs Biomed* 74  
43 (1): 29-46  
44 68 Duffull SB DN, Eccleston, J, Kimko, H (2008) WinPOPT User Guide Version  
45 1.2. School of Pharmacy, University of Otago, New Zealand.  
46 69 Retout S, Mentre F (2003) Optimization of individual and population designs  
47 using Splus. *J Pharmacokinet Pharmacodyn* 30 (6): 417-443  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 70 Duffull SB (2001) Design of clinical pharmacology trials. *Clin Exp Pharmacol*  
4 *Physiol* 28 (11): 905-912  
5 71 Knibbe CA, Melenhorst-de Jong G, Mestrom M, et al (2002) Pharmacokinetics  
6 and effects of propofol 6% for short-term sedation in paediatric patients following  
7 cardiac surgery. *Br J Clin Pharmacol* 54 (4): 415-422  
8 72 Breimer DD (2008) PK/PD modelling and beyond: impact on drug development.  
9 *Pharm Res* 25 (12): 2720-2722  
10 73 Alcorn J, McNamara PJ (2008) Using ontogeny information to build predictive  
11 models for drug elimination. *Drug Discov Today* 13 (11-12): 507-512  
12 74 Johnson TN (2005) Modelling approaches to dose estimation in children. *Br J*  
13 *Clin Pharmacol* 59 (6): 663-669  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1: Schematic representation of the relationship between dose and concentration (pharmacokinetics, PK) and between concentration and a pharmacological (side) effect (pharmacodynamics, PD). Important covariates which may affect both the PK and/or PD are bodyweight, age, disease status (e.g. critically ill versus healthy children) and genetics.

Figure 2: Simulation of propofol concentrations and response using COMFORT-B score versus time based on developed PK and PD models, after administration of different doses of propofol (0, 18, 30, and 36 mg/h) for a 10 kg and a 5 kg non-ventilated infant in the first night at the Intensive Care following craniofacial surgery. Target COMFORT-B scores are between 12 and 14 preferably. Reproduced from [Peeters MY, Prins SA, Knibbe CA, DeJongh J, van Schaik RH, van Dijk M, et al. Propofol pharmacokinetics and pharmacodynamics for depth of sedation in nonventilated infants after major craniofacial surgery. *Anesthesiology* 2006 Mar;104(3):466-74.]

Figure 3: Concentration-time profiles of the same study using two different approaches. In figure 3A the standard two-stage approach is applied to a rich dataset. 3B shows the population approach with mixed effect modeling applied to the same dataset using only two datapoints for each individual so a sparse dataset is created. In 3A, in each of the six individuals 10 samples are available. The different symbols correspond to different individuals. Each black line corresponds to a separate fit to the 10 data points of each individual.

In 3B, which uses the mixed effect modeling approach, two samples of the 10 per subject in 2A are used. The different symbols correspond to the six different individuals. The black line illustrates the concentration time plot based on the population mean values of the PK parameters (PRED). The grey lines show the plots of the individual patients, which are based on the population mean values together with the measured concentrations of the specific individual (IPRED).

Figure 4: In 4A, the inter-individual variability is shown between three individuals who received the same dose. 4B presents the intra- or residual variability by showing the concentration-time profile after repeated administration. Both these random variables are assumed to be normally distributed with a mean of zero and a variance of  $\omega^2$  or  $\sigma^2$  respectively.

**Figure 5:**

Two examples of a visual predictive check (VPC) are illustrated based on the same dataset (warfarin concentrations and prothrombin complex activity (PCA)) using two different models. In 5A the VPC of the effect compartment model is shown, while in 5B the VPC of the turnover model is demonstrated. The median (black thick line) and the 90% intervals (black thin lines) together with the observed data (PCA) (dots) are shown. Based on both graphics, the turn over model is the most appropriate model since 90% of the observations are lying within the prediction interval. Furthermore, unlike the effect compartment model, no bias is seen in the observations.


Reproduced from [Holford N, 2005. The visual predictive check – Superiority to Standard Diagnostic (Rorschach) Plots. PAGE 14, Abstr 738. (<http://www.page-meeting.org/?abstract=738>)].

Figure 6: Example of a normalized prediction distribution error (NPDE) analysis, which show the NPDE distributions for morphine. The normal distribution is presented by the solid line. The values for the mean and standard deviation of the observed NPDE distribution are given below the histogram, with \* indicating a significant difference of a mean of 0 and a variance of 1 at the  $p < 0.05$  level, as determined by the Wilcoxon signed rank test and Fisher test for variance. Additionally the distribution of NPDE vs time after the first dose and NPDE vs the log of the concentrations are also shown. The dotted lines represent the 90% distribution of the NPDE. Reproduced from [Knibbe CA, Krekels EH, van den Anker JN, DeJongh J, Santen GW, van Dijk M, et al. Morphine glucuronidation in preterm neonates, infants and children younger than 3 years. *Clin Pharmacokinet* 2009;48(6):371-85] with permission from Wolters Kluwer Health | Adis (© Adis Data Information BV [2006]. All rights reserved.)

Deleted: ¶

Deleted: 5


FIGURE 1



For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


FIGURE 2



Or Peer Review


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

FIGURE 3



For Peer Review


FIGURE 4



For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

FIGURE 5




For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

FIGURE 6 Deleted: 5



For Peer Review

Table 1: Factors influencing the required number of patients and/or samples per patient.

<b>Factor</b>	<b>Number of patients/samples</b>
Study of PK only	relatively small number of patients/samples
Study of PK-PD relationship	relatively high number of patients/samples
Even distribution of covariates (age, bodyweight)	↓ number of patients/samples
↑ Number of changes in dose	↑/- number of patients/samples (depending on other aspects of the study design)
↑ Number of samples/child	↓ number of patients
Use of optimal sampling strategies	↓ number of patients/samples
Different sampling windows (e.g. two or three sampling schemes)	↓ number of patients/samples

## The role of population PK-PD modeling in paediatric clinical research

*Roosmarijn F. W. De Cock<sup>1</sup>, Chiara Piana<sup>1</sup>, Elke H.J. Krekels<sup>1,2</sup>, Meindert Danhof<sup>1</sup>,  
Karel Allegaert<sup>4</sup>, Catherijne A.J. Knibbe<sup>1,2,3</sup>*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
Division of Pharmacology, Leiden/Amsterdam Centre for Drug Research,  
Leiden University,  
Leiden, the Netherlands

2  
Department of Pediatric Intensive Care  
Erasmus MC - Sophia Children's Hospital  
Rotterdam, the Netherlands

3  
Department of Clinical Pharmacy  
St. Antonius Hospital  
P.O. Box 2500  
3430 EM Nieuwegein, the Netherlands  
Email: [c.knibbe@antoniuziekenhuis.nl](mailto:c.knibbe@antoniuziekenhuis.nl)

4  
Neonatal Intensive Care Unit  
University Hospital Leuven, Belgium

RDC and CP equally contributed to this paper

1  
2  
3 Children differ from adults in their response to drugs. While this may be the result of  
4 changes in dose-exposure (pharmacokinetics (PK)) and/or exposure-response  
5 (pharmacodynamics (PD)) relationships, the magnitude of these changes may not be  
6 solely reflected by differences in bodyweight. As a consequence, dosing  
7 recommendations empirically derived from adults dosing regimens using linear  
8 extrapolations based on bodyweight, can result in therapeutic failure, occurrence of  
9 adverse effect or even fatalities. In order to define rational, patient tailored dosing  
10 schemes, population PK-PD studies in children are needed. For the analysis of the data,  
11 population modeling using non-linear mixed effect modeling is the preferred tool since  
12 this approach allows for the analysis of sparse and unbalanced datasets. Additionally it  
13 permits the exploration of the influence of different covariates such as bodyweight, age  
14 and other covariates, to explain the variability in drug response. Finally, using this  
15 approach, these PK-PD studies can be designed in the most efficient manner in order to  
16 obtain the maximum information on the PK-PD parameters with the highest precision.  
17 Once a population PK-PD model is developed, internal and external validations should be  
18 performed. If the model performs well in these validation procedures, model simulations  
19 can be used to define a dosing regimen which in turn needs to be tested and challenged in  
20 a prospective clinical trial. This methodology will improve the efficacy/safety balance of  
21 dosing guidelines which will be of benefit to the individual child.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

38 Keywords: Pharmacokinetics – Pharmacodynamics – Population modeling – Paediatric  
39 population – Validation  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Introduction:

Children differ from adults in their response to drugs. These differences may be caused by changes in the pharmacokinetics (PK) and/or pharmacodynamics (PD) between children and adults and may also vary between children of different ages. The PK of a drug includes processes of absorption, distribution, metabolism and elimination of a drug whereas the PD comprises the physiological and biological response to the administered drug and therefore may represent both efficacy and safety measures. While a child grows, enzyme pathways (involved in the PK), function and expression of receptors and proteins (involved in the PD) mature, which can be referred to as 'developmental changes' in childhood. The maturation rates of these developmental changes vary however between the pathways and receptors and often do not correlate solely with the increase in bodyweight of the child. The question is therefore how to obtain data in children that allow for the study of these developmental changes ultimately resulting in evidence based dosing regimens for drugs in children.

To date, only a small number of drugs used in children is licensed for use in this specific group. Up to 70% of the drugs in paediatric intensive care, and 90% of the drugs in neonatal intensive care, are prescribed in an off-label or unlicensed manner [1-4]. Paediatric dosing regimens are usually empirically derived from adult regimens using linear extrapolations based on bodyweight. Since these developmental changes are non-linear dynamic processes, this dosing paradigm may result in under or over-dosing particularly in specific age groups. This may cause therapeutic failure, occurrence of severe adverse effects or even fatalities such as fatalities occurring after long-term sedation with high doses of propofol [5, 6] and occurrence of the grey baby syndrome in neonates after treatment with chloramphenicol [7, 8]. As a result, dose adjustments in the younger age groups are often proposed. For vancomycin for example lower doses are administered in neonates younger than 1 week (20 mg/kg/day) compared to 1-4 week-old neonates (30 mg/kg/day) and children between 1 month and 18 years (40 mg/kg/day) [9].

Instead of the *a priori* use of bodyweight for dosing guidelines in children, detailed information on PK and potentially also the PD needs to be considered in order to define

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

effective and safe dosing regimens throughout the paediatric age range. The lack of PK and PD information on drugs in children has led to the European Regulation which entered into force in 2007. This law imposes pharmaceutical companies to perform research in the whole paediatric age-range for all drugs that are developed for the European market, by requiring the submission of a paediatric investigational plan (PIP) in the early stages of the development of a new drug. In this PIP, a full description has to be given of the studies and of drug formulation in the paediatric population. In case little information is available about efficacy and safety of a drug, studies in children are only performed after more information is obtained in the adult population to increase the safety of the paediatric study [10-12]. The main targets of introducing the paediatric Regulation were to facilitate development and availability of medicines in children between 0 and 17 years, to improve the availability of information about medicines used in children, to ensure that the medicines are of high quality, can be administered in a safe and effective way and that paediatric studies are performed in an ethically correct way [10]. The reward for this effort is a six month supplementary production certificate for the pharmaceutical company.

Both for industry and for academic researchers, performing (PK-PD) studies in children in order to develop rational dosing schemes is very challenging because of ethical and practical issues. Unlike studies in healthy adults, research in healthy children is considered to be unethical, so all paediatric studies are performed in the vulnerable group of children suffering from a disease. In all clinical trials, an informed consent has to be signed by the patient before he or she can be enrolled into a trial. In paediatric trials, this informed consent can not be obtained by the patient that participates in the trial, and is therefore replaced by the consent of the parents or guardians. In older age groups, in addition to this consent, an assent is used in which the aim of the study is explained in an age-appropriate language so that children can understand. [1, 4, 13].

Apart from ethical issues, practical challenges also occur when performing studies in children. There are limitations to the number and volume of samples that can be obtained, resulting in infrequent sampling possibilities and the need for advanced drug assay techniques with improved sensitivity. Another complicating factor is the limited available

1  
2  
3 number of subjects that suffers from the same disease. Finally, pharmacodynamic  
4 endpoints that measure the efficacy of the drug, and which are validated for children may  
5 be lacking. All these factors call for highly advanced study designs and analysis  
6 techniques so that the burden for each child can be kept to a minimum while still  
7 addressing all the study objectives.  
8  
9  
10  
11

12  
13  
14 This paper aims to inform clinical pharmacologists, paediatricians and pharmacists about  
15 population PK-PD modeling in paediatric drug research. Advanced statistical tools are  
16 discussed that can be used to develop rational dosing schemes based on the PK and PD of  
17 a drug in children, despite practical and ethical restrictions. Using these tools, covariates  
18 can be identified in order to define appropriate doses and dosing intervals based on  
19 individual characteristics of each child with minimal burden to each patient. The paper  
20 also describes how to evaluate the predictive performance of the models by different  
21 validation methods including a prospective clinical trial. Ultimately, the efforts result in  
22 an individualized dosing regimen based on the PK-PD relation through the paediatric  
23 age-range.  
24  
25  
26  
27  
28  
29  
30  
31

### 32 33 PK-PD in children 34

35 Developmental changes in childhood can affect all PK processes from absorption until  
36 elimination as well as the pharmacodynamic effects. For example, in neonates intra-  
37 gastric pH is elevated (>4) which may increase the bioavailability of acid-labile  
38 compounds (penicillin G) and decrease the bioavailability of weak acids (phenobarbital)  
39 when given orally [14]. Additionally, gastric emptying in neonates is delayed, which  
40 means that also the absorption of drugs e.g. paracetamol is slower in neonates [15, 16].  
41  
42 Other examples are changes in metabolizing enzyme capacity in children. Although most  
43 uridine 5'-diphosphate (UDP)-glucuronosyltransferases (UGTs) and P-450 cytochromes  
44 (CYPs) are expressed during the first week of life, the activity at birth in comparison with  
45 adults is often low, e.g. UGT2B7 activity at birth is around 10% of the adult level and  
46 maturation rates of different enzyme systems are known to mature at different rates [14,  
47 17-20].  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 In addition, renal function and liver flow are influenced by physiological changes  
4 depending on age, e.g. the glomerular filtration rate in mL/min/70kg in full term neonates  
5 is 35% of the adult value, while mL/min/70kg adult values are reached at approximately  
6 1 year old [21]). When using units of mL/min/70kg however, it should be realized that  
7 actual values of GFR in children are still very low compared to adult values because of  
8 correction for differences in total body weight between adults and infants.

9  
10 Furthermore the body composition of children changes continuously resulting in an age-  
11 dependent proportion of body water and fat, which influences the distribution of drugs.  
12 For example, the total amount of body water (80-90 % of the bodyweight) is higher in  
13 neonates compared to adults (55-60%). Hydrophilic drugs like aminoglycosides have a  
14 larger volume of distribution in neonates which can be explained by larger extra-cellular  
15 fluid (45% of the bodyweight) compared to adults (20%) [14, 22].

16  
17 In order to characterize the specific influence of developmental changes in childhood on  
18 the PK of a drug, concentration-time profiles are necessary, which require measurements  
19 of drug concentrations. For ethical reasons, in paediatric studies, discomfort, like pain  
20 and anxiety associated with venipuncture, must be restricted and practical issues limit the  
21 volume and amount of blood samples that can be obtained. Therefore, sensitive analysis  
22 techniques requiring only small blood samples should be used. While HPLC methods  
23 have reported to require only 50  $\mu$ L of blood [23], more recently LC-MS methods can  
24 measure up to ten different drugs in volumes as low as 50-100  $\mu$ L [24]. Additionally, also  
25 alternative matrices such as saliva should be explored as a non-invasive, more child-  
26 friendly alternative to measure a drug concentration. An example in this respect is a LC-  
27 MS/MS method which was developed and validated for the measurement of busulphan in  
28 saliva [24]. Also the use of a dried blood spot method e.g. for tacrolimus can facilitate the  
29 measurement of drugs in children [25]. Another method is capillary electrophoresis  
30 which requires only a low sample volume for the quantification of drugs in biological  
31 fluids [26].

32  
33 Changes between children and adults may also result from differences in the  
34 pharmacodynamics of a drug in children, e.g. by changes in the relative number and  
35 function of receptors. These age-related PD differences are until present rarely reported  
36 in literature, but one of the few examples is the increased sensitivity to d-tubocurarine, an  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 antagonist of nicotinic neuromuscular acetylcholine receptors, in neonates and infants  
4 compared to children and adults [27]. Other examples are the observed lower minimum  
5 alveolar concentration (MAC) of isoflurane in preterm neonates compared to full-term  
6 neonates and older children [28, 29] and the different sensitivity to bronchodilators  
7 because of the lack of smooth muscles in the airways in neonates [30].

8  
9  
10 To study the PD of a drug in children, the use of a PD endpoint which is validated for use  
11 in children is a prerequisite. An illustrative example is the measurement of pain in young  
12 children. Since they are not able to report their pain using a visual analogue scale, an  
13 observational scale has been developed. This comfort behavioral (COMFORT-B) scale  
14 was developed and validated for use in children under the age of three years [31]. The  
15 scale assesses six behavioral items: alertness, calmness, muscle tone, body movement,  
16 facial tension, and crying (non-ventilated children) or respiratory response (ventilated  
17 children). All items range from 1 (no distress) to 5 (severe distress), resulting in a total  
18 score of varying from 6 to 30. This validated scale can then be used as a PD endpoint for  
19 the development of PD models for pain and/or sedation in children of different ages [32-  
20 34].

21  
22  
23 The influence of covariates such as the developmental changes, disease status and  
24 genetics on the PK and PD of drugs in children is depicted in Figure 1.

25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
When both the PK and PD of a drug in children are characterized, the developed models  
can be used to derive rational dosing regimens with predictable efficacy and  
concentration profiles. An example of such a PK-PD model with a derived dosing  
regimen is an article published by Peeters *et al.* In this paper both the PK and the PD  
were characterized in children, the latter with the use of the COMFORT-B scale as  
pharmacodynamic endpoint [33]. Based on the model it was found that propofol  
clearance is two times higher in non-ventilated children compared to ventilated children  
and adults. For the PD, a model was derived in which an effect of propofol was  
characterized within a naturally occurring sleep pattern of children in the ICU. Both  
models (PK as well as PD) were used to simulate concentrations as well as the effects  
that could be expected using different dosing schemes (Figure 2). As a result, based on

1  
2  
3 this PK-PD model, a propofol dose of 30mg/h was recommended for a child of 10 kg  
4 which will result in adequate COMFORT-B scales in the night following craniofacial  
5 surgery.  
6  
7  
8  
9

#### 10 11 Methods to analyse data: standard two-stage or population approach

12  
13 When concentration-time and concentration-effect datasets obtained in children are  
14 considered for analysis, two different methods can be applied: the standard two-stage  
15 approach and the population approach using non-linear mixed effect models [35-38].  
16 When using the standard two-stage approach or classical approach, in a first step  
17 parameters are estimated in each individual based on individual concentration-time  
18 profiles (figure 3A). In a second step, these parameters are summarized by calculating the  
19 mean or median of the parameters and the variability between subjects (SE or IQR). A  
20 major drawback of this methodology is that this approach requires a relatively high  
21 number of samples in each individual patient (Figure 3A) while each patient has to  
22 contribute roughly the same number of samples. Moreover it is very difficult to  
23 distinguish between inter-individual (variability between subjects), intra-individual and  
24 residual variability (variability within one subject, measurement error, and model  
25 misspecification) and as a result inter-individual variability is often overestimated [39].  
26 Since usually only a limited number of observations can be obtained in paediatric  
27 subjects, the population approach using non-linear mixed effect modeling to obtain PK  
28 and PD parameters, is the preferred approach [37]. The population approach differs from  
29 the standard two stage-approach in the fact that the analysis is based on simultaneous  
30 analysis of all data of the entire population while still taking into account that different  
31 observations come from different patients (Figure 3B). Additionally the population  
32 approach allows not only for the analysis of dense data but also for sparse (limited  
33 number of observations per individual) and unbalanced data (unequal distribution of  
34 observations in various parts of the concentration-time profile in the individuals) or a  
35 combination of both. Finally both the interindividual and intra-individual variability are  
36 separately estimated in the dataset using this approach.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 As a result of this methodology, when designing a paediatric study of which the data will  
56 be analyzed using the population approach, it is advisable to collect samples at different  
57  
58  
59  
60

times (or time-windows) or to set alternating sampling schemes in subgroups of patients. This also means that (part of the) samples can be collected during routine clinical sampling. Consequently, the burden for the child that participates in the trial is reduced and the statistical power to develop a model describing the concentration-time or concentration-effect profile is not affected or improved.

The term ‘mixed’ in non-linear mixed effects modeling stands for a mixture of fixed and random effects. For the fixed effects, a structural model describing the PK or PD is chosen (e.g. a two-compartment model for PK or an  $E_{\max}$  model for PD). The random effects quantify the variability that is not explained by the fixed effects. These random effects include inter-subject and intra-subject and random variability (Figure 4), which are both simultaneously and separately estimated. It is often assumed that the variability between subjects follows a normal distribution with a mean of zero and variance  $\omega^2$ . Equation 1 is used to describe the relationship between individual and population parameter estimates.

$$\theta_i = \theta_{mean} \cdot e^{\eta_i} \quad (\text{equation 1})$$

where  $\theta_i$  represents the parameter of the  $i$ th subject,  $\theta_{mean}$  the population mean, and  $\eta_i$  the variability between subjects. The residual error is in generally described using a proportional error (error is dependent on the concentration, which means a higher absolute error at higher concentrations (equation 2)) or additive error (constant for all observations (equation 3)) or a combination of both. This means for the  $j$ th observed concentration of the  $i$ th individual the relation ( $Y_{ij}$ ):

$$Y_{ij} = c_{pred,ij} \cdot (1 + \varepsilon_{ij}) \quad (\text{equation 2})$$

$$Y_{ij} = c_{pred,ij} + \varepsilon_{ij} \quad (\text{equation 3})$$

where  $c_{pred}$  is predicted concentration and  $\varepsilon_{ij}$  is a random variable with mean zero and variance  $\sigma^2$ .

1  
2  
3  
4  
5 In general, model building requires three different steps. First a structural model (fixed  
6 effects) has to be designed, then a statistical sub-model (random effects) has to be  
7 developed and in the final step a covariate sub-model is identified.  
8  
9

10 The structural model describes the overall trend in the data. The choice of structural  
11 model (e.g. one, two or three-compartment model for PK and an  $E_{\max}$  model for PD) is to  
12 be based upon the best *a priori* information about the drug to be studied [40]. The  
13 structural model uses fixed effects parameters such as clearance and volume of  
14 distribution for PK or  $E_{\max}$  and  $EC_{50}$  for PD. The population values for these parameters  
15 are called typical values (TV).  
16  
17

18 After selecting the structural model, the statistical sub-model which accounts for the  
19 inter-individual as well as the residual variability is chosen and tested. Information on  
20 both inter- and intra- and residual variability is of clinical value, because it describes  
21 differences in clinical response between and within patients and may therefore provide  
22 guidance to rational dose adjustments. With the population approach, both these random  
23 effects are obtained, apart from estimates of both the population values (TV) and the  
24 individual values of PK and PD parameters (so called *post hoc* parameter estimates).  
25  
26

27 In the final step the covariate sub-model is determined which expresses relationships  
28 between covariates and parameters of the structural model (e.g. influence of bodyweight  
29 on volume of distribution or clearance). Covariates can be individual-specific (age,  
30 bodyweight, genetic profile, etc) or time-varying (renal function, hemodynamic  
31 parameters, body temperature etc). The covariate analysis will be explained more in  
32 details in the following section.  
33  
34

35 As these three models are interrelated, the choice of the structural (and statistical) model  
36 may affect the choice of the covariate model and vice versa. The process of finding a  
37 model that adequately describes the data is thus an elaborate task, where model  
38 checking/refining is performed in several steps. To assess model fit in relation to the  
39 observed concentrations or effect measures, scatter plots or the so called goodness-of-fit  
40 plots are created (see Validation of the PK-PD models). Free software packages (Xpose,  
41 PSN etc.) are available to generate these plots.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 The most commonly used software package for model building, which is also supported  
5 by the European Medicines Agency (EMA) is the nonlinear mixed-effect modeling  
6 program NONMEM (GloboMax/ ICON, Ellicott City, MD) [4, 41-43]. NONMEM  
7 estimates parameters (e.g. clearance, volume of distribution or  $EC_{50}$ ) via a maximum  
8 likelihood approach. This means that with the given data, the estimations of the  
9 parameters are the estimations which occur with the highest probability. Alternative  
10 software packages that can be used are for example Monolix, WinNonMix, USC\*PAC  
11 which uses nonparametric maximum likelihood methods [44] or ADAPT using maximum  
12 *a posteriori* (MAP) methods [45].  
13  
14  
15  
16  
17  
18  
19  
20

### 21 Covariate analysis

22 To determine the optimal dose based on the individual characteristics of a patient, a  
23 covariate analysis has to be performed [40, 46, 47]. The aim of the covariate analysis is to  
24 identify specific predictors (covariates) of PK and PD variability and can typically be  
25 studied in population models. Covariate analysis involves the modeling of the distribution  
26 of the individual parameter estimates as a function of covariates which can be of  
27 demographic (e.g. age, bodyweight, gender), patho-physiological (e.g. renal or hepatic  
28 function), and genetic/environmental origin and/or be the result of the concomitant use of  
29 other drugs, which may influence the PK and/or PD. The identification of predictive  
30 covariates for variability provides the scientific basis for rational and individualized,  
31 patient tailored dosing schemes.  
32  
33  
34  
35  
36  
37  
38  
39

40 The influence of developmental changes in childhood can be explored primarily by using  
41 size and/or age as covariates. Size (bodyweight) can be incorporated into the model using  
42 two different approaches. The first approach or 'allometric size approach' includes size *a*  
43 *priori* by using a bodyweight based exponential equation with a fixed exponent of 0.75  
44 for clearance and 1 for volume of distribution [48-52]. Once size is incorporated in the  
45 model using this fixed manner, the influence of age is investigated, being the difference  
46 between actual value of the PK parameter and the 0.75 allometric equation. When  
47 incorporating age as a covariate, different age descriptors may be used like postmenstrual  
48 age (PMA), gestational age (GA) or postnatal age (PNA) [53]. The choice for any of  
49 these age descriptors is based on the results of the systematic covariate analysis as  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 described below [50, 54]. In the second approach or ‘systematic covariate analysis’,  
4 bodyweight is regarded as a covariate as any other which means that the descriptive  
5 properties on the PK parameters are evaluated in a systematic covariate analysis as  
6 described below [55-57].  
7  
8  
9

10  
11  
12 In a systematic analysis, when studying the influence of covariates, scatter plots and  
13 summary plots of individual parameter estimates and/or weighted residuals *versus*  
14 covariates are used to screen for appropriate covariates to include in the covariate sub-  
15 model. Additionally these plots are used to explore the nature of the influence of the  
16 covariate (linear, exponential, allometric, subpopulations etc). Likely candidate  
17 covariates are then added to the model (forward inclusion). The influence of each  
18 covariate on the parameters is examined separately and compared to the simple model  
19 (no covariates). To assess whether the model with covariate statistically improved the fit  
20 to the data, the difference between their objective function value, referred to as log-  
21 likelihood ratio, is calculated. This ratio is assumed to be Chi-square distributed, which  
22 means that a reduction in objective function of 3.84 is considered to be significant  
23 ( $P < 0.05$ ) [43, 58]. Beside the reduction in objective function, goodness-of-fit plots of the  
24 simple model and covariate model are explored for diagnostic purposes. Furthermore, the  
25 confidence interval of the parameter estimates, the correlation matrix (indicates the  
26 relationship between two structural parameters) and visual improvement of the individual  
27 plots are used to evaluate the model. Finally, a superior model is expected to reduce the  
28 inter-subject variance and/or the residual error terms. This procedure of covariate  
29 modeling implies that each covariate is only implemented if this can be fully justified by  
30 the data and the results of the statistic evaluations.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

46 When two or more covariates are found to significantly improve the model the covariate  
47 that reduced the objective function most is included in the model after which the other  
48 covariates are tested again for their significance. After all covariates that significantly  
49 improved the objective function are added to the simple model, a backward deletion is  
50 performed, which means that each covariate is removed from the full model, one at a  
51 time (the one which causes the smallest increase in objective function first). Retaining or  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 removing the covariate is statistically tested by the use of the objective function (Chi-  
4 square test) until each covariate has been tested.  
5  
6  
7

8  
9 In datasets containing sparse data, there may not be enough information to accurately  
10 estimate inter- and intra-individual variability. This causes the values of these parameters  
11 to shrink to 0, resulting in individual parameter estimates that are closer to the population  
12 parameter estimates than they really are. This phenomenon is called shrinkage [59].  
13 Shrinkage may cause individual predictions, individual parameter estimates and  
14 diagnostics based on them to be less reliable. It can also hide, falsely introduce or distort  
15 the shape of covariate relationships.  
16  
17  
18  
19

20 Shrinkage is the result of properties of the data and is therefore difficult to avoid. One can  
21 only be aware of the presence of shrinkage, realize the influence it may have on the  
22 covariate analysis and use diagnostics other than those based on individual predictions or  
23 individual weighted residuals in the model building and model evaluation procedures.  
24  
25  
26  
27  
28  
29

### 30 Validation of PK-PD models

31 The objective of a PK or PK-PD modeling exercise is usually not just to describe the  
32 dataset of the sample of individuals that were studied. Generally, models are used to  
33 simulate which concentrations and/or effects and their variability can be expected when  
34 different doses are given to future patients. These simulations may therefore lead to  
35 optimized dosing recommendations or to optimization of new studies for the entire  
36 population where the sample of individuals belongs to. It is often said that 'all models are  
37 wrong, but some are useful' [60]. In order to define whether a model is useful and valid  
38 for clinical and trial simulations, thorough evaluation and validation of the model is  
39 necessary. Although validations of PK models are only performed in 17% of the  
40 published paediatric studies [4] and in 28% of the adults studies [61], proper model  
41 validations are an essential step in model building. For this purpose, different evaluation  
42 and validation methods are available. As described before [62], a proper validation and  
43 evaluation procedure includes an internal model evaluation followed by an external  
44 evaluation and a prospective clinical study.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The first evaluation method is the basic internal model validation used to assess whether the model is able to describe the learning dataset (dataset used to develop the model) accurately and without bias. This evaluation should actually be considered the final stage of the model building procedure. Subsequently, in the external evaluation it is assessed whether the model is able to describe one or more external datasets (datasets other than the one used to develop the model) adequately. Alternatively if a dataset is sufficiently large the original dataset may be split in two so that the model is developed using one part (about two thirds) of the dataset and evaluated externally using the other part (one third) of the dataset. In paediatric studies, it is then especially important to stratify the data correctly and ascertain that all age groups are represented in equal proportions in both datasets.

Various techniques are available for the validation and evaluation of population PK and PK-PD models (both for internal and external validation procedures).

- Basic goodness-of-fit plots ((1) individual predicted versus observed concentrations, 2) population predicted versus observed, 3) (conditional) weighted residuals versus time and 4) (conditional) weighted residuals versus dependent variable plots).

WRES and CWRES are calculated as the following:

$$WRES = \frac{\bar{y}_i - E_{FO}(\bar{y}_i)}{\sqrt{COV_{FO}(\bar{y}_i)}} \quad (\text{equation 4})$$

$$CWRES = \frac{\bar{y}_i - E_{FOCE}(\bar{y}_i)}{\sqrt{COV_{FOCE}(\bar{y}_i)}} \quad (\text{equation 5})$$

Where  $\bar{y}_i$  is the vector of measurements,  $E(\bar{y}_i)$  is the expectation of the data and  $Cov(\bar{y}_i)$  is the covariance matrix of the data [63].

These plots are used in model building, but can also be used to ascertain that there is no trend or bias in the model predictions of the final model. Furthermore, these plots can also be used for both the internal and external evaluation of the model.

- In a bootstrap analysis new datasets are generated by resampling from the original dataset and is therefore an internal validation of the model. The new datasets are



1  
2  
3 subsequently refitted to the original model, yielding mean values and standard  
4 errors for every model parameter.

5  
6  
7 A bootstrap analysis provides information on the stability of the model and its  
8 dependence on specific individuals in the learning dataset. With the freely available  
9 PSN or Wings for NONMEM software packages an automated bootstrap analysis  
10 can be performed.

- 11  
12  
13  
14 • In a visual predictive check (VPC) [64] a PK or PD profile is simulated a 100 to  
15 1000 times and lines for the median values and their 90% prediction interval are  
16 plotted in a graph. The observed values in the internal or external dataset are  
17 subsequently plotted on top of this. It can then be visually checked whether 90% of  
18 the observations are within the indicated prediction interval and whether there is no  
19 bias in the observations compared to prediction interval. In figure 5, two examples  
20 of a VPC are given, showing when a model does not work and when a model does  
21 work on the same data.  
22  
23  
24  
25  
26  
27

28 The VPC is a simulation-based diagnostic that can be used when the PK or PD  
29 profiles for all individuals in the dataset are similar and it allows for easy  
30 interpretation of the result. For this diagnostic tool, there are not statistical tests and  
31 all evaluations are based on visual evaluations. When the individual profiles are  
32 expected to deviate largely from one another because there is for instance a large  
33 variability in the time and amount of dose administrated, or when there are many  
34 covariates, the use of this diagnostic becomes more difficult.  
35  
36  
37  
38  
39

- 40 • Another simulation-based diagnostic which can be used for both internal and  
41 external validations is the normalized prediction distribution error (NPDE) [65]. An  
42 example of an NPDE published before is shown in figure 6 [55]. This method yields  
43 information on how accurate the model predicts the median value of the  
44 observations and the variability within them. The interpretation of this diagnostic is  
45 less straightforward than for the VPC, but the advantage of this method is that it can  
46 be used when the variability in dosing regimen (both in time, amounts and rates) is  
47 high or when there is a large number of covariates in the model. This can for  
48 instance be the case for data obtained during routine paediatric clinical practice.  
49 Software (e.g. NPDE add-on package for R) [66] to perform this analysis is freely  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 available. For the NPDE, beside visual evaluation of the plots, statistical tests are  
4 available. These statistical tests are however reported to be highly sensitive and  
5 powerful, so that decisions for the model should primarily be based on visual  
6 assessments. An example is the statistically significant deviation of zero of the  
7 mean value because of the large number of data, while the actual deviation is small  
8 (e.g. 0.074) and not of clinical relevance.  
9

10  
11  
12  
13  
14  
15  
16 If the model performed well in both the evaluation procedures, the dosing algorithm that  
17 results from the PK-PD model needs to be tested and challenged in a prospective  
18 (clinical) trial. If the predictive performance of the model is corroborated by the trial it  
19 can be used with confidence in clinical practice.  
20  
21  
22

### 23 24 25 Optimal design of paediatric studies

26 When new population PK-PD studies are performed, it is important to design these  
27 studies in the most efficient manner possible to obtain maximum information about the  
28 PK and PD parameters so that they can be determined with the highest precision [51, 67].  
29 When designing PK-PD studies in paediatrics certain factors need to be taken into  
30 account e.g. age-range of the paediatric group, therapeutic index, possibility to collect  
31 blood samples, availability of validated PD endpoints for children, and the availability of  
32 sensitive analytical methods.  
33  
34  
35  
36  
37

38 When optimizing a PK or PK-PD study design, using literature data from adults or  
39 children of different age-ranges or possible *in vitro* or pre-clinical data, a concentration-  
40 time or effect-time profile for a study can be simulated. This can help to identify possible  
41 shortcomings in the design or to perform a power-analysis. Alternatively software  
42 packages are available (WINPOPT [68], PopED [67] and PFIM [69]) that can help to  
43 identify the optimal number and time points of observations in a study based on the prior  
44 information on a drug [70] . To determine the appropriate sample size certain factors,  
45 which are summarized in Table 1, need to taken into account. Each of these factors can  
46 influence the required number of patients and/or samples in a positive or negative way. In  
47 a study of Peeters *et al.* [32] only 24 patients (aged between 3 and 24 months) were  
48 required to determine both the PK and PD since rich sampling was performed (median of  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 11 samples per child) and no covariates were found in the relatively homogenous  
4 population. This in contrast to a study performed by Knibbe *et al.* [55] in which 250  
5 children were included. This higher number was required because in addition to the large  
6 dispersion in age from (preterm) neonates up to toddlers of 3 years of age, only 1 to 4  
7 samples were available for each subject. Moreover infusion rates and additional bolus  
8 doses varied for each child during the study to obtain the desired analgesic effect. In the  
9 another example [71], only 6 patients (aged between 1 and 5 years) were required in  
10 which 7 samples per patient were collected. This lower number of patients (N=6)  
11 compared to the study of Peeters *et al.* (n=24) can be explained because there often exist  
12 a lower variability in PK than in PD which results in a lower required number of patients  
13 (Table 1).  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

#### 24 Conclusions and perspectives

25  
26 In view of the European Regulation which came into force in 2007, it seems now time to  
27 use the progress that has been made in the field of integrated PK-PD modelling [72] to  
28 develop rational and individualized dosing schemes for children. Because of the  
29 possibility to analyse sparse and unbalanced datasets thereby minimizing the burden for  
30 each child, population PK-PD modelling and simulation using non-linear mixed effect  
31 modelling has become the preferred tool to develop effective and safe dosing regimens  
32 for children. Specifically in paediatrics where the developmental changes have to be  
33 taken into account, which may influence the PK and/or the PD of the drugs, this  
34 advanced statistical tool is of critical value.  
35  
36

37 Before dosing regimens can be tested in clinical practice, proper validations of the  
38 models should be performed, for which recently adequate tools have been developed.  
39 Beside internal and external validations, prospective clinical trials, which allow for the  
40 evaluation of the model based dosing regimens, are needed, not only to adjust the  
41 proposed dosing regimen but also to convince paediatricians to use the information that  
42 has been generated using these modelling exercises.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 Furthermore, one of the future goals may be to explore possibilities for cross-validation  
54 of the models, in which the reported influences of developmental changes on a certain PK  
55 or PD parameter of one drug are evaluated for use in another drug that go through the  
56  
57  
58  
59  
60

1  
2  
3 same metabolic route or share the same mechanism of action. In this respect,  
4 physiologically-based pharmacokinetic (PBPK) models are needed. PBPK models  
5 consider the physiological and biochemical processes by using *in vitro* data to describe  
6 the PK of drugs [73, 74]. The combination of these two approaches may use the  
7 information that is already available in a more optimal way in defining effective and safe  
8 dosing regimens for every individual patient.  
9

10  
11  
12  
13  
14 In conclusion, analyses of paediatric data using population PK-PD modelling and  
15 covariate analysis will result in individualized dosing regimens for children of different  
16 age, bodyweight and genetic background. Thus population PK-PD modelling constitutes  
17 an innovative approach to the study of drug effects in this very special patient population,  
18 which is otherwise difficult to study.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Acknowledgements:  
4

5 The clinical research of K Allegaert is supported by the Fund for Scientific Research,  
6 Flanders (Belgium) (FWO Vlaanderen) by a Fundamental Clinical Investigatorship  
7 (1800209N)  
8

9  
10 The clinical research of C.A.J. Knibbe is supported by the Innovational Research  
11 Incentives Scheme (Veni grant, July 2006) of the Dutch Organization for Scientific  
12 Research (NWO).  
13

14  
15 The research of E.H.J. Krekels is supported by the Top Institute Pharma project number  
16 D2-104  
17

18  
19 The authors would like to thank Pieter De Cock, Hospital Pharmacist, for critically  
20 reading the manuscript.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References:

- 1 Ernest TB, Elder DP, Martini LG, et al (2007) Developing paediatric medicines: identifying the needs and recognizing the challenges. *J Pharm Pharmacol* 59 (8): 1043-1055
- 2 t Jong GW, Vulto AG, de Hoog M, et al (2001) A survey of the use of off-label and unlicensed drugs in a Dutch children's hospital. *Pediatrics* 108 (5): 1089-1093
- 3 Conroy S, Choonara I, Impicciatore P, et al (2000) Survey of unlicensed and off label drug use in paediatric wards in European countries. *European Network for Drug Investigation in Children. BMJ* 320 (7227): 79-82
- 4 Tod M, Jullien V, Pons G (2008) Facilitation of drug evaluation in children by population methods and modelling. *Clin Pharmacokinet* 47 (4): 231-243
- 5 Bray RJ (1998) Propofol infusion syndrome in children. *Paediatr Anaesth* 8 (6): 491-499
- 6 Parke TJ, Stevens JE, Rice AS, et al (1992) Metabolic acidosis and fatal myocardial failure after propofol infusion in children: five case reports. *BMJ* 305 (6854): 613-616
- 7 Sutherland JM (1959) Fatal cardiovascular collapse of infants receiving large amounts of chloramphenicol. *AMA J Dis Child* 97 (6): 761-767
- 8 McIntyre J, Choonara I (2004) Drug toxicity in the neonate. *Biol Neonate* 86 (4): 218-221
- 9 Nederland Kenniscentrum Farmacotherapie bij Kinderen. (<http://www.kinderformularium.nl> Accessed on 29 June 2009).
- 10 European Medicines Agency (EMA): Medicines for children. (<http://www.emea.europa.eu/htms/human/paediatrics/introduction.htm> Accessed on 17 June 2009).
- 11 Dunne J (2007) The European Regulation on medicines for paediatric use. *Paediatr Respir Rev* 8 (2): 177-183
- 12 Consultant PR (2006) EU paediatric regulation for testing medicines in children. (<http://www.child-medicines-research-info.com/paediatric-research-consultancy/eu-paediatric-regulation.asp> Accessed on 23 June 2009).
- 13 Saint Raymond A, Brasseur D (2005) Development of medicines for children in Europe: ethical implications. *Paediatr Respir Rev* 6 (1): 45-51
- 14 Kearns GL, Abdel-Rahman SM, Alander SW, et al (2003) Developmental pharmacology--drug disposition, action, and therapy in infants and children. *N Engl J Med* 349 (12): 1157-1167
- 15 Grand RJ, Watkins JB, Torti FM (1976) Development of the human gastrointestinal tract. A review. *Gastroenterology* 70 (5 PT.1): 790-810
- 16 Anderson BJ, van Lingen RA, Hansen TG, et al (2002) Acetaminophen developmental pharmacokinetics in premature neonates and infants: a pooled population analysis. *Anesthesiology* 96 (6): 1336-1345
- 17 Hines RN, McCarver DG (2002) The ontogeny of human drug-metabolizing enzymes: phase I oxidative enzymes. *J Pharmacol Exp Ther* 300 (2): 355-360
- 18 de Wildt SN, Kearns GL, Leeder JS, et al (1999) Glucuronidation in humans. Pharmacogenetic and developmental aspects. *Clin Pharmacokinet* 36 (6): 439-452

- 1  
2  
3  
4 19 Leeder JS (2004) Translating pharmacogenetics and pharmacogenomics into drug  
5 development for clinical pediatrics and beyond. *Drug Discov Today* 9 (13): 567-  
6 573
- 7 20 Krekels EH, van den Anker JN, Baiardi P, et al (2007) Pharmacogenetics and  
8 paediatric drug development: issues and consequences to labelling and dosing  
9 recommendations. *Expert Opin Pharmacother* 8 (12): 1787-1799
- 10 21 Rhodin MM, Anderson BJ, Peters AM, et al (2009) Human renal function  
11 maturation: a quantitative description using weight and postmenstrual age. *Pediatr*  
12 *Nephrol* 24 (1): 67-76
- 13 22 Bartelink IH, Rademaker CM, Schobben AF, et al (2006) Guidelines on paediatric  
14 dosing on the basis of developmental physiology and pharmacokinetic  
15 considerations. *Clin Pharmacokinet* 45 (11): 1077-1097
- 16 23 Knibbe CA, Koster VS, Deneer VH, et al (1998) Determination of propofol in  
17 low-volume samples by high-performance liquid chromatography with  
18 fluorescence detection. *J Chromatogr B Biomed Sci Appl* 706 (2): 305-310
- 19 24 Rauh M, Stachel D, Kuhlen M, et al (2006) Quantification of busulfan in saliva  
20 and plasma in haematopoietic stem cell transplantation in children : validation of  
21 liquid chromatography tandem mass spectrometry method. *Clin Pharmacokinet*  
22 45 (3): 305-316
- 23 25 Edelbroek PM, van der Heijden J, Stolk LM (2009) Dried blood spot methods in  
24 therapeutic drug monitoring: methods, assays, and pitfalls. *Ther Drug Monit* 31  
25 (3): 327-336
- 26 26 Jabeen R, Payne D, Wiktorowicz J, et al (2006) Capillary electrophoresis and the  
27 clinical laboratory. *Electrophoresis* 27 (12): 2413-2438
- 28 27 Fisher DM, O'Keefe C, Stanski DR, et al (1982) Pharmacokinetics and  
29 pharmacodynamics of d-tubocurarine in infants, children, and adults.  
30 *Anesthesiology* 57 (3): 203-208
- 31 28 LeDez KM, Lerman J (1987) The minimum alveolar concentration (MAC) of  
32 isoflurane in preterm neonates. *Anesthesiology* 67 (3): 301-307
- 33 29 Blusse van Oud-Alblas HJ PM, Brill MJE, Knibbe CAJ, Klein J, Tibboel D,  
34 Weber F (2009) The pharmacodynamics of isoflurane in children using Bispectral  
35 index and composite auditory evoked potentials. .([www.page-  
36 meeting.org/?abstract=1576](http://www.page-meeting.org/?abstract=1576) Accessed on
- 37 30 Stephenson T (2005) How children's responses to drugs differ from adults. *Br J*  
38 *Clin Pharmacol* 59 (6): 670-673
- 39 31 van Dijk M, de Boer JB, Koot HM, et al (2000) The reliability and validity of the  
40 COMFORT scale as a postoperative pain instrument in 0 to 3-year-old infants.  
41 *Pain* 84 (2-3): 367-377
- 42 32 Peeters MY, Prins SA, Knibbe CA, et al (2006) Pharmacokinetics and  
43 pharmacodynamics of midazolam and metabolites in nonventilated infants after  
44 craniofacial surgery. *Anesthesiology* 105 (6): 1135-1146
- 45 33 Peeters MY, Prins SA, Knibbe CA, et al (2006) Propofol pharmacokinetics and  
46 pharmacodynamics for depth of sedation in nonventilated infants after major  
47 craniofacial surgery. *Anesthesiology* 104 (3): 466-474
- 48 34 van Dijk M, Peters JW, van Deventer P, et al (2005) The COMFORT Behavior  
49 Scale: a tool for assessing pain and sedation in infants. *Am J Nurs* 105 (1): 33-36  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3 35 Jonsson EN, Karlsson MO (1999) Xpose--an S-PLUS based population  
4 pharmacokinetic/pharmacodynamic model building aid for NONMEM. *Comput*  
5 *Methods Programs Biomed* 58 (1): 51-64  
6  
7 36 FDA C (1999) Guidance for industry. Population  
8 pharmacokinetics. (<http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm072137.pdf> Accessed on  
9  
10 37 FDA C (1998) Guidance for industry. General considerations for pediatric  
11 pharmacokinetic studies for drugs and biological products (Draft  
12 Guidance). (<http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm072114.pdf> Accessed on 22 June 2009).  
13  
14 38 Beal SL, Sheiner LB (1982) Estimating population kinetics. *Crit Rev Biomed Eng*  
15 8 (3): 195-222  
16  
17 39 Ette EI, Williams PJ, Lane JR (2004) Population pharmacokinetics III: design,  
18 analysis, and application of population pharmacokinetic Studies. *Ann*  
19 *Pharmacother* 38 (12): 2136-2144  
20  
21 40 Wade JR, Beal SL, Sambol NC (1994) Interaction between structural, statistical,  
22 and covariate models in population pharmacokinetic analysis. *J Pharmacokinet*  
23 *Biopharm* 22 (2): 165-177  
24  
25 41 Aarons L (1999) Software for population pharmacokinetics and  
26 pharmacodynamics. *Clin Pharmacokinet* 36 (4): 255-264  
27  
28 42 Beal SL SL, Broeckman AJ (2006) NONMEM Users Guides. Icon Development  
29 Solutions, Ellicott City, Maryland, USA.  
30  
31 43 Broeckman AJ SL, Beal SL (1994) NONMEM Users Guide - Part V. University  
32 of California at San Francisco, San Francisco.  
33  
34 44 Jelliffe R SA, Van Guilder M, et al. (1995) User manual for version 10.7 of the  
35 USC\*PAC Collection of PC Programs, USC Laboratory of Applied  
36 Pharmacokinetics. USC School of Medicine, Los Angeles.  
37  
38 45 Bustad A, Terziivanov D, Leary R, et al (2006) Parametric and nonparametric  
39 population methods: their comparative performance in analysing a clinical dataset  
40 and two Monte Carlo simulation studies. *Clin Pharmacokinet* 45 (4): 365-383  
41  
42 46 Mentre F, Mallet A (1994) Handling covariates in population pharmacokinetics.  
43 *Int J Biomed Comput* 36 (1-2): 25-33  
44  
45 47 Mandema JW, Verotta D, Sheiner LB (1992) Building population  
46 pharmacokinetic--pharmacodynamic models. I. Models for covariate effects. *J*  
47 *Pharmacokinet Biopharm* 20 (5): 511-528  
48  
49 48 Anderson BJ, Holford NH (2008) Mechanism-based concepts of size and maturity  
50 in pharmacokinetics. *Annu Rev Pharmacol Toxicol* 48: 303-332  
51  
52 49 Anderson BJ, Allegaert K, Holford NH (2006) Population clinical pharmacology  
53 of children: general principles. *Eur J Pediatr* 165 (11): 741-746  
54  
55 50 Anderson BJ, Allegaert K, Holford NH (2006) Population clinical pharmacology  
56 of children: modelling covariate effects. *Eur J Pediatr* 165 (12): 819-829  
57  
58 51 Meibohm B, Laer S, Panetta JC, et al (2005) Population pharmacokinetic studies  
59 in pediatrics: issues in design and analysis. *AAPS J* 7 (2): E475-487  
60  
52 Anderson BJ, Holford NH (2009) Mechanistic basis of using body size and  
maturation to predict clearance in humans. *Drug Metab Pharmacokinet* 24 (1): 25-  
36



- 1  
2  
3 53 Engle WA (2004) Age terminology during the perinatal period. *Pediatrics* 114 (5):  
4 1362-1364  
5  
6 54 Allegaert K, Anderson BJ, Cossey V, et al (2006) Limited predictability of  
7 amikacin clearance in extreme premature neonates at birth. *Br J Clin Pharmacol*  
8 61 (1): 39-48  
9  
10 55 Knibbe CA, Krekels EH, van den Anker JN, et al (2009) Morphine  
11 glucuronidation in preterm neonates, infants and children younger than 3 years.  
12 *Clin Pharmacokinet* 48 (6): 371-385  
13  
14 56 Mahmood I (2007) Prediction of drug clearance in children: impact of allometric  
15 exponents, body weight, and age. *Ther Drug Monit* 29 (3): 271-278  
16  
17 57 Mahmood I (2009) Application of fixed exponent 0.75 to the prediction of human  
18 drug clearance: an inaccurate and misleading concept. *Drug Metabol Drug*  
19 *Interact* 24 (1): 57-81  
20  
21 58 Sheiner LB, Ludden TM (1992) Population pharmacokinetics/dynamics. *Annu*  
22 *Rev Pharmacol Toxicol* 32: 185-209  
23  
24 59 Karlsson MO, Savic RM (2007) Diagnosing model diagnostics. *Clin Pharmacol*  
25 *Ther* 82 (1): 17-20  
26  
27 60 Box G (1979) Robustness in the strategy of scientific model building. In: Launer  
28 RL Wilkinson GN, editors. *Robustness in Statistics*. New York: Academic Press,  
29 pp202.  
30  
31 61 Brendel K, Dartois C, Comets E, et al (2007) Are population pharmacokinetic  
32 and/or pharmacodynamic models adequately evaluated? A survey of the literature  
33 from 2002 to 2004. *Clin Pharmacokinet* 46 (3): 221-234  
34  
35 62 Ince I, de Wildt SN, Tibboel D, et al (2009) Tailor-made drug treatment for  
36 children: creation of an infrastructure for data-sharing and population PK-PD  
37 modeling. *Drug Discov Today* 14 (5-6): 316-320  
38  
39 63 Hooker AC, Staats CE, Karlsson MO (2007) Conditional weighted residuals  
40 (CWRES): a model diagnostic for the FOCE method. *Pharm Res* 24 (12): 2187-  
41 2197  
42  
43 64 Fang L, Holford NH, Hinkle G, et al (2007) Population pharmacokinetics of  
44 humanized monoclonal antibody HuCC49deltaCH2 and murine antibody CC49 in  
45 colorectal cancer patients. *J Clin Pharmacol* 47 (2): 227-237  
46  
47 65 Brendel K, Comets E, Laffont C, et al (2006) Metrics for external model  
48 evaluation with an application to the population pharmacokinetics of gliclazide.  
49 *Pharm Res* 23 (9): 2036-2049  
50  
51 66 Comets E, Brendel K, Mentre F (2008) Computing normalised prediction  
52 distribution errors to evaluate nonlinear mixed-effect models: the npde add-on  
53 package for R. *Comput Methods Programs Biomed* 90 (2): 154-166  
54  
55 67 Foracchia M, Hooker A, Vicini P, et al (2004) POPEd, a software for optimal  
56 experiment design in population kinetics. *Comput Methods Programs Biomed* 74  
57 (1): 29-46  
58  
59 68 Duffull SB DN, Eccleston, J, Kimko, H (2008) WinPOPT User Guide Version  
60 1.2. School of Pharmacy, University of Otago, New Zealand.  
Retout S, Mentre F (2003) Optimization of individual and population designs  
using Splus. *J Pharmacokinet Pharmacodyn* 30 (6): 417-443

- 1  
2  
3 70 Duffull SB (2001) Design of clinical pharmacology trials. Clin Exp Pharmacol  
4 Physiol 28 (11): 905-912  
5  
6 71 Knibbe CA, Melenhorst-de Jong G, Mestrom M, et al (2002) Pharmacokinetics  
7 and effects of propofol 6% for short-term sedation in paediatric patients following  
8 cardiac surgery. Br J Clin Pharmacol 54 (4): 415-422  
9  
10 72 Breimer DD (2008) PK/PD modelling and beyond: impact on drug development.  
11 Pharm Res 25 (12): 2720-2722  
12  
13 73 Alcorn J, McNamara PJ (2008) Using ontogeny information to build predictive  
14 models for drug elimination. Drug Discov Today 13 (11-12): 507-512  
15  
16 74 Johnson TN (2005) Modelling approaches to dose estimation in children. Br J  
17 Clin Pharmacol 59 (6): 663-669  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

1  
2  
3  
4 Figure 1: Schematic representation of the relationship between dose and concentration (pharmacokinetics, PK) and between concentration and a pharmacological (side) effect (pharmacodynamics, PD). Important covariates which may affect both the PK and/or PD are bodyweight, age, disease status (e.g. critically ill versus healthy children) and genetics.  
5  
6  
7

8  
9 Figure 2: Simulation of propofol concentrations and response using COMFORT-B score versus time based on developed PK and PD models, after administration of different doses of propofol (0, 18, 30, and 36 mg/h) for a 10 kg and a 5 kg non-ventilated infant in the first night at the Intensive Care following craniofacial surgery. Target COMFORT-B scores are between 12 and 14 preferably. Reproduced from [Peeters MY, Prins SA, Knibbe CA, DeJongh J, van Schaik RH, van Dijk M, et al. Propofol pharmacokinetics and pharmacodynamics for depth of sedation in nonventilated infants after major craniofacial surgery. *Anesthesiology* 2006 Mar;104(3):466-74.]  
10  
11  
12  
13  
14  
15  
16

17  
18 Figure 3: Concentration-time profiles of the same study using two different approaches. In figure 3A the standard two-stage approach is applied to a rich dataset. 3B shows the population approach with mixed effect modeling applied to the same dataset using only two datapoints for each individual so a sparse dataset is created. In 3A, in each of the six individuals 10 samples are available. The different symbols correspond to different individuals. Each black line corresponds to a separate fit to the 10 data points of each individual.  
19  
20  
21  
22  
23

24 In 3B, which uses the mixed effect modeling approach, two samples of the 10 per subject in 2A are used. The different symbols correspond to the six different individuals. The black line illustrates the concentration time plot based on the population mean values of the PK parameters (PRED). The grey lines show the plots of the individual patients, which are based on the population mean values together with the measured concentrations of the specific individual (IPRED).  
25  
26  
27  
28  
29

30  
31 Figure 4: In 4A, the inter-individual variability is shown between three individuals who received the same dose. 4B presents the intra- or residual variability by showing the concentration-time profile after repeated administration. Both these random variables are assumed to be normally distributed with a mean of zero and a variance of  $\omega^2$  or  $\sigma^2$  respectively.  
32  
33  
34  
35


36 Figure 5:

37 Two examples of a visual predictive check (VPC) are illustrated based on the same dataset (warfarin concentrations and prothrombin complex activity (PCA)) using two different models. In 5A the VPC of the effect compartment model is shown, while in 5B the VPC of the turnover model is demonstrated. The median (black thick line) and the 90% intervals (black thin lines) together with the observed data (PCA) (dots) are shown. Based on both graphics, the turn over model is the most appropriate model since 90% of the observations are lying within the prediction interval. Furthermore, unlike the effect compartment model, no bias is seen in the observations.  
38  
39  
40  
41  
42  
43

44 Reproduced from [Holford N, 2005. The visual predictive check – Superiority to Standard Diagnostic (Rorschach) Plots. PAGE 14, Abstr 738. (<http://www.page-meeting.org/?abstract=738>)].  
45  
46  
47

48 Figure 6: Example of a normalized prediction distribution error (NPDE) analysis, which show the NPDE distributions for morphine. The normal distribution is presented by the solid line. The values for the mean and standard deviation of the observed NPDE distribution are given below the histogram, with \* indicating a significant difference of a mean of 0 and a variance of 1 at the  $p < 0.05$  level, as determined by the Wilcoxon signed rank test and Fisher test for variance. Additionally the distribution of NPDE vs time after the first dose and NPDE vs the log of the concentrations are also shown. The dotted lines represent the 90% distribution of the NPDE. Reproduced from [Knibbe CA, Krekels EH, van den Anker JN, DeJongh J, Santen GW, van Dijk M, et al. Morphine glucuronidation in preterm neonates, infants and children younger than 3 years. *Clin Pharmacokinet* 2009;48(6):371-85] with permission from Wolters Kluwer Health | Adis (© Adis Data Information BV [2006]. All rights reserved.)  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


FIGURE 1



Developmental changes, disease status and genetics

For Peer Review

FIGURE 2



Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

FIGURE 3






FIGURE 4



For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


FIGURE 5



Peer Review



FIGURE 6



Or Peer Review

Table 1: Factors influencing the required number of patients and/or samples per patient.

<b>Factor</b>	<b>Number of patients/samples</b>
Study of PK only	relatively small number of patients/samples
Study of PK-PD relationship	relatively high number of patients/samples
Even distribution of covariates (age, bodyweight)	↓ number of patients/samples
↑ Number of changes in dose	↑/- number of patients/samples (depending on other aspects of the study design)
↑ Number of samples/child	↓ number of patients
Use of optimal sampling strategies	↓ number of patients/samples
Different sampling windows (e.g. two or three sampling schemes)	↓ number of patients/samples