

## Medium chain fatty acid feed supplementation reduces the probability of colonization in broilers

Twan van Gerwe, Annemarie Bouma, Don Klinkenberg, Jaap A. Wagenaar,

Wilma F. Jacobs-Reitsma, Arjan Stegeman

#### ▶ To cite this version:

Twan van Gerwe, Annemarie Bouma, Don Klinkenberg, Jaap A. Wagenaar, Wilma F. Jacobs-Reitsma, et al.. Medium chain fatty acid feed supplementation reduces the probability of colonization in broilers. Veterinary Microbiology, 2010, 143 (2-4), pp.314. 10.1016/j.vetmic.2009.11.029 . hal-00594814

#### HAL Id: hal-00594814 https://hal.science/hal-00594814

Submitted on 21 May 2011  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#### Accepted Manuscript

Title: Medium chain fatty acid feed supplementation reduces the probability of *Campylobacter jejuni* colonization in broilers

Authors: Twan van Gerwe, Annemarie Bouma, Don Klinkenberg, Jaap A. Wagenaar, Wilma F. Jacobs-Reitsma, Arjan Stegeman

PII:	S0378-1135(09)00578-1
DOI:	doi:10.1016/j.vetmic.2009.11.029
Reference:	VETMIC 4684
To appear in:	VETMIC
Received date:	28-8-2009
Revised date:	18-11-2009
Accepted date:	20-11-2009

Please cite this article as: van Gerwe, T., Bouma, A., Klinkenberg, D., Wagenaar, J.A., Jacobs-Reitsma, W.F., Stegeman, A., Medium chain fatty acid feed supplementation reduces the probability of *Campylobacter jejuni* colonization in broilers, *Veterinary Microbiology* (2008), doi:10.1016/j.vetmic.2009.11.029

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1	Medium chain fatty acid feed supplementation reduces the probability of Campylobacter jejuni
2	colonization in broilers
3	
4	
5	Twan van Gerwe <sup>1*</sup> , Annemarie Bouma <sup>2</sup> , Don Klinkenberg <sup>3</sup> , Jaap A. Wagenaar <sup>2,3,4</sup> , Wilma F. Jacobs-
6	Reitsma <sup>5</sup> , Arjan Stegeman <sup>6</sup>
7	
8	<sup>1</sup> Department of Farm Animal Health, Faculty of Veterinary Medicine,
9	Utrecht University, P.O. Box 80151, 3508 TD Utrecht, the Netherlands
10	<sup>2</sup> Department of Infectious Diseases and Immunology,
11	Faculty of Veterinary Medicine, Utrecht University, P.O. Box 80.165,
12	3508 TD Utrecht, the Netherlands.
13	<sup>3</sup> Central Veterinary Institute of Wageningen UR,
14	P.O. Box 65, 8200 AB Lelystad, the Netherlands.
15	<sup>4</sup> WHO Collaborating Center for <i>Campylobacter</i> /OIE Reference Laboratory for Campylobacteriosis,
16	the Netherlands.
17	<sup>5</sup> Rikilt, Institute of Food Safety, Wageningen UR,
18	P.O. Box 230, 6700 AE Wageningen, the Netherlands.
19	
	<sup>*</sup> Corresponding author. Current affiliation: Nutreco Poultry and Rabbit Research Center, Toledo (Spain), E-mail address: <u>t.vangerwe@nutreco.com</u> . Tel.: +34 925 53 52 86. Fax.: +34 925 53 53 30.

#### 20 Abstract

21

22 Campylobacteriosis in humans is associated with handling and consumption of contaminated broiler 23 meat. Reduction of the number of *Campylobacter*-colonized broiler flocks could potentially be 24 realized by decreasing their susceptibility for colonization. The aim of this study was to determine the 25 effect of feed supplementation with a mixture of medium chain fatty acids ( $C_8$ - $C_{12}$ ) on susceptibility of 26 broilers for Campylobacter colonization, feed conversion and body weight gain. Two experiments 27 were carried out with individually housed commercial broilers. The birds were fed with medium chain 28 fatty acids supplemented feed (n=227), or received feed without supplement (n=87). The birds were 29 inoculated with a dose of *Campylobacter jejuni* varying between log<sub>10</sub> 1.19 - 5.47 CFU. During 14 30 days after inoculation, cecal or fecal samples were collected, in which the presence of C. jejuni was 31 determined by bacterial culture. Beta-binomial dose response modeling of the colonization status at 14 32 days post-inoculation was performed to estimate the C. jejuni dose necessary to colonize 50% of 33 inoculated broilers, which was estimated to be 200 times higher in broilers fed with supplemented feed 34  $(\log_{10} 4.8 \text{ CFU})$  than in control broilers  $(\log_{10} 2.5 \text{ CFU})$ . Feed conversion was not affected by feed 35 supplementation, while body weight gain was 49 g higher in broilers fed with supplemented feed. 36 These findings indicate that susceptibility of broilers for *Campylobacter* colonization is decreased by 37 supplementation with medium chain fatty acids, and that feed supplemented with this mixture may be 38 a promising tool for the reduction of *Campylobacter* colonization in commercial broiler flocks. 39

40 Keywords: Campylobacter, broiler, feed supplementation, fatty acid, dose response

- 41
- 42

#### 43 Introduction

44

45 Meat products from broilers colonized with Campylobacter jejuni are considered an important source 46 of human campylobacteriosis (Friedman et al., 2004), and reduction of human exposure is an 47 important goal of public health programs (EFSA, 2005). One way to achieve this is to prevent 48 intestinal colonization of broilers with *Campylobacter* spp., either by prevention of exposure or by 49 reducing the susceptibility for colonization (Wagenaar et al., 2006). Because highly effective 50 biosecurity measures to prevent exposure might be difficult to implement, reduction of susceptibility 51 of broilers for colonization, here defined as the probability of colonization upon exposure, may be an 52 alternative way to reduce the number of colonized flocks. 53 One way to reduce the susceptibility is to change the micro-environment in the gastro-54 intestinal tract in such a way that ingested Campylobacter bacteria are inactivated or unable to reach 55 the lower intestines. It has been shown that short and medium chain fatty acids (SCFA and MCFA, 56 respectively) have in vitro anti-Campylobacter activity that is additional to the Campylobacter 57 inactivating effect of an acidified micro-environment (Chaveerach et al., 2002; Thormar et al., 2006). 58 In vivo, feed supplemented with high level SCFA results in a reduced chance of colonization in 59 broilers, but also in reduced body weight gain (BWG), which is economically undesirable (Heres et 60 al., 2004). Lower fatty acid concentrations might be necessary to prevent BWG loss, preferably 61 combined with improved effectivity at higher pH, to extend the anti-Campylobacter activity 62 throughout a larger part of the gastro-intestinal tract. 63 Whereas SCFA activity ceases at pH  $\geq$  5.5 (Chaveerach et al., 2002), *in vitro* studies showed 64 that 1-Monoglycerid of Capric acid, a MCFA, has anti-Campylobacter activity in feed mixed with a 65 buffer at pH 5.5 and in feed mixed with tap water at pH 7.0 (Thormar et al., 2006). Consequently, 66 MCFA might inactivate *Campylobacter* cells in the crop (pH 4.5) and the intestines, where the pH is 67 approximately 5.8-6.0 (Chang and Chen, 2000; Farner, 1942). A low concentration mixture of MCFA 68 could therefore be a promising tool to decrease the susceptibility without negative effects on weight

69 gain or feed conversion.

## CCEPTED M

70 The aim of this study was to determine whether an acidified feed containing 1% MCFA was

71 able to reduce the susceptibility of broilers for colonization with Campylobacter. jejuni. Additionally,

72 the effects of supplemented feed on BWG and feed conversion rate (FCR) were determined.

- 73
- 74

#### 75 Material and methods

76

#### 77 Experimental Design

78 Two experiments were carried out subsequently. For each experiment hatching eggs, originating from 79 one commercial broiler breeder flock (Ross 308) were purchased at 17 days of incubation. Chicks 80 were hatched at the experimental facilities of the Faculty of Veterinary Medicine of Utrecht 81 University. After hatching broilers were randomly assigned to control feed (CF) or supplemented feed 82 (SF), which they received throughout the experiments. CF was an antibiotic and anti-coccidia drug 83 free, mashed diet. SF was the same diet, with 1% soybean-oil substituted by 1% Lodestar<sup>TM</sup> C8-10 84 (Loders Croklaan, Wormerveer, the Netherlands), which is produced by fractional distillation of palm 85 kernel oil free fatty acids, and typically consists of 56% C<sub>10</sub>, 30% C<sub>8</sub>, 10% C<sub>12</sub>, <3% C<sub>6</sub>, and <3% 86 other lipids. The composition and calculated chemical analyses of both diets is shown in table 1. 87 In experiment 1 (exp. 1), a group of 150 day-old chicks was provided CF, based on random 88 selection of the birds, and a group of 47 chicks was provided SF. In experiment 2 (exp. 2), chicks were 89 placed in multiple groups: 192 day-old chicks were randomly divided into 32 groups of 6 chicks each, 90 22 groups receiving CF (132 chicks) and 10 groups SF (60 chicks). Birds were housed in a well-91 controlled facility, on litter floors. Groups were separated by walls. Water and feed was available ad 92 *libitum*, and from 7 days of age a daily dark period of 6 hours was applied. 93 At 14 days of age, by random selection 114 of the initially 150 CF and 42 of 47 SF broilers 94 (exp. 1), and 113 of 132 CF and 43 of 60 SF broilers (exp. 2) were weighed and housed individually in

95 wired cages of 40x40 cm, with closed, littered floors, which were situated in four identical

96 compartments. In exp. 1, broilers in both treatment groups were randomly divided over the cages in all

97 compartments. In exp. 2, broilers per treatment group were evenly divided over the compartments, and

- 98 within each compartment, broilers were randomly assigned to cages. The sides and back of each cage
- 99 were covered with plastic sheets. The distance between the cages was 40 cm at least.
- 100
- 101 Inoculation

102 One dose of 10<sup>9</sup> CFU of *C. ieiuni* C356, originating from a broiler flock (Jacobs-Reitsma et al., 1995) 103 and stored in glycerol at -80°C, was orally administered to 3 five-day-old broilers, which were housed 104 with 3 non-inoculated contact broilers. Three days post-inoculation (PI) C. jejuni was isolated from the 105 ceca of one contact broiler. This chicken-passaged strain, here referred to as C. jejuni C356P (C356P), 106 was stored in glycerol at  $-80^{\circ}$ C and used in experiments 1 and 2. 107 Before inoculation, C356P was freshly cultured in Heart Infusion Broth (micro-aerobically, 108 37°C, overnight) and diluted in saline to obtain the intended inoculation doses. Based on power 109 calculations, using results from a pilot study (data not shown), and presuming an additive effect on the 110 inoculation dose required to colonized 50% of the birds ( $CD_{50}$ ) of approx. log<sub>10</sub> 1.5 CFU, SF broilers 111 were inoculated with higher doses C356P than CF broilers (table 2). In each compartment, non-112 inoculated broilers were housed in 4 randomly selected cages (sentinel) to detect Campylobacter 113 transmission. Broilers, except sentinels, were orally inoculated with 0.25 ml of the C356P inoculation 114 suspensions at 14 days of age (exp. 1; n=140) or 18 days of age (exp. 2; n=139; one broiler died one 115 day prior to inoculation). The inoculation doses were randomly divided over the compartments. To 116 limit variation between individuals, all broilers were feed-deprived for 11 hours prior to inoculation, 117 and provided feed directly afterwards. The concentration of Campylobacter in the administered

- 118 inocula was determined by plating on modified charcoal cefoperazone deoxycholate agar (mCCDA)
- 119 (Biotrading Benelux B.V., Mijdrecht, the Netherlands).

120

121 Sampling and Testing

122 Six (exp. 1) or seven days (exp. 2) prior to inoculation, broilers were tested for the presence of

- 123 Campylobacter by culture of a fecal dropping on mCCDA. All broilers tested negative. After
- 124 inoculation, birds were sampled at 4, 8, and 11 days post inoculation (PI) by swabbing fresh cecal

## CCEPTED M

125 droppings if present, or otherwise a swab of a fresh fecal dropping. If neither could be obtained, a swab from cloacal content was taken. In addition, at 1, 2, 3, 7, and 9 days PI 1 sentinel per 126 127 compartment and 1 to 5 inoculated broilers per inoculation dose were placed in cardboard boxes with 128 wire floors for 4 hours, to collect cloacal swabs and fresh cecal droppings (or fecal droppings if not 129 present). Different material was sampled with the intention to compare the sensitivity of each of these 130 sampling methods. Because positivity was limited to one of two sample types only incidentally, 131 sensitivity of different sample mediums was considered equal. At the end of the trial (14 days PI), 132 cecal contents were sampled after euthanasia by cervical dislocation. 133 The person sampling the broilers was blinded to dose groups and could not distinguish 134 sentinels from inoculated broilers. Long-sleeved plastic gloves were changed for each broiler to avoid 135 cross-contamination. Samples were collected with sterile swabs and transported to the laboratory in 136 modified Amies transport medium without charcoal (Biotrading, Mijdrecht, the Netherlands) within 6 137 hours. Samples were streaked on mCCDA plates and incubated micro-aerobically at 42°C, and 138 examined for the presence of Campylobacter-suspect colonies after 24 and 48 hours. Microscopic 139 examination of morphology and motility was used as confirmation. Broilers were considered 140 *Campylobacter*-colonized and were excluded from further sampling when at least one sample tested 141 positive. Ethical aspects of the experiments were judged and approved by the animal ethical 142 committee of Utrecht University. 143 To assess the effect of MCFA feed supplementation on technical performance, BWG and feed 144

145

#### 146 Statistical Analyses

147 The effect of MCFA feed supplementation on susceptibility was assessed by fitting a Beta-binomial 148 dose response model (Teunis and Havelaar, 2000). According to this model, inoculation with dose D 149 results in a probability of colonization  $Pr_{inoc}(D)$ :

intake during the period of individual housing (>14 days of age) were recorded in exp. 2.

150 
$$\operatorname{Pr}_{inoc} D = 1 - \left(1 + \frac{D}{\beta}\right)^{-\alpha}$$

The underlying assumption of the model is that each bacterium can independently establish
colonization, but that hosts may differ in their susceptibility (Teunis and Havelaar, 2000). The two
parameters (instead of single probabilities for each dose) provide opportunities to assess the dose
response relation over a wide range of exposure doses, and to compare treatments tested with different
ranges of inoculation doses. Furthermore, the dose-response model could be used to calculate a CD<sub>50</sub>:
the colonization dose resulting in a 50% probability of infection.

157 Because sentinels were detected positive after day 4 PI in exp. 2 (table 2), colonization as a 158 result of transmission could not be excluded for the inoculated birds either. Therefore, in the main 159 analysis (using data up to day 14 PI) we corrected for transmission by estimating a transmission 160 probability  $Pr_{tr}$ , different for each room and treatment group. Consequently  $Pr_{+}(D)$ , the probability of 161 being colonized, was equal to

162 
$$Pr_{+} D = 1 - 1 - Pr_{tr} 1 - Pr_{inoc} D$$

163 Additionally, a more simple analysis was performed using the colonization status up to day 4 164 PI, without the correction for transmission. In the day-4 analysis,  $Pr_+(D)$  was equal to  $Pr_{inoc}(D)$ . 165 For both analyses (day-4 and day-14), four different models were fitted to see if there were 166 group and treatment effects: the first with separate dose response relations for each of the 4 treatment 167 groups (CF<sub>1</sub> vs. CF<sub>2</sub> vs. SF<sub>1</sub> vs. SF<sub>2</sub>), the second with combined CF groups (CF<sub>12</sub>) vs. combined SF 168 groups (SF<sub>12</sub>), the third with combined exp. 1 groups vs. exp. 2 groups, and the fourth with all groups 169 combined. All parameters were estimated by maximum likelihood. The corrected Akaike Information 170 Criterium (AIC<sub>c</sub>) (Hurvich and Tsai, 1989) was used to decide which model explained the data best 171 and whether different dose response relations should be adopted for (combined) treatment groups. 172 Prior to inoculation, some groups of broilers got diarrhea at 1 week of age. To study the 173 possible confounding effect of diarrhea on susceptibility, data of exp. 2 were analyzed as described 174 above, in 4 groups, with diarrhea status instead of experiment. AIC<sub>c</sub> was used to assess if the occurrence of this symptom was associated with an increased or decreased susceptibility to 175 176 *Campylobacter* colonization.

177	Linear regression analyses (SPSS 15.0.1.) were performed to assess if feed treatment affected
178	BWG and FCR in exp. 2. Sixty-nine broilers, which were detected Campylobacter-colonized before or
179	at 4 days PI, were weighed and sexed at 28 days of age, while the remaining 84 broilers (19 colonized
180	and 65 non colonized) were weighed and sexed at 32 days of age. To correct for this, age and final
181	colonization status were included as dependent variables, next to the variables sex and feed treatment,
182	resulting in the equations
183	$BWG_i = \beta_0 + \beta_1 Feed_{i1} + \beta_2 Sex_{i2} + \beta_3 Age_{i3} + \beta_4 Col_{i4} + \varepsilon_i,$
184	and,
185	$FCR_{j} = \beta_{0} + \beta_{1}Feed_{j1} + \beta_{2}Sex_{j2} + \beta_{3}Age_{j3} + \beta_{4}Col_{j4} + \varepsilon_{j}.$
186	
187	
188	Results
189	
190	Colonization
191	In exp. 1, 35 of 46 colonized birds, and in exp. 2, 72 of 89 colonized birds were detected
192	Campylobacter-positive in the first 4 days PI.(table 2). In exp. 1, the sentinels remained negative, but
193	in exp. 2, 4 of 16 sentinels were detected Campylobacter-positive, at days 7, 8, 8 and 14 days PI. Four
194	birds in exp. 1 and one bird in exp. 2 died after day 4 PI (table 2).
195	
196	Dose Response: Effect on Susceptibility
197	The best fitting day-14 model (lowest $AIC_c$ ) included separate dose response relation for $CF_{12}$
198	and $SF_{12}$ , indicating equal relations in both experiments. Dose response relations of $CF_{12}$ and $SF_{12}$
199	show parallel sigmoid shapes (figure 1). $CD_{50}$ for $CF_{12}$ was $log_{10}$ 2.5 CFU (95% CI: 2.2-2.8) and $CD_{50}$
200	for $SF_{12}$ was $log_{10}$ 4.8 CFU (95% CI: 4.4-5.2) (figure 1). The AIC <sub>c</sub> of alternative models were at least
201	3.95 higher.
202	The best day-4 model included separate dose response relations for all four treatment groups
203	$(CF_1, CF_2, SF_1, and SF_2)$ . $CD_{50}$ for $CF_1$ and $CF_2$ were $log_{10} 2.8$ CFU (95% CI: 2.6-3.3) and 2.1 CFU

204	(95% CI: 1.8-2.3), respectively. $CD_{50}$ for $SF_1$ and $SF_2$ were $log_{10}$ 8.5 CFU (95% CI: 4.8 - $\infty$ ) and 4.7
205	CFU (95% CI: 4.2-5.2), respectively. The AIC <sub>c</sub> of alternative models were at least 12.2 higher.

In exp. 2, diarrhea occurred around 1 week of age, with similar frequencies in both treatment groups (14/22 CF groups and 5/10 SF groups; Fisher exact test: P=0.70), suggesting that diarrhea was not feed related. In a separate analysis of data of exp. 2 inclusion of a variable describing whether a

209 broiler originated from a group with diarrhea did not result in a better fit, with AIC<sub>c</sub> being approx. 4

- 210 points higher in both the day-4 and day-14 models.
- 211

#### 212 Effect on technical performance

The broiler that died during the exp. 2 (table 2) and a broiler that was lame during the last few days of the experiment were excluded from this analysis. BWG was estimated  $49 \pm 24$  g higher in SF broilers compared to CF broilers (p=0.044) when correcting for the effect of sex (p<0.001), age (p<0.001), and final colonization status (p=0.398). FCR was estimated 0.061 ± 0.034 lower in broilers with were provided supplemented feed (p=0.075) when correcting for sex (p=0.179), age (p=0.183), and final colonization status (p=0.735).

219

220

#### 221 **Discussion**

222

223 In two experiments the effect of MCFA feed supplementation on susceptibility and technical 224 performance of broilers was studied. The effect on susceptibility was assessed by determining the 225 relation between the C. jejuni inoculation dose and the subsequent occurrence of colonization, 226 resulting in dose response curves of SF broilers shifted to the right compared to CF broilers, indicating 227 that SF broilers required a higher inoculation dose to become colonized than CF broilers. The effect of 228 MCFA feed supplementation could also have been assessed by comparing the percentage of colonized 229 broilers exposed to equal inoculation doses (log<sub>10</sub> 2.19 CFU in exp. 1, and log<sub>10</sub> 2.47 and 4.37 CFU in 230 exp. 2) at 14 days PI (table 2). Although this would also have illustrated the susceptibility reducing 231 effect of the treatment (Fisher exact test: all p < 0.05), we would not have been able to predict the

probability of colonization for other inoculation doses, nor would we have been able to correct fortransmission.

234 We used the Beta-binomial dose response model (Teunis and Havelaar, 2000) to estimate dose 235 response curves and CD<sub>50</sub>s. Besides the fact that this analysis resulted in less parameters to be 236 estimated than separate analyses for different doses, and in the possibility to use different doses for 237 different treatments, it also turned out useful to correct for transmission. We used the colonization 238 status at 14 days PI to estimate the dose response relations, but when the *Campylobacter* colonization 239 status at 4 days PI was used in the alternative (day-4) model, similar estimates were obtained. This 240 similarity suggests that most broilers colonized after 4 days PI were colonized by transmission. Based 241 on the results in this study, challenge experiments with *Campylobacter* might not necessarily have to 242 last longer than 4 days to estimate the dose response relation properly. 243 Analysis of technical performance showed that BWG was increased in SF broilers, while FCR

243 Analysis of technical performance showed that BwO was increased in SF orohers, while FCR 244 was not affected. The effect on BWG might have been caused by antimicrobial effects of the fatty 245 acids, as feed supplementation with Capric Acid and Lauric acid, two MCFAs, has been shown to 246 decrease the concentration of *Clostridium perfringens* in jejunum and ileum of *C. perfringens* 247 challenged broilers (Jansman et al., 2006). Although feed supplementation with antimicrobial agents 248 has the potential to improve feed efficiency (Dibner and Richards, 2005; Jansman et al., 2006), in this 249 study no significant effect on FCR was observed.

- 250
- 251

#### 252 Conclusion

253

The number of *C. jejuni* bacteria required to colonize 50% of inoculated broilers was estimated 200 times higher in broilers fed with supplemented feed than in control broilers. Although the working mechanism of supplemented feed remains to be elucidated, this effect on susceptibility is a promising finding for the implementation of MCFA feed supplementation as an intervention for reduction of susceptibility in broilers. As the *Campylobacter* exposure dose that broilers experience in the field is

- 259 unknown, field trials are necessary to determine to what extent MCFA supplementation reduces
- 260 *Campylobacter* colonization in the field.
- 261
- 262
- 263 **References**
- 264
- Chang, M.H., Chen, T.C., 2000. Reduction of *Campylobacter jejuni* in a simulated chicken digestive
  tract by *Lactobacilli* cultures. J. Food Prot. 63, 1594-1597.
- Chaveerach, P., Keuzenkamp, D.A., Urlings, H.A., Lipman, L.J., van Knapen, F., 2002. In vitro study
  on the effect of organic acids on *Campylobacter jejuni/coli* populations in mixtures of water
  and feed. Poult. Sci. 81, 621-628.
- Dibner, J.J., Richards, J.D., 2005. Antibiotic growth promoters in agriculture: history and mode of
  action. Poult. Sci. 84, 634-643.
- EFSA 2005. Opinion of the Scientific Panel on biological hazards (BIOHAZ) related to
- 273 *Campylobacter* in animals and foodstuffs. http://www.efsa.europa.eu/EFSA/efsa\_locale-
- 274 1178620753812\_1178620776955.htm
- Farner, D.S., 1942. The hydrogen ion concentration in avian digestive tracts. Poult. Sci. 21, 445-450.
- 276 Friedman, C.R., Hoekstra, R.M., Samuel, M., Marcus, R., Bender, J., Shiferaw, B., Reddy, S., Ahuja,
- S.D., Helfrick, D.L., Hardnett, F., Carter, M., Anderson, B., Tauxe, R.V., 2004. Risk factors
  for sporadic *Campylobacter* infection in the United States: A case-control study in FoodNet
- 279 sites. Clin. Infect. Dis. 38, S285-S296.
- Heres, L., Engel, B., Urlings, H.A., Wagenaar, J.A., van Knapen, F., 2004. Effect of acidified feed on
- susceptibility of broiler chickens to intestinal infection by *Campylobacter* and *Salmonella*.
- 282 Vet. Microbiol. 99, 259-267.
- Hurvich, C.M., Tsai, C.L., 1989. Regression and Time-Series Model Selection in Small Samples.
  Biometrika 76, 297-307.
- 285 Jacobs-Reitsma, W.F., van de Giessen, A.W., Bolder, N.M., Mulder, R.W., 1995. Epidemiology of
- 286 *Campylobacter* spp. at two Dutch broiler farms. Epidemiol. Infect. 114, 413-421.

287	Jansman, A.J.M., Wagenaars, C.M.F., Schonewille, A., Snel, H., Van der Klis, J.D. 2006. Bestrijding
288	van Clostridium en Campylobacter infecties in pluimvee via natuurlijke antimicrobiële
289	voedingscomponenten (Animal Sciences Group - Wageningen UR), p. 50.
290	Teunis, P.F.M., Havelaar, A.H., 2000. The Beta Poisson dose-response model is not a single-hit
291	model. Risk Anal. 20, 513-520.
292	Thormar, H., Hilmarsson, H., Bergsson, G., 2006. Stable Concentrated Emulsions of the 1-
293	Monoglyceride of Capric Acid (Monocaprin) with Microbicidal Activities against the Food-
294	Borne Bacteria Campylobacter jejuni, Salmonella spp., and Escherichia coli. Appl. Environ.
295	Microbiol. 72, 522-526.
296	
297	
220	

299 Table 1. Ingredient composition and calculated analysis of the diets

Item	Amount (%)	
Ingredients	Control Feed	Supplemented Feed
Wheat	36.53	36.53
Soybean meal	21.20	21.20
Corn	20.00	20.00
Peas	15.00	15.00
Soybean oil	3.95	2.95
Calcium carbonate	1.44	1.44
MCFA <sup>b</sup>	0.00	1.00
Monocalcium phosphate	0.57	0.57
Lysine 65%	0.35	0.35
Sodium chloride	0.26	0.26
Methionine	0.25	0.25
Premix	0.25	0.25
Sodium bicarbonate	0.10	0.10
Threonine	0.04	0.04
Choline chloride	0.03	0.03
Endoxylanase	0.02	0.02
Phytase	0.01	0.01
Calculated analysis		
СР	19.60	19.60
$ME_n (kcal/kg)^*$	2790	2790
$C8^{\dagger}$	0.00	0.30
C10 <sup>b</sup>	0.00	0.56
C12 <sup>b</sup>	0.00	0.10

#### 300

 $^{*}$  Energetic value of MCFA and soybean oil was assumed to be equal  $^{\dagger}$  1% Lodestar^{TM} C8-10

<sup>†</sup> 1% Lodestar<sup>TM</sup> C8-10

- **301 Table 2.** Colonization status of broilers (exp. 1 and exp. 2).
- 302 Rows represent different feed treatment groups (CF = control feed; SF = supplemented feed) and *C. jejuni* C356P inoculation
- 303 doses (log<sub>10</sub> CFU/broiler). Nominators in cells express the number of broilers detected *Campylobacter*-positive, and
- 304 denominators express the number of broilers sampled throughout 4 or 14 days post-inoculation (PI).

Received and the second second

305

Exp. 1				Exp. 2			X	
treatment	Dose	Time (days PI)		treatment	Dose	Time (day	Time (days PI)	
group	C356P	4	14	group	C356P	4	14	
CF	-	0/16	0/16	CF	-	0/16	4/16	
CF	1.19	0/33	1/32 <sup>a</sup>	CF	1.47	8/32	13/32	
	2.19	9/33	12/32 <sup>b</sup>		2.47	21/33	25/33	
	3.19	20/32	21/31 <sup>a</sup>		3.47	30/32	31/32	
SF	2.19	1/21	2/21	SF	2.47	0/10	2/10	
	4.49	5/21	9/20 <sup>a</sup>		3.47	2/11	2/11	
					4.47	2/10	3/9 <sup>a</sup>	
					5.47	9/11	9/11	

306 <sup>a</sup> A broiler died after being tested negative at 4 dpi.

307 <sup>b</sup> A broiler died after being tested positive at 4 dpi.

308

- **309** Figure 1. Dose response curves for the final model of the day-14 analysis
- 310 The relation between inoculation dose en the probability of colonization occurring as a result of inoculation, Pr<sub>inoc</sub>(D), up to
- 311 14 days PI, for combined experiments (1 and 2). Best fitted curves, with 95% confidence bounds, represent relations for
- 312 control feed (CF<sub>12</sub>) and MCFA supplemented feed (SF<sub>12</sub>) fed broilers. Dots represent raw data as observed in the
- 313 experiments, not corrected for the occurrence of transmission.
- 314 << Figure 1.tif >>
- 315

