



HAL
open science

On Schrodinger equations with modified dispersion

Rémi Carles

► **To cite this version:**

| Rémi Carles. On Schrodinger equations with modified dispersion. 2011. hal-00595434v2

HAL Id: hal-00595434

<https://hal.science/hal-00595434v2>

Preprint submitted on 25 May 2011 (v2), last revised 31 May 2011 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON SCHRÖDINGER EQUATIONS WITH MODIFIED DISPERSION

RÉMI CARLES

ABSTRACT. We consider the nonlinear Schrödinger equation with a modified spatial dispersion, given either by an homogeneous Fourier multiplier, or by a bounded Fourier multiplier. Arguments based on ordinary differential equations yield ill-posedness results which are sometimes sharp. When the Fourier multiplier is bounded, we infer that no Strichartz-type estimate improving on Sobolev embedding is available. Finally, we show that when the symbol is bounded, the Cauchy problem may be ill-posed in the case of critical regularity, with arbitrarily small initial data. The same is true when the symbol is homogeneous of degree one, where scaling arguments may not even give the right critical value.

1. INTRODUCTION

For $(t, x) \in \mathbf{R}_+ \times \mathbf{R}^d$, we consider

$$(1.1) \quad i\partial_t u + P(D)u = \lambda |u|^{2\sigma} u \quad ; \quad u|_{t=0} = u_0,$$

where $D = -i\nabla_x$, $P : \mathbf{R}^d \rightarrow \mathbf{R}$, $\lambda \in \mathbf{R}$ and $\sigma > 0$. Since the Fourier multiplier P is real-valued, the free flow ($\lambda = 0$) generates a unitary group on $\dot{H}^s(\mathbf{R}^d)$, $s \in \mathbf{R}$:

$$S(t) = e^{-itP(D)}.$$

We consider two cases:

- P is homogeneous, $P(\mu\xi) = \mu^m P(\xi)$, for all $\xi \in \mathbf{R}^d$, $\mu > 0$, with $m \geq 1$.
- P is bounded, $P \in L^\infty(\mathbf{R}^d)$.

The first case includes the standard Schrödinger operator ($P(\xi) = -|\xi|^2$), and the fourth-order Schrödinger operator ($P(\xi) = |\xi|^4$), studied for instance in [17, 18]. Smoothing effects have been established for rather general Fourier multipliers P in [13]. The case $d = 1$, $P(\xi) = \xi^{2j+1}$, $j \in \mathbf{N}$, has been studied initially in [14]. The case $m \in 2\mathbf{N}$, with P elliptic and $\nabla^2 P$ non-degenerate outside $\{\xi = 0\}$, appears as a particular case of the framework in [9]. It is shown there that if $s > \max(0, s_0)$, then the Cauchy problem (1.1) is locally well-posed in $H^s(\mathbf{R}^d)$, where

$$(1.2) \quad s_0 = \frac{d}{2} - \frac{m}{2\sigma}.$$

This index corresponds to the one given by scaling arguments: if u solves (1.1), then for any $k > 0$, $u_k : (t, x) \mapsto k^{m/(2\sigma)} u(k^m t, kx)$ solves the same equation. The value of s for which the $\dot{H}^s(\mathbf{R}^d)$ -norm is invariant under $u \mapsto u_k$ is $s = s_0$. We will see in §1.3 that this scaling argument may not yield the sharp Sobolev regularity: if $m = 1$, the Cauchy problem (1.1) may be strongly ill-posed in $H^s(\mathbf{R}^d)$ for all $s \leq d/2$.

In [9], the proof of local well-posedness uses dispersive and Strichartz estimates for S , established in [7] for $d = 1$, and in [8] for $d \geq 2$. Note that in the case $d = 1$, dispersive and Strichartz estimates for S are proved in [12] for $P(\xi) = |\xi|^m$ and $m \geq 2$

2010 *Mathematics Subject Classification*. Primary 35Q55; Secondary 35A01, 35B30, 35B33, 35B45.
This work was supported by the French ANR project R.A.S. (ANR-08-JCJC-0124-01).

(not necessarily an integer). By resuming arguments similar to those presented in [4, 6], we prove that in this framework, the index s_0 is sharp, in the sense that the nonlinear flow map fails to be uniformly continuous at the origin in $H^s(\mathbf{R}^d)$, if $s < s_0$. This property has been established in [18] for the case $P(\xi) = |\xi|^4$ with $(d, \sigma) = (3, 1)$. However, when P is not elliptic, the index s_0 may not correspond to the critical Sobolev regularity (see §1.3).

The second case, where P is bounded, is motivated by the results presented in [10], where $P(\xi) = -\frac{1}{h} \arctan(h|\xi|^2)$ is considered to construct numerical approximations of the solution to the linear Schrödinger equation

$$i\partial_t u + \Delta u = V(x)u,$$

and $0 < h \ll 1$ denotes the time step. We will see below that in such a framework, no Strichartz estimate is available, even if one is ready to pay some loss of derivative. Another example of bounded symbol one may think of is

$$i\partial_t u + \Delta(1 - \Delta)^{-1}u = \lambda|u|^{2\sigma}u.$$

1.1. Norm inflation. Our result in this direction is:

Theorem 1.1. *Let $d \geq 1$, $\lambda \in \mathbf{R}$, $\sigma > 0$. Assume that either σ is an integer, or that there exists an integer r such that $2\sigma \geq r > d/2$.*

1. Suppose that P is m -homogeneous, with $m \geq 1$, and denote $s_0 = d/2 - m/(2\sigma)$. Suppose that $s_0 > 0$ and let $0 < s < s_0$. There exists a family $(u_0^h)_{0 < h \leq 1}$ in $\mathcal{S}(\mathbf{R}^d)$ with

$$\|u_0^h\|_{H^s(\mathbf{R}^d)} \rightarrow 0 \text{ as } h \rightarrow 0,$$

and a solution u^h to (1.1) and $0 < t^h \rightarrow 0$, such that:

$$\|u^h(t^h)\|_{H^s(\mathbf{R}^d)} \rightarrow +\infty \text{ as } h \rightarrow 0.$$

2. If P is bounded, then the above conclusion remains true for any $0 < s < d/2$.

Theorem 1.1 is proved in §2, by adapting the ordinary differential equation mechanism used in, e.g., [4, 6]. However, the critical Sobolev regularity s_c may satisfy $s_c > s_0$ (see §1.3): in view of [9] and Theorem 1.1, we have $s_c = s_0$ at least when $P(\xi) = \mu|\xi|^m$, $\mu \in \mathbf{R} \setminus \{0\}$, $m \in 2\mathbf{N} \setminus \{0\}$.

1.2. Absence of Strichartz estimates. In this paragraph, we focus our discussion on the case where P is bounded. For $s > d/2$, $H^s(\mathbf{R}^d)$ being an algebra, local well-posedness in $H^s(\mathbf{R}^d)$ is straightforward, provided that the nonlinearity is sufficiently smooth (see e.g. [19]). Therefore, the critical threshold is $s_c = d/2$, and from Theorem 1.1, no dispersive property is present to decrease this number. More precisely, no Strichartz estimate is available for $S(\cdot)$, even if one is ready to pay some loss of regularity which is not worse than the result provided by Sobolev embedding. To state this property precisely, we recall the standard definition.

Definition 1.2. *A pair $(p, q) \neq (2, \infty)$ is admissible if $p \geq 2$, $q \geq 2$, and*

$$\frac{2}{p} = d \left(\frac{1}{2} - \frac{1}{q} \right).$$

By Sobolev embedding, for all (p, q) (not necessarily admissible) with $2 \leq q < \infty$, there exists $C > 0$ such that for all $u_0 \in H^{d/2-d/q}(\mathbf{R}^d)$, and all finite time interval I ,

$$\begin{aligned} \|S(\cdot)u_0\|_{L^p(I; L^q(\mathbf{R}^d))} &\leq C \|S(\cdot)u_0\|_{L^p(I; H^{d/2-d/q}(\mathbf{R}^d))} \\ &\leq C \|u_0\|_{L^p(I; H^{d/2-d/q}(\mathbf{R}^d))} = C |I|^{1/p} \|u_0\|_{H^{d/2-d/q}(\mathbf{R}^d)}. \end{aligned}$$

When P is bounded, this estimate cannot be improved:

Corollary 1.3. *Let $d \geq 1$, and $P \in L^\infty(\mathbf{R}^d; \mathbf{R})$. Suppose that there exist an admissible pair (p, q) , an index $k \in \mathbf{R}$, a time interval $I \ni 0$, $|I| > 0$, and a constant $C > 0$ such that*

$$\|S(\cdot)u_0\|_{L^p(I; L^q(\mathbf{R}^d))} \leq C\|u_0\|_{H^k(\mathbf{R}^d)}, \quad \forall u_0 \in H^k(\mathbf{R}^d).$$

Then necessarily, $k \geq 2/p = d/2 - d/q$.

The fact that no standard Strichartz estimate (with no loss) is available for $S(\cdot)$ is rather clear, since the dispersion relation is given by $\tau = P(\xi)$, and defines a characteristic variety which is bounded in τ . However, one could expect the existence of Strichartz estimates with loss of regularity, in the same fashion as in [1, 2, 3] (where the geometric framework — the space variable belongs to a compact manifold — rules out the existence of the standard dispersive properties). The fact that this is not so is a rather direct consequence of Theorem 1.1 (where $\sigma > 0$ is arbitrary), and of the argument given in [3] to prove Proposition 3.1. It may seem surprising to prove Corollary 1.3 as a consequence of a nonlinear analysis; we insist on the fact that the proof of Theorem 1.1 is rather simple, and the deduction of Corollary 1.3 involves another nonlinear result, whose proof is also quite short (see Proposition 3.2 below).

1.3. Critical cases. In [5], local well-posedness in $H^{s_c}(\mathbf{R}^d)$ for small data is established for Equation (1.1) in the case $P(\xi) = -|\xi|^2$, where

$$s_c = \frac{d}{2} - \frac{1}{\sigma}$$

coincides with s_0 in that case, since $m = 2$. In [16], local well-posedness in $H^{d/2}(\mathbf{R}^d)$ for small data is established for the same operator, with nonlinearities which are allowed to grow exponentially. In these two papers, the proof uses Strichartz estimates (in Besov spaces). On the other hand, when P is bounded, the Cauchy problem may be ill-posed in $H^{d/2}(\mathbf{R}^d)$, even for nonlinearities growing algebraically. Moreover, when P is m -homogeneous with $m = 1$, the critical Sobolev regularity may not be s_0 , but $s_c = d/2 > s_0$, with ill-posedness for $s = s_c$.

Proposition 1.4. *Let $\lambda \in \mathbf{R} \setminus \{0\}$, $\sigma > 0$. Assume that either σ is an integer, or that there exists an integer r such that $2\sigma \geq r > d/2$. In either of the two cases:*

- $P(\xi) = c \cdot \xi$, $c \in \mathbf{R}^d$, or
- P is constant,

for all $\delta > 0$, there exists $u_0 \in H^{d/2}(\mathbf{R}^d)$ with $\|u_0\|_{H^{d/2}(\mathbf{R}^d)} \leq \delta$ such that (1.1) has a unique solution $u \in C(\mathbf{R}_+; \mathcal{D}'(\mathbf{R}^d))$, and for any $t > 0$, $u(t, \cdot) \notin H^{d/2}(\mathbf{R}^d)$.

Unlike what happens in the presence of Strichartz estimates ([5, 16]), this result yields examples where local well-posedness fails in the critical case $s = s_c$, even for small data. We prove Proposition 1.4 in §4: we present the cases $d = 2$, $\sigma > 0$, and $d \geq 1$, $\sigma = 1/2$, only, for the convenience of the exposition, but the argument can be extended to any space dimension, up to more intricate computations.

In the case $m = 1$ (at least), the mere assumption that P is m -homogeneous is not enough to characterize the critical Sobolev regularity in (1.1), or the existence of Strichartz estimates. Indeed, the symbol $P(\xi) = |\xi|$ corresponds to the wave equation, for which Strichartz estimates are available when $d \geq 2$, and so $s_c < d/2$. See e.g. [11]. This suggests that also when $m > 1$, the value s_0 may not be sharp when P is not proportional to $|\xi|^m$, but for instance of the form $|\xi|^{m-1}c \cdot \xi$, $c \in \mathbf{R}^d$, or more generally when P is not elliptic; see also [13] for remarks in this direction.

2. PROOF OF THEOREM 1.1

The proof of Theorem 1.1 proceeds along the same lines as in [4, 6] (see also [19]). Fix s as in Theorem 1.1. Consider initial data of the form

$$u_0^h(x) = h^{s-d/2} \kappa^h a_0\left(\frac{x}{h}\right),$$

with $0 < h \ll 1$, $a_0(x) = e^{-|x|^2}$, and

$$\kappa^h = \left(\log \frac{1}{h}\right)^{-\theta}$$

for some $\theta > 0$ to be fixed later. We have $\|u_0^h\|_{H^s(\mathbf{R}^d)} \xrightarrow{h \rightarrow 0} 0$. Introduce the scaling

$$\psi(\tau, y) = h^{d/2-s} u^h(h^{2+\alpha}\tau, hy),$$

for some α to be precised later:

$$\|\psi(\tau)\|_{\dot{H}^s(\mathbf{R}^d)} = \|u^h(h^{2+\alpha}\tau)\|_{\dot{H}^s(\mathbf{R}^d)}.$$

Denote $\varepsilon = h^{2\sigma(d/2-s)-2-\alpha}$. The function ψ solves the Cauchy problem

$$(2.1) \quad i\varepsilon \partial_\tau \psi + h^{2\sigma(d/2-s)} P(h^{-1}D_y) \psi = \lambda |\psi|^{2\sigma} \psi; \quad \psi|_{\tau=0} = \kappa^h a_0.$$

2.1. Choice of α . When P is m -homogeneous, Equation (2.1) simplifies to

$$i\varepsilon \partial_\tau \psi + h^{2\sigma(d/2-s)-m} P(D_y) \psi = \lambda |\psi|^{2\sigma} \psi; \quad \psi|_{\tau=0} = \kappa^h a_0.$$

For $\omega > 0$, we set

$$2 + \alpha = \frac{1}{m + \omega} \left((m - 1 + \omega) 2\sigma \left(\frac{d}{2} - s \right) + m \right),$$

in which case we have:

$$\varepsilon = h^{2\sigma(s_0-s)/(m+\omega)}.$$

Therefore, $\varepsilon \rightarrow 0$ as $h \rightarrow 0$ since $s < s_0$. We also compute

$$h^{2\sigma(d/2-s)-m} = \varepsilon^{m+\omega}.$$

When P is bounded, we consider $2 + \alpha = \sigma(d/2 - s)$. Therefore,

$$2\sigma \left(\frac{d}{2} - s \right) - 2 - \alpha > 0 \quad (\text{hence } \varepsilon \rightarrow 0 \text{ as } h \rightarrow 0), \quad \text{and } 2 + \alpha > 0.$$

2.2. The ODE approximation. Introduce the solution to

$$i\varepsilon \partial_\tau \varphi = \lambda |\varphi|^{2\sigma} \varphi; \quad \varphi|_{\tau=0} = \kappa^h a_0.$$

It is given by

$$\varphi(\tau, y) = \kappa^h a_0(y) \exp\left(-i\lambda \frac{\tau}{\varepsilon} (\kappa^h)^{2\sigma} |a_0(y)|^{2\sigma}\right).$$

Since a_0 is a Gaussian, $\varphi \in C^\infty(\mathbf{R} \times \mathbf{R}^d)$ regardless of $\sigma > 0$, and for any $r \geq 0$,

$$(2.2) \quad \|\varphi(\tau)\|_{H^r(\mathbf{R}^d)} \lesssim (\kappa^h)^{1+2\sigma r} \left(\frac{\tau}{\varepsilon}\right)^r + \kappa^h.$$

Lemma 2.1. *Let $r > d/2$, with in addition $r \leq 2\sigma$ if $\sigma \notin \mathbf{N}$. In either of the cases of Theorem 1.1, we can find $\delta > 0$ independent of $\theta > 0$ such that*

$$\sup_{0 \leq \tau \leq \varepsilon (\log \frac{1}{\varepsilon})^\delta} \|\psi(\tau) - \varphi(\tau)\|_{H^r(\mathbf{R}^d)} \xrightarrow{\varepsilon \rightarrow 0} 0.$$

Proof. Denote by $w = \psi - \varphi$ the error. It solves

$$\begin{aligned} i\varepsilon\partial_\tau w + h^{2\sigma(d/2-s)}P(h^{-1}D_y)w &= h^{2\sigma(d/2-s)}P(h^{-1}D_y)\varphi \\ &+ \lambda(|w + \varphi|^{2\sigma}(w + \varphi) - |\varphi|^{2\sigma}\varphi), \end{aligned}$$

with $w|_{\tau=0} = 0$. Using the facts that P is real-valued, $z \mapsto |z|^{2\sigma}z$ is sufficiently smooth, and $H^r(\mathbf{R}^d)$ is an algebra, we have

$$\begin{aligned} \|w(\tau)\|_{H^r(\mathbf{R}^d)} &\lesssim \frac{1}{\varepsilon} \int_0^\tau \left\| h^{2\sigma(d/2-s)}P(h^{-1}D_y)\varphi(\tau') \right\|_{H^r(\mathbf{R}^d)} d\tau' \\ &+ \frac{1}{\varepsilon} \int_0^\tau \left(\|w(\tau')\|_{H^r(\mathbf{R}^d)}^{2\sigma} + \|\varphi(\tau')\|_{H^r(\mathbf{R}^d)}^{2\sigma} \right) \|w(\tau')\|_{H^r(\mathbf{R}^d)} d\tau'. \end{aligned}$$

In the case where P is m -homogeneous, we have

$$\begin{aligned} \left\| h^{2\sigma(d/2-s)}P(h^{-1}D_y)\varphi(\tau') \right\|_{H^r(\mathbf{R}^d)} &\lesssim h^{2\sigma(d/2-s)-m} \|\varphi(\tau')\|_{H^{r+m}(\mathbf{R}^d)} \\ &\lesssim h^{2\sigma(s_0-s)} \|\varphi(\tau')\|_{H^{r+m}(\mathbf{R}^d)}. \end{aligned}$$

In the case where P is bounded, we have

$$\begin{aligned} \left\| h^{2\sigma(d/2-s)}P(h^{-1}D_y)\varphi(\tau') \right\|_{H^r(\mathbf{R}^d)} &\lesssim h^{2\sigma(d/2-s)} \|\varphi(\tau')\|_{H^r(\mathbf{R}^d)} \\ &\lesssim h^{2\sigma(s_0-s)} \|\varphi(\tau')\|_{H^r(\mathbf{R}^d)}, \end{aligned}$$

where we set $s_0 = d/2$ in this case.

In both cases, we check that there exists $\beta > 0$ (independent of θ) such that

$$h^{2\sigma(s_0-s)} = \varepsilon^{1+\beta}.$$

It is given by the formula

$$\beta = \frac{2\sigma(s_0 - d/2) + 2 + \alpha}{2\sigma(d/2 - s) - 2 - \alpha}.$$

In the homogeneous case, this formula becomes $\beta = m - 1 + \omega$, and when P is bounded, $\beta = 1$. Therefore, in view of (2.2) and since $\kappa^h \leq 1$, there exist $\beta, \gamma > 0$ such that

$$\left\| h^{2\sigma(d/2-s)}P(h^{-1}D_y)\varphi(\tau') \right\|_{H^r(\mathbf{R}^d)} \leq \varepsilon^{1+\beta} \left(\left(\frac{\tau}{\varepsilon} \right)^\gamma + 1 \right).$$

So long as $\|w(\tau)\|_{H^r(\mathbf{R}^d)} \leq 1$, with τ as above, we infer from (2.2):

$$\|w(\tau)\|_{H^r(\mathbf{R}^d)} \lesssim \int_0^\tau \varepsilon^\beta \left(\left(\frac{\tau'}{\varepsilon} \right)^\gamma + 1 \right) d\tau' + \frac{1}{\varepsilon} \int_0^\tau \left(1 + \left(\frac{\tau'}{\varepsilon} \right)^\gamma \right) \|w(\tau')\|_{H^r(\mathbf{R}^d)} d\tau'.$$

Gronwall's Lemma yields

$$\|w(\tau)\|_{H^r(\mathbf{R}^d)} \lesssim \varepsilon^\beta \left(\left(\frac{\tau}{\varepsilon} \right)^\gamma + 1 \right) e^{C\tau/\varepsilon + C(\tau/\varepsilon)^{\gamma+1}} \lesssim \varepsilon^\beta e^{2C\tau/\varepsilon + C(\tau/\varepsilon)^{\gamma+1}}.$$

By choosing $\delta > 0$ sufficiently small, the right hand side is controlled by, say, $\varepsilon^{\beta/2}$, for all $0 \leq \tau \leq \varepsilon \left(\log \frac{1}{\varepsilon} \right)^\delta$. The condition $\|w(\tau)\|_{H^r(\mathbf{R}^d)} \leq 1$ is verified for such times τ , provided that ε is sufficiently small, and the lemma follows. \square

2.3. Conclusion. Let $r > d/2$ as in Lemma 2.1. With $\delta > 0$ given by Lemma 2.1, we have, since $s < d/2$:

$$\begin{aligned} \left\| u^h \left(h^{2+\alpha} \varepsilon \left(\log \frac{1}{\varepsilon} \right)^\delta \right) \right\|_{H^s(\mathbf{R}^d)} &\geq \left\| \varphi \left(\varepsilon \left(\log \frac{1}{\varepsilon} \right)^\delta \right) \right\|_{H^s(\mathbf{R}^d)} \\ &\quad - C \left\| \varphi \left(\varepsilon \left(\log \frac{1}{\varepsilon} \right)^\delta \right) - \psi \left(\varepsilon \left(\log \frac{1}{\varepsilon} \right)^\delta \right) \right\|_{H^r(\mathbf{R}^d)}. \end{aligned}$$

On the other hand, similar to (2.2), we have:

$$\|\varphi(\tau)\|_{H^s(\mathbf{R}^d)} \gtrsim (\kappa^h)^{1+2\sigma s} \left(\frac{\tau}{\varepsilon}\right)^s - C\kappa^h,$$

and so,

$$\left\| \varphi \left(\varepsilon \left(\log \frac{1}{\varepsilon} \right)^\delta \right) \right\|_{H^s(\mathbf{R}^d)} \geq C \left(\log \frac{1}{\varepsilon} \right)^{s\delta - \theta - 2\sigma\theta s} - o(1).$$

For $\theta > 0$ sufficiently small, $s\delta - \theta - 2\sigma\theta s > 0$, and Lemma 2.1 yields

$$\left\| u^h \left(h^{2+\alpha} \varepsilon \left(\log \frac{1}{\varepsilon} \right)^\delta \right) \right\|_{H^s(\mathbf{R}^d)} \xrightarrow{h \rightarrow 0} +\infty.$$

Theorem 1.1 follows, with

$$t^h = h^{2+\alpha} \varepsilon \left(\log \frac{1}{\varepsilon} \right)^\delta = Ch^{2\sigma(d/2-s)} \left(\log \frac{1}{h} \right)^\delta \xrightarrow{h \rightarrow 0} 0.$$

3. PROOF OF COROLLARY 1.3

We argue by contradiction, by using a slight generalization of [3, Proposition 3.1].

Definition 3.1 (From [15]). *Let $s \in \mathbf{R}$. The Cauchy problem (1.1) is well posed in $H^s(\mathbf{R}^d)$ if, for all bounded subset $B \subset H^s(\mathbf{R}^d)$, there exist $T > 0$ and a Banach space X_T continuously embedded into $C([0, T]; H^s(\mathbf{R}^d))$ such that for all $u_0 \in H^s(\mathbf{R}^d)$, (1.1) has a unique solution $u \in X_T$, and the mapping $u_0 \mapsto u$ is uniformly continuous from $(B, \|\cdot\|_{H^s})$ to $C([0, T]; H^s(\mathbf{R}^d))$.*

Proposition 3.2. *Let $d \geq 1$, $P : \mathbf{R}^d \rightarrow \mathbf{R}$. Suppose that there exist an admissible pair (p, q) , an index $k < 2/p = d/2 - d/q$, $T_0 > 0$, and a constant $C > 0$ such that*

$$(3.1) \quad \|S(\cdot)u_0\|_{L^p([0, T_0]; L^q(\mathbf{R}^d))} \leq C\|u_0\|_{H^k(\mathbf{R}^d)}, \quad \forall u_0 \in H^k(\mathbf{R}^d).$$

Then for all

$$k + \frac{d}{q} < s < \frac{d}{2}, \quad 0 < \sigma < \frac{p}{2},$$

the Cauchy problem for (1.1) is well posed in $H^s(\mathbf{R}^d)$.

Since in Theorem 1.1, we can always consider $\sigma = 1$, Theorem 1.1 and Proposition 3.2 imply Corollary 1.3 in the non-endpoint case $p > 2$. The endpoint case then follows by interpolation with the case $(p, q) = (\infty, 2)$: if an endpoint Strichartz estimate (with some loss) was available, then a non-endpoint would be as well.

Proof. For $0 < T \leq T_0$, introduce

$$X_T = C([0, T]; H^s(\mathbf{R}^d)) \cap L^p([0, T]; W^{\ell, q}(\mathbf{R}^d)),$$

where $\ell = s - k$. By assumption, $\ell > d/q$, so we have

$$X_T \subset L^p([0, T]; L^\infty(\mathbf{R}^d)).$$

This space is equipped with the norm

$$\|u\|_{X_T} = \sup_{0 \leq t \leq T} \|u(t)\|_{H^s(\mathbf{R}^d)} + \left\| (1 - \Delta)^{\ell/2} u \right\|_{L^p([0, T]; L^q(\mathbf{R}^d))}.$$

We construct the solution to (1.1) by a fixed point argument. Set

$$\Phi(u)(t) = S(t)u_0 - i\lambda \int_0^t S(t - \tau) (|u(\tau)|^{2\sigma} u(\tau)) d\tau.$$

We prove that for $T \in]0, T_0]$ sufficiently small, Φ is a contraction on some ball of X_T centered at the origin. In view of (3.1) and Minkowski inequality,

$$\begin{aligned} \|\Phi(u)\|_{X_T} &\lesssim \|u_0\|_{H^s(\mathbf{R}^d)} + \int_0^t \left\| |u(\tau)|^{2\sigma} u(\tau) \right\|_{H^s(\mathbf{R}^d)} d\tau \\ &\lesssim \|u_0\|_{H^s(\mathbf{R}^d)} + \int_0^t \|u(\tau)\|_{L^\infty(\mathbf{R}^d)}^{2\sigma} \|u(\tau)\|_{H^s(\mathbf{R}^d)} d\tau \\ &\lesssim \|u_0\|_{H^s(\mathbf{R}^d)} + T^\gamma \|u\|_{L^p([0, T]; L^\infty(\mathbf{R}^d))}^{2\sigma} \|u\|_{L^\infty([0, T]; H^s(\mathbf{R}^d))}, \end{aligned}$$

with $\gamma = 1 - 2\sigma/p > 0$. Therefore,

$$\|\Phi(u)\|_{X_T} \leq C \|u_0\|_{H^s(\mathbf{R}^d)} + CT^\gamma \|u\|_{X_T}^{2\sigma+1}.$$

Similarly,

$$\|\Phi(u) - \Phi(v)\|_{X_T} \leq CT^\gamma \left(\|u\|_{X_T}^{2\sigma} + \|v\|_{X_T}^{2\sigma} \right) \|u - v\|_{X_T}.$$

This yields the local well-posedness result stated in Proposition 3.2. \square

4. ILL-POSEDNESS

The key remark is that all the cases of Proposition 1.4 boil down to an ordinary differential equation mechanism. Denote by v the solution to

$$i\partial_t v = \lambda |v|^{2\sigma} v; \quad v|_{t=0} = u_0.$$

When $P(\xi) = c \cdot \xi$, we have

$$u(t, x) = v(t, x + ct),$$

and when $P(\xi) = c$, we have

$$u(t, x) = v(t, x) e^{ict},$$

so it suffices to prove Proposition 1.4 in the case $P = 0$. For fixed $x \in \mathbf{R}^d$, we have

$$(4.1) \quad v(t, x) = u_0(x) e^{-i\lambda t |u_0(x)|^{2\sigma}}.$$

The idea is then that $H^{d/2}(\mathbf{R}^d)$ is not an algebra.

Consider

$$(4.2) \quad u_0(x) = \delta \times \left(\log \frac{1}{|x|} \right)^\alpha \chi(|x|^2), \quad x \in \mathbf{R}^d,$$

with $\chi \in C_0^\infty(\mathbf{R})$, $\chi = 1$ near the origin, and $\text{supp} \chi \subset]-1, 1[$. We compute

$$\nabla v(t, x) = e^{-i\lambda t |u_0(x)|^{2\sigma}} \nabla u_0(x) - 2i\sigma \lambda t |u_0(x)|^{2\sigma} \nabla u_0(x).$$

We split the proof into three cases: for $d = 2$, the proof is straightforward, for $d \geq 4$ even, the proof is similar but we omit the details of computations, and for d odd, we simply sketch the argument.

Case $d = 2$. First, $u_0 \in H^1(\mathbf{R}^2)$ provided that $\alpha < 1/2$. Now Proposition 1.4 follows if we can choose $\alpha < 1/2$ so that $|u_0|^{2\sigma} \nabla u_0 \notin L^2(\mathbf{R}^2)$. Near the origin, we have, leaving out the constants,

$$\left| |u_0(x)|^{2\sigma} \nabla u_0(x) \right|^2 \approx \frac{1}{|x|^2} (\log |x|)^{4\alpha\sigma + 2\alpha - 2}.$$

The right hand side fails to be in $L_{\text{loc}}^1(\mathbf{R}^2)$ if we impose $4\alpha\sigma + 2\alpha \geq 1$. So Proposition 1.4 follows, with

$$\frac{1}{4\sigma + 2} \leq \alpha < \frac{1}{2}.$$

Case $d \geq 4$ even. The argument is the same as in the case $d = 2$, with more computations that we simply sketch. We check by induction that for $k \geq 1$, there exist coefficients $(\beta_{jk})_{1 \leq j \leq k}$ such that near the origin,

$$\partial_r^k u_0(x) = \frac{1}{r^k} \sum_{j=1}^k \beta_{jk} \left(\log \frac{1}{|x|} \right)^{\alpha-j}, \quad \text{with } \beta_{1k} = (-1)^{k-1} (k-1)! \alpha.$$

Therefore, the asymptotic behavior of $\partial_r^k u_0$ near the origin is given by:

$$\partial_r^k u_0(x) \underset{r \rightarrow 0}{\sim} (-1)^{k-1} (k-1)! \frac{\alpha}{r^k} \left(\log \frac{1}{|x|} \right)^{\alpha-1}.$$

Like in the case $d = 2$, $u_0 \in H^{d/2}(\mathbf{R}^d)$ provided that $\alpha < 1/2$. We compute, for $t > 0$, and as $x \rightarrow 0$:

$$(4.3) \quad |\partial_r^k v(t, x)| = \frac{1}{r^k} \left(c_k(t) \left(\log \frac{1}{|x|} \right)^{\gamma_k} + \mathcal{O} \left(\left(\log \frac{1}{|x|} \right)^{\gamma_k - \omega_k} \right) \right),$$

with $c_k(t) > 0$, $\omega_k > 0$, and

$$\gamma_k = \max((2\sigma k + 1)\alpha - k, (2\sigma + 1)\alpha - 1).$$

For $t > 0$, $v(t, \cdot) \notin H^{d/2}(\mathbf{R}^d)$ if, for $k = d/2$, the first term in (4.3) is almost in $L_{\text{loc}}^2(\mathbf{R}^d)$, but not quite: we choose α so that $2\gamma_{d/2} = -1$. We find (like for $d = 2$)

$$\alpha = \frac{1}{4\sigma + 2},$$

which is consistent with the requirement $\alpha < 1/2$. Thus, the first term in (4.3) is not in $L_{\text{loc}}^2(\mathbf{R}^d)$ due to a logarithmic divergence, while the remainder term is in $L_{\text{loc}}^2(\mathbf{R}^d)$, since $\omega_k > 0$.

The case when d is odd. We keep u_0 of the same form as in even dimensions, since we have found a value for α which does not depend on d even:

$$u_0(x) = \delta \times \left(\log \frac{1}{|x|} \right)^{1/(4\sigma+2)} \chi(|x|^2), \quad x \in \mathbf{R}^d.$$

Recall the characterization of $H^s(\mathbf{R}^d)$ when $s \in]0, 1[$: a function $f \in L^2(\mathbf{R}^d)$ belongs to $H^s(\mathbf{R}^d)$, $s \in]0, 1[$, if and only if

$$\iint_{\mathbf{R}^d \times \mathbf{R}^d} \frac{|f(x) - f(x+y)|^2}{|y|^{d+2s}} dx dy < \infty.$$

When $d = 1$, we check that $u_0 \in H^{1/2}(\mathbf{R})$. We can also check that for $t > 0$, $v(t, \cdot) \notin H^{1/2}(\mathbf{R})$.

When $d \geq 3$, we compute $\partial_r^k u_0$ and $\partial_r^k v$ in the same fashion as above, and check that

$$\nabla^{[d/2]} u_0 \in H^{1/2}(\mathbf{R}^d), \quad \text{and for } t > 0, \quad \nabla^{[d/2]} v(t, \cdot) \notin H^{1/2}(\mathbf{R}^d).$$

We leave out the details, since the technicalities are more involved than in the even dimensional case, and we believe that proving Proposition 1.4 in details is not worth such an effort.

Acknowledgement. The author is grateful to Valeria Banica for valuable comments on this work.

REFERENCES

- [1] R. ANTON, *Strichartz inequalities for Lipschitz metrics on manifolds and nonlinear Schrödinger equation on domains*, Bull. Soc. Math. France, 136 (2008), pp. 27–65.
- [2] M. BLAIR, H. SMITH, AND C. SOGGE, *On Strichartz estimates for Schrödinger operators in compact manifolds with boundary*, Proc. Amer. Math. Soc., 136 (2008), pp. 247–256.
- [3] N. BURQ, P. GÉRARD, AND N. TZVETKOV, *Strichartz inequalities and the nonlinear Schrödinger equation on compact manifolds*, Amer. J. Math., 126 (2004), pp. 569–605.
- [4] ———, *Multilinear eigenfunction estimates and global existence for the three dimensional nonlinear Schrödinger equations*, Ann. Sci. École Norm. Sup. (4), 38 (2005), pp. 255–301.
- [5] T. CAZENAVE AND F. WEISSLER, *The Cauchy problem for the critical nonlinear Schrödinger equation in H^s* , Nonlinear Anal., 14 (1990), pp. 807–836.
- [6] M. CHRIST, J. COLLIANDER, AND T. TAO, *Ill-posedness for nonlinear Schrödinger and wave equations*. Archived as <http://arxiv.org/abs/math/0311048>.
- [7] S. CUI, *Pointwise estimates for a class of oscillatory integrals and related L^p - L^q estimates*, J. Fourier Anal. Appl., 11 (2005), pp. 441–457.
- [8] ———, *Pointwise estimates for oscillatory integrals and related L^p - L^q estimates. II. Multidimensional case*, J. Fourier Anal. Appl., 12 (2006), pp. 605–627.
- [9] S. CUI AND C. GUO, *Well-posedness of higher-order nonlinear Schrödinger equations in Sobolev spaces $H^s(\mathbb{R}^n)$ and applications*, Nonlinear Anal., 67 (2007), pp. 687–707.
- [10] A. DEBUSSCHE AND E. FAOU, *Modified energy for split-step methods applied to the linear Schrödinger equation*, SIAM J. Numer. Anal., 47 (2009), pp. 3705–3719.
- [11] J. GINIBRE AND G. VELO, *Generalized Strichartz inequalities for the wave equation*, J. Funct. Anal., 133 (1995), pp. 50–68.
- [12] N. HAYASHI AND P. I. NAUMKIN, *Asymptotic properties of solutions to dispersive equation of Schrödinger type*, J. Math. Soc. Japan, 60 (2008), pp. 631–652.
- [13] C. KENIG, G. PONCE, AND L. VEGA, *Oscillatory integrals and regularity of dispersive equations*, Indiana Univ. Math. J., 40 (1991), pp. 33–69.
- [14] ———, *Higher-order nonlinear dispersive equations*, Proc. Amer. Math. Soc., 122 (1994), pp. 157–166.
- [15] ———, *On the ill-posedness of some canonical dispersive equations*, Duke Math. J., 106 (2001), pp. 617–633.
- [16] M. NAKAMURA AND T. OZAWA, *Nonlinear Schrödinger equations in the Sobolev space of critical order*, J. Funct. Anal., 155 (1998), pp. 364–380.

- [17] B. PAUSADER, *Global well-posedness for energy critical fourth-order Schrödinger equations in the radial case*, Dyn. Partial Differ. Equ., 4 (2007), pp. 197–225.
- [18] ———, *The cubic fourth-order Schrödinger equation*, J. Funct. Anal., 256 (2009), pp. 2473–2517.
- [19] T. TAO, *Nonlinear dispersive equations*, vol. 106 of CBMS Regional Conference Series in Mathematics, Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2006. Local and global analysis.

UNIV. MONTPELLIER 2 & CNRS, UMR 5149, MATHÉMATIQUES, CC 051, F-34095 MONTPELLIER
E-mail address: Remi.Carles@math.cnrs.fr