



HAL
open science

Mechanistic insights into antitumor effects of new dinuclear cis Pt complexes containing aromatic linkers

Lenka Zerzankova, Hana Kostrhunova, Marie Vojtiskova, Olga Novakova, Tereza Suchankova, Miaoxin Lin, Zijian Guo, Jana Kasparkova, Viktor Brabec

► **To cite this version:**

Lenka Zerzankova, Hana Kostrhunova, Marie Vojtiskova, Olga Novakova, Tereza Suchankova, et al.. Mechanistic insights into antitumor effects of new dinuclear cis Pt complexes containing aromatic linkers. *Biochemical Pharmacology*, 2010, 80 (3), pp.344. 10.1016/j.bcp.2010.04.013 . hal-00596277

HAL Id: hal-00596277

<https://hal.science/hal-00596277>

Submitted on 27 May 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Mechanistic insights into antitumor effects of new dinuclear cis Pt^{II} complexes containing aromatic linkers

Authors: Lenka Zerzankova, Hana Kostrhunova, Marie Vojtiskova, Olga Novakova, Tereza Suchankova, Miaoxin Lin, Zijian Guo, Jana Kasparikova, Viktor Brabec



PII: S0006-2952(10)00258-3
DOI: doi:10.1016/j.bcp.2010.04.013
Reference: BCP 10530

To appear in: *BCP*

Received date: 6-3-2010
Revised date: 7-4-2010
Accepted date: 9-4-2010

Please cite this article as: Zerzankova L, Kostrhunova H, Vojtiskova M, Novakova O, Suchankova T, Lin M, Guo Z, Kasparikova J, Brabec V, Mechanistic insights into antitumor effects of new dinuclear cis Pt^{II} complexes containing aromatic linkers, *Biochemical Pharmacology* (2008), doi:10.1016/j.bcp.2010.04.013

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Mechanistic insights into antitumor effects of new dinuclear cis

Pt^{II} complexes containing aromatic linkers

Lenka Zerzankova ^a, Hana Kostrhunova ^a, Marie Vojtiskova ^a, Olga Novakova ^a,

Tereza Suchankova ^b, Miaoxin Lin ^c, Zijian Guo ^c, Jana Kasparikova ^{a,*}

Viktor Brabec^{a,*}

^a *Institute of Biophysics, Academy of Sciences of the Czech Republic, v.v.i., CZ-61265 Brno, Czech Republic*

^b *Department of Experimental Physics, Faculty of Sciences, Palacky University, tr. Svobody 26, 77146 Olomouc, Czech Republic*

^c *State Key Laboratory of Coordination Chemistry, School of Chemistry and Chemical Engineering, Nanjing University, Nanjing 210093, PR China*

Classification: (1) Antibiotics and Chemotherapeutics

* Corresponding authors at: Institute of Biophysics, Academy of Sciences of the Czech Republic, v.v.i., Kralovopolska 135, CZ-61265 Brno, Czech Republic. Tel.: +420-541517148; fax: +420-541240499.

E-mail addresses: jana@ibp.cz (J. Kasparikova); brabec@ibp.cz (V. Brabec)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abbreviations: BBR3464, [$\{trans\text{-PtCl}(\text{NH}_3)_2\}_2\text{-}\mu\text{-}trans\text{-Pt}(\text{NH}_3)_2(\text{NH}_2(\text{CH}_2)_6\text{NH}_2)_2$] $^{4+}$;
BBR3610, [$\{trans\text{-PtCl}(\text{NH}_3)_2\}_2\text{-}\mu\text{-}\{trans\text{-}(\text{H}_2\text{N}(\text{CH}_2)_6\text{NH}_2(\text{CH}_2)_2\text{NH}_2(\text{CH}_2)_6\text{NH}_2)\}_2$] $^{4+}$; bp,
base pair; cisplatin, *cis*-diamminedichloroplatinum(II); CFE, cell-free extract; CL, crosslink;
complex **1**, $\{[cis\text{-Pt}(\text{NH}_3)_2\text{Cl}]_2(4,4'\text{-methylenedianiline})\}^{2+}$; complex **2**, $\{[cis\text{-}$
 $\text{Pt}(\text{NH}_3)_2\text{Cl}]_2(\alpha,\alpha'\text{-diamino-p-xylene})\}^{2+}$; CT, calf-thymus; DPP, differential pulse
polarography; EtBr, ethidium bromide; FAAS, flameless atomic absorption
spectrophotometry; HMG, high-mobility-group; IC₅₀, concentration inhibiting cell growth by
50%; MTT, [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]; PAGE,
polyacrylamide gel electrophoresis; r_b , the number of molecules of the metal complex bound
per nucleotide residue; r_i , the molar ratio of free metal complex to nucleotide-phosphates at
the onset of incubation with DNA; SDS, sodium dodecyl sulfate.

ABSTRACT

1
2 The primary objective was to understand more deeply the molecular mechanism underlying
3
4 different antitumor effects of dinuclear Pt^{II} complexes containing aromatic linkers of different
5
6 length, {[*cis*-Pt(NH₃)₂Cl]₂(4,4'-methylenedianiline)}²⁺ (**1**) and {[*cis*-Pt(NH₃)₂Cl]₂(α,α' -
7
8 diamino-p-xylene)}²⁺ (**2**). These complexes belong to a new generation of promising
9
10 polynuclear platinum drugs resistant to decomposition by sulfur nucleophiles which hampers
11
12 clinical use of bifunctional polynuclear trans Pt^{II} complexes hitherto tested. Results obtained
13
14 with the aid of methods of molecular biophysics and pharmacology reveal differences and
15
16 new details of DNA modifications by **1** and **2** and recognition of these modifications by
17
18 cellular components. The results indicate that the unique properties of DNA interstrand cross-
19
20 links of this class of polynuclear platinum complexes and recognition of these cross-links may
21
22 play a prevalent role in antitumor effects of these metallodrugs. Moreover, the results show
23
24 for the first time a strong specific recognition and binding of high-mobility-group domain
25
26 proteins, which are known to modulate antitumor effects of clinically used platinum drugs, to
27
28 DNA modified by a polynuclear platinum complex.
29
30
31
32
33
34
35
36
37

38 Key words: dinuclear platinum; antitumor; DNA damage; interstrand cross-link; recognition
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

1
2
3
4
5 The polynuclear platinum compounds represent a class of new antitumor metallodrugs
6
7 that is structurally distinct from conventional *cis*-diamminedichloridoplatinum(II) (cisplatin)
8
9 and its mononuclear analogues, and whose clinical profile and mechanism of action are
10
11 different from these established platinum mononuclear compounds [1-6]. DNA adducts of
12
13 polynuclear platinum complexes, whose formation is associated with antitumor effects of
14
15 these agents, differ significantly in structure and type from those of mononuclear platinum
16
17 complexes. Especially because of markedly more distant leaving groups, long-range intra- and
18
19 interstrand crosslinks (CLs) are formed in DNA which affect DNA conformation, are
20
21 recognized by specific proteins and repaired differently in comparison with the CLs of
22
23 cisplatin and its mononuclear analogues [7-11]. Importantly, cells with resistance to cisplatin
24
25 showed no cross-resistance to polynuclear platinum compounds [12-14]. The polynuclear
26
27 platinum complexes also exhibit significantly higher levels of cellular uptake very likely
28
29 because of their lipophilic linker chains and hydrophilic platinum-amine coordination spheres,
30
31 which may enhance membrane permeability [13, 15, 16].
32
33
34
35
36
37
38

39 The biological activity of polynuclear platinum complexes may be modulated by the
40
41 geometry and number of leaving groups in the coordination sphere of platinum atoms as well
42
43 as by the nature of linkers connecting the platinum centers. In contrast with the mononuclear
44
45 complexes, such as antitumor cisplatin and clinically ineffective transplatin, in the dinuclear
46
47 case both geometries are antitumor active [2], although DNA adducts (CLs) as well as local
48
49 conformational distortions on DNA and their recognition by cellular components are affected
50
51 by geometry [7, 10, 17]. The dinuclear *cis* isomer $[\{cis\text{-PtCl}(\text{NH}_3)_2\}_2(\text{H}_2\text{N}-(\text{CH}_2)_6\text{-NH}_2)]^{2+}$ is
52
53 kinetically more inert in its reactions with DNA and in double-stranded DNA produces more
54
55 interstrand crosslinks than its trans counterpart [10].
56
57
58
59
60
61
62
63
64
65

Also importantly, the general structure, such as [$\{trans\text{-PtCl}(\text{NH}_3)_2\}_2\text{-}\mu\text{-Y}\}^{n+}$ (Y = $\text{NH}_2(\text{CH}_2)_6\text{NH}_2$, spermidine, spermine (BBR3610), $trans\text{-Pt}(\text{NH}_3)_2(\text{H}_2\text{N}(\text{CH}_2)_6\text{NH}_2)_2$ (BBR3464), etc.), is susceptible to decomposition by the sulfur nucleophiles [18] since substitution of the Pt-Cl bond by a trans-influencing S donor results in bridge cleavage [19-21]. Hence, the antitumor polynuclear $trans$ Pt^{II} complexes, such as for instance trinuclear BBR3464 or dinuclear BR3610, undergo deactivation upon reaction with sulfur nucleophiles, which could result in decreased bioavailability, as well as in increased amounts of toxic metabolites in the host system, which may limit their clinical use [19]. On the other hand, the cis geometry as in [$\{cis\text{-PtCl}(\text{NH}_3)_2\}_2\text{-}\mu\text{-Y}\}^{n+}$ should preserve the main features of antitumor polynuclear Pt^{II} complexes but with enhanced stability to metabolic deactivation.

In an effort to design a polynuclear platinum compound that maintains the target (DNA) binding profile of the cytotoxic Pt^{II} polynuclear compounds and is less susceptible to metabolic decomposition, the long-chain bifunctional dinuclear cis platinum complexes [$\{\text{PtCl}(\text{DACH})\}_2\text{-}\mu\text{-H}_2\text{N}(\text{CH}_2)_6\text{NH}_2(\text{CH}_2)_2\text{NH}_2(\text{CH}_2)_6\text{NH}_2\}^{4+}$ (DACH = 1,2-diaminocyclohexane) and [$\{cis\text{-Pt}(\text{NH}_3)_2\text{Cl}\}_2(4,4'\text{-methylenedianiline})\}^{2+}$ (complex **1**, Fig. 1) were synthesized [22, 23]. Hence, both dinuclear cis Pt^{II} complexes exhibit enhanced stability to metabolic deactivation [22, 23].

Platinum–DNA binding, recognition of platinated DNA by proteins and DNA repair are important pharmacological features dictating a platinum drug's efficacy [2, 24, 25]. As regards molecular mechanisms of antitumor polynuclear Pt^{II} complexes, a great attention has been so far paid mostly to polynuclear $trans$ Pt^{II} complexes whose clinical application is limited due to their lowered stability to metabolic deactivation. Thus, the important aim of this study was to broaden theoretical background of biological (antitumor) effects of dinuclear cis Pt^{II} complexes, which exhibit enhanced stability to metabolic deactivation. The primary objective in the present study was to understand more deeply those aspects of the molecular

1 mechanism of action of the new dinuclear *cis* Pt^{II} complex **1** containing a semi-rigid linker
2 and for comparative purposes the molecular mechanism of action of a structurally cognate
3
4 dinuclear *cis* Pt^{II} complex containing a shorter, more rigid linker, {[*cis*-Pt(NH₃)₂Cl]₂(α,α' -
5 diamino-p-xylene)}²⁺ (complex **2**, Fig. 1) which are related to DNA modification by these
6
7 complexes, recognition of these modifications by proteins and DNA repair. Here, we describe
8
9 experiments revealing (i) new details of the DNA binding mode of **1** and **2** in cell-free media;
10
11 (ii) repair of DNA adducts of **1** and **2** (since the integrity of this process in human cells is a
12
13 key indicator of the sensitivity of a tumor to platinum-based therapy [26]); (iii) recognition of
14
15 DNA adducts of **1** and **2** by high-mobility-group (HMG)-domain proteins (since these
16
17 proteins are involved in the cisplatin mechanism of action [25]); (iv) sensitization of breast
18
19 cancer cells to **1** and **2** by steroid hormones.
20
21
22
23
24
25
26
27
28
29
30

31 **2. Materials and methods**

32 *2.1. Chemicals*

33
34
35
36
37
38
39
40
41 Cisplatin was obtained from Sigma (Prague, Czech Republic) (purity was $\geq 99.9\%$ based
42
43 on elemental and ICP trace analysis). Dinuclear Pt^{II} complexes **1** and **2** were prepared as
44
45 described previously [23, 27]. The stock solutions of platinum compounds were prepared in
46
47 10 mM NaClO₄ and stored at 20°C in the dark. The concentrations of platinum complexes in
48
49 the stock solutions were determined by flameless atomic absorption spectrometry (FAAS).
50
51 The concentrations of the platinum complexes indicated in the present work are related to the
52
53 whole compound (not to the Pt content in the case of the dinuclear complexes). Calf thymus
54
55 (CT) DNA (42% G + C, mean molecular mass ca. 20 000 kDa) was prepared and
56
57
58
59
60
61
62
63
64
65

1 characterized as described previously [28, 29]. Plasmids, pUC19 [2686 base pairs (bp)] and
2 pBR322 (4361 bp) were isolated according to standard procedures. Restriction endonucleases
3
4 were purchased from New England Biolabs (Beverly, MA). Deoxyribonucleoside
5
6 triphosphates were from Roche Diagnostics, GmbH (Mannheim, Germany). Agarose was
7
8 from FMC BioProducts (Rockland, ME). Electrophoresis-grade acrylamide, N,N'-
9
10 methylenebisacrylamide, ethidium bromide (EtBr), urea, and dithiothreitol were from Merck
11
12 KgaA (Darmstadt, Germany). Proteinase K and ATP were from Boehringer (Mannheim,
13
14 Germany). Spermine, EZBlue Gel Staining Reagent, β -estradiol and progesterone were from
15
16 Sigma (Prague, Czech Republic). Expression and purification of recombinant rat full-length
17
18 HMGB1 protein (HMG = high mobility group) and its domains A (residues 1-84 [30]) and B
19
20 (residues 85-180 [30]) (HMGB1a and HMGB1b, respectively) were carried out as described
21
22 [30-32]. Nonidet P-40 was from Fluka (Prague, Czech Republic). SDS was from Serva
23
24 (Heidelberg, Germany). MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]
25
26 was from Calbiochem (Darmstadt, Germany). Radioactive products were from Amersham
27
28 (Arlington Heights, IL, USA). A cell-free extract (CFE) was prepared from the repair
29
30 proficient HeLa S3 cell line as described [33, 34]. Streptavidin-coated magnetic beads
31
32 (Dynabeads) and Biotin-14-dATP were from Invitrogen Dynal AS (Oslo, Norway).
33
34
35
36
37
38
39
40
41
42

43 2.2. Cytotoxicity

44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Cisplatin, **1** and **2** were dissolved in DMSO. Stock solutions of the compounds were freshly prepared before use. The final concentration of DMSO in cell culture medium did not exceed 0.25%. The A2780 and A2780cisR human ovarian carcinoma cell lines (parent cisplatin sensitive and with acquired cisplatin resistance, respectively) were grown in RPMI 1640 medium (GIBCO, Carlsbad, CA) supplemented with gentamycin [50 μ g/mL, Serva

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

(Heidelberg, Germany)] and 10% heat inactivated fetal bovine serum (GIBCO, Carlsbad, CA). The acquired resistance of A2780cisR cells was maintained by supplementing the medium with 1 μM cisplatin every second passage. The cells were cultured in a humidified incubator at 37°C in a 5% CO_2 atmosphere and subcultured 2-3 times a week with an appropriate plating density.

Cell death was evaluated by using a system based on the tetrazolium compound MTT which is reduced by living cells to yield a formazan product that can be detected colorimetrically [35]. The cells were seeded in 96-well tissue cultured plates at a density of 10^4 cells/well in 100 μL of medium. After overnight incubation (16 hr), the cells were treated with the platinum compounds at the final concentrations in the range of 0 to 128 μM in a final volume of 200 μL /well. After additional 72 hr 10 μL of a freshly diluted MTT solution (2.5 mg/mL) was added to each well and the plate was incubated at 37°C in a humidified 5% CO_2 atmosphere for 4 hr. At the end of the incubation period the medium was removed and the formazan product was dissolved in 100 μL of DMSO. The cell viability was evaluated by measurement of the absorbance at 570 nm, using an Absorbance Reader SUNRICE TECAN SCHOELLER. IC_{50} values were calculated from curves constructed by plotting cell survival (%) versus drug concentration (μM). All experiments were made in triplicate. The reading values were converted to the percentage of control (% cell survival). Cytotoxic effects were expressed as IC_{50} (concentration inhibiting cell growth by 50%).

2.3. *Platination reactions in cell-free media*

If not stated otherwise, DNA was incubated with the platinum complex in 10 mM NaClO_4 at 37°C in the dark. After 24 hr, the samples were exhaustively dialyzed against the medium required for subsequent biochemical or biophysical analysis. An aliquot of these

1 samples was used to determine the value of r_b by FAAS. Alternatively, DNA binding of **1** and
2 **2** was studied using differential pulse polarography (DPP) [36].
3
4
5
6

7 *2.4. Recognition of DNA adducts by HMG-domain protein*

8
9

10
11 The EcoRI/PvuII restriction fragment (92 bp) from pUC19 plasmid was modified with
12 cisplatin or complexes **1** or **2** to desired r_b value in 0.01 M NaClO₄ at 37°C, in dark for 24 hr.
13
14 Following the incubation or the optional radioactive labeling (by γ -³²P ATP), the fragments
15 were 3'-end biotinylated using Biotin-14-dATP label and terminal deoxynucleotidyl
16 transferase in terminal transferase buffer (100 mM cacodylate buffer, pH 6.8; 5 mM CoCl₂,
17
18 0.5 mM dithiothreitol). Biotinylated DNA was attached to streptavidine coated magnetic
19 beads in 1 M NaCl, 5 mM Tris.HCl pH 7.4 and 0.5 mM EDTA. The beads were washed three
20 times in the same buffer. HMG protein was added to the beads carrying DNA and incubated
21 on ice in 10 mM HEPES, pH 7.5, 10 mM MgCl₂, 50 mM LiCl, 100 mM NaCl, 1 mM
22 spermidine and 0.05 % Nonidet P40, in the presence of poly(dI-dC) as a competitor for 1 hr.
23
24 The beads were then washed 2 times in the same buffer. Finally the protein was released with
25 0.04 % sodium dodecyl sulfate (SDS). Following the addition of loading buffer (0.1 M Tris-
26 HCl, pH 6.8; 2 % SDS; 0.1 M dithiothreitol, 10 % glycerol, 0.25 % bromphenol blue) the
27 samples were heated for five minutes at 90°C, electrophoresed on 15 % SDS-PAGE (PAGE =
28 polyacrylamide gel electrophoresis) and stained with EZBlue gel staining reagent.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 *2.5. DNA repair synthesis by human cell extract*

52
53
54
55

56 Repair DNA synthesis of CFEs was assayed using pUC19 plasmid. Each reaction of 50
57 μ L contained 600 ng of nonmodified pBR322 and 600 ng of nonmodified or platinated
58
59
60
61
62
63
64
65

1 pUC19, 2 mM ATP, 30 mM KCl, 0.05 mg/mL creatine phosphokinase (rabbit muscle), 20
2 mM each dGTP, dATP, and dTTP, 8 mM dCTP, 74 kBq of [α - 32 P]dCTP in the buffer
3
4 composed of 40 mM HEPES-KOH, pH 7.5, 5 mM MgCl₂, 0.5 mM dithiothreitol, 22 mM
5
6 creatine phosphate, 1.4 mg of bovine serum albumin/mL, and 20 μ g of CFE from the HeLa
7
8 S3 cells. Reactions were incubated for 3 hr at 30°C and terminated by adding EDTA to a final
9
10 concentration of 20 mM, SDS to 0.6%, and proteinase K to 250 μ g/mL and then incubating
11
12 for 20 min. The products were extracted with one volume of 1:1 phenol/chloroform. The
13
14 DNA was precipitated from the aqueous layer by the addition of 0.1 volume of 3 M sodium
15
16 acetate and 2.5 volumes of ethanol. After 30 min of incubation at -20°C and centrifugation at
17
18 12 000 g for 30 min at 4°C, the pellet was washed with 0.2 mL of 80% ethanol and dried in a
19
20 vacuum centrifuge. DNA was finally linearized by SspI before electrophoresis on a 1%
21
22 agarose gel. Resulting gel was stained by EtBr. The experiments were made in quadruplicate.
23
24
25
26
27
28
29
30

31 *2.6. Sensitization of cancer cells by steroid hormones to platinum complexes by clonogenic* 32 33 *assay* 34 35

36
37
38 MCF-7 cells were seeded on 6 cm Petri dishes at a density of 400 cells per dish. After 24
39
40 hr, a fresh stock solution containing steroid hormone(s), estradiol or estradiol plus
41
42 progesterone, was prepared in *N,N*-dimethylformamide and added to the dishes so that a final
43
44 concentration of each hormone was 0.2 μ M. Control dishes were treated with the same
45
46 volume of *N,N*-dimethylformamide without hormone. The hormone was added at the same
47
48 time as platinum complex (**1**, **2**, or cisplatin). After 4 hr treatment, the cells were washed with
49
50 PBS, and fresh medium was added. After 10 days, the cell colonies were stained with 1%
51
52 methylene blue and were then counted. Each point is an average of six independent
53
54
55
56
57
58
59
60
61
62
63
64
65 determinations.

2.7. Other physical methods

Absorption spectra were measured with a Beckmann DU-7400 spectrophotometer. FAAS measurements were carried out with a Varian AA240Z Zeeman atomic absorption spectrometer equipped with a GTA 120 graphite tube atomizer. For FAAS analysis DNA was precipitated with ethanol and dissolved in 0.1 M HCl. The gels were visualized by using the BAS 2500 FUJIFILM bioimaging analyzer, and the radioactivities associated with bands were quantitated with the AIDA image analyzer software (Raytest, Germany).

3. Results

3.1. Cytotoxicity

The cytotoxic activity of the **1** and **2** tested in the present work was determined against A2780 (parent cisplatin sensitive) and A2780cisR (with acquired cisplatin resistance) human ovarian carcinoma cell lines, commonly used to test cytotoxic activity of cisplatin analogues and other antitumor metallodrugs. The tumor cell lines were incubated for 72 hr with the platinum compounds and the cell survival in the culture treated with the platinum compounds was evaluated as described in the section Experimental Procedures. Results (Table 1) show that the cytotoxicity of **1** in cisplatin sensitive A2780 cells was comparable to the cytotoxicity of cisplatin, whereas **2** was slightly less cytotoxic. Importantly, both **1** and **2** were markedly more active in the cisplatin-resistant line A2780cisR and again **2** was less cytotoxic (Table 1).

3.2. Repair DNA synthesis by human cell extracts

To investigate “downstream” effects of DNA damage by adducts of **1** and **2**, we used *in vitro* systems to study repair of platinum lesions by CFE. DNA repair efficiency in pUC19 plasmid (2686 bp) globally modified by **1**, **2** or cisplatin (for comparative purposes) at $r_b = 0.03$ (r_b is defined as the number of molecules of the platinum complex bound per nucleotide residue) was tested using CFE of repair proficient HeLa cells. Repair activity was monitored by measurement of the amount of incorporated radiolabelled nucleotide. The incorporation of radioactive material was corrected for the relative DNA content in each band. As illustrated in Fig. 2, damage-induced DNA repair synthesis detected in the plasmid modified by **1** or **2** was approximately 38 or 36 %, respectively, of that found for the cisplatin at the same level of modification.

3.3. DNA modification in a cell-free medium

The cellular DNA repair response strongly depends on the character and extent of DNA damage. Therefore, it was of great interest to characterize DNA distortions induced by **1** and **2**. With the view of this we examined the DNA binding properties of these complexes and compared them with those of cisplatin. The experiments were aimed at quantification of the binding of **1** and **2** to mammalian DNA, determination of their preferential binding sites, and characterization of DNA lesions induced by **1** and **2** within natural DNA. The results of these experiments are described in detail in the Supplemental Information and are summarized in the following paragraph.

3.3.1. DNA binding

1
2 CT DNA was incubated with the platinum complexes at r_i values of 0.01, 0.03, 0.05, and
3
4 0.1) (r_i is defined as the molar ratio of free platinum complex to nucleotides at the onset of
5
6 incubation with DNA) in 10 mM NaClO₄ at 37°C for 24 or 48 hr. The amount of the platinum
7
8 compound bound to DNA increased with time. In this binding reaction the time at which the
9
10 binding reached 50% ($t_{50\%}$) was 14.9 ± 0.6 min and 39 ± 6 min for **1** and **2**, respectively, and
11
12 both complexes were quantitatively bound after 24 hr. This result indicates that the rate of
13
14 binding of both **1** and **2** to natural double-helical DNA is higher than that of cisplatin ($t_{50\%} =$
15
16 ca. 120 min [37]), although cisplatin is also quantitatively bound to DNA after 24 hr under
17
18 identical experimental conditions [37].
19
20
21
22
23
24
25

3.3.2. Transcription mapping of platinum-DNA adducts

26
27 *In vitro* RNA synthesis by T7 and SP6 RNA polymerases on DNA templates containing
28
29 adducts of **1** or **2** revealed (Fig. S1) that the major stop sites produced by these complexes
30
31 were at guanine residues, a few stop sites were also at adenine and cytosine residues. These
32
33 results also suggest that the adducts formed by **1** and **2** are able to inhibit RNA polymerase.
34
35
36
37
38
39
40

3.3.3. Interstrand cross-linking

41
42 We quantitated the interstrand cross-linking efficiency of **1** and **2** in linearized pUC19
43
44 plasmid (Fig. S2). The interstrand cross-linking efficiencies (defined as the number of
45
46 interstrand crosslinks per one adduct of the platinum complex [17]) of both **1** and **2** (48 ± 4 %
47
48 and 20 ± 2 %, respectively) were significantly higher than that of cisplatin (6 % [38]).
49
50
51
52
53
54
55

3.3.4. Characterization of DNA adducts by ethidium bromide fluorescence

1 The results of these experiments (Fig. S3) suggest that the conformational distortion
2 induced in DNA by the adducts of **1** is more delocalized and extends over more base pairs
3 around the platination sites than in the case of the adducts of **2** or mononuclear cisplatin.
4 Thus, these results are consistent with the formation of long-range intra- or interstrand CLs of
5 **1**, whereas **2** apparently forms in DNA much less long-range lesions. In aggregate,
6
7 characterization of DNA adducts of **1** and **2** by EtBr fluorescence supports the view that the
8 DNA binding modes of these dinuclear platinum compounds are distinctly different.
9
10

11 3.3.5. Unwinding of negatively supercoiled DNA

12 Electrophoresis in native agarose gel was used to quantify the unwinding induced in
13 pUC19 plasmid by adducts of **1** and **2** by monitoring the degree of supercoiling (Fig. S4). We
14 determined the DNA unwinding angle of 17° for the adduct of **1** and 12° for that of **2**. The
15 high level of unwinding induced by **1** and **2** is notable. Interestingly, the unwinding angle
16 produced by **1** is higher than that produced by **2**. One plausible explanation for this
17 observation might be consistent with a possibility that the additional contribution to
18 unwinding is associated with the interaction of the linker chain of **1** with the duplex upon
19 covalent binding of platinum units. Thus, the results of unwinding experiments represent
20 another support for the view that structure of DNA lesions formed by **1** is substantially
21 different from the structure of DNA adducts of its counterpart **2**.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 3.4. Recognition of DNA adducts by high-mobility-group (HMG)-domain proteins

50 The altered structures induced in DNA by cisplatin and its analogs attract HMG-domain
51 and other proteins [30, 39]. This binding of HMG-domain proteins to DNA modified by
52 cisplatin and its analogs has been postulated to mediate the antitumor properties of these
53
54
55
56
57
58
59
60
61
62
63
64
65

1 drugs.[39] As dinuclear platinum complexes **1** and **2** exhibit antitumor activity qualitatively
2 different from cisplatin, it was of interest to examine whether also the adducts formed within
3 DNA by **1** or **2** are able to be specifically recognized by HMG domain proteins.
4
5

6
7 To assess the ability of HMGB1 protein to recognize DNA modified by complexes **1** and
8
9 **2** we employed the system of DNA immobilized on streptavidine coated magnetic beads
10 suspended in HMG binding buffer containing HMGB1 protein. 10 μ g of DNA fragment was
11 immobilized on 1 mg of Dynabeads. Radioactively labeled DNA enabled us to ensure the
12 same quantity of DNA in various samples. First we determined whether HMGB1 protein
13 exhibits the ability to recognize DNA modified by the two new dinuclear compounds.
14
15 Immobilized DNA (nonmodified or modified by cisplatin, complex **1** and **2** to $r_b = 0.02$) was
16 incubated with HMGB1 protein on ice for 1 hr in the presence of poly(dI-dC) as a competitor.
17
18 Following the incubation, DNA was washed extensively and bound protein was released in
19
20 0.04 % SDS and electrophoresed on 15 % SDS-PAGE.
21
22
23
24
25
26
27
28
29
30

31 The HMGB1 protein was found to bind DNA modified by cisplatin, in agreement with
32 experiments published earlier [40, 41] that was evidenced by the presence of band migrating
33 near the 34 kDa marker corresponding to HMGB1 protein (Fig. 3A, lane cisPt, Table S1).
34
35 Interestingly, the band corresponding to HMGB1 was also clearly seen for the probe modified
36 by **1** (Fig. 3A, lane **1**); the intensity of this band was almost comparable (80%) to that found
37 for cisplatin (Table S1). On the other hand, much less effective binding of the HMGB1 to the
38 DNA containing the adducts of **2** was detected, the amount of HMGB1 protein bound to this
39 probe was only 15% of the amount of HMGB1 bound to DNA modified by cisplatin under the
40 same conditions (Fig. 3A, lane 2, Table S1). No binding of the proteins occurred under
41 identical experimental conditions if the same DNA probe was unplatinated (Fig. 3A, lane
42 NoPt). These results clearly showed that DNA modified by **1** is efficiently recognized by
43 HMGB1 with relatively high affinity, though slightly less than to the probe containing the
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

adducts of cisplatin. In contrast, DNA adducts of **2** are recognized by this protein markedly less.

To determine which of the two HMGB1 domains A or B predominates in recognition and binding of DNA lesions of **1**, the binding of HMGB1 domain A and HMGB1 domain B to the DNA modified by complexes **1** or **2** was investigated (Fig. 3 B,C). The experiments revealed that in agreement with previously published data [41], HMGB1a effectively bound to the probe containing adducts of cisplatin (Fig. 3B, lane cisPt). Considerably weaker binding of HMGB1a was noticed to the probes containing adducts of **1** or **2**; relative amount of HMGB1a protein bound to DNA probe containing adducts of cisplatin, **1** or **2** was 100%, 20%, or 14%, respectively (Fig. 3B, Table S1). On the other hand, the amount of HMGB1b bound to the DNA modified by **1** was two-fold higher compared to that bound to DNA modified by cisplatin (Fig. 3C, Table S1). These results clearly show that DNA damaged by complex **1** was recognized by HMGB1 protein, but in contrast to DNA globally modified by cisplatin, the full-length HMGB1 protein binds to DNA globally modified by **1** mainly through domain B.

Complex **1** modifies DNA so that it forms approximately equal amounts of interstrand and intrastrand adducts (see paragraph Interstrand cross-linking in this Section and Fig. S2). To distinguish which type of lesions formed by **1** within DNA is preferentially recognized by HMGB1 protein, the following experiment was performed. The 92 bp fragment of DNA was modified by **1** so that the fragment contained only one adduct per DNA fragment on average. After the modification, DNA was loaded into the denaturing agarose gel and two DNA fractions were purified from the gel, one contained only intrastrand adducts (no interstrand CLs) (F-IAA) and the other contained at least one interstrand CL (F-IEC). These two separated DNA fractions were then allowed to interact with HMGB1b, as described in section Experimental procedures. The amount of HMGB1b protein bound to each DNA fraction was

1 determined by SDS/PAGE electrophoresis. The inspection of the electrophoreogram (Fig. 4)
2 revealed that HMGB1b bound to DNA containing interstrand CLs of **1** whereas it bound
3 considerably less (4-fold) to DNA modified by **1** to the similar level but containing other
4 adducts, but not interstrand CLs (Table S2).
5
6
7
8
9

10 11 **3.5. Steroid hormones sensitize cancer cells to complex 1, but not to complex 2**

12
13
14
15

16 Previous studies have shown that HMGB1 protein sensitizes cells to cisplatin by
17 shielding its DNA adducts from nucleotide excision repair [25, 42]. Human breast cancer cells
18 MCF-7 contain estrogen and progesterone receptors and respond to these steroids up-
19 regulating HMGB1 protein [43]. Importantly, treatment of these cancer cells with estrogen or
20 progesterone significantly increases the potency of cisplatin apparently owing to the
21 overexpression of HMGB1 [44]. Since DNA adducts of **1** are recognized by HMGB1 protein
22 markedly more than those of **2** (Fig. 3), it was of interest to assess the involvement of HMG-
23 domain proteins in mediating cytotoxicity of the dinuclear Pt^{II} complexes tested in the present
24 work. Therefore, we investigated the effects of elevated HMGB1 levels due to steroid
25 hormone treatment in MCF-7 cells on the sensitivity of these cells to **1** or **2** (Fig. 5) using
26 clonogenic assay in the same way as described previously for cisplatin and carboplatin [44].
27
28 As anticipated, we found that elevated HMGB1 expression levels were paralleled by
29 increased sensitivity toward **1**, but markedly less toward **2** (Fig. 5 and Table S4). For instance,
30 in MCF-7 cells, estradiol or estradiol plus progesterone treatment increased sensitivity toward
31 **1** about 2.7- or 4.2-fold, respectively, when only 50% of the cells are viable (Fig. 5A);
32 estradiol or estradiol plus progesterone treatment also increased sensitivity toward cisplatin,
33 but only about 1.8- or 2.5-fold, respectively (Fig. 5C). Thus, sensitivity toward **1** was
34 enhanced due to steroid hormone treatment even considerably more than toward cisplatin.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Sensitivity of MCF-7 cells, treated with hormones, toward **1**, **2**, and cisplatin was also
2 determined by MTT assay (Table S3). The trend was similar to that observed when
3
4 clonogenic assay was used, although IC₅₀ data obtained by MTT assay were uniformly higher
5
6 than in clonogenic assay.
7
8
9

10 11 12 13 14 **4. Discussion** 15

16
17
18
19 We demonstrate in the present work that the dinuclear complexes **1** and **2** are quite toxic
20
21 to the ovarian tumor cells, with IC₅₀ values that were comparable to that observed for cisplatin
22
23 in the cisplatin sensitive cell line A2780, and noticeably much lower in the cisplatin-resistant
24
25 line A2780cisR (Table 1). Thus, these results complement those obtained in the previous
26
27 study [27] showing that complexes **1** and **2** exhibit the promising cytotoxicity to the human
28
29 nonsmall-cell lung cancer cell line (A549). Interestingly, the cytotoxicity of both **1** and **2** to
30
31 the ovarian tumor cells is characterized by remarkably low resistance factors, equal to or even
32
33 less than **1**, which indicates that the dinuclear platinum complexes examined in the present
34
35 work are capable of circumventing of cisplatin resistance in some types of the cisplatin-
36
37 resistant lines. In aggregate, no cross-resistance to cisplatin and high cytotoxic effects in
38
39 cisplatin resistant cells represent important pharmacological features of **1** and **2**.
40
41
42
43
44
45

46 An important feature of the mechanism underlying antitumor effects of DNA-binding
47
48 metal-based compounds is repair of their DNA adducts [40, 45]. A persistence of these DNA
49
50 adducts may potentiate their antitumor effects in the cells sensitive to these compounds [8, 39,
51
52 40]. DNA repair synthesis was investigated in the present work using the CFE from human
53
54 tumor cells and DNA substrates randomly modified by **1** or **2**. Importantly, the adducts of
55
56 both **1** and **2** induced a considerably lower level of repair synthesis than the adducts of
57
58
59
60
61
62
63
64
65

1
2 cisplatin suggesting a less efficient removal from DNA and enhanced persistence of the
3 adducts of the dinuclear complexes **1** and **2**. Thus, enhanced resistivity of DNA adducts of **1**
4 and **2** against repair processes appears to be an important factor contributing to ability of **1**
5 and **2** to overcome resistance towards cisplatin.
6
7

8
9 Structural features of DNA distortions are considered to be major factors playing a key
10 role in recognition by damaged-DNA recognition-proteins and DNA repair systems. The
11 analysis of DNA binding of **1** and **2** described in this report and summarized in Table 2
12 provides experimental support for the view that the binding of these dinuclear platinum
13 complexes modifies DNA in a way which is in several aspects considerably different from
14 modification of DNA by "conventional" mononuclear cisplatin. These differences may
15 suggest that in comparison with adducts of cisplatin the reduced repair of DNA adducts of the
16 dinuclear Pt^{II} complexes is a consequence of their different DNA binding mode.
17
18
19
20
21
22
23
24
25
26
27

28
29 The binding of **1** and **2** to DNA is considerably faster than the binding of dichloro
30 derivatives of mononuclear Pt^{II} complexes, such as cisplatin or transplatin ($t_{50\%} \sim 120$ min
31 [37]) likely reflecting the higher charge. In addition, the different rate of the binding of **1** and
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Consistent with this view is the observation that affinity of HMG-domain proteins to DNA adducts of **1** is markedly higher than that to DNA adducts of **2** (Fig. 3A). The full-length HMGB1 protein consists of two tandem HMG box domains (A and B) and a C-terminal acidic tail. Both A and B domains of HMGB1 protein share a common HMG box structure. Domain A has a higher binding affinity to distorted DNA than domain B, whereas domain B bends DNA more effectively [48]. Because of the importance of the HMG box interaction with DNA [25], the studies of the individual HMG domains and their DNA complexes are of great interest. It has been shown that full-length HMGB1 protein binds to DNA containing a site-specific 1,2-GG intrastrand CL of cisplatin mainly through domain A, whereas domain B alone binds cisplatinated DNA much more weakly than domain A alone [41]. In contrast to these data, binding of full-length HMGB1 to DNA modified by the dinuclear complex **1** is mediated preferentially by domain B (Fig. 3B, C). This result, in combination with our previous findings that the domain B of the HMGB1 protein with the A/B linker plays a crucial role in the interaction between full-length HMGB1 protein and DNA interstrand cross-linked by cisplatin [30], implies a possible role of the interstrand DNA adducts of **1** in the interaction with HMG-domain proteins. Therefore, we tested the contribution of DNA intrastrand and interstrand adducts formed by **1** to the binding of HMGB1 to DNA globally modified by **1**. The sample containing DNA fragment globally modified by **1** was divided into two DNA fractions, one containing only intrastrand adducts (no interstrand CLs) and the other containing interstrand CLs and the affinity of HMGB1b towards these two fractions was tested. The results unambiguously show the prevalent role of DNA interstrand CLs of **1** in recognition and binding of DNA modified by **1** to full-length HMGB1 protein. To our knowledge, this is the first evidence for strong specific recognition and binding of HMG-domain proteins to DNA modified by a polynuclear platinum complex. Previously, a weak recognition of DNA interstrand CLs of [$\{trans\text{-PtCl}(\text{NH}_3)_2\}\text{H}_2\text{N}(\text{CH}_2)_2$]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

${}^6\text{NH}_2\text{]Cl}_2$ complexes by full-length HMGB1 protein has only been reported [7], whereas a recognition of intrastrand CLs of these complexes has not been detected.

The results of the present work provide evidence that HMGB1 domain B controls a relatively strong binding of full-length HMGB1 to DNA modified by **1** when both domain A and B are present. It is quite conceivable that domain B is the dominating domain in HMGB1 protein that binds to the interstrand CL of **1**, while domain A only facilitates binding by providing additional protein-DNA interactions. A structure of full-length HMGB1 complexed with DNA containing interstrand CL of **1** will provide more insights into how HMG-domain proteins can mediate cytotoxicity of **1**. On the other hand, the affinities of full-length HMGB1 protein and its domain A to DNA modified by **2** are relatively weak and almost identical. This result is consistent with the view and supports the hypothesis that the binding of full-length HMGB1 protein to DNA modified by the dinuclear complex **2**, although relatively weak, is mediated preferentially by the domain A. Collectively, the results of this work demonstrate that not only DNA binding modes of **1** and **2** and resulting conformational alterations are noticeably different, but also “downstream” intracellular effects of DNA adducts of the two dinuclear complexes may be different, particularly in the cells in which the HMGB1 level is high and in which HMGB1 protein is not prevented from interacting with platinum-modified DNA.

It has been shown that specific interaction between adducts of "conventional" cisplatin and cellular HMG-domain proteins modulate the sensitivity of tumor cells to this drug. In particular, protein-DNA interactions induced by cisplatin inhibit replication,[49] shield the adducts from nucleotide excision repair [50] and interfere with transcription by recruiting transcription factors from their native binding sites [51, 52]. Therefore, it is reasonable to assume that such interactions may also differently modulate the cellular responses both to the

1 platinum dinuclear complexes **1** and **2** and also to the class of dinuclear complexes tested in
2 the present work and cisplatin.
3

4 In conclusion, the results of the present work are consistent with the hypothesis that the
5 unique properties of interstrand CLs of the class of bifunctional polynuclear platinum
6 complexes tested in the present work and resulting conformational alterations in DNA have
7 critical consequences for their antitumor effects. An interesting finding that might be also
8 translated to the clinical trials is that treatment leading to enhancement of level of HMG-
9 domain proteins in tumor cells should allow the regimens involving **1** to show increased
10 antitumor effects toward these cells.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

26 **Acknowledgements**

27 This research was supported by the Ministry of Education of the CR (MSMT LC06030,
28 6198959216, ME08017, ME10066, OC08003, and OC09018), the Academy of Sciences of
29 the Czech Republic (Grants KAN200200651, M200040901, AV0Z50040507, and
30 AV0Z50040702), the Grant Agency of the Academy of Sciences of the CR (IAA400040803),
31 and the Grant Agency of the CR (301/09/H004 and P301/10/0598). Financial support from
32 the National Natural Science Foundation of China (No. 90713001) is also acknowledged. J.K.
33 is an international research scholar of the Howard Hughes Medical Institute. The authors also
34 acknowledge that their participation in the EU COST Action D39 has enabled them to
35 exchange regularly their most recent ideas in the field of anticancer metallodrugs with several
36 European colleagues.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 **Appendix A. Supplementary data**

53 Supplementary data associated with this article can be found, in the online version, at doi: ...
54
55
56
57
58
59
60
61
62
63
64
65

References

- [1] Pratesi G, Perego P, Polizzi D, Righetti SC, Supino R, Caserini C, et al. A novel charged trinuclear platinum complex effective against cisplatin-resistant tumours: hypersensitivity of p53-mutant human tumour xenografts. *Br J Cancer* 1999;80:1912-9.
- [2] Farrell N. Polynuclear platinum drugs. In: Sigel A, Sigel H, editors. *Metal ions in biological systems*. New York, Basel: Marcel Dekker, Inc.; 2004, p. 251-96.
- [3] Brabec V, Kasparkova J. Modifications of DNA by platinum complexes: Relation to resistance of tumors to platinum antitumor drugs. *Drug Resist Updat* 2005;8:131-46.
- [4] Brabec V, Kasparkova J. DNA interactions of platinum anticancer drugs. Recent advances and mechanisms of action. In: Perez-Martin J-M, Fuertes MA, Alonso C, editors. *Metal compounds in cancer chemotherapy*. Trivandrum, Kerala (India): Research Signpost; 2005, p. 187-218.
- [5] Brabec V, Christofis P, Slamova M, Kostrhunova H, Novakova O, Najajreh Y, et al. DNA interactions of new cytotoxic tetrafunctional dinuclear platinum complex trans,trans-[PtCl₂(NH₃)₂(piperazine)]. *Biochem Pharmacol* 2007;73:1887-900.
- [6] Cossa G, Gatti L, Zunino F, Perego P. Strategies to improve the efficacy of platinum compounds. *Curr Med Chem* 2009;16:2355-65.
- [7] Kasparkova J, Farrell N, Brabec V. Sequence specificity, conformation, and recognition by HMG1 protein of major DNA interstrand cross-links of antitumor dinuclear platinum complexes. *J Biol Chem* 2000;275:15789-98.
- [8] Kasparkova J, Zehnulova J, Farrell N, Brabec V. DNA interstrand cross-links of the novel antitumor trinuclear platinum complex BBR3464. Conformation, recognition by

- 1 high mobility group domain proteins, and nucleotide excision repair. *J Biol Chem*
2 2002;277:48076-86.
3
- 4 [9] Brabec V, Kasparkova J, Vrana O, Novakova O, Cox JW, Qu Y, et al. DNA
5 modifications by a novel bifunctional trinuclear platinum Phase I anticancer agent.
6 *Biochemistry* 1999;38:6781-90.
7
- 8 [10] Kasparkova J, Novakova O, Vrana O, Farrell N, Brabec V. Effect of geometric
9 isomerism in dinuclear platinum antitumor complexes on DNA interstrand cross-
10 linking. *Biochemistry* 1999;38:10997-1005.
11
- 12 [11] Zehnulova J, Kasparkova J, Farrell N, Brabec V. Conformation, recognition by high
13 mobility group domain proteins, and nucleotide excision repair of DNA intrastrand
14 cross-links of novel antitumor trinuclear platinum complex BBR3464. *J Biol Chem*
15 2001;276:22191-9.
16
- 17 [12] Roberts JD, Peroutka J, Farrell N. Cellular pharmacology of polynuclear platinum anti-
18 cancer agents. *J Inorg Biochem* 1999;77:51-7.
19
- 20 [13] Roberts JD, Beggiolin G, Manzotti C, Piazzoni L, Farrell N. Comparison of cytotoxicity
21 and cellular accumulation of polynuclear platinum complexes in L1210 murine
22 leukemia cell lines. *J Inorg Biochem* 1999;77:47-50.
23
- 24 [14] Servidei T, Ferlini C, Riccardi A, Meco D, Scambia G, Segni G, et al. The novel
25 trinuclear platinum complex BBR3464 induces a cellular response different from
26 cisplatin. *Eur J Cancer* 2001;37:930-8.
27
- 28 [15] Farrell N. Polynuclear charged platinum compounds as a new class of anticancer agents:
29 Toward a new paradigm. In: Kelland LR, Farrell NP, editors. *Platinum-based drugs in*
30 *cancer therapy*. Totowa/NJ: Humana Press Inc.; 2000, p. 321-38.
31
- 32 [16] Harris AL, Yang X, Hegmans A, Povirk L, Ryan JJ, Kelland L, et al. Synthesis,
33 characterization, and cytotoxicity of a novel highly charged trinuclear platinum
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- compound. Enhancement of cellular uptake with charge. *Inorg Chem* 2005;44:9598-600.
- [17] Farrell N, Qu Y, Feng L, Van Houten B. Comparison of chemical reactivity, cytotoxicity, interstrand cross-linking and DNA sequence specificity of bis(platinum) complexes containing monodentate or bidentate coordination spheres with their monomeric analogues. *Biochemistry* 1990;29:9522-31.
- [18] Summa N, Maigut J, Puchta R, van Eldik R. Possible biotransformation reactions of polynuclear Pt(II) complexes. *Inorg Chem* 2007;46:2094-104.
- [19] Oehlsen ME, Qu Y, Farrell N. Reaction of polynuclear platinum antitumor compounds with reduced glutathione studied by multinuclear (^1H , ^1H - ^{15}N gradient heteronuclear single-quantum coherence, and ^{195}Pt) NMR spectroscopy. *Inorg Chem* 2003;42:5498 - 506.
- [20] Billecke C, Finniss S, Tahash L, Miller C, Mikkelsen T, Farrell NP, et al. Polynuclear platinum anticancer drugs are more potent than cisplatin and induce cell cycle arrest in glioma. *Neuro Oncol* 2006;8:215-26.
- [21] Mitchell C, Kabolizadeh P, Ryan J, Roberts JD, Yacoub A, Curiel DT, et al. Low-dose BBR3610 toxicity in colon cancer cells is p53-independent and enhanced by inhibition of epidermal growth factor receptor (ERBB1)-Phosphatidyl inositol 3 kinase signaling. *Mol Pharmacol* 2007;72:704-14.
- [22] Williams JW, Qu Y, Bulluss GH, Alvorado E, Farrell NP. Dinuclear platinum complexes with biological relevance based on the 1,2-diaminocyclohexane carrier ligand. *Inorg Chem* 2007;46:5820-2.
- [23] Fan D, Yang X, Wang X, Zhang S, Mao J, Ding J, et al. A dinuclear monofunctional platinum(II) complex with an aromatic linker shows low reactivity towards glutathione

- 1 but high DNA binding ability and antitumor activity. *J Biol Inorg Chem* 2007;12:655-
2 65.
3
4 [24] Jamieson ER, Lippard SJ. Structure, recognition, and processing of cisplatin-DNA
5 adducts. *Chem Rev* 1999;99:2467-98.
6
7 [25] Jung Y, Lippard SJ. Direct cellular responses to platinum-induced DNA damage. *Chem*
8 *Rev* 2007;107:1387-407.
9
10 [26] Brabec V, Kasparkova J. Role of DNA repair in antitumor effects of platinum drugs. In:
11 Hadjiliadis N, Sletten E, editors. *Metal complex - DNA interactions*. Chichester, UK:
12 Wiley; 2009, p. 175-208.
13
14 [27] Zhu J, Lin M, Fan D, Wu Z, Zhang J, Luc Y, et al. The role of bridging ligands in
15 determining DNA-binding ability and cross-linking patterns of dinuclear platinum(II)
16 antitumor complexes. *Dalton Trans* 2009:10889 -95.
17
18 [28] Brabec V, Palecek E. The influence of salts and pH on polarographic currents produced
19 by denatured DNA. *Biophysik* 1970;6:290-300.
20
21 [29] Brabec V, Palecek E. Interaction of nucleic acids with electrically charged surfaces. II.
22 Conformational changes in double-helical polynucleotides. *Biophys Chem* 1976;4:76-
23 92.
24
25 [30] Kasparkova J, Delalande O, Stros M, Elizondo-Riojas MA, Vojtiskova M, Kozelka J, et
26 al. Recognition of DNA interstrand cross-link of antitumor cisplatin by HMGB1
27 protein. *Biochemistry* 2003;42:1234-44.
28
29 [31] Stros M. DNA bending by the chromosomal protein HMG1 and its high mobility group
30 box domains. Effect of flanking sequences. *J Biol Chem* 1998;273:10355-61.
31
32 [32] Stros M. Two mutations of basic residues within the N-terminus of HMG-1 B domain
33 with different effects on DNA supercoiling and binding to bent DNA. *Biochemistry*
34 2001;40:4769-79.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [33] Reardon JT, Vaisman A, Chaney SG, Sancar A. Efficient nucleotide excision repair of cisplatin, oxaliplatin, and bis-aceto-ammine-dichloro-cyclohexylamine-platinum(IV) (JM216) platinum intrastrand DNA diadducts. *Cancer Res* 1999;59:3968-71.
- [34] Manley JL, Fire A, Cano A, Sharp PA, Gefter ML. DNA-dependent transcription of adenovirus genes in a soluble whole-cell extract. *Proc Natl Acad Sci USA* 1980;77:3855-9.
- [35] Alley MC, Scudiero DA, Monks A, Hursey ML, Czerwinski MJ, Fine DL, et al. Feasibility of drug screening with panels of human tumor cell lines using a microculture tetrazolium assay. *Cancer Res* 1988;48:589-601.
- [36] Kim SD, Vrana O, Kleinwächter V, Niki K, Brabec V. Polarographic determination of subnanogram quantities of free platinum in reaction mixture with DNA. *Anal Lett* 1990;23:1505-18.
- [37] Bancroft DP, Lepre CA, Lippard SJ. ^{195}Pt NMR kinetic and mechanistic studies of cis-diamminedichloroplatinum and trans-diamminedichloroplatinum(II) binding to DNA. *J Am Chem Soc* 1990;112:6860-71.
- [38] Brabec V, Leng M. DNA interstrand cross-links of trans-diamminedichloroplatinum(II) are preferentially formed between guanine and complementary cytosine residues. *Proc Natl Acad Sci USA* 1993;90:5345-9.
- [39] Brabec V. DNA modifications by antitumor platinum and ruthenium compounds: their recognition and repair. *Prog Nucleic Acid Res Mol Biol* 2002;71:1-68.
- [40] Cohen SM, Lippard SJ. Cisplatin: From DNA damage to cancer chemotherapy. *Prog Nucleic Acid Res Mol Biol* 2001;67:93-130.
- [41] Jung YW, Lippard SJ. Nature of full-length HMGB1 binding to cisplatin-modified DNA. *Biochemistry* 2003;42:2664-71.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [42] Huang JC, Zamble DB, Reardon JT, Lippard SJ, Sancar A. HMG-domain proteins specifically inhibit the repair of the major DNA adduct of the anticancer drug cisplatin by human excision nuclease. *Proc Natl Acad Sci USA* 1994;91:10394-8.
- [43] Chau KY, Lam HYP, Lee KLD. Estrogen treatment induces elevated expression of HMG1 in MCF-7 cells. *Exp Cell Res* 1998;241:269-72.
- [44] He Q, Liang CH, Lippard SJ. Steroid hormones induce HMG1 overexpression and sensitize breast cancer cells to cisplatin and carboplatin. *Proc Natl Acad Sci USA* 2000;97:5768-72.
- [45] Brabec V, Kasparkova J. Molecular aspects of resistance to antitumor platinum drugs. *Drug Resist Updat* 2002;5:147-61.
- [46] Hegmans A, Berners-Price SJ, Davies MS, Thomas D, Humphreys A, Farrell N. Long range 1,4 and 1,6-interstrand cross-links formed by a trinuclear platinum complex. Minor groove pre-association affects kinetics and mechanism of cross-link formation as well as adduct structure. *J Am Chem Soc* 2004;126:2166-80.
- [47] Berners-Price SJ, Davies MS, Cox JW, Thomas DS, Farrell N. Competitive reactions of interstrand and intrastrand DNA-Pt adducts: A dinuclear-platinum complex preferentially forms a 1,4-interstrand cross-link rather than a 1,2 intrastrand cross-link on binding to a GG 14-Mer duplex. *Chem Eur J* 2003;9:713-25.
- [48] Thomas JO. HMG 1 and 2: architectural DNA-binding proteins. *Biochem Soc Trans* 2001;29:395-401.
- [49] Comess KM, Burstyn JN, Essigmann JM, Lippard SJ. Replication inhibition and translesion synthesis on templates containing site-specifically placed cis-diamminedichloroplatinum(II) DNA adducts. *Biochemistry* 1992;31:3975-90.
- [50] Zamble DB, Mu D, Reardon JT, Sancar A, Lippard SJ. Repair of cisplatin-DNA adducts by the mammalian excision nuclease. *Biochemistry* 1996;35:10004-13.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [51] Zhai XQ, Beckmann H, Jantzen HM, Essigmann JM. Cisplatin-DNA adducts inhibit ribosomal RNA synthesis by hijacking the transcription factor human upstream binding factor. *Biochemistry* 1998;37:16307-15.
- [52] Vichi P, Coin F, Renaud JP, Vermeulen W, Hoeijmakers JHJ, Moras D, et al. Cisplatin- and UV-damaged DNA lure the basal transcription factor TFIID/TBP. *EMBO J* 1997;16:7444-56.
- [53] Fichtinger-Schepman AMJ, Van der Veer JL, Den Hartog JHJ, Lohman PHM, Reedijk J. Adducts of the antitumor drug cis-diamminedichloroplatinum(II) with DNA: Formation, identification, and quantitation. *Biochemistry* 1985;24:707-13.
- [54] Keck MV, Lippard SJ. Unwinding of supercoiled DNA by platinum ethidium and related complexes. *J Am Chem Soc* 1992;114:3386-90.

Table 1. *In vitro* growth inhibition of human ovarian cisplatin sensitive (A2780) and resistant (A2780cisR) cells, IC₅₀ (μM) ^{a,b}

| | A2780 | A2780cisR ^c |
|-----------|-----------|------------------------|
| cisplatin | 2.8 ± 0.7 | 18.6 ± 0.4 (6.6) |
| 1 | 2.7 ± 0.7 | 2.7 ± 0.1 (1) |
| 2 | 4.8 ± 2.7 | 3.6 ± 1.1 (0.75) |

^a Drug-treatment period was 72 hr.

^b Values shown in the table are the means (±SEM) of three separate experiments, each conducted with 6 replicates.

^c Resistance factor, defined as IC₅₀ (resistant) / IC₅₀ (sensitive), is given in parentheses.

Table 2. Summary of the effects of complex **1**, **2** and cisplatin on DNA ^a

| | 1 | 2 | cisplatin |
|---|----------|----------|---------------------|
| DNA binding ($t_{50\%}$) (min) ^b | 14.9±0.6 | 39±6 | 120 ^c |
| Preferential DNA binding sites ^d | G | G | GG, AG ^e |
| Reduction of EtBr fluorescence | strong | medium | medium |
| Plasmid DNA unwinding angle/adduct | 17° | 12° | 13° ^f |
| % interstrand cross-links/adduct | 48±4 | 20±2 | 6 ^g |

^a See also Figs. S1 - S4.

^b See the text for details.

^c Ref. [37].

^d Determined by transcription mapping.

^e Ref. [53].

^f Ref. [54].

^g Ref. [38].

Figure Captions

1
2
3
4
5 Fig. 1 - Structures of the platinum complexes.
6
7
8

9
10 Fig. 2 - *In vitro* DNA repair synthesis assay. Repair synthesis of the extract prepared from the
11 repair-proficient HeLa cell line used as substrates nonmodified pBR322 plasmid and pUC19
12 plasmid nonmodified or modified at $r_b = 0.03$ by cisplatin, complex **1**, and complex **2**. (A)
13
14 Results of a typical experiment. Top panel, a photograph of the EtBr stained gel; bottom
15
16 panel, autoradiogram of the gel showing incorporation of [α - 32 P]dCTP. Lanes: 1, nonmodified
17
18 pBR322 plus pUC19 plasmids; 2, nonmodified pBR322 plus pUC19 modified by cisplatin; 3,
19
20 nonmodified pBR322 plus pUC19 modified by **1**; 4, nonmodified pBR322 plus pUC19
21
22 modified by **2**. (B) Incorporation of dCTP into nonmodified or platinated pUC19 plasmid. For
23
24 all quantifications representing mean values of three separate experiments, incorporation of
25
26 radioactive material is corrected for the relative DNA content in each band. The radioactivity
27
28 associated with the incorporation of [α - 32 P]dCTP into DNA modified by cisplatin was taken
29
30 as 100%. Values shown in the graph are the means (\pm SEM) of three separate experiments,
31
32 each conducted with 4 replicates.
33
34
35
36
37
38
39
40
41
42

43 Fig. 3 - Recognition by HMG-domain proteins of platinated DNA. Affinity of full-length
44
45 HMGB1 (A) HMGB1 domain A (B), and HMGB1 domain B (C) to nonmodified (control)
46
47 DNA (lanes noPt) or DNA modified by cisplatin (lanes cisPt), complex **1** (lanes 1) or
48
49 complex **2** (lanes 2). Lanes M, a protein size marker; lane HMGB1, HMGB1 protein was
50
51 loaded; lane HMGB1 dom A, HMGB1 domain A was loaded.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig. 4 - Recognition by HMGB1 domain B of DNA fragments intrastrand or interstrand cross-linked by complex **1**. After reaction with **1**, the DNA was separated by gel electrophoresis so that the fraction F-IEC contained all interstrand cross-linked fragments and the fraction F-IAA contained exclusively platinum adducts formed within one strand of DNA (contained no interstrand CLs, for other details, see the text). Lanes: M, a protein size marker; noPt, control, nonplatinated; F-IAA, intrastrand adducts containing DNA incubated with HMGB1 domain B; F-IEC, interstrand cross-links containing DNA incubated with HMGB1 domain B.

Fig. 5 - The effects of estradiol or estradiol plus progesterone cotreatment on sensitivity of MCF-7 cells towards complexes **1** (A), **2** (B) or cisplatin (C) determined by clonogenic assay. MCF-7 cells were cotreated with estradiol (0.2 μ M) (\blacktriangle) or estradiol (0.2 μ M) plus progesterone (0.2 μ M) (\blacktriangledown) with platinum complex for 4 hr; (\square), no steroid hormone added. Values shown in the graphs are the means (\pm SEM) of six separate experiments, each conducted with 6 replicates; where error bars are not shown, they were smaller than the symbols.

Mechanistic insights into antitumor effects of new dinuclear cis Pt^{II} complexes containing aromatic linkers

Lenka Zerzankova ^a, Hana Kostrhunova ^a, Marie Vojtiskova ^a, Olga Novakova ^a,
Tereza Suchankova ^b, Miaoxin Lin ^c, Zijian Guo ^c, Jana Kasparikova ^{a,*}
Viktor Brabec^{a,*}

^a *Institute of Biophysics, Academy of Sciences of the Czech Republic, v.v.i., CZ-61265 Brno,
Czech Republic*

^b *Department of Experimental Physics, Faculty of Sciences, Palacky University, tr. Svobody
26, 77146 Olomouc, Czech Republic*

^c *State Key Laboratory of Coordination Chemistry, School of Chemistry and Chemical
Engineering, Nanjing University, Nanjing 210093, PR China*

Classification: (1) Antibiotics and Chemotherapeutics

* Corresponding authors at: Institute of Biophysics, Academy of Sciences of the Czech Republic, v.v.i., Kralovopolska 135, CZ-61265 Brno, Czech Republic. Tel.: +420-541517148; fax: +420-541240499.

E-mail addresses: jana@ibp.cz (J. Kasparikova); brabec@ibp.cz (V. Brabec)

Abbreviations: BBR3464, [$\{trans\text{-PtCl}(\text{NH}_3)_2\}_2\text{-}\mu\text{-}trans\text{-Pt}(\text{NH}_3)_2(\text{NH}_2(\text{CH}_2)_6\text{NH}_2)_2$] $^{4+}$;
BBR3610, [$\{trans\text{-PtCl}(\text{NH}_3)_2\}_2\text{-}\mu\text{-}\{trans\text{-}(\text{H}_2\text{N}(\text{CH}_2)_6\text{NH}_2(\text{CH}_2)_2\text{NH}_2(\text{CH}_2)_6\text{NH}_2)\}_2$] $^{4+}$; bp,
base pair; cisplatin, *cis*-diamminedichloroplatinum(II); CFE, cell-free extract; CL, crosslink;
complex **1**, [$\{cis\text{-Pt}(\text{NH}_3)_2\text{Cl}_2(4,4'\text{-methylenedianiline})\}_2$] $^{2+}$; complex **2**, [$\{cis\text{-}$
 $\text{Pt}(\text{NH}_3)_2\text{Cl}_2(\alpha,\alpha'\text{-diamino-p-xylene})\}_2$] $^{2+}$; CT, calf-thymus; DPP, differential pulse
polarography; EtBr, ethidium bromide; FAAS, flameless atomic absorption
spectrophotometry; HMG, high-mobility-group; IC₅₀, concentration inhibiting cell growth by
50%; MTT, [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]; PAGE,
polyacrylamide gel electrophoresis; r_b , the number of molecules of the metal complex bound
per nucleotide residue; r_i , the molar ratio of free metal complex to nucleotide-phosphates at
the onset of incubation with DNA; SDS, sodium dodecyl sulfate.

ABSTRACT

The primary objective was to understand more deeply the molecular mechanism underlying different antitumor effects of dinuclear Pt^{II} complexes containing aromatic linkers of different length, {[*cis*-Pt(NH₃)₂Cl]₂(4,4'-methylenedianiline)}²⁺ (**1**) and {[*cis*-Pt(NH₃)₂Cl]₂(α,α' -diamino-*p*-xylene)}²⁺ (**2**). These complexes belong to a new generation of promising polynuclear platinum drugs resistant to decomposition by sulfur nucleophiles which hampers clinical use of bifunctional polynuclear trans Pt^{II} complexes hitherto tested. Results obtained with the aid of methods of molecular biophysics and pharmacology reveal differences and new details of DNA modifications by **1** and **2** and recognition of these modifications by cellular components. The results indicate that the unique properties of DNA interstrand cross-links of this class of polynuclear platinum complexes and recognition of these cross-links may play a prevalent role in antitumor effects of these metallodrugs. Moreover, the results show for the first time a strong specific recognition and binding of high-mobility-group domain proteins, which are known to modulate antitumor effects of clinically used platinum drugs, to DNA modified by a polynuclear platinum complex.

Key words: dinuclear platinum; antitumor; DNA damage; interstrand cross-link; recognition

1. Introduction

The polynuclear platinum compounds represent a class of new antitumor metallodrugs that is structurally distinct from conventional *cis*-diamminedichloridoplatinum(II) (cisplatin) and its mononuclear analogues, and whose clinical profile and mechanism of action are different from these established platinum mononuclear compounds [1-6]. DNA adducts of polynuclear platinum complexes, whose formation is associated with antitumor effects of these agents, differ significantly in structure and type from those of mononuclear platinum complexes. Especially because of markedly more distant leaving groups, long-range intra- and interstrand crosslinks (CLs) are formed in DNA which affect DNA conformation, are recognized by specific proteins and repaired differently in comparison with the CLs of cisplatin and its mononuclear analogues [7-11]. Importantly, cells with resistance to cisplatin showed no cross-resistance to polynuclear platinum compounds [12-14]. The polynuclear platinum complexes also exhibit significantly higher levels of cellular uptake very likely because of their lipophilic linker chains and hydrophilic platinum-amine coordination spheres, which may enhance membrane permeability [13, 15, 16].

The biological activity of polynuclear platinum complexes may be modulated by the geometry and number of leaving groups in the coordination sphere of platinum atoms as well as by the nature of linkers connecting the platinum centers. In contrast with the mononuclear complexes, such as antitumor cisplatin and clinically ineffective transplatin, in the dinuclear case both geometries are antitumor active [2], although DNA adducts (CLs) as well as local conformational distortions on DNA and their recognition by cellular components are affected by geometry [7, 10, 17]. The dinuclear *cis* isomer $[\{cis\text{-PtCl}(\text{NH}_3)_2\}_2(\text{H}_2\text{N}-(\text{CH}_2)_6\text{-NH}_2)]^{2+}$ is kinetically more inert in its reactions with DNA and in double-stranded DNA produces more interstrand crosslinks than its trans counterpart [10].

Also importantly, the general structure, such as $[\{trans\text{-PtCl}(\text{NH}_3)_2\}_2\text{-}\mu\text{-Y}]^{n+}$ ($\text{Y} = \text{NH}_2(\text{CH}_2)_6\text{NH}_2$, spermidine, spermine (BBR3610), $trans\text{-Pt}(\text{NH}_3)_2(\text{H}_2\text{N}(\text{CH}_2)_6\text{NH}_2)_2$ (BBR3464), etc.), is susceptible to decomposition by the sulfur nucleophiles [18] since substitution of the Pt-Cl bond by a trans-influencing S donor results in bridge cleavage [19-21]. Hence, the antitumor polynuclear $trans$ Pt^{II} complexes, such as for instance trinuclear BBR3464 or dinuclear BR3610, undergo deactivation upon reaction with sulfur nucleophiles, which could result in decreased bioavailability, as well as in increased amounts of toxic metabolites in the host system, which may limit their clinical use [19]. On the other hand, the cis geometry as in $[\{cis\text{-PtCl}(\text{NH}_3)_2\}_2\text{-}\mu\text{-Y}]^{n+}$ should preserve the main features of antitumor polynuclear Pt^{II} complexes but with enhanced stability to metabolic deactivation.

In an effort to design a polynuclear platinum compound that maintains the target (DNA) binding profile of the cytotoxic Pt^{II} polynuclear compounds and is less susceptible to metabolic decomposition, the long-chain bifunctional dinuclear cis platinum complexes $[\{\text{PtCl}(\text{DACH})\}_2\text{-}\mu\text{-H}_2\text{N}(\text{CH}_2)_6\text{NH}_2(\text{CH}_2)_2\text{NH}_2(\text{CH}_2)_6\text{NH}_2]^{4+}$ (DACH = 1,2-diaminocyclohexane) and $\{[cis\text{-Pt}(\text{NH}_3)_2\text{Cl}]_2(4,4'\text{-methylenedianiline})\}^{2+}$ (complex **1**, Fig. 1) were synthesized [22, 23]. Hence, both dinuclear cis Pt^{II} complexes exhibit enhanced stability to metabolic deactivation [22, 23].

Platinum–DNA binding, recognition of platinated DNA by proteins and DNA repair are important pharmacological features dictating a platinum drug's efficacy [2, 24, 25]. As regards molecular mechanisms of antitumor polynuclear Pt^{II} complexes, a great attention has been so far paid mostly to polynuclear $trans$ Pt^{II} complexes whose clinical application is limited due to their lowered stability to metabolic deactivation. Thus, the important aim of this study was to broaden theoretical background of biological (antitumor) effects of dinuclear cis Pt^{II} complexes, which exhibit enhanced stability to metabolic deactivation. The primary objective in the present study was to understand more deeply those aspects of the molecular

mechanism of action of the new dinuclear *cis* Pt^{II} complex **1** containing a semi-rigid linker and for comparative purposes the molecular mechanism of action of a structurally cognate dinuclear *cis* Pt^{II} complex containing a shorter, more rigid linker, {[*cis*-Pt(NH₃)₂Cl]₂(α,α' -diamino-p-xylene)}²⁺ (complex **2**, Fig. 1) which are related to DNA modification by these complexes, recognition of these modifications by proteins and DNA repair. Here, we describe experiments revealing (i) new details of the DNA binding mode of **1** and **2** in cell-free media; (ii) repair of DNA adducts of **1** and **2** (since the integrity of this process in human cells is a key indicator of the sensitivity of a tumor to platinum-based therapy [26]); (iii) recognition of DNA adducts of **1** and **2** by high-mobility-group (HMG)-domain proteins (since these proteins are involved in the cisplatin mechanism of action [25]); (iv) sensitization of breast cancer cells to **1** and **2** by steroid hormones.

2. Materials and methods

2.1. Chemicals

Cisplatin was obtained from Sigma (Prague, Czech Republic) (purity was $\geq 99.9\%$ based on elemental and ICP trace analysis). Dinuclear Pt^{II} complexes **1** and **2** were prepared as described previously [23, 27]. The stock solutions of platinum compounds were prepared in 10 mM NaClO₄ and stored at 20°C in the dark. The concentrations of platinum complexes in the stock solutions were determined by flameless atomic absorption spectrometry (FAAS). The concentrations of the platinum complexes indicated in the present work are related to the whole compound (not to the Pt content in the case of the dinuclear complexes). Calf thymus (CT) DNA (42% G + C, mean molecular mass ca. 20 000 kDa) was prepared and

characterized as described previously [28, 29]. Plasmids, pUC19 [2686 base pairs (bp)] and pBR322 (4361 bp) were isolated according to standard procedures. Restriction endonucleases were purchased from New England Biolabs (Beverly, MA). Deoxyribonucleoside triphosphates were from Roche Diagnostics, GmbH (Mannheim, Germany). Agarose was from FMC BioProducts (Rockland, ME). Electrophoresis-grade acrylamide, N,N'-methylenebisacrylamide, ethidium bromide (EtBr), urea, and dithiothreitol were from Merck KgaA (Darmstadt, Germany). Proteinase K and ATP were from Boehringer (Mannheim, Germany). Spermine, EZBlue Gel Staining Reagent, β -estradiol and progesterone were from Sigma (Prague, Czech Republic). Expression and purification of recombinant rat full-length HMGB1 protein (HMG = high mobility group) and its domains A (residues 1-84 [30]) and B (residues 85-180 [30]) (HMGB1a and HMGB1b, respectively) were carried out as described [30-32]. Nonidet P-40 was from Fluka (Prague, Czech Republic). SDS was from Serva (Heidelberg, Germany). Radioactive products were from Amersham (Arlington Heights, IL, USA). A cell-free extract (CFE) was prepared from the repair proficient HeLa S3 cell line as described [33, 34]. Streptavidin-coated magnetic beads (Dynabeads) and Biotin-14-dATP were from Invitrogen Dynal AS (Oslo, Norway).

2.2. Cytotoxicity

Cisplatin, **1** and **2** were dissolved in DMSO. Stock solutions of the compounds were freshly prepared before use. The final concentration of DMSO in cell culture medium did not exceed 0.25%. The A2780 and A2780cisR human ovarian carcinoma cell lines (parent cisplatin sensitive and with acquired cisplatin resistance, respectively) were grown in RPMI 1640 medium (GIBCO) supplemented with gentamycin (50 μ g/mL, Serva) and 10% heat inactivated fetal bovine serum (GIBCO). The acquired resistance of A2780cisR cells was

maintained by supplementing the medium with 1 μM cisplatin every second passage. The cells were cultured in a humidified incubator at 37°C in a 5% CO_2 atmosphere and subcultured 2-3 times a week with an appropriate plating density.

Cell death was evaluated by using a system based on the tetrazolium compound MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] which is reduced by living cells to yield a formazan product that can be detected colorimetrically [35]. The cells were seeded in 96-well tissue cultured plates at a density of 10^4 cells/well in 100 μL of medium. After overnight incubation (16 hr), the cells were treated with the platinum compounds at the final concentrations in the range of 0 to 128 μM in a final volume of 200 μL /well. After additional 72 hr 10 μL of a freshly diluted MTT solution (2.5 mg/mL) was added to each well and the plate was incubated at 37°C in a humidified 5% CO_2 atmosphere for 4 hr. At the end of the incubation period the medium was removed and the formazan product was dissolved in 100 μL of DMSO. The cell viability was evaluated by measurement of the absorbance at 570 nm, using an Absorbance Reader SUNRICE TECAN SCHOELLER. IC_{50} values were calculated from curves constructed by plotting cell survival (%) versus drug concentration (μM). All experiments were made in triplicate. The reading values were converted to the percentage of control (% cell survival). Cytotoxic effects were expressed as IC_{50} (concentration inhibiting cell growth by 50%).

2.3. *Platination reactions in cell-free media*

If not stated otherwise, DNA was incubated with the platinum complex in 10 mM NaClO_4 at 37°C in the dark. After 24 hr, the samples were exhaustively dialyzed against the medium required for subsequent biochemical or biophysical analysis. An aliquot of these

samples was used to determine the value of r_b by FAAS. Alternatively, DNA binding of **1** and **2** was studied using differential pulse polarography (DPP) [36].

2.4. Recognition of DNA adducts by HMG-domain protein

The EcoRI/PvuII restriction fragment (92 bp) from pUC19 plasmid was modified with cisplatin or complexes **1** or **2** to desired r_b value in 0.01 M NaClO₄ at 37°C, in dark for 24 hr. Following the incubation or the optional radioactive labeling (by γ -³²P ATP), the fragments were 3'-end biotinylated using Biotin-14-dATP label and terminal deoxynucleotidyl transferase in terminal transferase buffer (100 mM cacodylate buffer, pH 6.8; 5 mM CoCl₂, 0.5 mM dithiothreitol). Biotinylated DNA was attached to streptavidine coated magnetic beads in 1 M NaCl, 5 mM Tris.HCl pH 7.4 and 0.5 mM EDTA. The beads were washed three times in the same buffer. HMG protein was added to the beads carrying DNA and incubated on ice in 10 mM HEPES, pH 7.5, 10 mM MgCl₂, 50 mM LiCl, 100 mM NaCl, 1 mM spermidine and 0.05 % Nonidet P40, in the presence of poly(dI-dC) as a competitor for 1 hr. The beads were then washed 2 times in the same buffer. Finally the protein was released with 0.04 % sodium dodecyl sulfate (SDS). Following the addition of loading buffer (0.1 M Tris-HCl, pH 6.8; 2 % SDS; 0.1 M dithiothreitol, 10 % glycerol, 0.25 % bromphenol blue) the samples were heated for five minutes at 90°C, electrophoresed on 15 % SDS-PAGE (PAGE = polyacrylamide gel electrophoresis) and stained with EZBlue gel staining reagent.

2.5. DNA repair synthesis by human cell extract

Repair DNA synthesis of CFEs was assayed using pUC19 plasmid. Each reaction of 50 μ L contained 600 ng of nonmodified pBR322 and 600 ng of nonmodified or platinated

pUC19, 2 mM ATP, 30 mM KCl, 0.05 mg/mL creatine phosphokinase (rabbit muscle), 20 mM each dGTP, dATP, and dTTP, 8 mM dCTP, 74 kBq of [α - 32 P]dCTP in the buffer composed of 40 mM HEPES-KOH, pH 7.5, 5 mM MgCl₂, 0.5 mM dithiothreitol, 22 mM creatine phosphate, 1.4 mg of bovine serum albumin/mL, and 20 μ g of CFE from the HeLa S3 cells. Reactions were incubated for 3 hr at 30°C and terminated by adding EDTA to a final concentration of 20 mM, SDS to 0.6%, and proteinase K to 250 μ g/mL and then incubating for 20 min. The products were extracted with one volume of 1:1 phenol/chloroform. The DNA was precipitated from the aqueous layer by the addition of 0.1 volume of 3 M sodium acetate and 2.5 volumes of ethanol. After 30 min of incubation at -20°C and centrifugation at 12 000 g for 30 min at 4°C, the pellet was washed with 0.2 mL of 80% ethanol and dried in a vacuum centrifuge. DNA was finally linearized by SspI before electrophoresis on a 1% agarose gel. Resulting gel was stained by EtBr. The experiments were made in quadruplicate.

2.6. Sensitization of cancer cells by steroid hormones to platinum complexes by clonogenic assay

MCF-7 cells were seeded on 6 cm Petri dishes at a density of 400 cells per dish. After 24 hr, a fresh stock solution containing steroid hormone(s), estradiol or estradiol plus progesterone, was prepared in *N,N*-dimethylformamide and added to the dishes so that a final concentration of each hormone was 0.2 μ M. Control dishes were treated with the same volume of *N,N*-dimethylformamide without hormone. The hormone was added at the same time as platinum complex (**1**, **2**, or cisplatin). After 4 hr treatment, the cells were washed with PBS, and fresh medium was added. After 10 days, the cell colonies were stained with 1% methylene blue and were then counted. Each point is an average of six independent determinations.

2.7. Other physical methods

Absorption spectra were measured with a Beckmann DU-7400 spectrophotometer. FAAS measurements were carried out with a Varian AA240Z Zeeman atomic absorption spectrometer equipped with a GTA 120 graphite tube atomizer. For FAAS analysis DNA was precipitated with ethanol and dissolved in 0.1 M HCl. The gels were visualized by using the BAS 2500 FUJIFILM bioimaging analyzer, and the radioactivities associated with bands were quantitated with the AIDA image analyzer software (Raytest, Germany).

3. Results

3.1. Cytotoxicity

The cytotoxic activity of the **1** and **2** tested in the present work was determined against A2780 (parent cisplatin sensitive) and A2780cisR (with acquired cisplatin resistance) human ovarian carcinoma cell lines, commonly used to test cytotoxic activity of cisplatin analogues and other antitumor metallodrugs. The tumor cell lines were incubated for 72 hr with the platinum compounds and the cell survival in the culture treated with the platinum compounds was evaluated as described in the section Experimental Procedures. Results (Table 1) show that the cytotoxicity of **1** in cisplatin sensitive A2780 cells was comparable to the cytotoxicity of cisplatin, whereas **2** was slightly less cytotoxic. Importantly, both **1** and **2** were markedly more active in the cisplatin-resistant line A2780cisR and again **2** was less cytotoxic (Table 1).

3.2. Repair DNA synthesis by human cell extracts

To investigate “downstream” effects of DNA damage by adducts of **1** and **2**, we used *in vitro* systems to study repair of platinum lesions by CFE. DNA repair efficiency in pUC19 plasmid (2686 bp) globally modified by **1**, **2** or cisplatin (for comparative purposes) at $r_b = 0.03$ (r_b is defined as the number of molecules of the platinum complex bound per nucleotide residue) was tested using CFE of repair proficient HeLa cells. Repair activity was monitored by measurement of the amount of incorporated radiolabelled nucleotide. The incorporation of radioactive material was corrected for the relative DNA content in each band. As illustrated in Fig. 2, damage-induced DNA repair synthesis detected in the plasmid modified by **1** or **2** was approximately 38 or 36 %, respectively, of that found for the cisplatin at the same level of modification.

3.3. DNA modification in a cell-free medium

The cellular DNA repair response strongly depends on the character and extent of DNA damage. Therefore, it was of great interest to characterize DNA distortions induced by **1** and **2**. With the view of this we examined the DNA binding properties of these complexes and compared them with those of cisplatin. The experiments were aimed at quantification of the binding of **1** and **2** to mammalian DNA, determination of their preferential binding sites, and characterization of DNA lesions induced by **1** and **2** within natural DNA. The results of these experiments are described in detail in the Supplemental Information and are summarized in the following paragraph.

3.3.1. DNA binding

CT DNA was incubated with the platinum complexes at r_i values of 0.01, 0.03, 0.05, and 0.1) (r_i is defined as the molar ratio of free platinum complex to nucleotides at the onset of incubation with DNA) in 10 mM NaClO₄ at 37°C for 24 or 48 hr. The amount of the platinum compound bound to DNA increased with time. In this binding reaction the time at which the binding reached 50% ($t_{50\%}$) was 14.9 ± 0.6 min and 39 ± 6 min for **1** and **2**, respectively, and both complexes were quantitatively bound after 24 hr. This result indicates that the rate of binding of both **1** and **2** to natural double-helical DNA is higher than that of cisplatin ($t_{50\%}$ = ca. 120 min [37]), **although cisplatin is also quantitatively bound to DNA after 24 hr under identical experimental conditions [37].**

3.3.2. Transcription mapping of platinum-DNA adducts

In vitro RNA synthesis by T7 and SP6 RNA polymerases on DNA templates containing adducts of **1** or **2** revealed (Fig. S1) that the major stop sites produced by these complexes were at guanine residues, a few stop sites were also at adenine and cytosine residues. These results also suggest that the adducts formed by **1** and **2** are able to inhibit RNA polymerase.

3.3.3. Interstrand cross-linking

We quantitated the interstrand cross-linking efficiency of **1** and **2** in linearized pUC19 plasmid (Fig. S2). The interstrand cross-linking efficiencies (**defined as the number of interstrand crosslinks per one adduct of the platinum complex [17]**) of both **1** and **2** (48 ± 4 % and 20 ± 2 %, respectively) were significantly higher than that of cisplatin (6 % [38]).

3.3.4. Characterization of DNA adducts by ethidium bromide fluorescence

The results of these experiments (Fig. S3) suggest that the conformational distortion induced in DNA by the adducts of **1** is more delocalized and extends over more base pairs around the platination sites than in the case of the adducts of **2** or mononuclear cisplatin. Thus, these results are consistent with the formation of long-range intra- or interstrand CLs of **1**, whereas **2** apparently forms in DNA much less long-range lesions. In aggregate, characterization of DNA adducts of **1** and **2** by EtBr fluorescence supports the view that the DNA binding modes of these dinuclear platinum compounds are distinctly different.

3.3.5. Unwinding of negatively supercoiled DNA

Electrophoresis in native agarose gel was used to quantify the unwinding induced in pUC19 plasmid by adducts of **1** and **2** by monitoring the degree of supercoiling (Fig. S4). We determined the DNA unwinding angle of 17° for the adduct of **1** and 12° for that of **2**. The high level of unwinding induced by **1** and **2** is notable. Interestingly, the unwinding angle produced by **1** is higher than that produced by **2**. One plausible explanation for this observation might be consistent with a possibility that the additional contribution to unwinding is associated with the interaction of the linker chain of **1** with the duplex upon covalent binding of platinum units. Thus, the results of unwinding experiments represent another support for the view that structure of DNA lesions formed by **1** is substantially different from the structure of DNA adducts of its counterpart **2**.

3.4. Recognition of DNA adducts by high-mobility-group (HMG)-domain proteins

The altered structures induced in DNA by cisplatin and its analogs attract HMG-domain and other proteins [30, 39]. This binding of HMG-domain proteins to DNA modified by cisplatin and its analogs has been postulated to mediate the antitumor properties of these

drugs.[39] As dinuclear platinum complexes **1** and **2** exhibit antitumor activity qualitatively different from cisplatin, it was of interest to examine whether also the adducts formed within DNA by **1** or **2** are able to be specifically recognized by HMG domain proteins.

To assess the ability of HMGB1 protein to recognize DNA modified by complexes **1** and **2** we employed the system of DNA immobilized on streptavidine coated magnetic beads suspended in HMG binding buffer containing HMGB1 protein. 10 μ g of DNA fragment was immobilized on 1 mg of Dynabeads. Radioactively labeled DNA enabled us to ensure the same quantity of DNA in various samples. First we determined whether HMGB1 protein exhibits the ability to recognize DNA modified by the two new dinuclear compounds. Immobilized DNA (nonmodified or modified by cisplatin, complex **1** and **2** to $r_b = 0.02$) was incubated with HMGB1 protein on ice for 1 hr in the presence of poly(dI-dC) as a competitor. Following the incubation, DNA was washed extensively and bound protein was released in 0.04 % SDS and electrophoresed on 15 % SDS-PAGE.

The HMGB1 protein was found to bind DNA modified by cisplatin, in agreement with experiments published earlier [40, 41] that was evidenced by the presence of band migrating near the 34 kDa marker corresponding to HMGB1 protein (Fig. 3A, lane cisPt, [Table S1](#)). Interestingly, the band corresponding to HMGB1 was also clearly seen for the probe modified by **1** (Fig. 3A, lane **1**); the intensity of this band was almost comparable (80%) to that found for cisplatin ([Table S1](#)). On the other hand, much less effective binding of the HMGB1 to the DNA containing the adducts of **2** was detected, the amount of HMGB1 protein bound to this probe was only 15% of the amount of HMGB1 bound to DNA modified by cisplatin under the same conditions (Fig. 3A, lane 2, [Table S1](#)). No binding of the proteins occurred under identical experimental conditions if the same DNA probe was unplatinated (Fig. 3A, lane NoPt). These results clearly showed that DNA modified by **1** is efficiently recognized by HMGB1 with relatively high affinity, though slightly less than to the probe containing the

adducts of cisplatin. In contrast, DNA adducts of **2** are recognized by this protein markedly less.

To determine which of the two HMGB1 domains A or B predominates in recognition and binding of DNA lesions of **1**, the binding of HMGB1 domain A and HMGB1 domain B to the DNA modified by complexes **1** or **2** was investigated (Fig. 3 B,C). The experiments revealed that in agreement with previously published data [41], HMGB1a effectively bound to the probe containing adducts of cisplatin (Fig. 3B, lane cisPt). Considerably weaker binding of HMGB1a was noticed to the probes containing adducts of **1** or **2**; relative amount of HMGB1a protein bound to DNA probe containing adducts of cisplatin, **1** or **2** was 100%, 20%, or 14%, respectively (Fig. 3B, Table S1). On the other hand, the amount of HMGB1b bound to the DNA modified by **1** was two-fold higher compared to that bound to DNA modified by cisplatin (Fig. 3C, Table S1). These results clearly show that DNA damaged by complex **1** was recognized by HMGB1 protein, but in contrast to DNA globally modified by cisplatin, the full-length HMGB1 protein binds to DNA globally modified by **1** mainly through domain B.

Complex **1** modifies DNA so that it forms approximately equal amounts of interstrand and intrastrand adducts (see paragraph Interstrand cross-linking in this Section and Fig. S2). To distinguish which type of lesions formed by **1** within DNA is preferentially recognized by HMGB1 protein, the following experiment was performed. The 92 bp fragment of DNA was modified by **1** so that the fragment contained only one adduct per DNA fragment on average. After the modification, DNA was loaded into the denaturing agarose gel and two DNA fractions were purified from the gel, one contained only intrastrand adducts (no interstrand CLs) (F-IAA) and the other contained at least one interstrand CL (F-IEC). These two separated DNA fractions were then allowed to interact with HMGB1b, as described in section Experimental procedures. The amount of HMGB1b protein bound to each DNA fraction was

determined by SDS/PAGE electrophoresis. The inspection of the electrophoreogram (Fig. 4) revealed that HMGB1b bound to DNA containing interstrand CLs of **1** whereas it bound considerably less (4-fold) to DNA modified by **1** to the similar level but containing other adducts, but not interstrand CLs (Table S2).

3.5. Steroid hormones sensitize cancer cells to complex **1**, but not to complex **2**

Previous studies have shown that HMGB1 protein sensitizes cells to cisplatin by shielding its DNA adducts from nucleotide excision repair [25, 42]. Human breast cancer cells MCF-7 contain estrogen and progesterone receptors and respond to these steroids up-regulating HMGB1 protein [43]. Importantly, treatment of these cancer cells with estrogen or progesterone significantly increases the potency of cisplatin apparently owing to the overexpression of HMGB1 [44]. Since DNA adducts of **1** are recognized by HMGB1 protein **markedly more than those of 2** (Fig. 3), it was of interest to assess the involvement of HMG-domain proteins in mediating cytotoxicity of the dinuclear Pt^{II} complexes tested in the present work. Therefore, we investigated the effects of elevated HMGB1 levels due to steroid hormone treatment in MCF-7 cells on the sensitivity of these cells to **1** or **2** (Fig. 5) using clonogenic assay in the same way as described previously for cisplatin and carboplatin [44]. As anticipated, we found that elevated HMGB1 expression levels were paralleled by increased sensitivity toward **1**, but markedly less toward **2** (Fig. 5 and Table S4). For instance, in MCF-7 cells, estradiol or estradiol plus progesterone treatment increased sensitivity toward **1** about 2.7- or 4.2-fold, respectively, when only 50% of the cells are viable (Fig. 5A); estradiol or estradiol plus progesterone treatment also increased sensitivity toward cisplatin, but only about 1.8- or 2.5-fold, respectively (Fig. 5C). Thus, sensitivity toward **1** was enhanced due to steroid hormone treatment even considerably more than toward cisplatin.

Sensitivity of MCF-7 cells, treated with hormones, toward **1**, **2**, and cisplatin was also determined by MTT assay (Table S3). The trend was similar to that observed when clonogenic assay was used, although IC₅₀ data obtained by MTT assay were uniformly higher than in clonogenic assay.

4. Discussion

We demonstrate in the present work that the dinuclear complexes **1** and **2** are quite toxic to the ovarian tumor cells, with IC₅₀ values that were comparable to that observed for cisplatin in the cisplatin sensitive cell line A2780, and noticeably much lower in the cisplatin-resistant line A2780cisR (Table 1). Thus, these results complement those obtained in the previous study [27] showing that complexes **1** and **2** exhibit the promising cytotoxicity to the human nonsmall-cell lung cancer cell line (A549). Interestingly, the cytotoxicity of both **1** and **2** to the ovarian tumor cells is characterized by remarkably low resistance factors, equal to or even less than **1**, which indicates that the dinuclear platinum complexes examined in the present work are capable of circumventing of cisplatin resistance in some types of the cisplatin-resistant lines. In aggregate, no cross-resistance to cisplatin and high cytotoxic effects in cisplatin resistant cells represent important pharmacological features of **1** and **2**.

An important feature of the mechanism underlying antitumor effects of DNA-binding metal-based compounds is repair of their DNA adducts [40, 45]. A persistence of these DNA adducts may potentiate their antitumor effects in the cells sensitive to these compounds [8, 39, 40]. DNA repair synthesis was investigated in the present work using the CFE from human tumor cells and DNA substrates randomly modified by **1** or **2**. Importantly, the adducts of both **1** and **2** induced a considerably lower level of repair synthesis than the adducts of

cisplatin suggesting a less efficient removal from DNA and enhanced persistence of the adducts of the dinuclear complexes **1** and **2**. Thus, enhanced resistivity of DNA adducts of **1** and **2** against repair processes appears to be an important factor contributing to ability of **1** and **2** to overcome resistance towards cisplatin.

Structural features of DNA distortions are considered to be major factors playing a key role in recognition by damaged-DNA recognition-proteins and DNA repair systems. The analysis of DNA binding of **1** and **2** described in this report and summarized in Table 2 provides experimental support for the view that the binding of these dinuclear platinum complexes modifies DNA in a way which is in several aspects considerably different from modification of DNA by "conventional" mononuclear cisplatin. These differences may suggest that in comparison with adducts of cisplatin the reduced repair of DNA adducts of the dinuclear Pt^{II} complexes is a consequence of their different DNA binding mode.

The binding of **1** and **2** to DNA is considerably faster than the binding of dichloro derivatives of mononuclear Pt^{II} complexes, such as cisplatin or transplatin ($t_{50\%} \sim 120$ min [37]) likely reflecting the higher charge. In addition, the different rate of the binding of **1** and **2** may be associated with the differences in the linking chains, such as flexibility and hydrophobicity, which may affect pre-association step of DNA binding [46, 47].

Importantly, the effect of the linker **character** and geometry results in enhanced extent of the long-range delocalized cross-linking by **1** as revealed by fluorescence measurements with EtBr (Fig. S3). Similarly, DNA interstrand cross-linking by **1** is enhanced compared with that by **2** (Fig. S2, Table 2). This enhanced interstrand crosslinking efficiency of **1** correlates with its higher potency compared with that of **2** so that interstrand CLs appear to play the prevalent role in cytotoxicity of **1**. In addition, compared with **2**, **1** distorts DNA secondary structure differently and unwinds DNA more (Fig. S4 and Table 2). Thus, it is reasonable to expect that several "downstream" intracellular effects of DNA adducts of **1** and **2** are different.

Consistent with this view is the observation that affinity of HMG-domain proteins to DNA adducts of **1** is markedly higher than that to DNA adducts of **2** (Fig. 3A). The full-length HMGB1 protein consists of two tandem HMG box domains (A and B) and a C-terminal acidic tail. Both A and B domains of HMGB1 protein share a common HMG box structure. Domain A has a higher binding affinity to distorted DNA than domain B, whereas domain B bends DNA more effectively [48]. Because of the importance of the HMG box interaction with DNA [25], the studies of the individual HMG domains and their DNA complexes are of great interest. It has been shown that full-length HMGB1 protein binds to DNA containing a site-specific 1,2-GG intrastrand CL of cisplatin mainly through domain A, whereas domain B alone binds cisplatinated DNA much more weakly than domain A alone [41]. In contrast to these data, binding of full-length HMGB1 to DNA modified by the dinuclear complex **1** is mediated preferentially by domain B (Fig. 3B, C). This result, in combination with our previous findings that the domain B of the HMGB1 protein with the A/B linker plays a crucial role in the interaction between full-length HMGB1 protein and DNA interstrand cross-linked by cisplatin [30], implies a possible role of the interstrand DNA adducts of **1** in the interaction with HMG-domain proteins. Therefore, we tested the contribution of DNA intrastrand and interstrand adducts formed by **1** to the binding of HMGB1 to DNA globally modified by **1**. The sample containing DNA fragment globally modified by **1** was divided into two DNA fractions, one containing only intrastrand adducts (no interstrand CLs) and the other containing interstrand CLs and the affinity of HMGB1b towards these two fractions was tested. The results unambiguously show the prevalent role of DNA interstrand CLs of **1** in recognition and binding of DNA modified by **1** to full-length HMGB1 protein. To our knowledge, this is the first evidence for strong specific recognition and binding of HMG-domain proteins to DNA modified by a polynuclear platinum complex. Previously, a weak recognition of DNA interstrand CLs of [$\{trans\text{-PtCl}(\text{NH}_3)_2\}\text{H}_2\text{N}(\text{CH}_2)_2$ -

${}^6\text{NH}_2\text{Cl}_2$ complexes by full-length HMGB1 protein has only been reported [7], whereas a recognition of intrastrand CLs of these complexes has not been detected.

The results of the present work provide evidence that HMGB1 domain B controls a relatively strong binding of full-length HMGB1 to DNA modified by **1** when both domain A and B are present. It is quite conceivable that domain B is the dominating domain in HMGB1 protein that binds to the interstrand CL of **1**, while domain A only facilitates binding by providing additional protein-DNA interactions. A structure of full-length HMGB1 complexed with DNA containing interstrand CL of **1** will provide more insights into how HMG-domain proteins can mediate cytotoxicity of **1**. On the other hand, the affinities of full-length HMGB1 protein and its domain A to DNA modified by **2** are relatively weak and almost identical. This result is consistent with the view and supports the hypothesis that the binding of full-length HMGB1 protein to DNA modified by the dinuclear complex **2**, although relatively weak, is mediated preferentially by the domain A. Collectively, the results of this work demonstrate that not only DNA binding modes of **1** and **2** and resulting conformational alterations are noticeably different, but also “downstream” intracellular effects of DNA adducts of the two dinuclear complexes may be different, particularly in the cells in which the HMGB1 level is high and in which HMGB1 protein is not prevented from interacting with platinum-modified DNA.

It has been shown that specific interaction between adducts of "conventional" cisplatin and cellular HMG-domain proteins modulate the sensitivity of tumor cells to this drug. In particular, protein-DNA interactions induced by cisplatin inhibit replication,[49] shield the adducts from nucleotide excision repair [50] and interfere with transcription by recruiting transcription factors from their native binding sites [51, 52]. Therefore, it is reasonable to assume that such interactions may also differently modulate the cellular responses both to the

platinum dinuclear complexes **1** and **2** and also to the class of dinuclear complexes tested in the present work and cisplatin.

In conclusion, the results of the present work are consistent with the hypothesis that the unique properties of interstrand CLs of the class of bifunctional polynuclear platinum complexes tested in the present work and resulting conformational alterations in DNA have critical consequences for their antitumor effects. An interesting finding that might be also translated to the clinical trials is that treatment leading to enhancement of level of HMG-domain proteins in tumor cells should allow the regimens involving **1** to show increased antitumor effects toward these cells.

Acknowledgements

This research was supported by the Ministry of Education of the CR (MSMT LC06030, 6198959216, ME08017, ME10066, OC08003, and OC09018), the Academy of Sciences of the Czech Republic (Grants KAN200200651, M200040901, AV0Z50040507, and AV0Z50040702), the Grant Agency of the Academy of Sciences of the CR (IAA400040803), and the Grant Agency of the CR (301/09/H004 and P301/10/0598). Financial support from the National Natural Science Foundation of China (No. 90713001) is also acknowledged. J.K. is an international research scholar of the Howard Hughes Medical Institute. The authors also acknowledge that their participation in the EU COST Action D39 has enabled them to exchange regularly their most recent ideas in the field of anticancer metallodrugs with several European colleagues.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi: ...

References

- [1] Pratesi G, Perego P, Polizzi D, Righetti SC, Supino R, Caserini C, et al. A novel charged trinuclear platinum complex effective against cisplatin-resistant tumours: hypersensitivity of p53-mutant human tumour xenografts. *Br J Cancer* 1999;80:1912-9.
- [2] Farrell N. Polynuclear platinum drugs. In: Sigel A, Sigel H, editors. *Metal ions in biological systems*. New York, Basel: Marcel Dekker, Inc.; 2004, p. 251-96.
- [3] Brabec V, Kasparkova J. Modifications of DNA by platinum complexes: Relation to resistance of tumors to platinum antitumor drugs. *Drug Resist Updat* 2005;8:131-46.
- [4] Brabec V, Kasparkova J. DNA interactions of platinum anticancer drugs. Recent advances and mechanisms of action. In: Perez-Martin J-M, Fuertes MA, Alonso C, editors. *Metal compounds in cancer chemotherapy*. Trivandrum, Kerala (India): Research Signpost; 2005, p. 187-218.
- [5] Brabec V, Christofis P, Slamova M, Kostrhunova H, Novakova O, Najajreh Y, et al. DNA interactions of new cytotoxic tetrafunctional dinuclear platinum complex trans,trans-[PtCl₂(NH₃)₂(piperazine)]. *Biochem Pharmacol* 2007;73:1887-900.
- [6] Cossa G, Gatti L, Zunino F, Perego P. Strategies to improve the efficacy of platinum compounds. *Curr Med Chem* 2009;16:2355-65.
- [7] Kasparkova J, Farrell N, Brabec V. Sequence specificity, conformation, and recognition by HMG1 protein of major DNA interstrand cross-links of antitumor dinuclear platinum complexes. *J Biol Chem* 2000;275:15789-98.
- [8] Kasparkova J, Zehnulova J, Farrell N, Brabec V. DNA interstrand cross-links of the novel antitumor trinuclear platinum complex BBR3464. Conformation, recognition by

- high mobility group domain proteins, and nucleotide excision repair. *J Biol Chem* 2002;277:48076-86.
- [9] Brabec V, Kasparikova J, Vrana O, Novakova O, Cox JW, Qu Y, et al. DNA modifications by a novel bifunctional trinuclear platinum Phase I anticancer agent. *Biochemistry* 1999;38:6781-90.
- [10] Kasparikova J, Novakova O, Vrana O, Farrell N, Brabec V. Effect of geometric isomerism in dinuclear platinum antitumor complexes on DNA interstrand cross-linking. *Biochemistry* 1999;38:10997-1005.
- [11] Zehnulova J, Kasparikova J, Farrell N, Brabec V. Conformation, recognition by high mobility group domain proteins, and nucleotide excision repair of DNA intrastrand cross-links of novel antitumor trinuclear platinum complex BBR3464. *J Biol Chem* 2001;276:22191-9.
- [12] Roberts JD, Peroutka J, Farrell N. Cellular pharmacology of polynuclear platinum anticancer agents. *J Inorg Biochem* 1999;77:51-7.
- [13] Roberts JD, Beggiolin G, Manzotti C, Piazzoni L, Farrell N. Comparison of cytotoxicity and cellular accumulation of polynuclear platinum complexes in L1210 murine leukemia cell lines. *J Inorg Biochem* 1999;77:47-50.
- [14] Servidei T, Ferlini C, Riccardi A, Meco D, Scambia G, Segni G, et al. The novel trinuclear platinum complex BBR3464 induces a cellular response different from cisplatin. *Eur J Cancer* 2001;37:930-8.
- [15] Farrell N. Polynuclear charged platinum compounds as a new class of anticancer agents: Toward a new paradigm. In: Kelland LR, Farrell NP, editors. *Platinum-based drugs in cancer therapy*. Totowa/NJ: Humana Press Inc.; 2000, p. 321-38.
- [16] Harris AL, Yang X, Hegmans A, Povirk L, Ryan JJ, Kelland L, et al. Synthesis, characterization, and cytotoxicity of a novel highly charged trinuclear platinum

- compound. Enhancement of cellular uptake with charge. *Inorg Chem* 2005;44:9598-600.
- [17] Farrell N, Qu Y, Feng L, Van Houten B. Comparison of chemical reactivity, cytotoxicity, interstrand cross-linking and DNA sequence specificity of bis(platinum) complexes containing monodentate or bidentate coordination spheres with their monomeric analogues. *Biochemistry* 1990;29:9522-31.
- [18] Summa N, Maigut J, Puchta R, van Eldik R. Possible biotransformation reactions of polynuclear Pt(II) complexes. *Inorg Chem* 2007;46:2094-104.
- [19] Oehlsen ME, Qu Y, Farrell N. Reaction of polynuclear platinum antitumor compounds with reduced glutathione studied by multinuclear (^1H , ^1H - ^{15}N gradient heteronuclear single-quantum coherence, and ^{195}Pt) NMR spectroscopy. *Inorg Chem* 2003;42:5498 - 506.
- [20] Billecke C, Finniss S, Tahash L, Miller C, Mikkelsen T, Farrell NP, et al. Polynuclear platinum anticancer drugs are more potent than cisplatin and induce cell cycle arrest in glioma. *Neuro Oncol* 2006;8:215-26.
- [21] Mitchell C, Kabolizadeh P, Ryan J, Roberts JD, Yacoub A, Curiel DT, et al. Low-dose BBR3610 toxicity in colon cancer cells is p53-independent and enhanced by inhibition of epidermal growth factor receptor (ERBB1)-Phosphatidyl inositol 3 kinase signaling. *Mol Pharmacol* 2007;72:704-14.
- [22] Williams JW, Qu Y, Bulluss GH, Alvorado E, Farrell NP. Dinuclear platinum complexes with biological relevance based on the 1,2-diaminocyclohexane carrier ligand. *Inorg Chem* 2007;46:5820-2.
- [23] Fan D, Yang X, Wang X, Zhang S, Mao J, Ding J, et al. A dinuclear monofunctional platinum(II) complex with an aromatic linker shows low reactivity towards glutathione

- but high DNA binding ability and antitumor activity. *J Biol Inorg Chem* 2007;12:655-65.
- [24] Jamieson ER, Lippard SJ. Structure, recognition, and processing of cisplatin-DNA adducts. *Chem Rev* 1999;99:2467-98.
- [25] Jung Y, Lippard SJ. Direct cellular responses to platinum-induced DNA damage. *Chem Rev* 2007;107:1387-407.
- [26] Brabec V, Kasparikova J. Role of DNA repair in antitumor effects of platinum drugs. In: Hadjiladis N, Sletten E, editors. *Metal complex - DNA interactions*. Chichester, UK: Wiley; 2009, p. 175-208.
- [27] Zhu J, Lin M, Fan D, Wu Z, Zhang J, Luc Y, et al. The role of bridging ligands in determining DNA-binding ability and cross-linking patterns of dinuclear platinum(II) antitumor complexes. *Dalton Trans* 2009:10889 -95.
- [28] Brabec V, Palecek E. The influence of salts and pH on polarographic currents produced by denatured DNA. *Biophysik* 1970;6:290-300.
- [29] Brabec V, Palecek E. Interaction of nucleic acids with electrically charged surfaces. II. Conformational changes in double-helical polynucleotides. *Biophys Chem* 1976;4:76-92.
- [30] Kasparikova J, Delalande O, Stros M, Elizondo-Riojas MA, Vojtiskova M, Kozelka J, et al. Recognition of DNA interstrand cross-link of antitumor cisplatin by HMGB1 protein. *Biochemistry* 2003;42:1234-44.
- [31] Stros M. DNA bending by the chromosomal protein HMG1 and its high mobility group box domains. Effect of flanking sequences. *J Biol Chem* 1998;273:10355-61.
- [32] Stros M. Two mutations of basic residues within the N-terminus of HMG-1 B domain with different effects on DNA supercoiling and binding to bent DNA. *Biochemistry* 2001;40:4769-79.

- [33] Reardon JT, Vaisman A, Chaney SG, Sancar A. Efficient nucleotide excision repair of cisplatin, oxaliplatin, and bis-aceto-ammine-dichloro-cyclohexylamine-platinum(IV) (JM216) platinum intrastrand DNA diadducts. *Cancer Res* 1999;59:3968-71.
- [34] Manley JL, Fire A, Cano A, Sharp PA, Gefter ML. DNA-dependent transcription of adenovirus genes in a soluble whole-cell extract. *Proc Natl Acad Sci USA* 1980;77:3855-9.
- [35] Alley MC, Scudiero DA, Monks A, Hursey ML, Czerwinski MJ, Fine DL, et al. Feasibility of drug screening with panels of human tumor cell lines using a microculture tetrazolium assay. *Cancer Res* 1988;48:589-601.
- [36] Kim SD, Vrana O, Kleinwächter V, Niki K, Brabec V. Polarographic determination of subnanogram quantities of free platinum in reaction mixture with DNA. *Anal Lett* 1990;23:1505-18.
- [37] Bancroft DP, Lepre CA, Lippard SJ. ^{195}Pt NMR kinetic and mechanistic studies of cis-diamminedichloroplatinum and trans-diamminedichloroplatinum(II) binding to DNA. *J Am Chem Soc* 1990;112:6860-71.
- [38] Brabec V, Leng M. DNA interstrand cross-links of trans-diamminedichloroplatinum(II) are preferentially formed between guanine and complementary cytosine residues. *Proc Natl Acad Sci USA* 1993;90:5345-9.
- [39] Brabec V. DNA modifications by antitumor platinum and ruthenium compounds: their recognition and repair. *Prog Nucleic Acid Res Mol Biol* 2002;71:1-68.
- [40] Cohen SM, Lippard SJ. Cisplatin: From DNA damage to cancer chemotherapy. *Prog Nucleic Acid Res Mol Biol* 2001;67:93-130.
- [41] Jung YW, Lippard SJ. Nature of full-length HMGB1 binding to cisplatin-modified DNA. *Biochemistry* 2003;42:2664-71.

- [42] Huang JC, Zamble DB, Reardon JT, Lippard SJ, Sancar A. HMG-domain proteins specifically inhibit the repair of the major DNA adduct of the anticancer drug cisplatin by human excision nuclease. *Proc Natl Acad Sci USA* 1994;91:10394-8.
- [43] Chau KY, Lam HYP, Lee KLD. Estrogen treatment induces elevated expression of HMG1 in MCF-7 cells. *Exp Cell Res* 1998;241:269-72.
- [44] He Q, Liang CH, Lippard SJ. Steroid hormones induce HMG1 overexpression and sensitize breast cancer cells to cisplatin and carboplatin. *Proc Natl Acad Sci USA* 2000;97:5768-72.
- [45] Brabec V, Kasparkova J. Molecular aspects of resistance to antitumor platinum drugs. *Drug Resist Updat* 2002;5:147-61.
- [46] Hegmans A, Berners-Price SJ, Davies MS, Thomas D, Humphreys A, Farrell N. Long range 1,4 and 1,6-interstrand cross-links formed by a trinuclear platinum complex. Minor groove pre-association affects kinetics and mechanism of cross-link formation as well as adduct structure. *J Am Chem Soc* 2004;126:2166-80.
- [47] Berners-Price SJ, Davies MS, Cox JW, Thomas DS, Farrell N. Competitive reactions of interstrand and intrastrand DNA-Pt adducts: A dinuclear-platinum complex preferentially forms a 1,4-interstrand cross-link rather than a 1,2 intrastrand cross-link on binding to a GG 14-Mer duplex. *Chem Eur J* 2003;9:713-25.
- [48] Thomas JO. HMG 1 and 2: architectural DNA-binding proteins. *Biochem Soc Trans* 2001;29:395-401.
- [49] Comess KM, Burstyn JN, Essigmann JM, Lippard SJ. Replication inhibition and translesion synthesis on templates containing site-specifically placed cis-diamminedichloroplatinum(II) DNA adducts. *Biochemistry* 1992;31:3975-90.
- [50] Zamble DB, Mu D, Reardon JT, Sancar A, Lippard SJ. Repair of cisplatin-DNA adducts by the mammalian excision nuclease. *Biochemistry* 1996;35:10004-13.

- [51] Zhai XQ, Beckmann H, Jantzen HM, Essigmann JM. Cisplatin-DNA adducts inhibit ribosomal RNA synthesis by hijacking the transcription factor human upstream binding factor. *Biochemistry* 1998;37:16307-15.
- [52] Vichi P, Coin F, Renaud JP, Vermeulen W, Hoeijmakers JHJ, Moras D, et al. Cisplatin- and UV-damaged DNA lure the basal transcription factor TFIID/TBP. *EMBO J* 1997;16:7444-56.
- [53] Fichtinger-Schepman AMJ, Van der Veer JL, Den Hartog JHJ, Lohman PHM, Reedijk J. Adducts of the antitumor drug cis-diamminedichloroplatinum(II) with DNA: Formation, identification, and quantitation. *Biochemistry* 1985;24:707-13.
- [54] Keck MV, Lippard SJ. Unwinding of supercoiled DNA by platinum ethidium and related complexes. *J Am Chem Soc* 1992;114:3386-90.

Table 1. *In vitro* growth inhibition of human ovarian cisplatin sensitive (A2780) and resistant (A2780cisR) cells, IC₅₀ (μM) ^{a,b}

| | A2780 | A2780cisR ^c |
|-----------|-----------|------------------------|
| cisplatin | 2.8 ± 0.7 | 18.6 ± 0.4 (6.6) |
| 1 | 2.7 ± 0.7 | 2.7 ± 0.1 (1) |
| 2 | 4.8 ± 2.7 | 3.6 ± 1.1 (0.75) |

^a Drug-treatment period was 72 hr.

^b Values shown in the table are the means (±SEM) of three separate experiments, each conducted with 6 replicates.

^c Resistance factor, defined as IC₅₀ (resistant) / IC₅₀ (sensitive), is given in parentheses.

Table 2. Summary of the effects of complex **1**, **2** and cisplatin on DNA ^a

| | 1 | 2 | cisplatin |
|---|----------|----------|---------------------|
| DNA binding ($t_{50\%}$) (min) ^b | 14.9±0.6 | 39±6 | 120 ^c |
| Preferential DNA binding sites ^d | G | G | GG, AG ^e |
| Reduction of EtBr fluorescence | strong | medium | medium |
| Plasmid DNA unwinding angle/adduct | 17° | 12° | 13° ^f |
| % interstrand cross-links/adduct | 48±4 | 20±2 | 6 ^g |

^a See also Figs. S1 - S4.

^b See the text for details.

^c Ref. [37].

^d Determined by transcription mapping.

^e Ref. [53].

^f Ref. [54].

^g Ref. [38].

Figure Captions

Fig. 1 - Structures of the platinum complexes.



Fig. 2 - *In vitro* DNA repair synthesis assay. Repair synthesis of the extract prepared from the repair-proficient HeLa cell line used as substrates nonmodified pBR322 plasmid and pUC19 plasmid nonmodified or modified at $r_b = 0.03$ by cisplatin, complex **1**, and complex **2**. (A) Results of a typical experiment. Top panel, a photograph of the EtBr stained gel; bottom panel, autoradiogram of the gel showing incorporation of [α - 32 P]dCTP. Lanes: 1, nonmodified pBR322 plus pUC19 plasmids; 2, nonmodified pBR322 plus pUC19 modified by cisplatin; 3, nonmodified pBR322 plus pUC19 modified by **1**; 4, nonmodified pBR322 plus pUC19 modified by **2**. (B) Incorporation of dCTP into nonmodified or platinated pUC19 plasmid. For all quantifications representing mean values of three separate experiments, incorporation of radioactive material is corrected for the relative DNA content in each band. The radioactivity associated with the incorporation of [α - 32 P]dCTP into DNA modified by cisplatin was taken as 100%. Values shown in the graph are the means (\pm SEM) of three separate experiments, each conducted with 4 replicates.

Fig. 3 - Recognition by HMG-domain proteins of platinated DNA. Affinity of full-length HMGB1 (A) HMGB1 domain A (B), and HMGB1 domain B (C) to nonmodified (control) DNA (lanes noPt) or DNA modified by cisplatin (lanes cisPt), complex **1** (lanes 1) or complex **2** (lanes 2). Lanes M, a protein size marker; lane HMGB1, HMGB1 protein was loaded; lane HMGB1 dom A, HMGB1 domain A was loaded.

Fig. 4 - Recognition by HMGB1 domain B of DNA fragments intrastrand or interstrand cross-linked by complex **1**. After reaction with **1**, the DNA was separated by gel electrophoresis so that the fraction F-IEC contained all interstrand cross-linked fragments and the fraction F-IAA contained exclusively platinum adducts formed within one strand of DNA (contained no interstrand CLs, for other details, see the text). Lanes: M, a protein size marker; noPt, control, nonplatinated; F-IAA, intrastrand adducts containing DNA incubated with HMGB1 domain B; F-IEC, interstrand cross-links containing DNA incubated with HMGB1 domain B.

Fig. 5 - The effects of estradiol or estradiol plus progesterone cotreatment on sensitivity of MCF-7 cells towards complexes **1** (A), **2** (B) or cisplatin (C) determined by clonogenic assay. MCF-7 cells were cotreated with estradiol (0.2 μ M) (\blacktriangle) or estradiol (0.2 μ M) plus progesterone (0.2 μ M) (\blacktriangledown) with platinum complex for 4 hr; (\square), no steroid hormone added. Values shown in the graphs are the means (\pm SEM) of six separate experiments, each conducted with 6 replicates; where error bars are not shown, they were smaller than the symbols.

Figure 1







Figure 3

