



**HAL**  
open science

## Ochratoxin A levels in plasma from healthy blood donors from Valencia and estimation of exposure degree: comparison with previous national Spanish data

Angel Medina-Vayá, Eva M. Mateo, Roberto Roig, Armando Blanquer, Misericordia Jiménez

### ► To cite this version:

Angel Medina-Vayá, Eva M. Mateo, Roberto Roig, Armando Blanquer, Misericordia Jiménez. Ochratoxin A levels in plasma from healthy blood donors from Valencia and estimation of exposure degree: comparison with previous national Spanish data. *Food Additives and Contaminants*, 2010, 27 (09), pp.1273-1284. 10.1080/19440049.2010.487876 . hal-00597455


**HAL Id: hal-00597455**

**<https://hal.science/hal-00597455>**

Submitted on 1 Jun 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Ochratoxin A levels in plasma from healthy blood donors from Valencia and estimation of exposure degree: comparison with previous national Spanish data**

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2010-036.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	19-Apr-2010
Complete List of Authors:	Medina-Vayá, Angel; University of Valencia., Microbiology and Ecology Mateo, Eva; University of Valencia, Dpt. Microbiology and Ecology Roig, Roberto; Centre de trasnfusion de la Generalitat Blanquer, Armando; Centre de trasnfusion de la Generalitat Jiménez, Misericordia; University of Valencia, Microbioloby and Ecology
Methods/Techniques:	Chromatography - HPLC, Statistical analysis
Additives/Contaminants:	Ochratoxin A, Mycotoxins – ochratoxin A
Food Types:	

SCHOLARONE™  
Manuscripts

1  
2  
3 **Ochratoxin A levels in plasma from healthy blood donors from**  
4 **Valencia and estimation of exposure degree: comparison with previous**  
5 **national Spanish data**  
6  
7  
8  
9

10  
11  
12  
13 Á. Medina<sup>1,2\*</sup>, E.M. Mateo<sup>1</sup>, R.J. Roig<sup>3</sup>, A. Blanquer<sup>3</sup>, M. Jiménez<sup>1</sup>  
14  
15

16  
17 <sup>1</sup> Departamento de Microbiología y Ecología. Universidad de Valencia. Dr. Moliner 50, E-  
18 46100 Burjassot, Valencia, Spain.  
19

20 <sup>2</sup> Applied Mycology Group, Cranfield Health, Vincent Building, Cranfield University. College  
21 Road, Cranfield, Bedford MK43 0AL, U.K.  
22

23 <sup>3</sup> Centre de Transfusió de la Comunitat Valenciana, Agència Valenciana de Salut. Conselleria  
24 de Sanitat, Generalitat Valenciana, Avda. del Cid, 65-acc. E-46014, Valencia, Spain  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

35 \* Corresponding author

36 Dr. Angel Medina-Vayá

37 Telf: +34 963543144

38 Fax: +34 96 3544570

39 E-mail: [angel.medina@uv.es](mailto:angel.medina@uv.es)  
40  
41  
42  
43  
44

45 **Running title:** OTA levels in Valencian blood donors

46 **Keywords:** Ochratoxin A; Dietary intake; Plasma; Human exposure; Valencian Community;  
47 Spain  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abstract**

Blood plasma levels of ochratoxin A, a toxic secondary metabolite of several fungal species belonging to the genera *Aspergillus* and *Penicillium*, were determined in 168 blood donors from Valencian Community (Spain), using LC-FLD. In conjunction with the blood collection, detailed information about the diet was obtained by using a questionnaire that included a wide range of products potentially contaminated with the toxin. The investigation revealed a detection frequency of 100%. Mean level was 1.09 µg OTA/l of plasma and concentrations ranged between 0.15 and 5.71 µg OTA/l of plasma. Men's levels were slightly higher than levels observed in women. Results were analysed by Spearman rank correlation test and Gamma correlation statistic. There was no strong correlation between individual consumption of 26 food groups, described as possibly contaminated with OTA and plasma level of OTA. Multiple regression models and factorial regression models were obtained explaining 14.9% and 64.4% respectively, but they could not explain the overall of the variability. The dietary daily intake of ochratoxin A was calculated on the basis of plasma toxin levels using Breitholtz and Klassen formulas. For the overall population, mean values were 1.47±1.25 and 2.16±1.88 ng/kg b.w/day respectively. All results have been compared with other Spanish data.

## Introduction

Ochratoxin A (OTA) is a toxic secondary metabolite of several fungal species belonging to the genera *Aspergillus* and *Penicillium* (van der Merwe et al., 1965; Pitt, 1987; Medina et al., 2005a).

OTA contaminates agricultural products and due to its long half life it accumulates in food, representing a serious threat to human and animal health throughout the world since it has been implicated in a diverse range of toxicological effects (O'Brien and Dietrich, 2005). OTA produces an inhibition of protein synthesis and lipid peroxidation by oxidative processes (Gautier et al., 2001; Omar et al., 1996). These mechanisms may generate nephrotoxic, neurotoxic and immunotoxic effects (Luhe et al., 2003; Schaaf et al., 2002; Álvarez et al., 2004). Teratogenic and carcinogenic effects have been described in some animal species (Veselá et al., 1983; Lioi et al., 2004). The IARC has classified OTA as a "possible human carcinogen (Group 2B)" based on sufficient evidence for carcinogenicity in experimental animal studies and inadequate evidence in humans (IARC, 1993). The toxic-kinetic parameters of OTA have not been well established in humans, but OTA showed high availability after oral administration in some animal studies (Fuchs and Hult, 1992; Omar et al., 1996).

OTA binds rapidly and with high affinity to plasma proteins with increased  $T_{1/2}$  values (Mally et al., 2005). The metabolism in humans is basically produced through cytochrome P-450 (Dortant et al., 2001) generating less toxic metabolites (Fazecas et al., 2005) which are eliminated in urine and faeces (Li et al., 2000, 1997). OTA can cross the placenta and is also excreted in animal milk (Valenta and Goll, 1996; Breitholtz-Emanuelsson et al., 1993).

As a consequence of the possible health hazards related to the ingestion of OTA, efforts to assess OTA levels in foods have been made by researchers worldwide and by governmental authorities in many countries. Several authors have described contamination of OTA in a wide range of products as cereal, beans, roasted coffee and cocoa, malt and beer, bread and bakery products, wines and grape juices, spices, poultry meat and kidney, pig kidneys and pork sausages (Van Egmond and Speijers, 1994; Jorgensen, 1998; Vrabcheva et al., 1998, Otteneder and Majerus 2000, 2001; Castellari et al., 2000; Thirumala-Devi et al., 2001; Thuvander et al., 2001; Visconti et

1  
2  
3 al., 2000; Leitner et al., 2002; Medina et al., 2005b, 2006; Mateo et al., 2007). It has also  
4  
5 been found in animal tissues, muscle, and organs, eggs (Marquardt and Frolich, 1992;  
6  
7 Valenta, H., 1998; Moreno Guillamont et al., 2005), cow's and human milk (Zimmerli  
8  
9 and Dick, 1995; Valenta, H., 1998; Breitholtz-Emanuelson et al., 1993).

10  
11 There are numerous reports from many countries in the world describing high  
12  
13 frequencies of human blood contamination by ochratoxin A, although at low levels,  
14  
15 showing widespread human exposure to this mycotoxin in Europe (Petkova-Bocharova  
16  
17 et al., 1988; Breitholtz et al., 1991; Goliński et al., 1991; Hald, 1991; Petkova-Bocharova  
18  
19 and Castegnaro, 1991; Creppy et al., 1993; Hadlok, 1993; Ruprich and Ostrý, 1993;  
20  
21 Breitholtz-Emanuelson et al., 1994; Kovács et al., 1995; Zimmerli and Dick, 1995; Solti  
22  
23 et al., 1997; Domijan et al., 1999; Palli et al., 1999; Peraica et al., 1999; Ozcelik et al.,  
24  
25 2001, Thuvander et al., 2001), in North America (Kuiper-Goodman et al., 1993; Scott et  
26  
27 al., 1998), in Japan (Ueno et al., 1998), in Africa (Kane et al., 1991; Singare-Tigori et al.,  
28  
29 2006) and in Asia (Assaf et al., 2004). Roughly, ochratoxin A levels were found to be at  
30  
31 the  $\mu\text{g/l}$  range, often with frequencies among population extending from 6.6% to 100%  
32  
33 (Marquardt and Frolich, 1992; Thuvander et al., 2001; Zimmerli and Dick, 1995) which  
34  
35 also depends on the level of the limit of detection of the method carried out for  
36  
37 analysis in sera or plasma samples.

38  
39 The evaluation of OTA levels in different types of populations is unquestionable  
40  
41 to provide data on exposure and evaluate the possible human health risk to OTA.

42  
43 Over the last 12 years four studies have been published analysing levels of OTA  
44  
45 in human blood. Burdaspal and Legarda and Jimenez et al. in 1998, Perez de Obanos et  
46  
47 al. in 2001 and recently Coronel et al. in 2009.

48  
49 Moreover the scarce of works done in Spain, generally, most of the studies  
50  
51 published regarding other countries just show the OTA incidence in human blood  
52  
53 serum or plasma but the relationship between the plasma levels of OTA, the estimated  
54  
55 dietary intake and the possible correlation with the consumption of different food and  
56  
57 beverages has been rarely studied.

58  
59 In the present paper, we developed i) an evaluation of the consumption of  
60  
foods and beverages considered as potentially contaminated with OTA in a wide  
number of healthy blood donors from the Valencian Community population  
(Mediterranean region) in Spain, ii) quantification of OTA plasma levels from these

1  
2  
3 healthy blood donors iii) a study of the potential correlation between the OTA levels in  
4 plasma and the diet, iv) the calculated dietary intake and v) a comparison with data  
5 obtained in previous studies made in different Spanish areas.  
6  
7  
8

## 9 **Materials and methods**

### 10 *Sampling and description of populations*

11  
12 Blood plasma samples were collected from healthy blood donors at the “Centro de  
13 Transfusiones de la Comunidad Valenciana” (CTCV) Valencian Community Transfusion  
14 Centre, Valencia, Spain, from 1st July to 14th of November 2008. The blood samples  
15 were taken in connection with the ordinary collection of blood, and all blood donors  
16 were informed and invited to participate in the survey after giving their informed  
17 consent. Ethylenediaminetetrachloroacetic acid (EDTA) was used as an anticoagulant.  
18 After centrifugation, the 3ml plasma samples were immediately frozen at -70°C.  
19 Different tests were made with the original plasma as routine in the CTCV to ensure  
20 absence of infectious agents. Once the results were obtained, samples which were  
21 found to be free of any infectious agents were then forwarded on dry ice to the  
22 Valencia University Fungi and Mycotoxin Research Laboratory, for OTA analysis.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

### 35 *Collection of individual food consumption data*

36  
37 In conjunction with the blood collection, detailed information about the diet of the  
38 healthy blood donors was obtained. The questionnaire included information about the  
39 consumption of a wide range of products such as meat (chicken, turkey, beef, pork, ...),  
40 beverages (beer, wine, fruit juices, coffee), cereal derived products (bakery products,  
41 pasta,...), spices, cheese, chocolate, etc. Additional information about the typical  
42 amount eaten of each item, consumption frequency or origin (supermarket, retails  
43 shops...) of all food items was collected. Furthermore some question regarding the  
44 age, sex, job, consumption of diet and bio-products, meals in restaurants... were done  
45 to all the healthy blood donor subjects that agreed to participate in the study.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

### 56 *Standards and reagents*

57  
58 The ochratoxin A standard was purchased from Sigma (Sigma-Aldrich, Alcobendas,  
59 Spain). A stock solution of about 500 mg/l was prepared by solving 1mg of OTA in 2ml  
60

1  
2  
3 of toluene–acetic acid (99:1, v/v). A series of working standards was prepared by  
4 evaporation of known aliquots of the stock solution and dissolution in filtered LC  
5 mobile phase. They were used to calibrate the detector response.  
6  
7

8  
9 The concentration of the stock solution was determined by measuring absorbance at  
10 333 nm of a diluted solution (20-30 mg/l) of OTA in toluene-acetic acid (99:1, v/v).  
11 Acetonitrile, chloroform, and methanol (all LC grade), acetic acid (A.R.) and phosphoric  
12 acid (85%, A.R.) were from J.T. Baker (Deventer, the Netherlands). Sodium chloride and  
13 dimethyl sulfoxide were purchased from Sigma (Sigma-Aldrich, Alcobendas, Spain).  
14 Pure water was obtained from a Milli-Q apparatus (Millipore, Billerica, MA, USA) and  
15 was used when water was required.  
16  
17  
18  
19  
20  
21  
22  
23

#### 24 *Extraction, clean-up and analysis of ochratoxin A*

25  
26 Extraction was made following the protocol used by Lino et al. (2008) with some  
27 modifications. Briefly, two mL of plasma were added of 1 mL of a solution containing  
28 33.7 mL of 85% orthophosphoric acid and 118 g sodium chloride per liter, and mixed  
29 for 1 min. After, 2 x 4 mL of chloroform were added and mixed for 20 min using a  
30 rotary mixer. The organic extracts were reunified and concentrated to dryness at 45°C  
31 under gentle stream of nitrogen. The residue was dissolved in 0.25ml of water:  
32 dimethyl sulfoxide (70:30, v/v) and transferred to an amber glass vial. One hundred µl  
33 were injected into the LC system.  
34  
35  
36  
37  
38  
39  
40

41 The LC system consisted of a Waters 600E system controller, a Millipore Waters 717  
42 Plus autosampler and a Waters 470 scanning fluorescence detector (Waters, Milford,  
43 Ma, USA). Excitation and emission wavelengths were 330 and 460 nm, respectively.  
44 The samples were separated using a C<sub>18</sub> Zorbax Eclipse Plus<sup>®</sup> (150 x 4.6 mm, 3.5µm)  
45 (Agilent Technologies, Waldbronn, Germany), with a guard column of the same  
46 material. Run time for samples was 20 min with OTA being detected at about 12 min.  
47 The flow rate of the mobile phase (water-acetonitrile-acetic acid; 59:39:2, v/v/v) was 1  
48 mL min<sup>-1</sup>.  
49  
50  
51  
52  
53  
54  
55

56 Analysis of the results was carried out on a computer with Millennium<sup>®</sup> 4.0 software  
57 (Waters).  
58

59 Calibration lines were constructed using blood plasma spiked with different amounts  
60 of OTA standard in the range 0.05-10 µg OTA/l of medium. Since OTA free samples

were not found, calibration lines were constructed with the standard addition method using a 50ml sample supplied by CTCV. The interval trend line obtained fitted perfectly with an  $R^2=0.997$ . For each point three replicates were made.

#### *Method performance*

The limit of detection (LOD) was calculated using the IUPAC formula (Long and Winefordner, 1983):

$$LOD = \bar{X}_n + 3 \cdot S_{n-1}$$

Where,  $\bar{X}$  is the concentration mean and  $S_n$  the standard deviation. The limit of quantification (LOQ) was calculated according the next equation:

$$LOQ = \bar{X}_n + 10 \cdot S_{n-1}$$

The results have shown a LOD of 0.01  $\mu\text{g}$  OTA/l blood plasma, and a LOQ 0.07  $\mu\text{g}$  OTA/l. All these determinations were carried out in triplicate.

The OTA recovery rate from blood plasma was calculated (using another 50 ml sample as in the previous section) by addition of known amounts of OTA in the range of 0.1 to 5  $\mu\text{g}$  OTA/l obtaining a mean recovery of  $81.6 \pm 10.9$  %. All the results presented have been corrected according to this value.

Repeatability was tested in the same day on five replicates at the spiking level of 0.5  $\mu\text{g}$  OTA/l, and presented a relative standard deviation of 11%. Reproducibility was assessed in three different days also at 0.5  $\mu\text{g}$  OTA/l, and the relative standard deviation was 10%.

#### *LC-MS/MS confirmation of OTA in plasma samples*

Once the LC-FLD data were collected, medium-high concentration samples were selected and the extract was dried in low volume chromatograph vials, using a gentle stream of  $\text{N}_2$  at room temperature. Once they were near to be dry, 75  $\mu\text{L}$  of methanol: water (1:1 v/v) were added and 50  $\mu\text{L}$  were injected in the LC-ESI-MS/MS system.

LC-triple quadrupole mass spectrometry (LC-MS/MS) was used to confirm the presence of OTA in plasma samples. Transition between 403.8  $m/z$  ion and 358.2  $m/z$  ion was investigated. OTA was detected in all concentrated samples containing more than 15

1  
2  
3 ng OTA/ml (LOD) or above, confirming the results obtained previously by LC-FLD for  
4 samples containing more than 0.40-0.45 µg OTA/l of blood.  
5

6  
7 The analysis was carried out on a Shimadzu LC-10AD (Shimadzu Corporation, Kyoto,  
8 Japan), and an automatic injector Shimadzu SIL-10AS equipped with a C<sub>18</sub> Teknokroma  
9 Mediterranea Sea18 (50 mm x 2.1 mm, 5 µm) (Teknokroma, Sant Cugat del Valles,  
10 Spain) at room temperature and coupled to a Micromass Quattro™ triple-quadrupole  
11 mass spectrometer (Waters, Beverly, MA, USA). The mobile phase was programmed  
12 following a linear gradient at a flow rate of 0.3 mL/min. Solvent A was water with 0.5%  
13 formic acid, and solvent B was methanol with 0.5% formic acid. The gradient program  
14 was as follows: 0-5 min, 60% A; 2 min, 10% A; 8 min, 2 % A; 0.5 min, 60% A. Injection  
15 volume was 20 µL.  
16

17  
18 Micromass Quattro™ triple-quadrupole mass spectrometer was equipped with a Z-  
19 spray electrospray ionization source. Electrospray ionization method was performed  
20 using the following ionization source parameters: Capillary: 3.5 KV, skimmer: 2 V, RF  
21 Lens: 0.5 V, source temperature: 120 °C, desolvation temperature: 150 °C. MS1  
22 parameters were LM resolution, 13; HM resolution, 13; ion energy, 1. MS2 parameters  
23 were LM resolution, 13; HM resolution, 13; ion energy, 1; multiplier 650 V.  
24

25  
26 Spectra were obtained in positive ionization multiple reaction monitoring (MRM)  
27 mode.  
28

29  
30 Confirmation by ESI-MS/MS was based on the transition between parent ion [M + H]<sup>+</sup>  
31 and the daughter ion [(M + H)<sup>+</sup> - HCOOH], whose *m/z* ratios are 403.8 and 358.2,  
32 respectively.  
33

#### 34 *Calculation of ochratoxin A intake based on plasma levels*

35  
36 The dietary daily intake of ochratoxin A was calculated on the basis of plasma toxin  
37 levels using the equation  $k_0 = Cl_p \cdot C_p / A$ , where  $k_0$  is the dietary intake (ng/kg body weight  
38 and day) (Breitholtz et al. 1991).  $Cl_p$  is the plasma clearance (mL/kg body weight and  
39 day),  $C_p$  is the plasma concentration of ochratoxin A (ng/mL), and  $A$  is the  
40 bioavailability of the toxin.  $Cl_p$  and  $A$  were estimated at 0.67 mL/kg body weight/day  
41 and 0.5 (i.e. 50%), respectively, by Breitholtz et al. (1991), and these figures were used  
42 in the present calculations.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Moreover, the dietary daily intake was calculated according to the Klassen equation  
4 (Directorate General-Health and consumer protection, 2002; Thuvander et al., 2004).  
5  
6 In these later studies  $Cl_p$  was estimated in 0.485 mL/kg body weight/day; a value lesser  
7  
8 than the obtained by Breitholtz et al. (1991).  
9

### 10 11 12 *Statistical analysis*

13  
14 Statistical analysis was performed using the package STATISTICA 8 (StatSoft® Inc.,  
15  
16 2007. Tulsa, OK, USA).  
17

18  
19 Data on ochratoxin A in plasma/BMI were tested for normality using standardized  
20  
21 skewness and standardized Kurtosis tests. In both cases values were not within the  
22  
23 range expected for data from a normal distribution. The correlation between plasma  
24  
25 levels of the toxin and food consumption was investigated, due to non-normality, using  
26  
27 the Spearman rank correlation test and *Gamma* statistic. Spearman R is useful  
28  
29 assessing whether the two quantities we measure actually are varying together as  
30  
31 predicted by our hypothesis, or whether any apparent correlation produced is just due  
32  
33 to the amount of variation in results we could expect from chance circumstance. In  
34  
35 short, *Gamma* is a probability; specifically, it is computed as the difference between  
36  
37 the probability that the rank ordering of the two variables agree minus the probability  
38  
39 that they disagree, divided by 1 minus the probability of ties.

40  
41 STATISTICA 8 General Regression Module (GRM) was also used to build mathematical  
42  
43 models to extract additional information from the data. This module allowed us to  
44  
45 build models for designs with single-degree-of-freedom effects for continuous  
46  
47 predictor variables using Sigma restricted parameterization. GRM implements stepwise  
48  
49 and best-subset model-building techniques and uses the least squares methods of the  
50  
51 general linear model to build models and to estimate and test hypotheses about  
52  
53 effects included in the final model.

54  
55 Multiple regression models, that do not study the effects between the different  
56  
57 predictors, and factorial models, analyzing the above effects, were built.

## 58 **Results**

### 59 *Dietary intake questionnaire*

60

1  
2  
3 Meat product consumption, cereal and derived cereal consumption; beverages and  
4 other food items are shown in tables 1 to 4, respectively.

5  
6  
7 The main differences between men and woman food consumption were found on the  
8 breakfast cereals, beer and wine.

9  
10 Women, among all age ranges, use to eat more breakfast cereals than men. In the case  
11 of the alcoholic beverages (wine and beer) the frequency of consumption is higher in  
12 men but also the average quantity is approximately 2-fold bigger in men group.

#### 13 14 15 16 17 18 *Ochratoxin A levels in plasma samples*

19  
20 A total of 168 samples of plasma were analyzed under the described conditions and  
21 OTA was detected in all of them at concentrations ranging between 0.15 and 5.71  $\mu\text{g}$   
22 OTA/l of plasma. Mean level was 1.09  $\mu\text{g}$  OTA/l of plasma. Table 5 shows the Body  
23 Mass Index (B.M.I.), OTA level ( $\mu\text{g}/\text{l}$  plasma), Standard Deviation (S.D.) and the OTA  
24 ranges in the blood samples from the populations of men and women studied. In order  
25 to simplify the interpretation, data have been split in 3 different groups according to  
26 the age (18-35, 36-45 and over 45 years old). As it is shown in Table 5, level of OTA in  
27 men's blood plasma was slightly higher than levels observed in woman's blood plasma.  
28 Due to the non-normality of OTA level, data normalization was made by calculating the  
29 logarithm and then ANOVA was performed in the transformed variable using sex as  
30 categorical factor. No significant statistical differences were found in log(OTA blood  
31 levels) ( $P=0.216$ ) between men and women populations.

#### 32 33 34 35 36 37 38 39 40 41 42 43 44 45 *Evaluation of daily intake*

46 The dietary daily intake of ochratoxin A was calculated on the basis of plasma toxin  
47 levels using Breitholtz and Klassen formulas. For the overall population, mean values  
48 were  $1.47 \pm 1.25$  and  $2.16 \pm 1.88$  ng/kg b.w/day respectively.

49 Splitting the cases by sex and using both formula, Breitholtz and Klassen, the results  
50 were  $1.52 \pm 1.29$  and  $2.24 \pm 1.90$  ng/kg b.w/day for men and  $1.31 \pm 1.23$  and  $1.93 \pm 1.81$   
51 ng/kg b.w/ day for women.

52 Performing calculations with the Breitholtz's formula, 3.0% of cases showed values  
53 higher than 5 ng/kg b.w/day. With Klassen formula, 6.7% of cases showed higher  
54 values.

1  
2  
3  
4  
5 *Correlation between consumption of individual foods and plasma levels of ochratoxin*

6  
7 A.

8  
9 Correlations between individual consumption of each 26 food groups described as  
10 possibly contaminated with OTA and plasma level of OTA/BMI, and some data about  
11 habits, were examined. This transformation was done in order to relate the amount of  
12 OTA in plasma with the possible total volume of blood that is between 2800 and 3800  
13 ml/m<sup>2</sup> body surface.  
14

15  
16 Results for OTA/BMI variable analyzed by Spearman rank correlation test and *Gamma*  
17 correlation statistic are shown in Table 6.  
18

19  
20 In the overall population, weak correlations were seen between plasma levels of OTA  
21 and beef, turkey, cold meat and bread consumptions. Results were the same in both  
22 tests performed.  
23

24  
25 Also a weak correlation was found between the plasma levels of OTA and people that  
26 use to have meals outside home.  
27

28  
29 Regarding men population, Spearman rank correlation test showed weak correlations  
30 with turkey, cold meat and bread consumptions. In this case, beef consumption wasn't  
31 statistically significant and neither outside home meals. Results of *Gamma* correlation  
32 statistic, that is preferable to Spearman *R* when the data contain many tied  
33 observations, were slightly different in this case and showed the same results obtained  
34 for the overall population.  
35

36  
37 Study of women population showed different results than the obtained for men. In  
38 these case Spearman rank correlation test and *Gamma* correlation statistic showed  
39 weak correlations between plasma levels of OTA and wine and cheese consumption.  
40

41  
42 Multiple regression models were obtained to extract more information about the data.  
43 As first step multiple regression methods were created using *Backward stepwise*  
44 model building (P to enter: .1; P to remove: .2). Using that methodology, main factors  
45 affecting the variability of OTA/BMI were found to be: wine, beer, bread and turkey  
46 factors, and also bun, fruit juice and Italian pasta consumption, despite they had less  
47 influence.  
48

49  
50 With this first approximation and building the whole model, the adjusted R<sup>2</sup>  
51 (determination factor) was 0.1493. Therefore, only 14.9% of the total variation in OTA  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 level/BMI can be explained by the linear relationship between the factors described  
4 above.  
5

6  
7 Next step was the building of factorial regression models. In this case, interaction  
8 between factors was considered. Five different set of 20% of the raw data were  
9 obtained randomly and were used for cross-validation. When models were built these  
10 5 new variables were used to perform the model validation.  
11

12  
13 Models built included the main affecting factors obtained in multiple regressions.  
14 Other factors were also included and tested using *Backward stepwise* model building  
15 changing enter and remove condition in order trying to improve the model fitting.  
16

17  
18 Finally a third order, cross validated, whole model was built with factors: overall meat  
19 consumption, bread, bun, Italian pasta, beer, wine, coffee, fruit juices and dried fruit  
20 consumptions.  
21

22  
23 The adjusted  $R^2$  obtained in this case was 0.6438, meaning that from the overall  
24 variation in plasma OTA level/BMI, only 64.4% of variability could be explained using  
25 the model built with the above factors. The other 35.6% of the total variation in OTA  
26 level/BMI remained unexplained.  
27  
28  
29  
30  
31  
32  
33

### 34 35 Discussion

36  
37 In this work, original results regarding consumption of foods considered as ochratoxin  
38 A vectors in the Valencian Community, wide area in the east of Spain where the  
39 Mediterranean diet is common, are described. Moreover, the OTA levels of a large  
40 sample of healthy individuals from this population are shown.  
41

42  
43 Incidence of OTA in plasma has been reported in subjects from different geographical  
44 parts of Spain, but the dietary habits and climate in many regions are very different.  
45

46  
47 The mean level we found in the Valencian Community (1.09  $\mu\text{g}$  OTA/l) is lower than  
48 levels found in Madrid (1.19 ng/mL) (Burdaspal and Legarda, 1998) and higher than  
49 levels found in Lleida (0.86  $\mu\text{g}$  OTA/l) (Coronel et al., 2009), in Navarra (0.71 $\mu\text{g}$  OTA/l)  
50 (Jimenez et al., 1998) and Granada (0.63 ng/mL) (Pérez de Obanos et al., 2001).  
51

52  
53 These differences could be well explained attending to diet differences between these  
54 areas. However it is impossible to compare and discuss it because diet parameters of  
55 the analysed subjects were not studied in the other studies carried out in Spain or data  
56 used have been extracted from literature (Coronel et al., 2009).  
57  
58  
59  
60

1  
2  
3 Studies carried out in other countries as Portugal, Chile and Croatia also have shown  
4 differences in the OTA levels of populations belonging to different areas in the same  
5 country (Peraica et al., 2001; Muñoz et al., 2006; Lino et al., 2008).  
6  
7

8  
9 Comparing our results with others from different countries, the obtained levels in  
10 plasma samples of healthy blood donors from the Valencian Community are similar to  
11 samples from Canada, United Kingdom (0.88 and 1.09 respectively), lower than values  
12 reported in Tunisia and Denmark (Hald, 1991; Hassen et al., 2004 ) and higher than  
13 levels reported in other countries as Italy, Morocco, Lebanon, Croatia, Sweden, Poland  
14 and Portugal (Assaf et al., 2004; Brera et al., 1999; Fillali et al., 2002; Hald, 1991; Lino  
15 et al., 2008; Peraica et al., 1999, 2001; Thuvander et al., 2001).  
16  
17

18  
19 Also we found a higher frequency of positive samples (100%) in comparison with other  
20 studies carried out in Spain. Jimenez et al. (1998) found 53% of positives. Pérez de  
21 Obanos et al. (2001) found 72% and recently Coronel et al., (2009) have found 98.6%  
22 of positive samples. Our results agree with percentage found by Burdaspal and  
23 Legarda (1998) that found 100% of positives. These differences could be explained  
24 attending to the different analytical method limits of detection and quantification in  
25 the studies. These authors reported a limit of detection of 0.075 µg OTA/l (Coronel et  
26 al., 2009), 0.21 µg OTA/l (Pérez de Obanos et al., 2001) and 0.52 µg OTA/l (Jimenez et  
27 al., 1998), whereas the limit of detection of Burdaspal and Legarda (1998) was 0.020  
28 µg OTA/l and in this study was 0.010 µg OTA/l of blood plasma.  
29  
30

31  
32 In the literature, frequencies between 6.6% and 100% of samples with presence of  
33 OTA have been reported and explanations for these differences have been found in  
34 the different LOD and LOQ as well (Lino et al., 2008; Jimenez et al., 1998; Thuvander et  
35 al., 2001).  
36  
37

38  
39 Lack of differences between OTA plasma levels in men and women agrees with results  
40 found in Lleida by Coronel et al. (2009), and with other studies in other countries  
41 (Breitholtz et al., 1991; Thuvander et al., 2001; Ueno et al., 1998). Even though other  
42 authors described significant differences in OTA blood plasma levels regarding sex  
43 (Muñoz et al., 2006).  
44  
45

46  
47 We did not find differences between age groups and these results do not agree with  
48 those obtained by Coronel et al. (2009) in Spain and in other countries as Portugal  
49 (Lino et al., 2008). Our results agree with the study carried out by Jimenez et al. (1998)  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 in Navarra (Spain) and other studies carried out overseas (Ueno et al., 1998; Palli et al.,  
4 1999; Gareis et al., 2000; Grosso et al., 2003; Skaug, 2003; Assaf et al., 2004) that  
5 showed no differences among age groups. Although these results should be  
6 interpreted with care due to in each case the age groups were defined differently.  
7

8  
9  
10 The dietary daily intake was calculated from plasma levels using two different  
11 formulas. Basically the calculation is the same but the figure of the plasma clearance  
12 ( $Cl_p$ ) is lesser using the Klassen approach. This leads to higher calculated ochratoxin  
13 daily intakes. Despite the fact the means obtained with both formulas ( $1.47 \pm 1.25$  and  
14  $2.16 \pm 1.88$  ng/kg b.w/ day, respectively) were below the tolerable daily intake, 5 ng/kg  
15 b.w, estimated by the Scientific Committee on Foods (SCF) of the European Union in  
16 2002, individual values higher than the limits have been found. In the case of  
17 Breitholtz's formula, 3.0% of cases showed higher values and with Klassen formula,  
18 6.7%. That high levels found may cause concern, once OTA is classified as possible  
19 carcinogen for human (group 2B) by International Agency for Research on Cancer  
20 (IARC, 1993).  
21

22  
23 Although the study involved a high number of subjects and a high number of food  
24 items that often are contaminated with ochratoxin A, the relation between the OTA  
25 plasma levels and these food items were weak.  
26

27  
28 Correlations tests performed did not show stronger correlation between the OTA level  
29 in plasma and some variables, despite of some of them resulted statistically significant,  
30 the correlation coefficients or probabilities were very small. Clear differences were  
31 observed between the men and women groups. These differences show that maybe  
32 each individual or group of individuals has different main sources of OTA.  
33

34  
35 Investigation on possible effects between variables was done building regression  
36 models and in this case some variables and their combinations explained around the  
37 60% of the variability among the data.  
38

39  
40 Models could not explain the overall of the variability and when some samples were  
41 removed from the model in order to perform cross validation samples some of them  
42 were not well predicted by the model, despite of most of them were well predicted.  
43 These results mean that some variability sources are being ignored in the model. Also  
44 it should be taken into account the different sources of uncertainty that this kind of  
45 studies has. Apart from the analytical uncertainty typical in all mycotoxin analysis it is  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

possible that there are some mistakes in data from the questionnaires, possible bias during the sampling as only volunteer people was included and because there are seasonal differences regarding the food consumption; we did not consider other sources of ochratoxin A, for example via airborne particles and spores from ochratoxigenic fungi (Brera et al., 2002).

### Acknowledgements

The authors wish to thank financial support from FEDER and Spanish Government "Ministerio de Ciencia e Innovación" (Project AGL2007-66416-C05-01/ALI, and two research grants) and the Generalitat Valenciana (Project ACOMP2009/371). A. Medina wants to acknowledge funding from the Spanish Ministry of Science and Innovation (Programa Nacional de Movilidad de Recursos Humanos del Plan nacional de I+D+I 2008-2011). The authors are also indebted to Ms Silvia Sarrión, Mara Argente, Carmina Mas, M. Angeles Arnau and Dr. Vicente Mirabet for their helpful collaboration concerning blood sampling and data collection from donors, and with F.M. Valle-Algarra for his help with LC-MS/MS analysis.

### References

- Álvarez L, Gil AG, Ezpeleta O, García Jalón JA, López de Cerain A. 2004. Immunotoxic effects of Ochratoxin A in wistar rats after oral administration. *Food Chem Toxicol* 42: 825-834.
- Assaf H, Betbeder A-M, Creppy EE, Pallardy M, Azouri H. 2004. Ochratoxin A levels in human plasma and foods in Lebanon. *Hum Exper Toxicol* 23: 495-501.
- Brera C, Caputi R, Miraglia M, Iavicoli I, Salerno A, Carelli G. 2002. Exposure assessment to mycotoxins in workplaces: aflatoxins and ochratoxin A occurrence in airborne dusts and human sera. *Microchem J* 73: 167-173.
- Breitholtz A, Olsen M, Dahlback A, Hult K. 1991. Plasma ochratoxin A levels in three Swedish populations surveyed using an ion-pair HPLC technique. *Food Addit Contam* 8: 183-192.
- Breitholtz-Emanuelsson A, Olsen M, Oskarsson A, Palminger I, Hult K. 1993. Ochratoxin A in cow's milk and in human milk with corresponding human blood samples. *J AOAC Int* 76: 842-846.
- Breitholz-Emmanuelson A, Minervini F, Hult K, Visconti A. 1994. Ochratoxin A in human serum samples collected in southern Italy from healthy individuals and individuals suffering from different kidneys disorders. *Natural Toxins* 2: 366-370.
- Burdaspal PA, Legarda TM. 1998. Datos sobre presencia de ocratoxina A en plasma humano en España. *Alimentaria* 292, 103-109.

- 1  
2  
3 Castellari M, Fabbri S, Fabiani A, Amati A, Galassi S. 2000. Comparison of different  
4 immunoaffinity clean-up procedures for high-performance liquid chromatographic  
5 analysis of ochratoxin A in wines. *J Chromatogr A* 888: 129-136.
- 6  
7 Coronel MB, Sanchis V, Ramos AJ, Marin S. 2009. Assessment of the exposure to ochratoxin A  
8 in the province of Lleida, Spain. *Food Chem Toxicol* 47: 2847-2852
- 9  
10 Creppy E, Castegnaro M, Grosse Y, Meriaux J, Manier C, Moncharmont P, Waller C. 1993. Etude  
11 de l'ochratoxicose humaine dans trois regions de France : Alsace, Aquitaine, et region  
12 Rhone-Alpes. In: Creppy, E., Castegnaro, M., Dirheimer, G. (Eds.), *Human Ochratoxicosis*  
13 *and its Pathologies*, INSERM vol 231. John Libbey Eurotext Ltd, Paris, pp. 147-158.
- 14  
15 Domijan AM, Peraica M, Fuchs R, Lucia A, Radic B, Balija M, Bosanac I, Gricevic D. 1999.  
16 Ochratoxin A in blood of healthy population in Zagreb. *Archives of Industrial Hygiene*  
17 *and Toxicology* 50: 263-271.
- 18  
19 Dortant PM, Peters-Volleberg GWM, Van Loveren H, Marquardt RR, Speijers GJA. 2001. Age-  
20 related differences in the toxicity of Ochratoxin A in female rats. *Food Chem Toxicol* 39:  
21 55-65.
- 22  
23 European Commission, 2002. Assessment of dietary intake of Ochratoxin A by the population  
24 of European Union Members states. Directorate General-Health and consumer  
25 protection. Report on tasks for scientific cooperation. Report of experts participating in  
26 Task 3.2.7, pp. 18-19.
- 27  
28 Fazecas B, Tar A, Kovacs M. 2005. Ochratoxin a content of urine samples of healthy humans in  
29 Hungary. *Acta Vet Hung* 53: 35-44.
- 30  
31 Fuchs R, Hult K. 1992. Ochratoxin A in blood and its pharmacokinetic properties. *Food Chem*  
32 *Toxicol* 30: 201-204.
- 33  
34 Gautier JC, Holzhaeuser D, Markovic J, Gremaud E, Schilter B, Turesky RJ. 2001. Oxidative  
35 damage and stress response from Ochratoxin A exposure in rats. *Free Radical Biol Med*  
36 30 (10): 1089-1098.
- 37  
38 Gareis M, Rosner H, Ehrhardt S. 2000. Blood serum levels of ochratoxin A and nutrition habits.  
39 *Arch Lebensmittelhyg* 51: 108-110.
- 40  
41 Goliński P, Grabatkiewicz-Szczena J, Chelkowski J, Hult K, Kostecki M. 1991. Possible sources of  
42 ochratoxin A in human blood in Poland. In: Castegnaro M, Plestina R, Dirheimer G,  
43 Chernozemsky IN, Bartsch H. (Eds.), *Mycotoxins, Endemic Nephropathy and Urinary*  
44 *Tract Tumours*. IARC scientific publications 115, Lyon, pp. 153-157.
- 45  
46 Grosso F, Said S, Mabrouk I, Fremy JM, Castegnaro M, Jemmali M, Dragacci S. 2003. New data  
47 on the occurrence of ochratoxin A in human sera from patients affected or not by renal  
48 diseases in Tunisia. *Food Chem Toxicol* 41: 1133-1140.
- 49  
50 Hadlok RM. 1993. Human ochratoxicosis in Germany, updating 1993. In: Creppy EE, Castegnaro  
51 M, Dirheimer G. (Eds.), *Human Ochratoxicosis and its Pathologies*, INSERM vol 231. John  
52 Libbey Eurotext Ltd, Paris, pp. 141-145.
- 53  
54 Hald B, 1991. Ochratoxin A in human blood in European countries. In: Castegnaro M, Plestina  
55 R, Dirheimer G, Chernozemsky IN, Bartsch H. (Eds.), *Mycotoxins, Endemic Nephropathy*  
56 *and Urinary Tract Tumours*. IARC scientific publications 115, Lyon, pp. 159-164.
- 57  
58 IARC, 1993. Ochratoxin A. Some naturally occurring substances: food items and constituents,  
59 heterocyclic aromatic amines and mycotoxins. IARC Monograph on the Evaluation of  
60 Carcinogenic Risk to Humans. Lyon, vol. 56, pp. 489-521.

- 1  
2  
3 Jimenez AM, Cerain AL, González-Penas E, Bello J, Betbeder AM, Creppy EE. 1998. Exposure to  
4 ochratoxin A in Europe: comparison with a region of northern Spain. *J Toxicol-Toxin*  
5 *Reviews* 17: 479-491.  
6  
7 Jorgensen K. 1998. Survey of pork, poultry, coffee, beer and pulses for ochratoxin A. *Food*  
8 *Additives and Contaminants* 15: 550-554.  
9  
10 Kane A, Diop N, Diack TS. 1991. Natural occurrence of ochratoxin A in food and feed in  
11 Senegal. *IARC Monographs on mycotoxins, endemic nephropathy and urinary tract*  
12 *tumours*. IARC, Lyon, 93-96.  
13  
14 Kovács F, Sándor G, Ványi A, Domány S, Zomborszky-Kovács M. 1995. Detection of ochratoxin  
15 A in human blood and colostrum. *Acta Veterinaria Hungarica* 43: 393-400.  
16  
17 Kuiper-Goodman T, Ominski K, Marquardt RR, Malcom S, McMullen E, Lombaert GA, Morton T.  
18 1993. Estimating human exposure to ochratoxin A in Canada. In: Creppy E, Castegnaro  
19 M, Dirheimer G. (Eds.), *Human Ochratoxicosis and its Pathologies*, INSERM vol 231. John  
20 Libbey Eurotext Ltd, Paris, pp. 167-174.  
21  
22 Leitner A, Zoller P, Paolillo A, Stroka J, Papadopoulou-Bouraoui A, Jaborek Anklam E, Lindner  
23 W. 2002. Comparison of methods for the determination of ochratoxin A in wine. *Anal.*  
24 *Chim. Acta* 453: 33-41.  
25  
26 Li S, Marquardt RR, Frohlich AA. 2000. Identification of ochratoxins and some of their  
27 metabolites in bile and urine of rats. *Food Chem Toxicol* 38: 141-152.  
28  
29 Li S, Marquardt RR, Frohlich AA, Vitti TG, Crow G. 1997. Pharmacokinetics of Ochratoxin A and  
30 its metabolites in rats. *Toxicol Appl Pharmacol* 145: 82-90.  
31  
32 Lino CM, Baeta ML, Henri M, Dinis AMP, Pena AS, Silveira MIN. 2008. Levels of ochratoxin A in  
33 serum from urban and rural Portuguese populations and estimation of exposure degree.  
34 *Food Chemical Toxicol* 46: 879-885.  
35  
36 Lioi MB, Santoro A, Barbieri R, Salzano S, Ursini MV. 2004. Ochratoxin A and zearalenone: a  
37 comparative study on genotoxic effects and cell death induced in bovine lymphocytes.  
38 *Mutat Res* 557: 19-27.  
39  
40 Long GL and Winefordner JD. 1983. Limit of detection: a closer look at the IUPAC definition.  
41 *Anal. Chem.* 55, 713A.  
42  
43 Luhe A, Hildebrand H, Bach U. 2003. A new approach to studying ochratoxin A (OTA)-induced  
44 nephrotoxicity: expression profiling in vivo and in vitro employing cDNA microarrays.  
45 *Toxicol. Sci.* 73 (2): 315-328.  
46  
47 Mally A, Keim-Heusler H, Amberg A, Kurz M, Zepnik H, Mantle P, Völkel W, Hard G, Dekant W,  
48 2005. Biotransformation and nephrotoxicity of ochratoxin B in rats. *Toxicol Appl*  
49 *Pharmacol* 206: 43-53.  
50  
51 Marquardt RR, Frolich AA. 1992. A review of recent advances in understanding ochratoxicosis.  
52 *J. Anim. Sci.* 70, 3968-3988.  
53  
54 Mateo R, Medina Á, Mateo EM, Mateo F, Jiménez M. 2007. An overview of ochratoxin A in  
55 beer and wine. *Int J Food Microbiol* 119: 79-83.  
56  
57 Medina A, Jiménez M, Gimeno-Adelantado JV, Valle-Algarra FM, Mateo. 2005b. Determination  
58 of ochratoxin A in beer marketed in Spain by liquid chromatography with fluorescence  
59 detection using lead hydroxyacetate as a clean-up agent. *J Chromatogr A* 1083: 7-13.  
60  
61 Medina A, Mateo R, López-Ocaña L, Valle-Algarra FM, Jiménez M. 2005a. Study of Spanish  
62 grape mycobiota and ochratoxin A production by isolates of *Aspergillus tubingensis* and  
63 other members of *Aspergillus* Section *Nigri*. *Appl Environ Microbiol* 71: 4696-4702.

- 1  
2  
3 Medina A, Valle-Algarra FM, Gimeno-Adelantado JV, Mateo R, Mateo F, Jiménez M. 2006. New  
4 method for determination of ochratoxin A in beer using zinc acetate and solid-phase  
5 extraction silica cartridges. *J Chromatography A* 1121: 178-183.
- 6  
7 Moreno Guillamont E, Lino CM, Baeta ML, Pena AS, Silveira MIN, V Mañes J. 2005. A  
8 comparative study of extraction apparatus in HPLC analysis of ochratoxin A in muscle.  
9 *Anal Bioanal Chem* 383: 570-575.
- 10  
11 Muñoz K, Vega M, Rios G, Muñoz S, Madariaga R. 2006. Preliminary study of Ochratoxin A in  
12 human plasma in agricultural zones of Chile and its relation to food consumption. *Food*  
13 *Chem Toxicol* 44: 1884–1889.
- 14  
15 O'Brien E and Dietrich DR. 2005. Ochratoxin A: The Continuing Enigma. *Crit Rev Toxicol* 35: 33-  
16 60.
- 17  
18 Omar RF, Gelboin HV, Rahimtula AD. 1996. Effect of cytochrome P450 induction on the  
19 metabolism and toxicity of ochratoxin A. *Biochem Pharmacol* 51: 207–216.
- 20  
21 Ottener H, Majerus P. 2000. Occurrence of ochratoxin A in wines: influence of the type of  
22 wine and its geographical origin. *Food Addit Contam* 17: 793–798.
- 23  
24 Ottener H, Majerus P. 2001. Ochratoxin A in coffee: nation-wide evaluation of data  
25 collected by German Food Control (1995–1999). *Food Addit Contam* 18: 431–435.
- 26  
27 Ózcelik N, Kosar A, Soysal D. 2001. Ochratoxin A in human serum samples collected in Isparta-  
28 Turkey from healthy individuals and individuals suffering from different urinary  
29 disorders. *Toxicology Letter* 121: 9–13.
- 30  
31 Palli D, Miraglia M, Saieva C, Masala G, Cava E, Colatosti M, Corsi AM, Russo A, Brera C. 1999.  
32 Serum levels of ochratoxin A in healthy adults in Tuscany: correlation with individual  
33 characteristics and between repeat measurements. *Cancer Epidemiology Biomarkers*  
34 *and Prevention* 8: 265–269.
- 35  
36 Peraica M, Domijan AM, Fuchs R, Lucia A, Radia B. 1999. The occurrence of ochratoxin A in  
37 blood in general population of Croatia. *Toxicology Letter* 110: 105–112.
- 38  
39 Pérez de Obanos A, López de Ceraín A, Jiménez AM, González E, Bello J. 2001. Ochratoxin A in  
40 human plasma: new data of exposition in Spain. *Rev Toxicol* 18: 19–23.
- 41  
42 Petkova-Bocharova T, Castegnaro M. 1991. Ochratoxin A in human blood in relation to  
43 endemic nephropathy and urinary tract tumours in Bulgaria. In: Castegnaro M, Plestina  
44 R, Dirheimer G, Chernozemsky IN, Bartsch H. (Eds.), *Mycotoxins, Endemic Nephropathy*  
45 *and Urinary Tract Tumours*. IARC scientific publications 115, Lyon, pp. 135–137.
- 46  
47 Petkova-Bocharova T, Chernozemsky IN, Castegnaro M. 1988. Ochratoxin A in human blood in  
48 relation to Balkan endemic nephropathy and urinary system tumours in Bulgaria. *Food*  
49 *Addit. Contam.* 5: 299–301.
- 50  
51 Pitt JI. 1987. *Penicillium viridicatum*, *Penicillium verrucosum* and production of ochratoxin A.  
52 *Appl Environ Microbiol* 53: 266–269.
- 53  
54 Ruprich, J., Ostrý, V., 1993. Study of human exposure to ochratoxin A and assessment of  
55 possible sources. *Central European Journal of Public Health* 1, 46–48.
- 56  
57 Schaaf GJ, Nijmeijer SM, Maas RFM, Roestenberg P, De Groene EM, Fink-Gremmels J. 2002.  
58 The role of oxidative stress in the ochratoxin A-mediated toxicity in proximal tubular  
59 cells. *Biochim Biophys Acta* 1588: 149–158.
- 60  
61 Scott PM, Kanhere SR, Lau BPY, Lewis DA, Hayward S, Ryan JJ, Kuiper-Goodman T. 1998. Survey  
62 of Canadian human blood for ochratoxin A. *Food Addit Contam* 15: 555–562.

- 1  
2  
3 Skaug MA. 2003. Levels of ochratoxin A and IgG against conidia of *Penicillium verrucosum* in  
4 blood samples from healthy farm workers. *Ann Agric Environ Med* 10: 73-77.
- 5  
6 Singare-Tigori B, Moukha S, Kouadio JH, Dano DS, Betbeder A-M, Achour A, Creppy EE. 2006.  
7 Ochratoxin A in human blood in Abidjan, Côte d'Ivoire. *Toxicon* 47: 894-900.
- 8  
9 Solti L, Salomon F, Barna-Vetro I, Gyongyosi A, Szabo E, Wolfling A. 1997. Ochratoxin A content  
10 of human sera determined by a sensitive ELISA. *J Anal Toxicol* 21: 44-48.
- 11  
12 Thirumala-Devi K, Mayo MA, Reddy G, Emmanuel KE, Larondelle Y, Reddy DV. 2001.  
13 Occurrence of ochratoxin A in black pepper, coriander, ginger and turmeric in India.  
14 *Food Addit Contam* 18: 830-835.
- 15  
16 Thuvander A, Paulsen JE, Axberg K, Johansson N, Videnes A, Enghardt-Barbieri H, Tryg K, Lund-  
17 Larsen K, Jahrl S, Widenfalk A, Bosnes V, Alexander J, Hult K, Olsen M. 2001. Levels of  
18 ochratoxin A in blood from Norwegian and Swedish blood donors and their possible  
19 correlation with food consumption. *Food Addit Contam* 39, 1145-1151.
- 20  
21 Thuvander A, Paulsen JE, Axberg K, Johansson N, Vidnes A, Enghardt-Barbieri H, Tressou J,  
22 Leblanc J-Ch, Feinberg M, Bertail P. 2004. Statistical methodology to evaluate food  
23 exposure to a contaminant and influence of sanitary limits: application to ochratoxin A.  
24 *Regul Toxicol Pharmacol* 40: 252-263.
- 25  
26 Ueno Y, Maki S, Lin J, Furuya M, Sugimura Y, Kawamura O. 1998. A 4-year study of plasma  
27 ochratoxin A in a selected population in Tokyo by immunoassay and immunoaffinity  
28 columnlinked HPLC. *Food Chem Toxicol* 36: 445-449.
- 29  
30 Valenta H. 1998. Chromatographic methods for the determination of ochratoxin A in animal  
31 and human tissues and fluids. *J Chromatogr A* 815: 72-92.
- 32  
33 Valenta H, Goll M. 1996. Determination of ochratoxin A in regional samples of cow's milk from  
34 Germany. *Food Addit Contam* 13: 669-676.
- 35  
36 van der Merwe KJ, Steyn PS, Fourie L. 1965. Ochratoxin A, a toxic metabolite produced by  
37 *Aspergillus ochraceus*. *Wilh. Nature*, 205 (4976): 1112-1113.
- 38  
39 van Egmond HP, Speijers GJA. 1994. Survey of data on the incidence and levels of ochratoxin A  
40 in food and animal feed worldwide. *Natural Toxins* 3: 125-144.
- 41  
42 Veselá D, Veselý D, Jelínek R. 1983. Toxic effects of ochratoxin A and citrinin, alone and in  
43 combination, on chicken embryos. *Appl Environ Microbiol* 45 (1): 91-93.
- 44  
45 Visconti A, Pascale M, Centonze G. 2000. Determination of ochratoxin A in domestic and  
46 imported beers in Italy by immunoaffinity clean-up and liquid chromatography. *J.*  
47 *Chromatogr. A* 888: 321-326.
- 48  
49 Vrabcheva T, Gareis M, Bresch H, Bedectel C, Engel G, Majerus P, Rosner H, Wolff J. 1998.  
50 Occurrence of ochratoxins A and B in spices and herbs. *Revue de Médecine Vétérinaire*  
51 (special issue June 1998) 533.
- 52  
53 Zimmerli B and Dick R. 1995. Determination of ochratoxin A at ppb level in human blood,  
54 serum, milk and some foodstuffs by HPLC with enhanced fluorescence detection and  
55 immunoaffinity column cleanup: methodology and Swiss data. *J Chromatogr B Biomed.*  
56 *Appl.* 666: 85-99.
- 57  
58  
59  
60

Table 1- Meat and meat-derived products frequency of consumption, from healthy blood donors living in the Valencian Community, in times per week. Data collected from 168 questionnaires are shown in overall and also split by sex and age.

	Age range	Number of cases	Overall meat consumption (times/week)	Pork consumption (times/week)	Beef consumption (times/week)	Chicken consumption (times/week)	Turkey consumption (times/week)	Other meats consumption (times/week)	Animal entrails (% of Yes) [average frequency among positives]	Cold meat consumption (times/week)	Sausages consumption (times/week)
All data		168	4.88	1.55	1.26	2.02	0.54	0.48	27.9 [2 or 3 days/month ]	1.98	1.27
Men											
	Overall	120	4.93	1.66	1.34	1.93	0.43	0.53	28.3 [2 or 3 days/month]	2.19	1.54
	18-35	42	5.85	2.07	1.38	2.19	0.62	0.67	33.3 [2 or 3 days/month]	2.55	2.09
	36-45	53	4.77	1.62	1.36	1.77	0.42	0.38	20.8 [3 or + days/month]	2.11	1.36
	46-	25	3.72	0.95	1.21	1.71	0.17	0.67	36.0 [1 day/month]	1.79	0.92
Women											
	Overall	48	4.77	1.27	1.06	2.27	0.79	0.33	27.1 [2 or 3 days/month ]	1.46	0.60
	18-35	24	5.00	1.54	1.04	2.08	1.17	0.25	29.2 [3 or + days/month ]	2.04	0.58
	36-45	13	4.30	1.00	1.08	2.53	0.39	0.15	23.1 [2 or 3 days/month ]	1.46	0.69
	46-	11	4.81	1.00	1.09	2.36	0.46	0.73	27.3 [1 day/month ]	0.18	0.55

Table 2- Cereals and derivates frequency of consumption, from healthy blood donors living in the Valencian Community, in times per week and average quantity of amount consumed each time. Data collected from 168 questionnaires are shown in overall and also split by sex and age.

	Age range	Number of cases	Days of bread consumption / week [Avg. quantity in grams]	Days of bun consumption / week [Avg. quantity in grams]	Days of Italian pasta consumption / week [Avg. quantity in grams]	Days of breakfast cereal consumption/ week [Avg. quantity in grams]	Days of cookie consumption/ week [Avg. number of cookies]	Days of sweet corn consumption/ week [Avg. quantity in grams]	Days of snacks and crisp consumption/ week	% people that selected whole-grain flour [% people that selected both plain and wholegrain]
All data		168	6.06 [129.75]	1.27 [32.6]	2.02 [99.82]	1.36 [11.6]	2.19 [2.2]	1.28	0.90	14.3 [4.2]
Men										
	Overall	120	6.09 [150.5]	1.39 [35.8]	2.08 [103.1]	0.98 [10.0]	2.23 [2.9]	1.26	0.87	14.2 [4.2]
	18-35	42	5.98 [147.9]	1.71 [46.7]	2.09 [105.4]	1.48 [13.6]	2.21 [3.1]	1.43	1.33	4.16 [1.7]
	36-45	53	6.02 [153.1]	1.58 [40.2]	2.26 [103.8]	0.60 [7.1]	2.11 [3.1]	1.09	0.79	5.83 [1.7]
	46-	25	6.42 [145.3]	0.46 [8.3]	1.67 [97.9]	0.96 [10.7]	2.54 [2.5]	1.29	0.25	4.16 [0.83]
Women										
	Overall	48	5.98 [78.4]	0.98 [36.9]	1.88 [91.38]	2.33 [16.1]	2.08 [2.0]	1.29	1.0	14.6 [4.2]
	18-35	24	5.41 [75.5]	1.42 [45.7]	2.13 [86.0]	3.0 [19.6]	2.0 [1.9]	1.75	1.46	6.3 [2.1]
	36-45	13	6.69 [100.0]	0.77 [30.8]	2.08 [106.2]	2.77 [16.2]	2.31 [1.6]	1.15	0.92	6.3 [2.1]
	46-	11	6.36 [59.0]	0.27 [25.0]	1.09 [82.0]	0.36 [8.0]	2.0 [2.4]	0.46	0.1	2.1 [0.0]

Table 3- Beverages frequency of consumption, from healthy blood donors living in the Valencian Community, in times per week and average quantity of amount consumed each time. Data collected from 168 questionnaires are shown in overall and also split by sex and age.

	Age range	Number of cases	Days of beer consumption /week [Avg. quantity in mL]	Days of wine consumption / week [Avg. quantity in mL]	Days of coffee consumption / week [Avg. quantity coffees/day]	Days of fruit juice consumption/ week [Avg. quantity in mL]	Days of infusion consumption/ week [Avg. quantity infusions/day]
All data		168	2.31 [303.6]	1.22 [100.6]	5.10 [1.6]	2.98 [164.1]	1.26 [0.43]
Men							
	Overall	120	2.85 [348.3]	1.38 [114.6]	5.02 [1.5]	3.31 [182.7]	1.07 [0.4]
	18-35	42	2.36 [361.9]	0.71 [86.9]	4.95 [1.5]	3.43 [198.6]	1.07 [0.5]
	36-45	53	3.13 [292.5]	1.34 [116.0]	4.89 [1.5]	3.51 [176.9]	0.87 [0.3]
	46-	25	2.92 [450.0]	2.38 [147.9]	5.33 [1.4]	2.67 [166.7]	1.54 [0.5]
Women							
	Overall	48	0.96 [186.9]	0.83 [63.3]	5.29 [1.7]	2.17 [115.0]	1.73 [0.5]
	18-35	24	0.54 [173.9]	0.33 [39.1]	4.71 [1.5]	3.33 [183.8]	2.17 [0.6]
	36-45	13	1.92 [238.5]	1.85 [129.2]	6.46 [1.9]	0.08 [15.4]	1.31 [0.3]
	46-	11	0.73 [150.0]	0.73 [40.0]	5.18 [1.9]	2.10 [100.0]	1.27 [0.5]

Table 4- Frequency of consumption in times per week of different products described as contaminated with OTA and typical in the Mediterranean diet. In some cases the amount consumed each time is also shown. Data collected from 168 questionnaires are shown in overall and also split by sex and age.

	Age range	Number of cases	Days of chocolate consumption / week [Avg. quantity in grams]	Days of dried fruits consumption / week	Days of cheese consumption / week [Avg. quantity in grams]	Days of spices consumption/ week	Days of olive consumption/ week [Avg. quantity of olives]
All data		168	1.69 [13.7]	1.78	3.88 [59.4]	1.86	1.68 [4.7]
Men							
	Overall	120	1.83 [14.3]	1.81	3.74 [60.5]	1.97	1.93 [5.2]
	18-35	42	2.14 [19.5]	2.43	4.29 [72.0]	2.38	1.95 [5.1]
	36-45	53	1.75 [11.1]	1.25	3.62 [56.0]	1.83	1.70 [4.9]
	46-	25	1.50 [12.5]	1.96	3.00 [48.5]	1.58	2.38 [6.0]
Women							
	Overall	48	1.33 [11.9]	1.67	4.25 [56.6]	1.60	1.04 [3.4]
	18-35	24	1.58 [16.9]	1.46	4.50 [57.8]	1.58	1.08 [3.3]
	36-45	13	1.08 [8.4]	1.38	4.00 [42.3]	1.77	1.08 [3.5]
	46-	11	1.09 [5.4]	2.45	4.00 [70.9]	1.45	0.91 [3.5]

Table 5- Ochratoxin A levels found in blood plasma and body mass index from 168 healthy donors. Cases are split by sex and age.

	Age range	Number of cases	B.M.I. <sup>a</sup>	OTA $\mu\text{g/L}$ plasma $\pm$ S.D. <sup>b</sup>	Range OTA $\mu\text{g/L}$ plasma
All data		168	25.92	1.09 $\pm$ 0.95	0.15 - 5.71
Men					
	Overall	120	26.2	1.14 $\pm$ 0.97	0.15 - 5.71
	18-35	42	25.13	1.13 $\pm$ 0.97	0.15 - 5.71
	36-45	53	26.43	1.35 $\pm$ 1.11	0.22 - 5.10
	46-	25	27.5	0.76 $\pm$ 0.43	0.21 - 2.00
Women					
	Overall	48	25.24	0.98 $\pm$ 0.92	0.26 - 5.62
	18-35	24	25.43	0.81 $\pm$ 0.62	0.26 - 2.45
	36-45	13	24.32	1.38 $\pm$ 1.41	0.37 - 5.62
	46-	11	25.93	0.92 $\pm$ 0.74	0.37 - 2.89

a: Body Mass Index

b: Standard Deviation

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Table 6. Correlation coefficients (Spearman rank) and correlation probabilities (Gamma correlations) between food consumption and blood plasma levels of ochratoxin A. Only food items and habits data for which statistically significant correlation coefficients (P<0.05) were obtained in one or more of the populations are shown.

Variables	Overall population		Men		Women	
	Spearman rank correlation test	<i>Gamma</i> correlation statistic	Spearman rank correlation test	<i>Gamma</i> correlation statistic	Spearman rank correlation test	<i>Gamma</i> correlation statistic
Beef consumption	<b>0.199<sup>a</sup></b>	<b>0.181</b>	0.173	<b>0.155</b>	0.214	0.205
Turkey consumption	<b>0.163</b>	<b>0.183</b>	<b>0.184</b>	<b>0.220</b>	0.237	0.249
Cold meat consumption	<b>0.220</b>	<b>0.182</b>	<b>0.287</b>	<b>0.235</b>	-0.103	-0.084
Bread consumption	<b>0.211</b>	<b>0.149</b>	<b>0.206</b>	<b>0.146</b>	0.150	0.046
Wine consumption	-0.052	-0.047	-0.161	-0.148	<b>0.332</b>	<b>0.405</b>
Cheese consumption	0.114	0.082	0.043	0.031	<b>0.336</b>	<b>0.237</b>
Habits						
Meals outside home	<b>0.202</b>	<b>0.240</b>	0.179	<b>0.208</b>	0.237	0.314

<sup>a</sup> Figures given in bold indicate that the correlation is statistically significant (P<0.05)