



HAL
open science

Modeling water exchange between Baltimore Harbor and Chesapeake Bay using artificial tracers: seasonal variations

Bo Hong, Nauth Panday, Jian Shen, Harry V. Wang, Wenping Gong, Anna Soehl

► **To cite this version:**

Bo Hong, Nauth Panday, Jian Shen, Harry V. Wang, Wenping Gong, et al.. Modeling water exchange between Baltimore Harbor and Chesapeake Bay using artificial tracers: seasonal variations. *Marine Environmental Research*, 2010, 70 (1), pp.102. 10.1016/j.marenvres.2010.03.010 . hal-00598203

HAL Id: hal-00598203

<https://hal.science/hal-00598203>

Submitted on 5 Jun 2011


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Modeling water exchange between Baltimore Harbor and Chesapeake Bay using artificial tracers: seasonal variations

Authors: Bo Hong, Nauth Panday, Jian Shen, Harry V. Wang, Wenping Gong, Anna Soehl



PII: S0141-1136(10)00050-4

DOI: [10.1016/j.marenvres.2010.03.010](https://doi.org/10.1016/j.marenvres.2010.03.010)

Reference: MERE 3437

To appear in: *Marine Environmental Research*

Received Date: 22 October 2009

Revised Date: 12 March 2010

Accepted Date: 24 March 2010

Please cite this article as: Hong, B., Panday, N., Shen, J., Wang, H.V., Gong, W., Soehl, A. Modeling water exchange between Baltimore Harbor and Chesapeake Bay using artificial tracers: seasonal variations, *Marine Environmental Research* (2010), doi: 10.1016/j.marenvres.2010.03.010

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Modeling water exchange between Baltimore Harbor and Chesapeake Bay**
2 **using artificial tracers: seasonal variations**

3

4 Bo Hong^{1*}, Nauth Panday², Jian Shen¹, Harry V. Wang¹, Wenping Gong³, and Anna Soehl²

5 1. Virginia Institute of Marine Science, College of William and Mary, Gloucester Point, VA 23062, USA

6 2. Maryland Department of Environmental Quality, Baltimore, MD 21230

7 3. School of Marine Science, Sun Yat-Sen University, 135West Xingang Rd., Guangzhou 510275, China

8

9

10

11

12

13

14

15

16

17

18

19 To be submitted to Marine Environmental Research

20

21 *Corresponding author:

22 Dr. Bo Hong

23 Virginia Institute of Marine Science, College of William and Mary

24 Rt. 1208, Greate Road, P. O. Box 1346

25 Gloucester Point, VA 23062, USA

26

27 Phone: 804-684-7897

28 Fax: 804-684-7899

29 Email: andyhong@vims.edu

30

31

32

33

34

1 **Abstract** Understanding the dynamics of water exchange between Baltimore Harbor and the
2 Chesapeake Bay is essential when evaluating transport and fate of dissolved substances in both of these
3 systems. Conservative artificial tracers are used in this study to investigate transport processes through a
4 three-dimensional hydrodynamic model (CH3D). The model well reproduced the three-layered
5 circulation pattern in Baltimore Harbor. Several numerical experiments are performed to trace the water
6 mass coming from different sources. The results indicate that both the upper and lower layers of the
7 Harbor are the dominant pathways of transporting dissolved substances from Susquehanna River to the
8 Harbor. Such inward transport is intensified (suppressed) during the high-discharge (low-discharge)
9 period. The upper layer inflow transports water mass with high concentrations of dissolved substances
10 while the inflow from the lower layer transports water mass with low concentrations of dissolved
11 substances. The bottom layer is the dominant pathway for transporting dissolved substances from the
12 lower Bay to the Harbor. Lower river discharge and stronger along-Bay pressure gradient (resulting in
13 stronger landward residual flow in the bottom layer of the Bay) facilitate the bottom intrusion of dissolved
14 substances from lower Bay to the Harbor. Once contaminants are transported into the Harbor, they
15 usually stay for a longer time in the mid-depth of the Harbor than those in other layers due to the three-
16 layer circulation in the Harbor. The time needed for the contaminants being transported out of the Harbor
17 during a typical low-discharge period is about 1 month longer than that needed during a typical high-
18 discharge period. The results, from the environmental perspective, provide new insights for quantitative
19 evaluation on the transport processes of the dissolved biogeochemical substances between Baltimore
20 Harbor and Chesapeake Bay.

21
22 **Keywords:** Water exchange; three-dimensional model; Tracer; Age; Baltimore Harbor; Chesapeake Bay

1 **1. Introduction**

2
3 Baltimore Harbor is a tributary embayment in the upper western portion of Chesapeake Bay
4 draining a highly urbanized and industrialized Patapsco River basin (Fig. 1). Water quality and
5 ecosystem health in the Harbor is susceptible to the industrial, municipal, and stormwater runoff
6 from the surrounding watershed as well as the inflow from the adjacent Bay. Excessive nutrient
7 runoff from the Chesapeake Bay drainage basin is known to contribute to the hypoxia or anoxia
8 events in deepwater (referred to as “dead zone”) in the mainstem of the Bay during much of the
9 summer (Kemp et al., 2005). These deepwater are not only harmful to the aquatic life in the Bay
10 itself, but can also spread into most of the nearby embayments (Kuo and Neilson, 1987). Owing
11 to the different source and transport pathways of dissolved substances, the quantitative
12 interpretation of water exchange between Baltimore Harbor and the Bay is necessary for better
13 protecting and utilizing the environmental resources in these areas.

14 The stratification in upper Chesapeake Bay is maintained by dynamic processes associated
15 with freshwater discharged into the system primarily by the Susquehanna River, resulting in a
16 two-layered circulation (Schubel and Pritchard, 1986). Baltimore Harbor has relatively small
17 drainage areas. The combined average daily discharge of the tributaries to the Harbor represents
18 only about 1/315 of the volume of the Harbor (Schubel and Pritchard, 1986). The mean tidal
19 range of the Harbor is about 0.35 m and the tidal currents in the Harbor are relatively weak
20 (<http://tidesandcurrents.noaa.gov>). The estimated maximum tidal excursion is about 2.63 km.
21 Given the insignificant freshwater discharge and the low tidal energy input to the Harbor, the
22 unique three-layered circulation developed in the Harbor and the stratification are mainly
23 controlled by the density conditions in the adjacent Bay.

24 Although the three-layered circulation in Baltimore Harbor has been extensively investigated
25 by many previous studies (e.g., Carpenter, 1960; Stroup et al., 1961; Cameron and Pritchard,
26 1963; Chao et al., 1996), quantitative evaluation of the water exchange and pollutant transport
27 processes between the Harbor and the Bay from the environmental prospective is still very
28 limited. The questions that need to be further addressed include: (i) what amount of dissolved
29 pollutants in the Baltimore Harbor is associated with the inflow from the main Bay? (ii) how
30 long does it take for these substances to be transported from the source area to the area of

1 concern? and (iii) how does the transport time-scale change spatially and temporally under
2 different hydrodynamic conditions?

3 Transport of water and dissolved substances in estuary is strongly influenced by numerous
4 factors such as river discharge, tides, wind, bottom topography, Coriolis effect, etc. It is usually
5 difficult to trace water mass (or nutrients, dissolved pollutants, etc.) coming from different
6 source regions in *in-situ* observations since hydrodynamic processes usually result in the mixture
7 of them. However, the fate and transport timescales of these biogeochemical substances are
8 essential to both oceanographic research and environmental assessment. Using artificial tracers
9 in numerical modeling is one of the feasible methods that can be used to address the problem.

10 Previous studies have demonstrated the advantages of using artificial tracers in numerical
11 simulations to determine the fate and transport timescales of dissolved substances since they left
12 the source region. For example, Hirst (1999) examined the penetration and fate of North Atlantic
13 Deep Water in a global ocean model; Delhez and Deleersnijder (2002) simulated the advection-
14 dispersion of tracers discharged at the nuclear fuel reprocessing plant of Cap de La Hague; Shen
15 and Haas (2004) used a three-dimensional numerical model to calculate the age distribution for
16 the substance released from the head waters of the York River estuary under different
17 hydrodynamic conditions. In addition, both particle trajectories and passive tracers were used in
18 a three-dimensional circulation model by Döös and Engqvist (2007) to estimate the potential fate
19 and distribution of radio-nuclides released in the coastal region of the Baltic Sea. Through age
20 tracer modeling, Gustafsson and Bendtsen (2007) quantified the timescales of downward
21 (upward) mixing of surface (bottom) water in a shallow fjord. In these studies, artificial tracers
22 are integrated simultaneously with the hydrodynamic fields and provide powerful tools that can
23 help us to quantify the transport processes of dissolved substances and understand the
24 mechanisms that control their temporal and spatial variations.

25 For the purpose of quantifying water exchange between Baltimore Harbor and the mainstem
26 of the Chesapeake Bay, we add conservative artificial tracers in an extensively calibrated three-
27 dimensional hydrodynamic model. The hydrodynamic model and the method of quantifying
28 water exchange are introduced in Section 2. Section 3 simply describes the hydrodynamic
29 background of the concerned area. The diagnostic and quantitative analyses of water exchange
30 between Baltimore Harbor and main Bay are presented in Section 4. Discussion and conclusions
31 are provided in Section 5.

1

2 **2. Model Description**

3

4 2.1 Hydrodynamic Model

5 A three-dimensional numerical hydrodynamic model (Curvilinear Hydrodynamics in Three
6 Dimensions, CH3D) was used in this study. CH3D was originally developed by Sheng (1986),
7 and subsequently modified extensively by the US Waterways Experiment Station for application
8 to Chesapeake Bay (Johnson et al., 1991; 1993; 1995; Wang and Chapman, 1995). The detailed
9 description and theoretical aspects of CH3D can be found in Johnson et al. (1991). Second order
10 k- ϵ turbulence closure model (Kundu, 1980) is used to calculate the eddy viscosity and
11 diffusivity coefficient. The background horizontal diffusivity is set to $1 \times 10^{-5} \text{m}^2 \text{s}^{-1}$. A particular
12 feature of CH3D is the solution of transformed equations on a boundary-fitted curvilinear
13 coordinates in the horizontal plane. This allows the model grid to cope with the complicated
14 shoreline configuration and keep a good representation of the deep channel orientation in the
15 Chesapeake Bay and its tributaries, whereas high-resolution grids are required in other models
16 with orthogonal curvilinear coordinates to meet the needs of keeping reasonable
17 orthogonalization and good representation of complex estuarine bathymetry.

18 CH3D has been successfully used to investigate the hydrodynamic features and transport
19 processes of dissolved substances in the Chesapeake Bay (e.g., Hood et al., 1999; Xu et al.,
20 2002; Shen and Wang, 2007). It is also extended to include the sediment transport (Lin et al.,
21 2004; Park et al., 2008) and biogeochemical sub-model (Xu and Hood, 2006), in which the
22 physical model provides hydrodynamic background for the sediment and biogeochemical
23 constituents in three dimensions. The model grid of this application (developed by Wang et al.,
24 2004) is shown in Fig. 2. There are 19 layers in the vertical with a uniform layer thickness of
25 1.52 m, except for the top layer whose thickness fluctuates with water level. The model is
26 initialized by the Chesapeake Bay Program (CBP) Monitoring data
27 (<http://www.chesapeakebay.net/dataandtools.aspx/>). Daily fresh water inflow
28 (<http://waterdata.usgs.gov/nwis/dv/>) with zero salinity and time-varying temperature was
29 prescribed for the major tributaries, which include Susquehanna River, C&D Canal, Chester
30 River, Choptank River, and Patapsco River (see Fig. 1 for the locations of these tributaries). At
31 the southern open boundary the tidal elevations are prescribed by hourly data observed at

1 Solomons Island, MD (<http://tidesandcurrents.noaa.gov/>, Station ID: 8577330). The upstream
2 schemes are adopted for the temperature and salinity fields at the southern open boundary
3 condition in which the observed vertical salinity and temperature profiles at Station CB4.4 (see
4 Fig. 2 for its location) are advected into the model domain for the inflow conditions. Since the
5 observed salinity and temperature data are collected at approximately monthly intervals, the data
6 were first linearly interpolated into daily intervals to obtain the annual cycle. Then the station
7 data were interpolated onto the model grid as open boundary forcings. The radiation boundary
8 condition was used for the velocity field. The velocity at the open boundary was calculated in the
9 model based on the observed surface elevation field and the density profile. The daily averaged
10 surface heat exchange and equilibrium temperature are computed from the corresponding
11 meteorological data (NOAA, 2007).

12 Lin et al. (2004) used this model configuration to simulate hydrodynamic fields covering both
13 very wet and very dry hydrological years. Modeled bottom velocity fields were compared with
14 observations at a mooring station inside the Baltimore Harbor in spring of 2000. The simulated
15 along-Harbor velocity agrees very well with the observations. Park et al. (2008) also used this
16 model configuration to simulate the hydrodynamic field in 1996. By using the two surveys in
17 October followed a peak discharge of $5,267 \text{ m}^3 \text{ s}^{-1}$ on October 21 and the two surveys in July
18 under the low discharge conditions (data presented in Sanford et al. (2001)), the model
19 performance were validated over a wide range of freshwater discharge values. The spatial
20 distribution characteristics of observed along-Bay salinity profiles were reproduced well by the
21 model. The model-data comparisons at a mooring station indicated that the model could catch
22 the observed vertical structure variations and the vertical mixing scheme worked properly. These
23 model-data comparisons indicated the success of model performance on tidal timescales. Liu et
24 al. (2009) presented a good example of model-data comparisons. They evaluated model skill not
25 only for tidal properties but also for subtidal properties. Since we are focusing on the seasonal
26 variations in this study, further model validation has been conducted to confirm the model
27 performance in modeling seasonal estuarine circulation. Whereas simulations were conducted for
28 multiple years, only the results in 2000 are presented here (similar model performances were
29 obtained in other years). Comparisons between the modeled salinity and CBP monitoring data
30 are shown in Fig. 3a. Model results show consistent seasonal variations with the observations. It
31 is shown that the model can catch the seasonal salinity variations in the upper Chesapeake Bay.

1 Fig. 3b presents the comparisons of water surface elevation at NOAA observation stations. Both
 2 the modeled and observed results are 15-day low-pass filtered before plotting. It can be seen that
 3 model results reproduced the seasonal water surface elevations very well. The correlation
 4 coefficients (CC) at each station are also presented in Fig. 3b, which are at the 95% confidence
 5 level. The CC at Tolchester is the lowest (= 0.93). It can be seen that this model reproduces the
 6 upper Bay seasonal variations very well. The present model configuration is robust and suitable
 7 for this study.

8

9 2.2 Methods

10 Water coming from a specific origin is identified by the passive tracer, which also represents
 11 the dissolved substances (contaminants, etc.) in the water. There are two aspects that need to be
 12 addressed for quantifying the exchange of water coming from different origins. One is the
 13 amount of the dissolved substances, which is represented by the tracer concentration. The other
 14 is the transport timescale of the dissolved substances, which is represented by the tracer age (the
 15 tracer age is defined as the time elapsed since the tracer under consideration is released into the
 16 water body where its age is prescribed to be zero (Delhez et al., 1999)). The concentration and
 17 the age of the tracer are calculated in the numerical model using transport equations
 18 (Deleersnijder et al., 2001). The tracer concentration equation is

$$19 \quad \frac{\partial c(t, \vec{x})}{\partial t} + \nabla \cdot (\vec{v}c(t, \vec{x}) - K \nabla c(t, \vec{x})) = 0 \quad (1)$$

20 Where $c(t, \vec{x})$ is the tracer concentration, \vec{v} is the velocity vector, K is the diffusivity tensor, t is
 21 the time, and \vec{x} is the spatial coordinate. There is no sink of the tracer within the study area.
 22 According to Deleersnijder et al. (2001), the evolution of age concentration ($\alpha(t, \vec{x})$) is described
 23 as

$$24 \quad \frac{\partial \alpha(t, \vec{x})}{\partial t} + \nabla \cdot (\vec{v}\alpha(t, \vec{x}) - K \nabla \alpha(t, \vec{x})) = c(t, \vec{x}) \quad (2)$$

25 Then the mean age ($a(t, \vec{x})$) can be calculated as

$$26 \quad a(t, \vec{x}) = \frac{\alpha(t, \vec{x})}{c(t, \vec{x})} \quad (3)$$

27 At the releasing site, the tracer concentration is 1 (arbitrary unit) and the tracer age is zero. The
 28 tracer age at any other location is representative of the timescale for water or a dissolved

1 substance to be transported from its source to this location. The spatial and temporal variations of
2 tracer concentration and mean age (hereafter referred to “water age”) will be used in the
3 following to quantify the water exchange under different hydrodynamic conditions.

4 5 2.3 Numerical Experiments Design

6 In this study, tracers are released from three different locations in three independent
7 experiments for quantifying water exchange between the Baltimore Harbor and Chesapeake Bay.
8 Previous studies have indicated that the circulation in the upper Chesapeake Bay is driven
9 primarily by the discharge of Susquehanna River (Schubel and Pritchard 1986). In order to
10 obtain more general results of seasonal variation, we use the climatological daily river runoff
11 data (1985-2007) instead of using the data from a single year. In reality, the perturbation/mixing
12 induced by wind forcing is more complicated (MacCready et al., 2009). We exclude the wind
13 effect in this study for simplicity and only focus on the effects of river discharge, which is the
14 dominant physical process determining the seasonal variation in the upper Bay. We are aware of
15 the importance of wind mixing as it affects the estuarine stratification. Wind effects are more
16 dominant during synoptic events. The influences of short-term wind events on long-term
17 stratification and long-term transport warrant more observations and studies and will be studied
18 separately. The three tracer simulation experiments are hereafter referred to as Exp. 1, Exp. 2,
19 and Exp. 3. In Exp. 1, tracers are continuously released from the entrance of Susquehanna River,
20 which are used to track the water masses (contaminants, etc.) coming from Susquehanna River.
21 In Exp. 2, tracers are continuously released from the mid-Bay (along the southern open boundary
22 of the model), which are used to track the water masses coming from the lower Bay. In Exp. 3,
23 tracers are instantaneously released over the entire Baltimore Harbor, which are used to track the
24 dissolved substances over the Harbor. There is no other source of tracer in each experiment.
25 Hydrodynamic equations and passive tracer equations are integrated simultaneously. In order to
26 reach the dynamic equilibrium condition and eliminate transition errors resulting from model
27 spin-up, the model is run repeatedly for 2 years with the annual cycle forcing fields. Only outputs
28 of the last year are used for the following analyses. Note that the spin-up time in Chesapeake Bay
29 is relatively longer than most estuaries, for example, the spin-up time is less than two weeks for
30 Columbia River estuary hindcast modeling (Liu et al., 2009). This is due to the relatively slow
31 adjustment of baroclinic fields in the highly stratified Chesapeake Bay.

1

2 **3. Circulation patterns**

3

4 To understand the long-term variability of water exchange between the Harbor and the Bay,
5 two fundamental processes need to be addressed. The first one is the variation of the
6 Susquehanna River discharge. Susquehanna River has a seasonal flow pattern typical of mid-
7 latitude rivers: high discharge in spring produced by snow melting and spring rains followed by
8 low-to-moderate flow throughout most of the remainder of the year (Carpenter, 1960; Schubel
9 and Pritchard, 1986). This variation greatly impacts the vertical stratification of the main Bay,
10 horizontal salinity gradients between the Bay and Baltimore Harbor, and the amount of dissolved
11 substances transported by river discharge. The second one is the impact of deep water coming
12 from lower Chesapeake Bay, in which the high-salinity water is advected up-Bay mainly through
13 non-tidal gravitational circulation (Pritchard, 1956). Because of the persistence of stratification
14 and strong two-layered circulation developed in the main Bay, the flow in the bottom layer not
15 only plays a role of conveying saline water from the lower Bay to the upper Bay, but also
16 conveys biogeochemical substances that mix into the deep water during various destratification
17 events.

18 The Susquehanna River discharge accounts for 87% of the total river runoff in the upper
19 Chesapeake Bay. The 23-year (1985-2007) daily mean river runoff of Susquehanna River (Fig.
20 4) indicates that the maximum (minimum) river discharge usually appears in March-April (July-
21 October), which can be defined as the typical high-discharge (low-discharge) period. Baltimore
22 Harbor is famous on its three-layered circulation pattern driven by differences in the vertical
23 salinity structure in the Harbor and in the adjacent Chesapeake Bay (Carpenter, 1960; Stroup et
24 al., 1961; Boicourt and Olson, 1982; Chao et al., 1996). The typical three-layered circulation in
25 Baltimore Harbor is shown in Fig. 5a (after Cameron and Pritchard, 1963). The characteristic
26 salinity distribution of the Harbor has the conventional positive seaward gradient in the lower
27 layer, but a negative seaward gradient nearer the surface. Thus, there is flow into the Harbor both
28 at the surface and at the bottom driven by the pressure gradient. These inflows return to the
29 Chesapeake Bay at mid-depth. Figs. 5b and 5c present the typical three-layered circulation
30 (along the navigation channel of the Harbor, transect I in Fig. 2) simulated in this study. The
31 model results show similar pattern with that demonstrated by Cameron and Pritchard (1963): in

1 the upper water column, the less saline, less dense surface waters of the Bay tend to overflow the
2 more saline, denser surface waters in the Harbor. Near the bottom, the more saline, denser waters
3 of the Bay tend to flow under the less saline, less dense waters of the Harbor. The mid-depth
4 water flows outward to offset the inward flow.

5 The mean salinity of Chesapeake Bay is a product of two competing processes, i.e., freshwater
6 influx into the Bay through rivers and exchange with the adjacent ocean (Austin, 2002). The
7 modeled surface salinity fields in the upper Chesapeake Bay averaged during the typical high-
8 discharge and low-discharge periods are shown in Fig. 6a. Competitions between the river runoff
9 and lower Bay saline water intrusion are clearly reflected. During the high-discharge period,
10 water in the upper Bay is greatly diluted. The intrusion of the freshet into Baltimore Harbor can
11 be clearly discerned from the isohalines. During the low-discharge period, the saline water takes
12 over much of the domain gradually and the surface fresher water intrusion from the main Bay to
13 the Harbor is greatly weakened and mainly confined to the area along the northern bank of the
14 Harbor. The residual velocity profile across the mouth of Baltimore Harbor (facing west, transect
15 II in Fig. 2) is presented in Fig. 6b. Without the wind forcing, three-layered circulation appears
16 throughout the year, but both the spatial distribution and the strength of the flow vary with time.
17 Uniform surface inward flow usually occurs during the high-discharge period, whereas surface
18 inward flow only occurs near the northern bank of the Harbor during the low-discharge period
19 resulting in the two-layered circulation near the southern bank. The mid-depth outward flow is
20 strengthened when the surface inflow is fully developed during the high-discharge period, and
21 then weakened during the low-discharge period. The inward flow in the bottom layer shows less
22 variation compared with those in the upper layers. This is consistent with the findings of Chao et
23 al. (1996) that indicated the existence of persistent and dominant intrusion into the bottom of the
24 Harbor against all adversities.

25 26 **4. Results**

27
28 The introduction of artificial tracers in a numerical model is a convenient way to quantify the
29 timescale for the transport process of dissolved substances relative to its source region. It also
30 can reveal the spatial distribution of tracer concentrations of dissolved substances coming from a

1 specific source. These are usually difficult to measure in observations. Water exchange between
2 Baltimore Harbor and Chesapeake Bay are investigated below from three different aspects.

3

4 4.1 Water Mass Coming from Susquehanna River (Exp. 1)

5 Exp. 1 tracks the dissolved substances coming from Susquehanna River. The selected
6 snapshots (daily mean results sampled at Day 91 and Day 255, respectively) of surface water age
7 distribution are shown in Figs. 7a and 7b, which correspond to the very high-discharge and very
8 low-discharge conditions (see Fig. 4), respectively. The water age at Day 91 is much lower than
9 that at Day 255, indicating that the transport time of the water mass coming from Susquehanna
10 River during a high-discharge period is much less than that during a low-discharge period. The
11 water age difference between Days 91 and 255 is approximately 70 (40) days at the mouth (head)
12 of the Harbor. The corresponding surface velocity field (Figs. 7e and 7f) shows that the flow at
13 Day 91 is much stronger than that at Day 255. There is strong and consistent surface inflow in
14 the Harbor at Day 91, whereas weak and divergent flow appears in the Harbor at Day 255. These
15 demonstrate that Susquehanna River discharge has substantial influence on water exchange
16 between the Harbor and the main Bay.

17 Fig. 8 presents the monthly mean water age profiles along the navigation channel in
18 Baltimore Harbor. Since the tracers are continuously released at the entrance of Susquehanna
19 River, the age distribution in Fig. 8 represents the transport timescale needed for the dissolved
20 substances to reach the Harbor. The common feature of the age distribution is that the water age
21 at either the surface or bottom layer of the Harbor is usually lower than that in the mid-depth of
22 the water column. The largest vertical age difference can be about 50 days and appears in winter.
23 After the high-discharge period starts, such vertical difference decreases to about 20 days. The
24 lowest water age appears in April when Susquehanna River is at its maximum discharge. The age
25 starts to increase from May and reaches its maximum in November.

26 More detailed structure of the water exchange process can be observed from the time evolution
27 of tracer concentration and water age at selected stations (Fig. 9). Stations C, D, and E are
28 located in the mouth, the exterior, and the interior of the Harbor respectively, and Station B is in
29 the main channel of the Bay (see Fig. 2 for these locations). For the water mass coming from the
30 Susquehanna River, the tracer concentration decreases with depth in the Harbor (Figs. 9a-c). The
31 highest tracer concentration (>80% at the mouth of the Harbor, >60% in the interior of the

1 Harbor) appears in April. The time required for the tracer concentration in the lower layer to
2 reach its maximum value ($>30\%$) is delayed approximately a half-month compared with that in
3 the surface layer. When examining the tracer concentration in the main channel of the Bay
4 (Station B, Fig. 9d), we can find such temporal and spatial variations of tracer concentration also
5 occur in the main Bay. The corresponding water age at each station in the Harbor (Figs. 9e-g)
6 show significant temporal variations. In general, the water age during the high-discharge period
7 is about 2 months longer than that during the low-discharge period. As the inflow is greatly
8 weakened during the low-discharge period (Fig. 6 and Fig. 7f), the bottom layer water age in the
9 inner Harbor is approximately 1 month larger than that at the surface layer due to reduced
10 gravitation circulation. The largest water age appears at about 6 m (4 m) depth during the high-
11 discharge (low-discharge) period, corresponding to where the mid-depth outflow exists. Such
12 pattern disappears in the main channel of the Bay (Station B, Fig. 9h) where the water age
13 always increases with depth due to the stratification effect (Shen and Lin, 2006).

14 4.2 Water Mass Coming from Lower Chesapeake Bay (Exp. 2)

15 Exp. 2 tracks the water mass coming from the lower Chesapeake Bay. Since the transport
16 timescale of water mass coming from the lower Bay highly relies on the releasing site, the mean
17 age can be different for tracers released at different sites. However, using the same reference site,
18 the results in Exp. 2 are still efficient for studying the variations of transport timescale for
19 dissolved substances coming from the lower Bay. The selected snapshots (daily mean results
20 sampled at Day 91 and Day 255, respectively) of surface age distribution are shown in Figs. 7c
21 and 7d, which correspond to the very high-discharge and very low-discharge conditions,
22 respectively. A notable feature existing in these two snapshots is that the water age near the head
23 of the Harbor is lower than that in the inner Harbor. Such difference tends to be augmented at
24 Day 255 compared with that at Day 91. In general, the water age near the northern bank of the
25 Harbor is lower than that near the southern bank. Since the water mass coming from the lower
26 Bay mainly occupies the lower layer of the Bay body and enters Baltimore Harbor mainly
27 through the bottom inflow, the detailed vertical structure needs to be examined to reveal the
28 water exchange processes of the lower Bay and the Harbor.

29 The monthly mean water age profiles along the navigation channel in Baltimore Harbor are
30 shown in Fig. 10. Unlike the results in Exp.1, the highest water age does not manifest in the mid-
31 depth of the Harbor in Exp. 2, except near the Harbor mouth. The water age decreases

1 consistently with depth. The contours are quite flat at the mid-depth. The water age near the
2 bottom of the Harbor is usually less than that near the surface by 40-50 days. During the high-
3 discharge period, bottom water age increases by approximately 20 days. This indicates that the
4 bottom water exchange between the Harbor and the Bay is faster during the low-discharge period
5 than during the high-discharge period.

6 Fig. 11 presents the depth-time profiles of tracer concentration and water age at selected
7 stations. Tracer concentrations in the Harbor (Figs.11a-c) increase with depth throughout the
8 year. The lowest tracer concentration appears during the high-discharge period (at both the upper
9 and the lower layers of the water column), and then increases gradually as the low-discharge
10 period begins. The tracer concentration during the low-discharge period is usually higher than
11 that during the high-discharge period by 20-30%, and the vertical difference of tracer
12 concentration during the high-discharge period is higher than that during the low-discharge
13 period. Similar patterns can be seen in the main Bay (Fig. 11d). The time evolutions of vertical
14 age profiles at selected stations are shown in Figs. 11e-h. The highest vertical age variations
15 appear at around 6 m, where the haloclines fluctuate with the dynamics. The bottom layer water
16 age is almost the same between the Harbor and the Bay. But in the upper layer, the water age in
17 the Harbor is about 30 days larger than that in the main Bay.

18

19 4.3 Influence of Water Mass in Baltimore Harbor to the upper Bay (Exp. 3)

20 The highly urbanized and industrialized watershed draining to the Harbor contributes various
21 contaminants. In order to better understand the potential impact of the Baltimore Harbor
22 watershed on downstream water quality, tracers were released instantaneously in the Harbor
23 (Exp.3). The spatial and temporal variations of these tracers were used to quantify the influences
24 of the contaminants in the Harbor to the Bay. At each grid cell in the Harbor, the tracers were
25 released over the entire water column. Two model runs were conducted under the typical high-
26 discharge and low-discharge conditions, respectively. For the high-discharge simulation, a flow
27 rate of $1900 \text{ m}^3 \text{ s}^{-1}$ from the Susquehanna River was specified. For the low-discharge simulation,
28 this flow rate specification was $360 \text{ m}^3 \text{ s}^{-1}$. The temporal evolution of the total tracer
29 concentration for the entire Harbor is shown in Fig. 12. Following the instantaneous release, the
30 total tracer concentration decreases quickly. The decrease of total tracer concentration under the
31 high-flow condition is much faster than that under the low-flow condition, which indicates the

1 high-flow condition facilitates the renewal of water in the Harbor. According to Prandle (1984),
 2 the turnover time is defined as the time needed for the total mass of a material within the area of
 3 interest to be reduced to a factor of e^{-1} , i.e.

$$4 \quad R(\tau) = R_0 e^{-\tau/\tau_i} \quad (4)$$

5 where R is the total tracer concentration and τ_i is the turn over time. So the total tracer
 6 concentration will be reduced by a factor of e^{-1} at the time $\tau = \tau_i$. The turnover time equals the
 7 residence time if the concentration in an embayment decrease exponentially (Zimmerman, 1976).
 8 The turnover time of the Harbor is 21.7 days (39.5 days) under the high-discharge (low-
 9 discharge) condition. Fig. 13 shows the evolution of tracer concentration in the Harbor under the
 10 typical high-discharge condition (a similar pattern develops under the low-discharge condition).
 11 It can be shown that dissolved substances in the Harbor usually remain for a longer time in the
 12 mid-depth of the Harbor (and upper layer of the inner Harbor) than those in other areas. Thus
 13 when organic contaminants are discharged into the Harbor from the drainage basin, the mid-
 14 depth of the Harbor (and the upper layer of the inner Harbor) usually sustains water quality stress
 15 over a longer period than other areas.

16 Examining the conditions at Stations f, g, and h can provide additional insight about the
 17 influence of Baltimore Harbor outflow on water quality in the main Bay. Stations f and h lie in
 18 the south and north of the Harbor, respectively, while Station g lies near the mouth of the Harbor
 19 (see Fig. 2). These stations are used to represent different areas in the Bay. The depth-averaged
 20 tracer concentrations at each station are shown in Figs. 14a and 14b. Under the high-flow
 21 condition, the tracer concentrations at Station g are zero at the beginning and start to increase
 22 after the first week. The peak tracer concentration appears at around Day 27, at which point it
 23 starts to decrease gradually with small perturbations. Similar temporal variations can be observed
 24 at other stations, with peaks occurring at different times at each station (Day 35 and Day 45 for
 25 Stations f and h, respectively). The concentration distribution shows the age density function.
 26 The calculated mean age values suggest that contaminants in Baltimore Harbor can be
 27 transported to the selected stations f, g, and h at around 66, 58, and 73 days, respectively, after
 28 being released in the Harbor. The mean age under the low-flow condition is about 1 month
 29 longer than that under the high-flow condition (around 97, 82, and 105 days for Station f, g, and
 30 h, respectively). The appearance of peak tracer concentration under the low-flow condition is

1 also later than that under the high-flow condition. The time evolution of the vertical tracer
2 concentration profile helps to reveal the distribution of tracer concentration for the entire water
3 column at each station. As an example, the results under the low-flow condition (a similar
4 pattern appears in the high-flow condition) are shown in Fig. 14c. These results suggest that,
5 after the dissolved substances are transported out of the Harbor, they show different vertical
6 occupation patterns in different areas of the main Bay. At the station south (north) of the Harbor,
7 the tracer concentration decreases (increases) with depth. For the station close to the mouth of
8 the Harbor, the highest tracer concentration occurs at around 6 m depth. The difference in the
9 vertical distribution of tracer concentration at different locations suggests that the contaminants
10 in the Harbor may influence different habitats in the mainstem of the Bay.

11

12 **5. Discussion**

13

14 The results presented in the previous section (summarized in Table 1) reveal that tracers
15 coming from different origins have different pathways and different transport timescales under
16 the three-layered circulation condition in the Baltimore Harbor. These reflect the difference of
17 water exchange processes with respect to different source origins.

18

19 5.1 Dominant Pathway of Water Exchange for Different Water Sources

20 The dissolved substances that originate from Susquehanna River (lower Bay) mainly occupy
21 the upper (lower) layer of the water column both in the upper Bay and in the Harbor and their
22 concentrations decrease (increase) consistently with depth (Figs. 9 and 11). However, this kind
23 of symmetric distribution does not appear in the vertical age distribution. In the case of releasing
24 tracers from the Susquehanna River (Exp. 1), the water age profile along the navigation channel
25 of the Harbor shows a 'sandwich' structure (high water age in the mid-depth and low water age
26 in both the upper and the lower layers, Fig. 8). For the case of releasing tracers from the lower
27 Bay (Exp. 2), the water age within the Harbor shows a monotonic vertical structure (decreasing
28 consistently with depth, Fig. 10). The results in Exp. 1 indicate that both the upper and the lower
29 layer inflow of the Harbor are the dominant pathways of transporting dissolved substances from
30 Susquehanna River to the Harbor. The upper layer inflow, associated with the high concentration
31 of dissolved substances, dominates the mass transport during a typical high-discharge period. But

1 the mass transportation through the bottom inflow becomes dominant during a typical low-
2 discharge period. Although the concentrations of dissolved substances associated with the
3 bottom inflow is less than those in the surface inflow, the transportation by bottom inflow is
4 faster than that by the surface inflow during a typical low-discharge period. In Exp. 2, on the
5 other hand, only the inflow from the lower layer is the dominant pathway for transporting
6 dissolved substances from the lower Bay into the Harbor. Because of the persistence of
7 stratification in the main Bay, which prevents the vertical mixing of bottom water to the surface,
8 the transport of water that originates from the lower Bay through the upper layer is still much
9 slower than that in the lower layer, even during a typical high-discharge period with strong
10 surface inflow to the Harbor.

11

12 5.2 Comparison with theoretical scales

13 Examining the age distribution for Exp. 2 (Fig. 11), the low age distribution near the bottom
14 does not follow the freshwater discharge, which can be attributed to the influence of the lower
15 Bay through the gravitational circulation. The observed salinity values at Station CB4.4 (see Fig.
16 2 for its location) is presented in Fig. 15a, which were used as the model open boundary
17 conditions in this study. The temporal vertical salinity distribution shows that there are several
18 high-salinity water intrusion events (i.e., salinity > 20 ppt) in the bottom layer of the Bay. The
19 depth-time profiles of water age in Exp. 2 are shown in Figs. 15b and 15c, which are sampled at
20 Stations B and C, respectively. It can be seen that the low water age events (and the high tracer
21 concentration events, (Fig. 11)) are usually associated with the high-salinity water intrusion
22 events. Although Station CB4.4 is miles away from Stations B and C, the bottom layer salinity at
23 Stations B and C (figures not shown) have consistent temporal variations with salinity at Station
24 CB4.4. In order to further examine the underlying transport mechanisms, the depth-time profile
25 of modeled along-Bay velocity at Station V (see Fig. 2) is presented in Fig. 15d. A 15-day low-
26 pass filter was applied to the velocity data before the plotting. The results clearly show the
27 two-layered gravitational circulation development in the Bay and the bottom residual flow
28 variations are highly in phase with the bottom salinity field. Velocity profiles at other stations in
29 the main channel of the Bay show similar temporal variations to those at Station V. Those results
30 indicate that the higher bottom salinity field usually corresponds to the stronger landward flow
31 resulting in a decrease of water age in the bottom layer (Figs. 15b-c). The gravitational

1 circulation is critical in changing the transport processes. When the return flow is stronger, tracer
 2 transportation is faster, which results in smaller water age. In general, the water age near the
 3 bottom of the main Bay decreases about 20 days when the landward residual velocity increases
 4 by about 8 cm s^{-1} (Fig. 15).

5 The gravitational circulation in the estuary is influenced by the combined effect of river
 6 discharge and bottom saline water intrusion from the ocean. The competition between freshwater
 7 inflow from the head and high-salinity water intrusion from the mouth results in the variation of
 8 the along-Bay pressure gradient. The Hansen and Rattray (1965) solution (HR65 hereafter) for
 9 the along-channel velocity provides a classical way to quantify the contribution of along-channel
 10 pressure gradient and river discharge to the gravitational circulation. Progress has been made
 11 since HR65 and several assumptions in HR65 have been relaxed now (e.g., Geyer et al., 2000;
 12 MacCready, 2004; 2007). MacCready and Geyer (2010) provide an excellent review on the
 13 recent advances of the theoretical solution and formulae for calculating adjustment timescales
 14 under perturbations. Using the solution technique of MacCready (2007), the along-channel
 15 velocity $u(x, z, t)$ can be divided into depth-averaged (overbar) and deviations from depth-
 16 averaged (prime) components so that $u = \bar{u} + u'$. The quantity $\bar{u} = Q_r/A$ where Q_r is the river
 17 discharge and A is the cross-sectional area. The quantity u' has the standard vertical profile as
 18 described in MacCready (2004):

$$19 \quad u' = \underbrace{\bar{u}P_1}_{(a)} + \underbrace{u_E P_2}_{(b)}$$

20 Where

$$21 \quad P_1 = \frac{1}{2} - \frac{3}{2} \left(\frac{z}{h} \right)^2 \quad \text{and} \quad P_2 = 1 - 9 \left(\frac{z}{h} \right)^2 - 8 \left(\frac{z}{h} \right)^3 \quad (5)$$

22 The magnitude of the exchange flow is $u_E = g\beta\bar{s}_x H^3 / (48K_m)$, H is the channel depth, g is the
 23 acceleration of gravity, and K_m is the vertical eddy viscosity. For simplicity, the density anomaly
 24 in the solution is assumed to depend only on salinity ($\rho = \rho_0(1 + \beta S)$, where $\beta = 0.77 \times 10^{-3}$).
 25 The depth-averaged along-estuary salinity gradient \bar{s}_x is set to be constant in the HR65 solution.
 26 Here we use the solution of MacCready (2004) (Eq. 7) which allows \bar{s}_x to vary along the estuary.
 27 Following MacCready (2007), we assume that $K_m = a_0 C_d U_r H$, where C_d is a drag coefficient

1 $(= 2.6 \times 10^{-3})$ and a_0 is a tuning coefficient (0.0325 in MacCready (2007), but 0.0227 in this
 2 study). Part (a) in Eq. (5) represents the circulation contributed by river discharge (RV), and part
 3 (b) represents the circulation contributed by pressure gradient (PG). Channel depth and width are
 4 prescribed based on the realistic conditions in the upper Chesapeake Bay. In the following, we
 5 use s_{ocn} to represent the oceanic salinity, which are based on observations at Station CB4.4 (Fig.
 6 15.1). Based on our 3D model configuration, we choose river discharge values of $2400 \text{ m}^3 \text{ s}^{-1}$
 7 and $800 \text{ m}^3 \text{ s}^{-1}$, which largely represent the conditions of high-discharge and low-discharge
 8 periods, respectively. Using the linear adjustment time theory of MacCready (2007), the salt
 9 intrusion length can be scaled by $L \propto 2 \int_R^0 \frac{\bar{s}}{s_{ocn}} dx$ and the freshwater replacement time can be
 10 approximated by $\frac{1}{2} \frac{L}{\bar{u}}$. When river discharge is $2400 \text{ m}^3 \text{ s}^{-1}$ ($800 \text{ m}^3 \text{ s}^{-1}$), the freshwater
 11 replacement time is about 15.2 (56.0) days, and the salt intrusion length is about 104 km (128
 12 km). The estimated freshwater replacement timescales are quite consistent with the water ages at
 13 the mouth calculated by Exp.1 which represents the mean transport timescale of the freshwater
 14 from the Susquehanna River. As the freshwater moves downstream, it replaces the old water.
 15 For transport of water from the lower Bay, the transport timescale depends on the exchange flow.
 16 The Knudsen relation provides a suitable estimation of the exchange flow from a given
 17 stratification. For $\bar{u} \ll u_E$, the dimensionless stratification is estimated by
 18 $\Phi = \frac{s_{bot} - s_{top}}{s_{ocn}} = \frac{3}{20} u_E \frac{H^2}{K_s} \frac{\bar{s}_x}{s_{ocn}}$ (K_s is the vertical eddy diffusivity, which has the same definition
 19 as Eq. 3.9 in MacCready (2007)). A dimensionless expression for the exchange flow can be
 20 written as $\frac{u_E}{c} = \frac{1}{48} \frac{cH^2}{K_M} \frac{1}{L}$, where $c = \sqrt{g\beta s_{ocn} H}$ is twice the speed of the fastest internal wave
 21 that could be supported in the estuary given the density difference between ocean and freshwater.
 22 Following the methodology of MacCready and Geyer (2010) and using the theoretical solution,
 23 the graphical presentation of these scaling results is shown in the plane of $(u_E/c, \Phi)$ (Fig. 16).
 24 The Knudsen relation is purely diagnostic. It tells us only which exchange flow is consistent with
 25 a given stratification. On its own, it cannot predict both stratification and exchange. Meanwhile,
 26 our 3D model results are sampled to calculate the dimensionless exchange flow and stratification

1 under various real forcing conditions. The 3D model results largely match the theoretical results
2 (Fig. 16). It is indicated that in the upper Chesapeake Bay stronger exchange flow corresponds to
3 stronger stratification. During the high salinity water intrusion events, the horizontal salinity
4 gradients increase and both the vertical stratifications and exchange flow becomes stronger.
5 Therefore, the age of the water mass coming from the lower Bay is relatively smaller.

6 5.3 Renewal of Baltimore Harbor

7 Persistent low DO during the summer in the deep-water (dead zone) of the main Bay
8 remains a serious concern for water quality managers. Observations indicate that the hypoxic or
9 anoxic water in the deep channel of the main Bay can enter most of the major tributaries due to
10 gravitational circulation (Kuo and Neilson, 1987). This phenomenon has been observed in the
11 Baltimore Harbor, which greatly degrades the ecosystem in the inner Harbor. In the experiment
12 of tracing deep water coming from the lower Bay, we find the upper layer tracer concentration
13 near the head of the Harbor is higher than that in the inner and outer Harbor (figure not shown).
14 The age distribution (Figs. 7c-d and Fig. 10) suggests that, in the upper layer of the water
15 column, the time needed for the deep water of the lower Bay to reach the head of the Harbor is
16 less than that to the inner and outer Harbor. According to Cameron and Pritchard (1963), the
17 characteristic salinity distribution of the Harbor has a positive seaward gradient in the lower
18 layer, but a negative seaward gradient nearer the surface. The resulting stratification is weakest
19 near the head of the Harbor, where vertical mixing is stronger than in other areas of the Harbor.
20 The increased vertical mixing (or upwelling) near the head of the Harbor may facilitate the
21 upward transport of dissolved substances in that area.

22 Although the concentrations of riverine pollutants entering the Harbor are higher during the
23 high-discharge period than that during the low-discharge period, the renewal of water in the
24 Harbor is faster during the high-discharge period than that during the low-discharge period.
25 Thus, the impact of dissolved riverine pollutants on the Harbor is not severe during the high-
26 discharge period as large amounts of substances will be transported out of the Harbor quickly.
27 The worst situation can occur at the end of the high-discharge period when the concentrations in
28 the Harbor are still high but the water renewal is slowing down.

30 6. Conclusions

1 The water exchange between the Baltimore Harbor and Chesapeake Bay is investigated using
2 artificial tracer in conjunction with a calibrated realistic 3D model. The tracer concentration is
3 representative of water or dissolved substances coming from the concerned source region, while
4 the water age can indicate the transport timescale of these dissolved substances. These features
5 are important factors that need to be examined when trying to protect the tidal environmental
6 resources and manage pollutant transport in estuaries.

7 The results suggest that the temporal and spatial variations of water exchange between the
8 Harbor and the Bay have obvious seasonal features, which are highly correlated with the
9 Susquehanna River discharge. The hydrodynamic condition during the high-discharge period
10 facilitates the intrusion of dissolved substances coming from Susquehanna River to the Harbor
11 (mainly via the upper layer of the water column). For the case of releasing tracers from
12 Susquehanna River (Exp. 1), the tracer concentrations increase approximately 40%, but the water
13 age decreases approximately 2 months during a high-discharge period compared to the results
14 during a low-discharge period. Both the upper and lower layers are dominant pathways for
15 transporting dissolved substances from the Susquehanna River to the Harbor, but the upper layer
16 tends to transport water masses with higher concentrations of dissolved substances.

17 During a high-discharge period, the salinity decreases in the upper Bay and the baroclinic
18 pressure between the Bay and the Harbor decreases, which suppresses the saline water intrusion
19 to the Harbor thus decreases the transport time of water coming from the lower Bay (Exp. 2).
20 Compared to a high-discharge period, the tracer concentration decreases about 20% and the
21 water age increases about 20 days during a low-discharge period. Only the lower layer is the
22 dominant pathway of transporting dissolved substances coming from the lower Bay into the
23 Harbor.

24 The transport in the main channel is highly related to the dynamic conditions in the lower Bay.
25 A high-salinity water intrusion event in the bottom layer of the Bay usually corresponds to a low
26 water age event near the bottom both in the Bay and in the Harbor. The transport time depends
27 on the strength of the exchange flow. The water age decreases by approximately 20 days when
28 the landward residual velocity in the main Bay increases by approximately 8 cm/s. Diagnostic
29 analyses suggest that the low river discharge and strong along-Bay pressure gradient facilitate
30 the up-Bay transport of dissolved substance coming from the lower Bay.

1 The time needed for the contaminants in the Harbor to be transported to the main Bay
2 during a typical low-discharge period is about 1 month longer than during a typical high-
3 discharge period. Because of the development of the three-layered circulation, dissolved
4 substances usually remain the longest in the mid-depth of the Harbor. The maximum tracer
5 concentration occurs at around 6 m depth in the area close to the mouth of the Harbor. As soon
6 as the dissolved substances are transported out of the Harbor, water mass follows two different
7 pathways. When being transported toward the downstream of the Bay, high tracer concentrations
8 occur near the surface. Conversely, when being transported toward the upstream of the Bay, high
9 tracer concentrations occur near the bottom.

10 The long-term mean transport processes, which mainly depend on the low frequency and mean
11 motion of the water, are essential to evaluate water exchange between the Harbor and the Bay.
12 This study indicates that the water age and tracer concentration in the Harbor are sensitive to the
13 perturbations of the Susquehanna River discharge. In reality, the perturbation can also be
14 induced by the wind forcing, and affect the water exchange over shorter timescales. The
15 incorporation of the wind effect will be covered in our future work. As the freshwater discharge
16 dominates stratification in the upper Bay, the results presented in this study represents the
17 seasonal density-driven exchange flow between the Harbor and the Bay. The approach used here
18 can be used to study the water exchange processes in other estuaries and harbors.

20 **Acknowledgements**

21 The funding of the project is provided by the Maryland Department of Environment. This is the
22 contribution number #xxx of Virginia Institute of Marine Sciences, College of William and Mary.
23 This work was performed using computational facilities at the College of William and Mary which were
24 enabled by grants from Sun Microsystems, the National Science Foundation, and Virginia's
25 Commonwealth Technology Research Fund.

27 **References**

- 28
29 Austin, J.A., 2002. Estimating the mean ocean-bay exchange rate of the Chesapeake Bay. *Journal of*
30 *Geophysical Research* 110 (C11), 3192.
31
32 Boicourt, W.C., and Olson, P., 1982. A hydrodynamic study of the Baltimore Harbor system. Tech. Rep.
33 82-10, Chesapeake Bay Institute, The Johns Hopkins University, 131 pp.

- 1
2 Cameron, W.M., and Pritchard, D.W., 1963. Estuaries, p.306-324. *In* The Sea – Vol. 2. Ed. By M.N. Hill.
3 Interscience, New York, N.Y., 757 pp.
4
- 5 Carpenter, J. H. 1960. The Chesapeake Bay Institute study of the Baltimore Harbor, p. 62-78. In Proc.
6 33rd Ann. Conf. Maryland-Delaware Water and Sewage Assoc.
7
8
- 9 Chao, S.-Y., Boicourt, W.C., Wang, H. V., 1996. Three layered circulation in reverse estuaries.
10 Continental Shelf Research 16, 1379-1397.
11
- 12 Deleersnijder, E., Campin, J., Delhez, E., 2001. The concept of age in marine modeling I. Theory and
13 preliminary model results. Journal of Marine Systems 28, 229-267.
14
- 15 Delhez, E. J. M., J.M. Campin, A.C. Hirst, E. Deleersnijder, 1999. Toward a general theory of the age in
16 ocean modeling. Ocean Modeling 1, 17-27.
17
- 18 Delhez, E. J. M., Deleersnijder, E., 2002. The concept of age in marine modeling II. Concentration
19 distribution function in the English Channel and the North Sea. Journal of Marine Systems 31, 279-297.
20
- 21 DÖÖs, K., Engqvist, A., 2007. Assessment of water exchange between a discharge region and the open
22 sea – A comparison of different methodological concepts. Estuarine, Coastal and Shelf Science 74,709-
23 721.
24
- 25 Geyer, W.R., Trowbridge, J.H., and Bowen, M.M., 2000. The dynamics of a partially mixed estuary.
26 Journal of Physical Oceanography 30, 2035-2048.
27
- 28 Gustafsson, K. E., Bendtsen, J., 2007. Elucidating the dynamics and mixing agents of a shallow fjord
29 through age tracer modeling. Estuarine, Coastal and Shelf Science 74, 641-654.
30
- 31 Hansen, D.V., and Rattray, Jr. M., 1965. Gravitational circulation in straits and estuaries. Journal of
32 Marine Research 23, 104-122.
33
- 34 Hirst, A. C., 1999. Determination of water component age in ocean models: application to the fate of
35 North Atlantic Deep Water. Ocean Modelling 1, 81-94.
36
- 37 Hood, R.R., Wang, H. V., Purcell, J.E., Houde, E.D., Harding Jr., L.W., 1999. Modeling particles and
38 pelagic organisms in Chesapeake Bay: Convergent features control plankton distributions. Journal of
39 Geophysical Research 104, 1223-1243.
40
- 41 Johnson, B.H. et al., 1991. User's guide for a three-dimensional numerical hydrodynamic model of
42 Chesapeake Bay. Department of the army, Waterways experiment station, Corps of Engineers.
43
- 44 Johnson, B. H., Kim, K.W., Heath, R.E., Hsieh, B.B., Butler, H. L., 1993. Validation of three-
45 dimensional hydrodynamic model of Chesapeake Bay. Journal of Hydraulic Engineer 119, 2-20.
46
- 47 Johnson, B. H., Wang, H. V., Kim, K. W., 1995. Can numerical estuarine models be driven at the estuary
48 mouth, in: ASCE Estuarine and Coastal Modeling, pp. 255-267, Am. Soc. Of Civ. Eng., New York.
49
- 50 Kemp, W.M., W.R. Boynton, J.E. Adolf, D.F. Boesch, W.C. Boicourt, G. Brush, J.C. Cornwell, T.R.
51 Fisher, P.M. Gilbert, J.D. Hagy, L. W. Harding, E.D. Houde, D.G. Kimmel, W.D. Miller, R.I.E. Newell,

- 1 M.R. Roman, E.M. Smith, and J.C. Stevenson, 2005. Eutrophication of Chesapeake Bay: historical trends
2 and ecological interactions. *Marine Ecology Progress Series* 30321: 1–29.
3
- 4 Kundu, P. K., 1980. A numerical investigation of mixed-layer dynamics, *Journal of Physical*
5 *Oceanography* 10, 220-236.
6
- 7 Kuo, A.Y., and B.J. Neilson, 1987. Hypoxia and salinity in Virginia Estuaries. *Estuaries* 104: 277–283,
8 doi:10.2307/1351884.
9
- 10 Lin, J., Wang, H. V., Sisson, G. M., Shen, J., 2004. Toxic modeling in an industrial Harbor – A case study
11 for Baltimore Harbor. *Proceedings of the Eight International Conference of Estuarine and coastal*
12 *modeling*, November 3-5, 2003, Monterey, CA, 455-474.
13
- 14 Liu, Y., P. MacCready, B. M. Hickey, E. P. Dever, P.M. Kosro, and N. S. Banas, 2009. Evaluation of a
15 coastal ocean circulation model for the Columbia River plume in 2004. *Journal of Geophysical Research*
16 114, C00B04, doi: 10.1029/2008JC004929.
17
- 18 MacCready, P., 2004. Toward a Unified Theory of Tidally-Averaged Estuarine Salinity
19 Structure. *Estuaries* 27, 561-570.
20
- 21 MacCready, P., 2007. Estuarine Adjustment. *Journal of Physical Oceanography* 37, 2133-2145.
22
- 23 MacCready, P., N. S. Banas, B. H. Jickey, E. P. Dever, Y. Liu, 2009. A model study of tide- and wind-
24 induced mixing in the Columbia River Estuary and plume. *Continental Shelf Research* 29, 278-291.
25
- 26 MacCready, P. and W.R. Geyer, 2010. Advances in Estuarine Physics. *Annual Review of Marine Science*
27 2, 35-58, 10.1146/annurev-marine-120308-081015.
28
- 29 Park, K., H. V. Wang, S. Kim, 2008. A model study of the estuarine turbidity maximum along the main
30 channel of the Upper Chesapeake Bay, *Estuaries and Coasts* 31, 115–133.
31
- 32 Prandle, D. A., 1984. A modeling study of the mixing of ¹³⁷Cs in the seas of European continent shelf.
33 *Philosophical Transactions of Royal Society of London* 310, 407-436.
34
- 35 Pritchard, D.W., 1956. The dynamic structure of a coastal plain estuary. *Journal of Marine Research*
36 15(I), 33-42.
37
- 38 Sanford, L.P., S.E. Suttles, J. P. Halka, 2001. Reconsidering the physics of the Chesapeake Bay estuarine
39 turbidity maximum. *Estuaries* 24, 655-669.
40
- 41 Schubel, J.R., Pritchard, D.W., 1986. Responses of upper Chesapeake bay to variations in discharge of
42 the Susquehanna River. *Estuaries* 9, 236-249.
43
- 44 Shen, J., Haas, L., 2004. Calculating age and residence time in the tidal York River using three-
45 dimensional model experiments. *Estuarine, Coastal and Shelf Science* 61, 449-461.
46
- 47 Shen, J., Wang, H. V., 2007. Determining the age of water and long-term transport timescale of the
48 Chesapeake Bay. *Estuarine, Coastal and Shelf Science* 74, 750-763.
49
- 50 Sheng, Y.P., 1986. A three-dimensional mathematical model of coastal estuarine and lake currents using a
51 boundary fitted grid, Rep. No. 585, ARAR Group of Titan Syst., Princeton, N.J..

1
2 Stroup, E. E., D. W. Pritchard, J. H. Carpenter, 1961. Final report, Baltimore Harbor Study. The John
3 Hopkins University Tech. Rep. XXVI, Ref. 61-5. Chesapeake Bay Institute, Baltimore, MD. 79p.

4
5 Wang, H. V., Chapman, R. S., 1995. Application of vertical turbulence closure Schemes in the
6 Chesapeake Bay Circulation Model – A comparative study, in ASCE Estuarine and Coastal Modeling, pp.
7 283-297, Am. Soc. Civ. Eng., New York.

8
9 Wang, H.V., H. Liu, K. Park. 2004. *The development of a water quality model for Baltimore Harbor,*
10 *Back River, and the adjacent upper Chesapeake Bay. A report to Maryland Department of the*
11 *Environment, SRAMSOE 386.* Gloucester Point, VA: Virginia Institute of Marine Science.

12
13 Xu, J., Chao, S.-Y., Hood, R.R., Wang, H., Boicourt, W.C., 2002. Assimilating high-resolution salinity
14 data into a model for a partially mixed estuary. *Journal of Geophysical Research* 107 (C7), 3074, doi:
15 10.1029/2000JC000626.

16
17 Xu, J., and Hood, R.R., 2006. Modeling biogeochemical cycles in Chesapeake Bay with a couple
18 physical-biological model. *Estuarine, Coastal and Shelf Science* 69, 19-46.

19
20 Zimmerman, J. T. F., 1976. Mixing and flushing of tidal embayments in the western Dutch Wadden sea,
21 Part I: distribution of salinity and calculation of mixing time scales. *Netherlands Journal of Sea Research*
22 10, 149-191.

23 24 25 **Figure Captions**

26
27 Figure 1. Bathymetry of upper Chesapeake Bay. The color gradient represents water depth in meters.

28
29 Figure 2. Model domain and boundary-fitted computational grids (101 X 153). The red bold lines denote
30 transects along the navigational channel of the Harbor (transect I) and across the mouth of the Harbor
31 (transect II) respectively. The star symbols represent stations where time series analyses are conducted.
32 The rectangle (Station V) in the main channel of the Bay is the sample site of modeled velocity profile.
33 The triangles represent CBP monitoring stations in the Bay.

34
35 Figure 3. Model-data comparisons. (a) Salinity results. The subscripts s and b represent surface
36 salinity and bottom salinity, respectively ; (b) low-pass filtered water elevation. The observations
37 (black line) are missing in some time intervals. The correlation coefficients (CC) in (b) are at the 95%
38 confidence level.

39
40 Figure 4. Long-term mean (1985-2007) daily freshwater discharge ($\times 10^3 \text{ m}^3 \text{ s}^{-1}$) from Susquehanna River.

41
42 Figure 5. (a) Typical longitudinal-vertical section of the salinity (ppt) distribution along the navigational
43 channel of the Baltimore Harbor, with the Chesapeake Bay lies in the right entrance of the figure. Arrows
44 indicate the schematic depiction of the three-layered circulation pattern in the Harbor (after Cameron and
45 Pritchard, 1963). (b) Modeled salinity (ppt) profile along transect I. (c) Modeled velocity (cm s^{-1}) profile
46 along section I. The positive (negative) values represent the outward (inward) flow. The abscissa is
47 consistent with model grid coordinates in Figure 2.

1 Figure 6. Surface salinity (ppt) in the upper Chesapeake Bay averaged during (a) the typical high-
 2 discharge period and (b) the typical low-discharge period. Zonal velocity (cm s^{-1}) profiles across the
 3 mouth of Baltimore Harbor (transect II in Fig. 2) averaged during (c) the typical high-discharge period
 4 and (d) the typical low-discharge period. The positive (negative) values represent the outward (inward)
 5 flow. The contour interval is 1 ppt for salinity field and 0.5 cm s^{-1} for the velocity field.

6
 7 Figure 7. Snapshot of surface water age (day) (Exp. 1 (a-b) and Exp. 2 (c-d)) and velocity (cm s^{-1}) (e-f) at
 8 Day 91 (left) and Day 255 (right). The contour interval is 4 days.

9
 10 Figure 8. Monthly mean vertical profile of water age (day) resulting from Exp.1 along the navigation
 11 channel of Baltimore Harbor (transect I in Fig. 2). The contour interval is 5 days. The abscissa is
 12 consistent with the model abscissa in Figure 2.

13
 14 Figure 9. Temporal vertical tracer concentration (arbitrary unit) (left, a-d) and water age (day) (right, e-h)
 15 profiles resulting from Exp. 1 at Stations C, D, E and B (from top to bottom), respectively. The contour
 16 interval for the tracer concentration and water age are 0.1 and 10 days, respectively.

17 Figure 10. Same as Fig. 8, but for the results of Exp. 2.

18
 19 Figure 11. Same as Fig. 9, but for the results of Exp. 2.

20
 21 Figure 12. Temporal evolution of total tracer concentration (arbitrary unit) in Baltimore Harbor under
 22 high-flow (thin line) and low-flow (bold line) condition, respectively. The dashed lines indicate where the
 23 total tracer concentration is reduced to a factor of e^{-1} (i.e., measure of the turnover time of the Harbor
 24 waters).

25
 26 Figure 13. Evolution of tracer concentration (arbitrary unit) in the case of releasing tracers instantly in
 27 Baltimore Harbor under typical high-discharge condition (Exp. 3). Chesapeake Bay lies to the right of the
 28 figure. The abscissa is consistent with the model abscissa in Figure 2.

29
 30 Figure 14. Time series of depth-averaged tracer concentration under (a) typical high-discharge condition
 31 and (b) typical low-discharge condition. The temporal vertical profiles of tracer concentration under
 32 typical low-discharge condition are shown in (c). In each row, the results at Stations f, g, and h are
 33 presented from the left to the right.

34
 35 Figure 15. The temporal vertical profile of (a) observed salinity (ppt) at Station CB4.4; (b) water age
 36 (day) at Station C (Exp. 2); (c) water age at Station B (Exp. 2); and (d) residual velocity (cm s^{-1}) at Station
 37 V. The negative (positive) velocity represents the seaward (landward) flow.

38
 39 Figure 16. The plane ($uE/c, \Phi$) presentation of all possible estuary states. Φ and uE/c denotes the
 40 dimensionless stratification and exchange flow, respectively. The solid line represents the results
 41 calculated using theoretical solution, the triangle represents the results calculated using the 3D model.

42 43 44 **Table**

45 Table 1. A summary of the tracer age results in the Baltimore Harbor.

Description	Exp. 1	Exp. 2	Exp. 3
Vertical age distribution pattern	Low age occurs at both surface and bottom, and	Decreases consistently with depth	Larger in the mid-depth than those in other areas

	high age occurs in mid-depth		
Vertical age difference (VAD) in the water column	The highest (lowest) VAD can be about 50 (20) days and appears in winter (April)	VAD is about 40-50 days. larger than bottom layer	Not available
Age variations from high-discharge condition to low-discharge condition	Increases about 2 months	Decreases about 20 days	The turnover time of the Harbor is 21.7 days (39.5 days) under the high-discharge (low-discharge) condition
Tracer concentration (TC) distribution pattern	Decrease with depth. Highest TC appears in April	Increase with depth. Lowest TC appears in April	After instantaneous release, high TC remains for a longer time in the mid-depth than those in other areas

1




Figure 1. Bathymetry of upper Chesapeake Bay. The color gradient represents water depth in meters.




Figure 2. Model domain and boundary-fitted computational grids (101 X 153). The red bold lines denote transect along the navigational channel of the Harbor (transect I) and across the mouth of the Harbor (transect II) respectively. The star symbols represent stations where time series analyses are conducted. The rectangle (Station V) in the main channel of the Bay is the sample site of modeled velocity profile. The triangles represent CBP monitoring stations in the Bay.




Figure 3. Model-data comparisons. (a) Salinity results. The subscripts *s* and *b* represent surface salinity and bottom salinity, respectively; (b) low-pass filtered water elevation. The observations (black line) are missing in some time intervals. The correlation coefficients (CC) in (b) are at the 95% confidence level.




Figure 4. Long-term mean (1985-2007) daily freshwater discharge ($\times 10^3 \text{ m}^3 \text{ s}^{-1}$) from Susquehanna River.




Figure 5. (a) Typical longitudinal-vertical section of the salinity (ppt) distribution along the navigational channel of the Baltimore Harbor; Chesapeake Bay lies to the right of the figure. Arrows indicate the schematic depiction of the three-layered circulation pattern in the Harbor (after Cameron and Pritchard, 1963). (b) Modeled salinity (ppt) profile along transect I. (c) Modeled velocity (cm s⁻¹) profile along transect I. The positive (negative) values represent the outward (inward) flow. The abscissa in (b) and (c) is consistent with the model abscissa in Figure 2.




Figure 6. Surface salinity (ppt) in the upper Chesapeake Bay averaged during (a) the typical high-discharge period and (b) the typical low-discharge period. Zonal velocity (cm s⁻¹) profiles across the mouth of Baltimore Harbor (transect II in Fig. 2) averaged during (c) the typical high-discharge period and (d) the typical low-discharge period. The positive (negative) values represent the outward (inward) flow. The contour interval is 1 ppt for salinity field and 0.5 cm s⁻¹ for the velocity field.




Figure 7. Snapshot of surface water age (day) (Exp. 1 (a-b) and Exp. 2 (c-d)) and velocity (cm s^{-1}) (e-f) at Day 91 (left) and Day 255 (right). The contour interval is 4 days.




Figure 8. Monthly mean vertical profile of water age (day) resulting from Exp.1 along the navigational channel of Baltimore Harbor (transect I in Fig. 2). The contour interval is 5 days. The abscissa is consistent with the model abscissa in Figure 2.




Figure 9. Temporal vertical tracer concentration (arbitrary unit) (left, a-d) and water age (day) (right, e-h) profiles resulting from Exp. 1 at Stations C, D, E, and B (from top to bottom), respectively. The contour intervals for the tracer concentration and water age are 0.1 and 10 days, respectively.




Figure 10. Same as Fig. 8, but for the results of Exp.2.




Figure 11. Same as Fig. 9, but for the results of Exp.2.




Figure 12. Temporal evolution of total tracer concentrations (arbitrary unit) in Baltimore Harbor under high-flow (thin line) and low-flow (bold line) condition, respectively. The dashed lines indicate where the total tracer concentration is reduced to a factor of e^{-1} (i.e., measure of the turnover time of the Harbor waters).




Figure 13. Evolution of tracer concentrations (arbitrary unit) in the case of releasing tracers instantly in Baltimore Harbor under typical wet condition (Exp. 3). Chesapeake Bay lies to the right of the figure. The abscissa is consistent with the model abscissa in Figure 2.




Figure 14. Time series of depth-averaged tracer concentration (arbitrary unit) under (a) typical wet and (b) typical dry conditions. The temporal vertical profiles of tracer concentrations under typical dry condition (the same pattern occurs during the typical wet condition) are shown in (c). In each row, the results at Stations f, g, and h are presented from the left to the right.




Figure 15. The temporal vertical profile of (a) observed salinity (ppt) at Station CB4.4; (b) water age (day) at Station C (Exp. 2); (c) water age at Station B (Exp. 2); and (d) residual velocity (cm s^{-1}) at Station V. The negative (positive) velocity represents the seaward (landward) flow.




Figure 16. The plane (u_E/c , Φ) presentation of all possible estuary states. Φ and u_E/c denotes the dimensionless stratification and exchange flow, respectively. The solid line represents the results calculated using theoretical solution, the triangle represents the results calculated using the 3D model.