



**HAL**  
open science

## On multivariate dispersion orderings based on the standard construction

Félix Belzunce, José M. Ruiz, Alfonso Suárez-Llorens

► **To cite this version:**

Félix Belzunce, José M. Ruiz, Alfonso Suárez-Llorens. On multivariate dispersion orderings based on the standard construction. *Statistics and Probability Letters*, 2009, 78 (3), pp.271. 10.1016/j.spl.2007.07.001 . hal-00598961

**HAL Id: hal-00598961**

**<https://hal.science/hal-00598961>**

Submitted on 8 Jun 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Author's Accepted Manuscript

On multivariate dispersion orderings based on the standard construction

Félix Belzunce, José M. Ruiz, Alfonso Suárez-Llorens


PII: S0167-7152(07)00238-6  
DOI: doi:10.1016/j.spl.2007.07.001  
Reference: STAPRO 4704

To appear in: *Statistics & Probability Letters*

Received date: 10 November 2006  
Revised date: 2 April 2007  
Accepted date: 24 July 2007

Cite this article as: Félix Belzunce, José M. Ruiz and Alfonso Suárez-Llorens, On multivariate dispersion orderings based on the standard construction, *Statistics & Probability Letters* (2007), doi:10.1016/j.spl.2007.07.001

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



[www.elsevier.com/locate/stapro](http://www.elsevier.com/locate/stapro)

# On multivariate dispersion orderings based on the standard construction

Félix Belzunce, José M. Ruiz  
Dept. Estadística e I. O., Univ., Murcia (Spain).  
Alfonso Suárez-Llorens \*  
Dept. Estadística e I.O., Univ., Cádiz (Spain).

April 2, 2007

## Abstract

We study the relationship between the multivariate dispersive orders based on the standard construction. In particular those given by Shaked and Shanthikumar (1998) and Fernández-Ponce and Suárez-Llorens (2003). In order to reach our objective we define a new weaker multivariate dispersive notion. Random vectors with a common copula and positive dependence properties are analyzed.

**AMS Subject Classification:** 60E15, 60E05, 60K10.

**Key words and phrases:** univariate and multivariate dispersive orders, st:icx order, convolutions, models of ordered random variables.

## 1 Introduction

In the context of stochastic orders, several orders have been defined to compare two univariate random quantities in terms of their variability or dispersion. In particular, one of the most widely used in the literature is the dispersive order introduced by Lewis and Thompson (1981), see Shaked and Shanthikumar (1994) as an excellent handbook to study all its properties.

Based on the properties and characterizations of the univariate dispersive order several authors have proposed different multivariate extensions, namely Shaked and Shanthikumar (1998) and Fernández-Ponce and Suárez-Llorens (2003) propose multivariate dispersive orders based on the standard construction.

In this paper we analyzed the relationship between these multivariate dispersion notions. The organization is the following. First in section 2 we present some well known characterizations of the univariate dispersion order and discuss how they have been used to define several multivariate dispersion orderings under the standard construction. Second in Section 3 we propose a new multivariate dispersion order which will be used later in Section 3 to relate the multivariate dispersion orders defined in Section 2. Third in Section 4 we show the relationship among the multivariate dispersion concepts introduced in the previous sections. In particular some interesting results are obtained for random vectors with a

---

\*Corresponding author: Tel: (+34) 956015481, fax: (+34) 956015378 <alfonso.suarez@uca.es>

common copula and positive dependence properties. Finally, in Section 5 we present how those relationships can be applied and used to simplify some known results in the literature.

In this paper for any random variable  $X$  and an event  $A$ , we let  $\{X|A\}$  denote any random variable whose distribution is the conditional distribution of  $X$  given  $A$ . Expected values are assumed to exist whenever they are mentioned. By  $=_{st}$  we denote equality in law.

## 2 The univariate dispersion order: characterizations and generalizations

A formal definition of the univariate dispersive order is based on the notion of quantile. Given a random variable  $X$  with distribution function  $F$ , we define the *univariate quantile* as  $Q_X(u) \equiv F_X^{-1}(u) = \inf\{x : F_X(x) \geq u\}$ , for all real value  $u \in (0, 1)$  (for definition and characterizations of the dispersive order see Section 2.B.2 in Shaked and Shanthikumar (1994)). Given two random variables  $X$  and  $Y$ , we say that  $X$  is less in the dispersive order than  $Y$ , denoted by  $X \leq_{disp} Y$ , if

$$Q_X(v) - Q_X(u) \leq Q_Y(v) - Q_Y(u) \text{ for all } 0 < u \leq v < 1. \quad (1)$$

Clearly this is a dispersive order notion because it requires the distance between any two quantiles of  $X$  to be less separated than the corresponding quantiles of  $Y$ .

An interesting characterization is given in terms of expansion functions. A real valued function  $\phi$  is said to be an expansion function if  $\phi(x') - \phi(x) \geq x' - x$  whenever  $x' \geq x$ . If  $\phi$  is differentiable then  $\phi$  is an expansion function if  $d\phi(x)/dx \geq 1$  for all  $x$ . We have that  $X \leq_{disp} Y$  if, and only if,

$$Y =_{st} \phi(X) \text{ for some expansion function } \phi. \quad (2)$$

Also, from the definition, it is easy to see that  $X \leq_{disp} Y$  if, and only if,

$$\phi(x) = Q_Y(F_X(x)) \text{ is an expansion function.} \quad (3)$$

In addition, if the distribution functions of  $X$  and  $Y$  are strictly increasing, then the function  $\phi$  defined in (3), is the only one that satisfies (2). To finalize we present another characterization which we will use later. Given two random variables  $X$  and  $Y$ , with absolutely continuous distribution functions, and density functions  $f_X$  and  $f_Y$ , then  $X \leq_{disp} Y$  if, and only if,

$$f_X(Q_X(u)) \geq f_Y(Q_Y(u)), \text{ for all } u \in (0, 1). \quad (4)$$

Based on these characterizations several authors have proposed different extensions to the multivariate case. Important contributions in this case have been made by Oja (1983) and Giovagnoli and Wynn (1995). Clearly inspired in (2), those authors define multivariate dispersion orders through the existence of a multivariate function  $k$  which maps stochastically a random vector  $\mathbf{X}$  to another one  $\mathbf{Y}$ , that is  $\mathbf{Y} =_{st} k(\mathbf{X})$ . It is well known that there are different transformations that map a multivariate random vector to another one. For this reason Shaked and Shanthikumar (1998) and Fernández-Ponce and Suárez-Llorens (2003) consider a particular one based on the standard construction. First we need some definitions. From now on, we will assume that the multivariate distribution function is an absolutely continuous function.

Let  $\mathbf{X}$  be a random vector and let  $\mathbf{u} = (u_1, \dots, u_n)$  in  $[0, 1]^n$ . The standard construction for  $\mathbf{X}$ , denoted by

$$\hat{x}(\mathbf{u}) = (\hat{x}_1(u_1), \hat{x}_2(u_1, u_2), \dots, \hat{x}_n(u_1, \dots, u_n)),$$

is defined as follows

$$\begin{aligned} \hat{x}_1(u_1) &= Q_{X_1}(u_1) \\ \hat{x}_i(u_1, \dots, u_i) &= Q_{\{X_i | \bigcap_{j=1}^{i-1} X_j = \hat{x}_j(u_j)\}}(u_i), \text{ for } i = 2, \dots, n. \end{aligned}$$

This known construction is widely used in simulation theory and plays the role of the quantile in the multivariate case. The following result, it is well known (see Li, Scarsini and Shaked (1996))

$$\hat{x}(\mathbf{U}) =_{st} \mathbf{X}, \quad (5)$$

where  $\mathbf{U}$  is a random vector with  $n$  independent uniform distributed components on  $[0, 1]$ .

By ‘‘inverting’’ the standard construction, we can express the independent uniform random variables  $U_i$ ’s as functions of the  $X_i$ ’s. Let us denote

$$\hat{x}^*(\mathbf{x}) = (\hat{x}_1^*(x_1), \dots, \hat{x}_n^*(x_1, \dots, x_n))$$

as the vector given by

$$\begin{aligned} \hat{x}_1^*(x_1) &= F_{X_1}(x_1), \\ \hat{x}_i^*(x_1, \dots, x_i) &= F_{\{X_i | \bigcap_{j=1}^{i-1} X_j = x_j\}}(x_i), \text{ for } i = 2, \dots, n. \end{aligned}$$

It is also well known, see for instance Shaked and Shantikumar (1998) that

$$\hat{x}^*(\mathbf{X}) =_{st} \mathbf{U}. \quad (6)$$

Let us consider the  $n$ -dimensional function  $\phi(\mathbf{x}) = (\phi_1(x_1), \dots, \phi_n(x_1, \dots, x_n))$  defined as

$$\begin{aligned} \phi_1(x_1) &= (\hat{y}_1 \circ \hat{x}_1^*)(x_1) = Q_{Y_1}(F_{X_1}(x_1)) \quad (7) \\ \phi_i(x_1, \dots, x_i) &= (\hat{y}_i \circ \hat{x}_i^*)(x_1, \dots, x_i) = Q_{\{Y_i | \bigcap_{j=1}^{i-1} Y_j = \phi_j(x_1, \dots, x_j)\}}(F_{\{X_i | \bigcap_{j=1}^{i-1} X_j = x_j\}}(x_i)), \quad (8) \end{aligned}$$

for  $i = 2, \dots, n$ .

It is clear from (5) and (6) that  $\phi$  maps stochastically  $\mathbf{X}$  to  $\mathbf{Y}$ , that is  $\phi(\mathbf{X}) =_{st} \mathbf{Y}$ . It is also apparent that the standard construction can be seen as a generalization of the univariate quantile function.

Let  $\mathbf{X}$  and  $\mathbf{Y}$  be two random vectors on  $\mathbb{R}^n$  and let  $\hat{x}(\cdot)$ ,  $\hat{y}(\cdot)$  the corresponding standard constructions. Shaked and Shantikumar (1998) considered the following condition

$$\hat{y}(\mathbf{u}) - \hat{x}(\mathbf{u}) \text{ is increasing in } \mathbf{u} \in (0, 1)^n, \quad (9)$$

as a multivariate generalization of (1). To maintain a coherent notation, we will say that  $\mathbf{X}$  is less than  $\mathbf{Y}$  in variability if (9) holds, and we will denote it as  $\mathbf{X} \leq_{var} \mathbf{Y}$ . Note that (9) can be rewritten in terms of the function  $\phi$  defined in (7) and (8) as

$$\phi(\hat{x}(\mathbf{u})) - \hat{x}(\mathbf{u}) \text{ is increasing in } \mathbf{u} \in (0, 1)^n.$$

Another multivariate generalization based on the standard construction was given recently by Fernández-Ponce and Suárez-Llorens (2003). Given two  $n$ -dimensional random vectors  $\mathbf{X}$  and  $\mathbf{Y}$ , we say that  $\mathbf{X}$  is less in the *multivariate dispersive order* than  $\mathbf{Y}$ , denoted by  $\mathbf{X} \leq_{disp} \mathbf{Y}$ , if, and only, if

$$\|\hat{x}(\mathbf{v}) - \hat{x}(\mathbf{u})\|_2 \leq \|\hat{y}(\mathbf{v}) - \hat{y}(\mathbf{u})\|_2$$

for all  $\mathbf{u}, \mathbf{v} \in [0, 1]^n$ , where  $\|\cdot\|_2$  means the Euclidean distance. Note that the last inequality can be considered as a multivariate extension of (1). Also Fernández-Ponce and Suárez-Llorens (2003) proved that the  $\leq_{disp}$  order is equivalent to check if the function  $\phi$  defined in (7) and (8) is a multivariate expansion function, where multivariate expansion means that  $\|\phi(\mathbf{x}) - \phi(\mathbf{x}')\|_2 \geq \|\mathbf{x} - \mathbf{x}'\|_2$  for all  $\mathbf{x}, \mathbf{x}'$  in  $\mathbb{R}^n$ , and therefore can be also seen as a multivariate generalization of (3).

### 3 The Conditional Dispersive Order

In this section we present a new multivariate dispersion order as a generalization of (3) for the univariate case. The purpose of this order is to relate the  $\leq_{disp}$  and  $\leq_{var}$  orders as we will see later on in Section 4. On the other hand it will also give a meaningful interpretation in terms of dispersion to both orders.

**Definition 1.** Let  $\mathbf{X}$  and  $\mathbf{Y}$  be two  $n$ -dimensional random vectors. We consider the function  $\phi$  defined in (7) and (8) which maps  $\mathbf{X}$  onto  $\mathbf{Y}$ . We say that  $\mathbf{X}$  is less in the *conditional dispersive order* than  $\mathbf{Y}$ , denoted by  $\mathbf{X} \leq_{c-disp} \mathbf{Y}$ , if  $\phi_i(x_1, \dots, x_i)$  is an expansion function in  $x_i$  in the univariate sense, for all  $i = 1, \dots, n$ .

Roughly speaking, the conditional dispersive order is defined through a function which maps stochastically a vector to another one and satisfies that  $i$ -th component is an expansion function in  $x_i$  when  $x_1, \dots, x_{i-1}$  remains fixed. We observe that from (7), (8) and (3) we have that  $\phi_1(x_1)$  is an expansion function in  $x_1$  if, and only if,

$$X_1 \leq_{disp} Y_1 \tag{10}$$

and for fixed  $(x_1, \dots, x_{i-1}) \in \mathbb{R}^{i-1}$ ,  $\phi_i(x_1, \dots, x_i)$  is an expansion function in  $x_i$ , if, and only if,

$$\left\{ X_i \left| \bigcap_{j=1}^{i-1} X_j = \hat{x}_j(u_1, \dots, u_j) \right. \right\} \leq_{disp} \left\{ Y_i \left| \bigcap_{j=1}^{i-1} Y_j = \hat{y}_j(u_1, \dots, u_j) \right. \right\}, \tag{11}$$

for  $i = 2, \dots, n$  and for all  $u_i$  such that  $0 < u_i < 1$ ,  $i = 1, \dots, n$ .

Therefore the conditional dispersive order can be checked by the univariate dispersive order. The conditions (10) and (11) provide a geometrical interpretation for the  $\leq_{c-disp}$  order which we present in the bivariate case.

Fernández-Ponce and Suárez-Llorens (2003) provided the notion of corrected orthant associated with the standard construction and studied the accumulated probability in all of them. Let  $\mathbf{X} = (X_1, X_2)$  be a bivariate random vector and let  $(u_1, u_2)$  in  $[0, 1]^2$ . The intersection between the curve  $x_2 = \hat{x}_2(\cdot, u_2)$ , where  $u_2$  is fixed, and the line  $x_1 = \hat{x}_1(u_1)$  is achieved at the point  $\hat{x}(u_1, u_2) = (\hat{x}_1(u_1), \hat{x}_2(u_1, u_2))$  and this intersection provides four corrected orthants ( $2^n$  on  $\mathbb{R}^n$ ) which we represent in Figure 1.

In order to clarify the understanding of the corrected orthant concept, we show the definition of the left lower corrected orthant one

$$\{(x_1, x_2) : x_1 \leq \hat{x}_1(u_1), x_2 \leq \hat{x}_2(F_{X_1}(x_1), u_2)\}.$$

The result states that the accumulated probability in all corrected orthants depend on  $u_1$  and  $u_2$  (see Figure 1). Note that the conditional function  $x_2 = \hat{x}_2(\cdot, u_2)$  represents the extreme behavior of the dependent variable  $X_2$  conditional on the explanatory variable  $X_1$ .




Figure 1: The four corrected orthants and accumulated probabilities.

Let us consider now  $(u_1, u_2)$  and  $(v_1, v_2) \in [0, 1]^2$  where  $u_i < v_i$ ,  $i = 1, 2$ . According to the previous discussion is easy to see that Figure 2 represents a bivariate central region for the random vector  $\mathbf{X}$  given by the conditional function. Let consider now  $\mathbf{Y} = (Y_1, Y_2)$  such that  $\mathbf{X} \leq_{c-disp} \mathbf{Y}$ . Then it holds by (10) and (11) that

$$\begin{aligned} \hat{x}_1(v_1) - \hat{x}_1(u_1) &\leq \hat{y}_1(v_1) - \hat{y}_1(v_1) \\ \hat{x}_2(u, v_2) - \hat{x}_2(u, u_2) &\leq \hat{y}_2(u, v_2) - \hat{y}_2(u, u_2), \end{aligned}$$

for all  $0 \leq u_i \leq v_i \leq 1$ ,  $i = 1, 2$ ,  $u \in [0, 1]$ . Then is easy to note that the central region for  $\mathbf{X}$  represented in Figure 2 is less widely separated than the corresponding for  $\mathbf{Y}$ . Therefore the interpretation of  $\leq_{c-disp}$  in terms of dispersion is clear: we have found a wider region that accumulates the same probability.




Figure 2: Central Region for  $\mathbf{X}$

To continue with the interpretation we observe that given an  $n$ -dimensional random

vector  $\mathbf{X}$  with density function  $f_{\mathbf{X}}$  then

$$f_{\mathbf{X}}(\hat{x}(\mathbf{u})) = f_1(\hat{x}(u_1)) \prod_{i=2}^n f_{i|1,\dots,i-1}(\hat{x}_i(u_1, \dots, u_{i-1})),$$

for all  $\mathbf{u} \in [0, 1]^n$ , where  $f_1$  is the density function of  $X_1$  and  $f_{i|1,\dots,i-1}$  is the density function of  $\left( X_i \left| \bigcap_{j=1}^{i-1} X_j = \hat{x}_j(u_1, \dots, u_j) \right. \right)$ . Therefore from (10), (11) and (4), it is clear that if  $\mathbf{X} \leq_{c-disp} \mathbf{Y}$  then

$$f_{\mathbf{X}}(\hat{x}(\mathbf{u})) \geq f_{\mathbf{Y}}(\hat{y}(\mathbf{u})), \forall \mathbf{u} \in [0, 1]^n.$$

Hence the multivariate density functions evaluated at standard constructions are ordered as a clear generalization of (4).

We observe that  $\leq_{c-disp}$  is not invariant under the permutation of the marginal distributions because the standard construction depends on the choice of the ordering of the marginal distributions. This is also the case of the  $\leq_{disp}$  and  $\leq_{var}$  orders. On the other hand is easily to prove that  $\leq_{c-disp}$  is location invariant.

The  $\leq_{c-disp}$  order can be also interpreted in terms of local volume elements. Let  $\mathbf{X}$  and  $\mathbf{Y}$  two  $n$ -dimensional random vectors. Oja (1983) defined that  $\mathbf{Y}$  is more scattered than  $\mathbf{X}$ , denoted by  $\mathbf{X} \leq_{\Delta} \mathbf{Y}$ , if there is a function  $k: \mathbb{R}^n \rightarrow \mathbb{R}^n$  such that  $\mathbf{Y} =_{st} k(\mathbf{X})$  and for all  $\{\mathbf{x}_1, \dots, \mathbf{x}_{n+1}\} \subset \mathbb{R}^n$  it holds that

$$\Delta(k(\mathbf{x}_1), \dots, k(\mathbf{x}_{n+1})) \geq \Delta(\mathbf{x}_1, \dots, \mathbf{x}_{n+1}), \quad (12)$$

where  $\Delta(\mathbf{x}_1, \dots, \mathbf{x}_{n+1})$  is the volume of the ‘‘simplex’’ with vertices at  $\mathbf{x}_1, \dots, \mathbf{x}_{n+1}$ .

Note that although condition (12) seems to be a strict one, in practice it is only necessary to check if the absolute value of the determinant of the Jacobian matrix of  $k$  is bigger than 1 (see Giovagnolo and Wynn (1995)), i.e.

$$\text{abs}(\text{Det}(J_k(\mathbf{x}))) \geq 1, \text{ for all } \mathbf{x} \in \mathbb{R}^n. \quad (13)$$

Let us consider two  $n$ -dimensional random vectors  $\mathbf{X}$  and  $\mathbf{Y}$  such that  $\mathbf{X} \leq_{c-disp} \mathbf{Y}$ . From the characterization of the univariate expansion functions it easily holds that  $\partial\phi_i/\partial x_i \geq 1$  for  $i = 1, \dots, n$ . Due to the fact that  $\phi$  has a lower triangular Jacobian matrix it is apparent that  $\text{Det}(J_{\phi}) \geq 1$ . Hence from (13) it holds that  $\leq_{c-disp} \Rightarrow \leq_{\Delta}$ .

Fernández-Ponce and Suárez-Llorens (2003) proved in Theorem 3.1 that if we take a function  $k$  such that  $\mathbf{Y} =_{st} k(\mathbf{X})$  and  $k$  has a lower triangular matrix with diagonal elements strictly positive then  $k$  has the form of the function  $\phi$ . From this result there are many possible distributions which can be compared in the  $\leq_{c-disp}$  ordering. If we consider a random vector  $\mathbf{X} = (X_1, \dots, X_n)$  and a transformation  $\mathbf{Y} = \phi(\mathbf{X})$ , such that  $Y_i = f_i(X_i) + g_i(X_1, \dots, X_{i-1})$ , where  $f_i$  is an strictly increasing differentiable expansion function and  $g_i$  is any differentiable function for all  $i = 1, \dots, n$ , then it is apparent that  $\mathbf{X} \leq_{c-disp} \mathbf{Y}$ . As a particular case, this result holds for  $f_i(x) = a_i x$  where  $a_i > 1$ , for  $i = 1, \dots, n$ .

To finish this section we present a result for the preservation of the conditional dispersive order under conjunction of independent random vectors ordered in the conditional dispersive order.



**Theorem 1.** Let  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_m$  be a set of independent random vectors where the dimension of  $\mathbf{X}_i$  is  $n_i$ ,  $i = 1, 2, \dots, m$  and let  $\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_m$  be another set of independent random vectors where the dimension of  $\mathbf{Y}_i$  is  $n_i$ ,  $i = 1, 2, \dots, m$ . If  $\mathbf{X}_i \leq_{c\text{-disp}} \mathbf{Y}_i$  for  $i = 1, 2, \dots, m$  then

$$(\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_m) \leq_{c\text{-disp}} (\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_m).$$

## 4 The relationships among $\leq_{disp}$ , $\leq_{var}$ and $\leq_{c\text{-disp}}$ orders

In this section we study the relationships among the  $\leq_{disp}$ ,  $\leq_{var}$  and  $\leq_{c\text{-disp}}$  orders. The relationship among the  $\leq_{disp}$  and conditions (10) and (11) was given by Fernández-Ponce and Suárez-Llorens (2003) (see Corollary 3.1), and therefore the following result holds.

**Theorem 2.** Let  $\mathbf{X}$  and  $\mathbf{Y}$  be two  $n$ -dimensional random vectors. If  $\mathbf{X} \leq_{disp} \mathbf{Y}$  then  $\mathbf{X} \leq_{c\text{-disp}} \mathbf{Y}$ .

We want to emphasize that Fernández-Ponce and Suárez-Llorens (2003) did not consider (10) and (11) as a multivariate dispersive order and they just studied the relationship among these conditions and the  $\leq_{disp}$  order.

Next we show that the variability order is stronger than the conditional dispersive order.

**Theorem 3.** Let  $\mathbf{X}$  and  $\mathbf{Y}$  be two  $n$ -dimensional random vectors. If  $\mathbf{X} \leq_{var} \mathbf{Y}$  then  $\mathbf{X} \leq_{c\text{-disp}} \mathbf{Y}$ .

*Proof.* Let  $\mathbf{X}$  and  $\mathbf{Y}$  be two  $n$ -dimensional random vectors and let  $\hat{x}(\cdot)$ ,  $\hat{y}(\cdot)$  the corresponding standard constructions. In particular, condition (9) implies that

$$\hat{y}_i(u_1, u_2, \dots, u_i) - \hat{x}_i(u_1, u_2, \dots, u_i) \text{ is increasing in } u_i \in (0, 1), \text{ for } i = 1, 2, \dots, n.$$

Note that if we fix  $u_1, \dots, u_{i-1}$  then  $\hat{y}_i(u_1, \dots, u_i)$  and  $\hat{x}_i(u_1, \dots, u_i)$  are the univariate quantile functions of  $(X_i \mid \bigcap_{j=1}^{i-1} X_j = \hat{x}_j(u_1, \dots, u_j))$  and  $(Y_i \mid \bigcap_{j=1}^{i-1} Y_j = \hat{y}_j(u_1, \dots, u_j))$  respectively. Hence the proof follows from (10) and (11).  $\square$

Next we show that previous implications are strict and also we show the  $\leq_{disp}$  and  $\leq_{var}$  orders, in general, are not related.

**Example 1.** Let  $\mathbf{X} \sim N((0, 0), \Sigma_{\mathbf{X}})$  and  $\mathbf{Y} \sim N((0, 0), \Sigma_{\mathbf{Y}})$  be two bivariate normal distributions where

$$\Sigma_{\mathbf{X}} = \begin{pmatrix} \sigma_{1\mathbf{X}} & \sigma_{12\mathbf{X}} \\ \sigma_{12\mathbf{X}} & \sigma_{2\mathbf{X}} \end{pmatrix} \text{ and } \Sigma_{\mathbf{Y}} = \begin{pmatrix} \sigma_{1\mathbf{Y}} & \sigma_{12\mathbf{Y}} \\ \sigma_{12\mathbf{Y}} & \sigma_{2\mathbf{Y}} \end{pmatrix}.$$

Note that without lack of generality the mean vectors can be considered with components equal to zero. It is well known that the conditional distributions for multivariate normal distribution are also normal distributions. Then, the distribution of  $X_2$  conditioned to  $X_1$  is given by

$$(X_2 \mid X_1 = x_1) \sim N\left(x_1 \frac{\sigma_{12\mathbf{X}}}{\sigma_{1\mathbf{X}}}, \sigma_{2\mathbf{X}} \sqrt{(1 - \rho_{\mathbf{X}}^2)}\right),$$

where  $\rho_{\mathbf{X}}$  represents the correlation coefficient of  $X_1$  and  $X_2$ . Analogously for  $Y_2$  conditioned to  $Y_1$ .

First, we consider the  $\leq_{c-disp}$  ordering. It is a well known result that two univariate normal distributions are ordered in the univariate dispersion sense if, and only if their variances are ordered. Hence using (10) and (11) it easily holds that  $\mathbf{X} \leq_{c-disp} \mathbf{Y}$  if, and only if

$$\begin{aligned} \sigma_{1\mathbf{Y}} &\geq \sigma_{1\mathbf{X}} \\ \sigma_{2\mathbf{Y}}\sqrt{1-\rho_{\mathbf{Y}}^2} &\geq \sigma_{2\mathbf{X}}\sqrt{1-\rho_{\mathbf{X}}^2}. \end{aligned} \quad (14)$$

Now we consider the  $\leq_{disp}$  order. Fernández-Ponce and Suárez-Llorens (2003) studied the Jacobian matrix of the function  $\phi$  given in (7) and (8) for the bivariate normal distribution and they obtained a  $2 \times 2$  lower triangular matrix of the form

$$J_\phi = \begin{pmatrix} \frac{\sigma_{1\mathbf{Y}}}{\sigma_{1\mathbf{X}}} & 0 \\ \frac{\sigma_{2\mathbf{Y}}}{\sigma_{1\mathbf{X}}} \left( \rho_{\mathbf{Y}} - \rho_{\mathbf{X}} \frac{\sqrt{(1-\rho_{\mathbf{Y}}^2)}}{\sqrt{(1-\rho_{\mathbf{X}}^2)}} \right) & \frac{\sigma_{2\mathbf{Y}}\sqrt{(1-\rho_{\mathbf{Y}}^2)}}{\sigma_{2\mathbf{X}}\sqrt{(1-\rho_{\mathbf{X}}^2)}} \end{pmatrix}$$

Giovagnoli and Wynn (1995) characterized a multivariate expansion function in terms of its Jacobian matrix. Let  $k(\cdot)$  be a continuously differentiable function. Then  $k(\cdot)$  is a multivariate expansion function if, and only if the matrix  $J_k(\mathbf{x})^t J_k(\mathbf{x}) - I_n$  is nonnegative definite  $\forall \mathbf{x} \in \mathbb{R}^n$ , where  $J_k = \left\{ \frac{\partial k_i}{\partial x_j} \right\}$  is the Jacobian matrix of  $k(\cdot)$  and  $I_n$  is the identity matrix of order  $n$ . Under a straightforward computation of the eigenvalues of the matrix  $J_\phi(\mathbf{x})^t J_\phi(\mathbf{x}) - I_2$  we obtain that this one is nonnegative definite if, and only if the inequalities given by (14) hold and

$$\sigma_{2\mathbf{Y}}^2 \left( \rho_{\mathbf{Y}}\sqrt{1-\rho_{\mathbf{X}}^2} - \rho_{\mathbf{X}}\sqrt{1-\rho_{\mathbf{Y}}^2} \right)^2 \leq (\sigma_{1\mathbf{Y}}^2 - \sigma_{1\mathbf{X}}^2)(\sigma_{2\mathbf{Y}}^2(1-\rho_{\mathbf{Y}}^2) - \sigma_{2\mathbf{X}}^2(1-\rho_{\mathbf{X}}^2)). \quad (15)$$

Let us consider now the  $\leq_{var}$  order. Given that we deal with conditional normal distributions the standard construction for  $\mathbf{X}$  is easily given by  $\hat{x}_1(u_1) = Q_Z(u_1)\sigma_{1\mathbf{X}}$  and  $\hat{x}_2(u_1, u_2) = Q_Z(u_1)\rho_{\mathbf{X}}\sigma_{2\mathbf{X}} + Q_Z(u_2)\sigma_{2\mathbf{X}}\sqrt{(1-\rho_{\mathbf{X}}^2)}$  where  $Z$  is distributed as  $N(0, 1)$  and analogously for  $\mathbf{Y}$ . Then  $\mathbf{X} \leq_{var} \mathbf{Y}$  holds if, and only if the inequalities given by (14) hold and

$$\sigma_{2\mathbf{Y}}\rho_{\mathbf{Y}} \geq \sigma_{2\mathbf{X}}\rho_{\mathbf{X}}. \quad (16)$$

Note that from (16) the  $\leq_{var}$  ordering depends on the signs of the correlation coefficients.

Now we present some detailed cases in order to study the strict implications among the different dispersion orders. If we take  $\sigma_{i\mathbf{X}} = \sigma_{i\mathbf{Y}}$ , for  $i = 1, 2$ , then (14) holds if, and only if  $|\rho_{\mathbf{X}}| \geq |\rho_{\mathbf{Y}}|$ . If we take  $\rho_{\mathbf{X}} = -0,7$  and  $\rho_{\mathbf{Y}} = 0,4$ , then  $\mathbf{X} \leq_{c-disp} \mathbf{Y}$  holds and from (16)  $\mathbf{X} \leq_{var} \mathbf{Y}$  also holds. However from (15)  $\mathbf{X} \leq_{disp} \mathbf{Y}$  does not hold. Under the same assumption for the marginal variances and taking  $\rho_{\mathbf{X}} = 0,7$  and  $\rho_{\mathbf{Y}} = 0,4$ ,  $\mathbf{X} \leq_{c-disp} \mathbf{Y}$  holds. However from (15) and (16) neither  $\mathbf{X} \leq_{disp} \mathbf{Y}$  nor  $\mathbf{X} \leq_{var} \mathbf{Y}$  hold respectively. On the other hand if we take  $\rho_{\mathbf{X}} = \rho_{\mathbf{Y}}$  and  $\sigma_{i\mathbf{X}} < \sigma_{i\mathbf{Y}}$ , for  $i = 1, 2$ . Then from (14) and (15), both  $\mathbf{X} \leq_{c-disp} \mathbf{Y}$  and  $\mathbf{X} \leq_{disp} \mathbf{Y}$  hold respectively. However the  $\leq_{var}$  ordering depends on the signs of the correlation coefficients. In particular if we consider  $\rho_{\mathbf{X}} = \rho_{\mathbf{Y}} < 0$  does not hold.

To summarize we have the following chains of strict implications:

$$\begin{array}{c} \leq_{disp} \Rightarrow \leq_{c-disp} \Rightarrow \leq_{\Delta} \\ \uparrow \\ \leq_{var} \end{array}$$

Following Müller and Scarsini (2001), Khakedi and Kochar (2005) and Arias-Nicolás, Fernández-Ponce, Luque-Clavo and Suarez-Llorens (2005), a natural question arises: Whether the  $\leq_{c-disp}$  order holds under the dispersive order of the marginals, for random vectors with the same copula?

A copula  $C$  is a cumulative distribution function with uniform margins on  $[0, 1]$ . Furthermore, it has been shown that if  $H$  is a  $n$ -dimensional distribution function, with marginal distribution functions  $F_1, \dots, F_n$  then there exists a  $n$ -copula  $C$  such that for all  $(x_1, \dots, x_n) \in \mathbb{R}^n$ , it holds that  $H(x_1, \dots, x_n) = C(F_1(x_1), \dots, F_n(x_n))$ . Moreover, if  $F_1, \dots, F_n$  are continuous then  $C$  is unique (for details about copulas see Nelsen (1999)). It follows that if  $\mathbf{X} = (X_1, \dots, X_n)$  and  $\mathbf{Y} = (Y_1, \dots, Y_n)$  are two  $n$ -dimensional random variables, then they have the same copula if, and only if  $(X_1, \dots, X_n) =_{st} (Q_{X_1}(F_{Y_1}(Y_1)), \dots, Q_{X_n}(F_{Y_n}(Y_n)))$ . To summarize the copula allows us to separate the effect of the dependence from effects of the marginal distributions. With this settings, we can formulate the following theorem.

**Theorem 4.** *Let  $\mathbf{X} = (X_1, \dots, X_n)$  and  $\mathbf{Y} = (Y_1, \dots, Y_n)$  be two  $n$ -dimensional random vectors such that they have the same copula. Then  $\mathbf{X} \leq_{c-disp} \mathbf{Y}$  if and only if  $X_i \leq_{disp} Y_i$  for all  $i = 1, \dots, n$ .*

*Proof.* Arias-Nicolás, Fernández-Ponce, Luque-Calvo and Suárez-Llorens (2005) showed that, for two random vectors with the same copula, the function  $\phi$  can be expressed as

$$\phi_i(x_1, \dots, x_i) = Q_{Y_i}(F_{X_i}(x_i)). \quad (17)$$

for  $i = 1, \dots, n$ . Hence in light of (3) the result holds.  $\square$

Therefore given two  $n$ -dimensional random vectors  $\mathbf{X}$  and  $\mathbf{Y}$ , with the same copula, then

$$\begin{array}{c} \mathbf{X} \leq_{disp} \mathbf{Y} \Leftrightarrow \mathbf{X} \leq_{c-disp} \mathbf{Y} \Leftrightarrow X_i \leq_{disp} Y_i \quad \forall i = 1, 2, \dots, n. \\ \uparrow \\ \mathbf{X} \leq_{var} \mathbf{Y} \end{array} \quad (18)$$

Looking at the last discussion of the Example 1 and from the well known fact that two bivariate normal distributions have a common copula if, and only if they have the same correlation coefficient then it is apparent a strict implication for  $\leq_{var}$  holds. As a special case of distribution functions having the same structure dependence are two random vectors  $\mathbf{X}$  and  $\mathbf{Y}$  having independent components.

The strict implication in (18) for the  $\leq_{var}$  ordering it is not surprising if we take in account that the  $\leq_{var}$  ordering seems to be associated with the positive dependence structure of the random vectors. Those considerations lead us to assume some dependence properties for the random vectors. To simplify the notation we will say that the random vector  $(X_1, \dots, X_n)$  is *conditionally increasing in quantile*, denoted by CIQ in order to simplify, if the standard construction  $\hat{x}(\mathbf{u})$  is increasing in  $\mathbf{u} \in (0, 1)^n$ .

The CIQ notion is related to the following property. The random vector  $(X_1, \dots, X_n)$  is said to be *conditionally increasing in sequence* (CIS) if (see Barlow and Proschan (1975)),  $X_i \uparrow_{st} (X_1, \dots, X_{i-1})$ ,  $i = 2, \dots, n$ , that is, if

$$[X_i | X_1 = x_1, \dots, X_{i-1} = x_{i-1}] \leq_{st} [X_i | X_1 = x'_1, \dots, X_{i-1} = x'_{i-1}]$$

whenever  $x_j \leq x'_j$ ,  $j = 1, 2, \dots, i-1$ .

Rubinstein, Samorodnitsky and Shaked (1985) proved that the CIS property implies the CIQ property. Also the CIQ property is preserved by strictly increasing transformations of each component.

**Theorem 5.** *Let  $\mathbf{X}$  and  $\mathbf{Y}$  be two  $n$ -dimensional random vectors with the same copula. If  $\mathbf{X}$  is CIQ then  $\mathbf{X} \leq_{var} \mathbf{Y}$  if, and only if  $X_i \leq_{disp} Y_i$  for all  $i = 1, \dots, n$ .*

*Proof.* Note that from (18) it is only necessary to show the sufficient condition. Under the expression of  $\phi$  for random vectors with the same copula given by equation (17) and the fact that  $\phi$  maps the standard construction of  $\mathbf{X}$  to the corresponding of  $\mathbf{Y}$  it easily holds that  $Q_{Y_i}(F_{X_i}(\hat{x}_i(u_1, \dots, u_i))) = \hat{y}_i(u_1, \dots, u_i)$  for all  $i = 1, \dots, n$ . Hence

$$F_{X_i}(\hat{x}_i(u_1, \dots, u_i)) = F_{Y_i}(\hat{y}_i(u_1, \dots, u_i)) \text{ for all } i = 1, \dots, n. \quad (19)$$

From (19) if  $\mathbf{X}$  is CIQ is apparent that  $\mathbf{Y}$  is also CIQ. If we take  $(v_1, \dots, v_i) \geq (u_1, \dots, u_i)$  then

$$\begin{aligned} \hat{y}_i(v_1, \dots, v_i) &\geq \hat{y}_i(u_1, \dots, u_i) \\ \hat{x}_i(v_1, \dots, v_i) &\geq \hat{x}_i(u_1, \dots, u_i) \end{aligned}$$

for  $i = 1, \dots, n$ . On the other hand, from (19) it holds that  $\hat{x}_i(u_1, \dots, u_i)$  and  $\hat{y}_i(u_1, \dots, u_i)$  provide the same univariate quantile for the marginal variables  $X_i$  and  $Y_i$  respectively. By hypothesis assumption the marginal distributions are ordered in dispersion. Then

$$\hat{y}_i(v_1, \dots, v_i) - \hat{y}_i(u_1, \dots, u_i) \geq \hat{x}_i(v_1, \dots, v_i) - \hat{x}_i(u_1, \dots, u_i)$$

for all  $(v_1, \dots, v_i) \geq (u_1, \dots, u_i)$ . Then we obtain condition (9). Hence  $\mathbf{X} \leq_{var} \mathbf{Y}$ .  $\square$

Therefore from (18) and Theorem 5, given two random vectors  $\mathbf{X}$  and  $\mathbf{Y}$ , with the same copula and with the CIQ property, then we have the following chain of equivalences:

$$\mathbf{X} \leq_{disp} \mathbf{Y} \Leftrightarrow \mathbf{X} \leq_{c-disp} \mathbf{Y} \Leftrightarrow \mathbf{X} \leq_{var} \mathbf{Y} \Leftrightarrow X_i \leq_{disp} Y_i \quad \forall i = 1, \dots, n. \quad (20)$$

We would like to emphasize that the equivalences given by (20) simplify the conditions where the  $\leq_{var}$  ordering can be verified which, from a practical viewpoint, can be very complicated to compute. Specifically, we can take advantage of (20) in the applications of the  $\leq_{var}$  ordering.

## 5 Applications

### 5.1 Models of ordered random variables

An interesting application of the implications given by (20) can be given for several models of ordered random variables with applications in statistics and reliability. For instance,

the order statistics from a sample of i.i.d. random variables, the random vector of the first  $n$  epoch times of a nonhomogeneous Poisson process,  $k$  records and order statistics under multivariate imperfect repair. A general concept where the previous models are included is the concept of generalized order statistics see Kamps (1995). Formally, let  $n \in \mathbb{N}$ ,  $k \geq 1$ ,  $m_1, \dots, m_{n-1} \in \mathbb{R}$ ,  $M_r = \sum_{j=r}^{n-1} m_j$ ,  $1 \leq r \leq n-1$ , be parameters such that  $\gamma_r = k + n - r + M_r \geq 1$  for all  $r \in 1, \dots, n-1$ , and let  $\tilde{m} = (m_1, \dots, m_{n-1})$ , if  $n \geq 2$  ( $\tilde{m} \in \mathbb{R}$  arbitrary, if  $n = 1$ ). We call uniform generalized order statistics to the random vector  $(U_{(1,n,\tilde{m},k)}, \dots, U_{(n,n,\tilde{m},k)})$  with joint density function

$$h(u_1, \dots, u_n) = k \left( \prod_{j=1}^{n-1} \gamma_j \right) \left( \prod_{j=1}^{n-1} (1 - u_j)^{m_j} \right) (1 - u_n)^{k-1}$$

on the cone  $0 \leq u_1 \leq \dots \leq u_n \leq 1$ . Now given a distribution function  $F$  we call generalized order statistics (GOS) based on  $F$  to the random vector

$$(X_{(1,n,\tilde{m},k)}, \dots, X_{(n,n,\tilde{m},k)}) \equiv (F^{-1}(U_{(1,n,\tilde{m},k)}), \dots, F^{-1}(U_{(n,n,\tilde{m},k)})). \quad (21)$$

Note that if  $F$  is an absolutely continuous distribution then  $F^{-1}$  is strictly increasing. Therefore is apparent from (21) that two random vectors of generalized order statistics, with the same parameters, and possibly, based on different absolutely continuous distributions  $F$  and  $G$ , have the same copula. In addition a random vector of generalized order statistics have the CIS property (this follows from the Markovian property of GOS and the transition probabilities), so it also has the CIQ property.

Under the previous arguments and using (20) it holds that two random vectors of generalized order statistics  $(X_{(1,n,\tilde{m},k)}, \dots, X_{(n,n,\tilde{m},k)})$  and  $(Y_{(1,n,\tilde{m},k)}, \dots, Y_{(n,n,\tilde{m},k)})$ , based on absolutely continuous distribution functions  $F$  and  $G$ , respectively, are ordered in the  $\leq_{disp}$ ,  $\leq_{c-disp}$  and  $\leq_{var}$  multivariate dispersive orderings if, and only if the marginals are ordered in the univariate dispersive order, that is if  $X_{(r,n,\tilde{m},k)} \leq_{disp} Y_{(r,n,\tilde{m},k)}$ , for all  $r : 1, \dots, n$ . The last condition holds under the assumption  $F \leq_{disp} G$  (see Theorem 3.12 in Belzunce, Mercader and Ruiz (2005)). With a simpler argument, this result simplifies Theorem 3.11 by Belzunce, Mercader and Ruiz (2005), Theorem 2.3 by Belzunce, Ruiz and Ruiz (2003) and Theorem 3.1 by Belzunce and Ruiz (2002).

## 5.2 Variability ordering of convolutions

Convolutions appear naturally in several context such as risk theory, reliability and statistics. For example consider an insurance company with a number  $n$  of clients, with individual risks  $X_1, \dots, X_n$  then the company bears the risk  $S = \sum_{i=1}^n X_i$ . In reliability theory convolutions appear when a failed unit is replaced by a new one and the total life is obtained by the addition of the two lifelength. Also several statistics are linear combinations of random variables. In the literature one can find several results on variability comparisons of convolutions. Most of these results are given for some parametric models (uniform and gamma distributions) of independent random variables see for example Kochar and Ma (1999a) and (1999b), Korwar (2002), Khaledi and Kochar (2002) and (2004) in the case of the dispersive order. For dependent components a simple but elegant result is provided by Bäuerle (1997), Bäuerle and Rieder (1997) and Müller (1997), which prove that the convolution of the components of two random vectors, ordered in the supermodular order, are ordered in the increasing convex order.

One of the most interesting properties of the  $\leq_{var}$  ordering is given by the study of conditions for the variability ordering of increasing directionally transformations of the random vectors. Recall from Rüschendorf (1983) that a real function  $\varphi$  on  $\mathbb{R}^n$  is said to be *directionally convex* if for any  $\mathbf{x}_i \in \mathbb{R}^n$ ,  $i = 1, 2, 3, 4$ , such that  $\mathbf{x}_1 \leq \mathbf{x}_2 \leq \mathbf{x}_4$ ,  $\mathbf{x}_1 \leq \mathbf{x}_3 \leq \mathbf{x}_4$  and  $\mathbf{x}_1 + \mathbf{x}_4 = \mathbf{x}_2 + \mathbf{x}_3$ , one has

$$\varphi(\mathbf{x}_2) + \varphi(\mathbf{x}_3) \leq \varphi(\mathbf{x}_1) + \varphi(\mathbf{x}_4). \quad (22)$$

Shaked and Shanthikumar (1998) in Theorem 4.2 proved that given two nonnegative  $n$ -dimensional random vectors  $\mathbf{X}$  and  $\mathbf{Y}$ , with the CIS property, if  $\mathbf{X} \leq_{var} \mathbf{Y}$  then  $\varphi(\mathbf{X}) \leq_{st:icx} \varphi(\mathbf{Y})$ , for all increasing directionally convex functions  $\varphi$ . Where the  $\leq_{st:icx}$  ordering means that  $E[h(\varphi(\mathbf{X}))] \leq E[h(\varphi(\mathbf{Y}))]$  for all increasing functions  $h$  for which the expectations exists (that is, if  $\varphi(\mathbf{X}) \leq_{st} \varphi(\mathbf{Y})$ ) and  $\text{Var}[h(\varphi(\mathbf{X}))] \leq \text{Var}[h(\varphi(\mathbf{Y}))]$  for all increasing convex functions  $h$  for which the variances exist. The  $\leq_{st:icx}$  ordering is of interest because it lets you to compare the variances.

Note that Shaked and Shantikumar (1998) assume the  $\leq_{var}$  ordering between two non-negative random vectors and the CIS property. If we consider random vectors with a common copula, using Theorem 5 and the mentioned fact that CIS implies CIQ, in this case the  $\leq_{var}$  ordering can be easily checked just comparing in dispersion the marginal distributions. Hence from the fact that the function  $\varphi(\mathbf{x}) = \sum_{i=1}^n x_i$  is increasing directionally convex we can state the following general result.

**Corollary 1.** *Let  $\mathbf{X} = (X_1, \dots, X_n)$  and  $\mathbf{Y} = (Y_1, \dots, Y_n)$  be two  $n$  dimensional nonnegative CIS random vectors with a common copula. If  $X_i \leq_{disp} Y_i$  for every  $i = 1, \dots, n$ , then  $\sum_{i=1}^n X_i \leq_{st:icx} \sum_{i=1}^n Y_i$ .*

In particular we have that  $\text{Var}[h(\sum_{i=1}^n X_i)] \leq \text{Var}[h(\sum_{i=1}^n Y_i)]$ , for every increasing convex function  $h$ , provided the previous variances exist, and therefore we can compare the variability of convolutions of random variables not necessarily independent.

## References

- [1] Arias-Nicolás, J. P.; Fernández-Ponce, J. M.; Luque-Calvo, P. and Suárez-Llorens, A.(2005). Multivariate dispersion order and the notion of copula applied to the multivariate  $t$ -distribution. *Probability in the Engineering and Informational Sciences* **19**, 363–375.
- [2] Barlow, R.E. and Proschan, F. (1975). *Statistical Theory of Reliability and Life Testing*. Holt, Rinehart and Winston, Inc., New York.
- [3] Bäuerle, N. (1997). Monotonicity results for MR/GI/1 queues. *Journal of Applied Probability* **34**, 514–524.
- [4] Bäuerle, N. and Rieder, U. (1997). Comparison results for Markov-modulated recursive models. *Probability in the Engineering and Informational Sciences* **11**, 203–217.
- [5] Belzunce, F., Mercader, J.A., and Ruiz, J.M. (2005). Stochastic comparisons of generalized order statistics, *Probability in the Engineering and Informational Sciences* **19**, 99–120.

- [6] Belzunce, F. and Ruiz, J.M. (2002). Multivariate dispersive ordering of epoch times of nonhomogeneous Poisson processes. *Journal of Applied Probability* **39**, 637–643.
- [7] Belzunce, F., Ruiz, J. M. and Ruiz, M.C. (2003), Multivariate properties of random vectors of order statistics, *Journal of Statistical Planning and Inference*, **115**, 413–424.
- [8] Fernández-Ponce, J. M. and Suárez-Llorens, A.(2003). A multivariate dispersion ordering based on quantiles more widely separated. *Journal of Multivariate Analysis* **85**, 40–53.
- [9] Giovagnoli, A. and Wynn, H.P. (1995). Multivariate dispersion orderings. *Statistics and Probability Letters* **22**, 325–332.
- [10] Kamps, U. (1995), *A Concept of Generalized Order Statistics*, B.G. Teubner, Stuttgart.
- [11] Khaledi, B.-E. and Kochar, S.C. (2002). Dispersive ordering among linear combinations of uniform random variables. *Journal of Statistical Planning and Inference* **100**, 13–21.
- [12] Khaledi, B.-E. and Kochar, S.C. (2004). Ordering convolutions of gamma random variables. *Sankhyā* **66**, 466–473.
- [13] Khaledi, B.-E. and Kochar, S.C. (2005). Dependence, dispersiveness, and multivariate hazard rate ordering. *Probability in the Engineering and Informational Sciences* **19**, 427–446.
- [14] Kochar, S.C. and Ma, C. (1999a). Dispersive ordering of convolutions of exponential random variables. *Statistics and Probability Letters* **43**, 321–324.
- [15] Kochar, S.C. and Ma, C. (1999b). Erratum: "Dispersive ordering of convolutions of exponential random variables". *Statistics and Probability Letters* **45**, 283.
- [16] Korwar, R.M.(2002). On stochastic orders for sums of independent random variables. *Journal of Multivariate Analysis* **80**, 344–357.
- [17] Lewis, T. and Thompson, J. W. (1981). Dispersive distributions, and the connection between dispersivity and strong unimodality. *Journal of Applied Probability* **18**, 76–90.
- [18] Li, H.; Scarsini, M. and Shaked, M, (1996). Linkages: A tool for the construction of multivariate distributions with given nonoverlapping multivariate marginals. *Journal of Multivariate Analysis* **56**, 20–41.
- [19] Müller, A. (1997). Stop-loss order for portfolios of dependent risks. *Insurance: Mathematics and Economics* **21**, 219–223.
- [20] Müller, A. and Scarsini, M. (2001). Stochastic comparison of random vectors with a common copula. *Mathematics of Operation Reserach* **26**, 723–740.
- [21] Nelsen, R.B. (1999). *An Introduction to Copulas*. Lectures Notes in Statistics **139**. Springer-Verlag, New York.
- [22] Oja, H. (1983). Descriptive statistics for multivariate distributions. *Statistics and Probability Letters* **1**, 327–332.

- [23] Rubinstein, R.Y.; Samorodnitsky, G. and Shaked, M. (1985). Antithetic variates, multivariate dependence and simulation of stochastic systems. *Management Science* **31**, 66–77.
- [24] Rüschemdorff, L. (1983). Solution of a statistical optimization problem by rearrangement methods. *Metrika* **30**, 55–61.
- [25] Shaked, M. and Shanthikumar, J.G. (1994). *Stochastic Orders and Their Applications*. Academic Press, San Diego.
- [26] Shaked, M. and Shanthikumar, J.G. (1998). Two variability orders. *Probability in the Engineering and Informational Sciences* **12**, 1–23.

Accepted manuscript