



**HAL**  
open science

## Modelling human diseases in *Caenorhabditis elegans*

Maria Markaki, Nektarios Tavernarakis

► **To cite this version:**

Maria Markaki, Nektarios Tavernarakis. Modelling human diseases in *Caenorhabditis elegans*. *Biotechnology Journal*, 2010, 5 (12), pp.1261. 10.1002/biot.201000183 . hal-00599455

**HAL Id: hal-00599455**

**<https://hal.science/hal-00599455v1>**

Submitted on 10 Jun 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Modelling human diseases in *Caenorhabditis elegans***

Journal:	<i>Biotechnology Journal</i>
Manuscript ID:	biot.201000183.R1
Wiley - Manuscript type:	Review
Date Submitted by the Author:	22-Oct-2010
Complete List of Authors:	Markaki, Maria; Foundation for Research and Technology - Hellas, Institute of Molecular Biology and Biotechnology Tavernarakis, Nektarios; Foundation for Research and Technology - Hellas, Institute of Molecular Biology and Biotechnology
Primary Keywords:	Genetics
Secondary Keywords:	Diseases
Keywords:	Ageing, Neurodegeneration, Model Organisms

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6 Review

7  
8 **Modelling human diseases in *Caenorhabditis elegans***  
9

10  
11  
12 **Maria Markaki and Nektarios Tavernarakis**  
13

14  
15  
16  
17 Institute of Molecular Biology and Biotechnology, Foundation for Research and  
18  
19 Technology - Hellas, Heraklion 71110, Crete, Greece

20  
21  
22  
23 **Keywords:** Ageing, Cell death, Model Organisms, Neurodegeneration, Protein  
24  
25 aggregation

26  
27  
28 **Correspondence:**  
29

30 Dr. Nektarios Tavernarakis

31  
32 Institute of Molecular Biology and Biotechnology,

33  
34 Foundation for Research and Technology - Hellas

35  
36 N. Plastira 100, Vassilika Vouton, PO Box 1385, Heraklion 70013, Crete, Greece

37  
38 Tel: +30 2810 391066, Fax: +30 2810 391067

39  
40 E-mail address: [tavernarakis@imbb.forth.gr](mailto:tavernarakis@imbb.forth.gr)  
41  
42

43  
44 **Abbreviations:** A $\beta$ ,  $\beta$ -amyloid peptide; AD, Alzheimer's disease; ALS, Amyotrophic  
45  
46 Lateral Sclerosis; APP, Amyloid Precursor Protein; ASPs, aspartyl proteases;  $\alpha$ -syn,  $\alpha$ -  
47  
48 synuclein; CFP, cyan fluorescent protein; CHIP, carboxyl-terminus of Hsc70 interacting  
49  
50 protein; CLPs, calpain proteases; CNS, Central Nervous System; DA, dopaminergic  
51  
52 neurons; DAPC, Dystrofin-Associated Protein Complex; DAT-1, Dopamine transporter -1;  
53  
54 DGC, Dystrofin-Glycoprotein Complex; DMD, Duchenne Muscular Dystrophy; DR, Dietary  
55  
56 restriction; Drp-1, Dynamin-related protein 1; dVAP; Drosophila VAP; ER, endoplasmic  
57  
58 reticulum; FALS, Familiar Amyotrophic Lateral Sclerosis; FTDP-17 tau, Fronto-Temporal  
59  
60

1  
2  
3  
4  
5  
6 Dementia and Parkinsonism tau linked to chromosome 17; HIF-1, Hypoxia-Inducible  
7 Transcription factor -1; HSF-1, Heat Shock Transcription Factor 1; Htt, Huntingtin; HD,  
8 Huntington's disease; hVAP, human VAP; ILS, insulin-like signaling pathway; LRRK2,  
9 leucine-rich repeat kinase 2; MAPK, Mitogen-Activated Protein Kinase; miRNAs,  
10 microRNAs; MSP, Major Sperm Protein; NTFs, intracellular neurofibrillary tangles;  
11 OPMD, Oculopharyngeal Muscular Dystrophy; PABPN1, Poly(A) Binding Protein Nuclear  
12 1; PD, Parkinson's disease; PHP tau, pseudohyperphosphorylated tau; ROS, Reactive  
13 Oxygen Species; RTK, Receptor Tyrosine Kinase ; SMA, Spinal Muscular Atrophy; SMN,  
14 survival motor neuron gene; SOD1, Cu/Zn Superoxide dismutase 1; snb-1, synaptobrevin  
15 gene -1; TGF $\beta$ , Transforming growth factor- $\beta$ ; ts, temperature-sensitive; VAMP,  
16 (Synaptobrevin) Associated Protein B (VAPB); V- ATPase, vacuolar-ATPase; YFP, yellow  
17 fluorescent protein  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Abstract

Genes linked to human diseases often function in evolutionary conserved pathways, which can be readily dissected in simple model organisms. Because of its short lifespan and the well known biology coupled with a completely sequenced genome that shares extensive homology with that of mammals, *Caenorhabditis elegans* is one of the most versatile and powerful model organisms. Research in *C. elegans* has been instrumental for the elucidation of molecular pathways implicated in many human diseases. In this review, we introduce *C. elegans* as a model organism for biomedical research and we survey recent relevant findings that shed light on the basic molecular determinants of human disease pathophysiology. The nematode holds promise of providing clear leads towards the identification of potential targets for the development of new therapeutic interventions against human diseases.

## 1 Introduction to *Caenorhabditis elegans* biology

The roundworm *Caenorhabditis elegans* (*C. elegans*) was selected as a simple metazoan model in the early 1960's by Sydney Brenner for pursuing research in developmental biology and neurobiology. Since its introduction as a model organism, *C. elegans* has been widely used to investigate important biological processes most of which have remained essentially unchanged during evolution [1]. *C. elegans* is a free-living, non-parasitic nematode, with a life cycle of 3.5 days at 20°C and a lifespan of about 2 to 3 weeks under suitable living conditions. The adult is about 1mm in length and 80µm in diameter; it feeds on bacteria such as *Escherichia coli* in liquid medium or on agar plates and can be easily cultivated in large numbers. *C. elegans* has five pairs of autosomes and one pair of sex chromosomes. It has two sexes, hermaphrodites and males; the ratio of sex chromosomes to autosomes determines its sex. A single hermaphrodite produces ~300 progeny by self-fertilization and more if it mates with males which arise occasionally at a frequency of 0.1%. Wild-type individuals consist of 959 somatic cells, 302 of which are neurons. The animal body is transparent so it is easy to track cells and follow cell lineages and biological processes [2]. The genome size of *C. elegans* is about a hundred million base pairs, is completely sequenced and surprisingly similar to that of humans; it is estimated that 60-80% of the genes have a human counterpart [3]. Viable mutant strains, strains that overexpress a gene or lack a gene function can be efficiently generated and the resulting phenotypes can be rapidly identified [4]. Comprehensive information concerning gene structure, expression patterns, protein-protein interactions, mutant or RNAi phenotypes and microarray data, is available in Wormbase, the online resource for nematode-related information (<http://www.wormbase.org/>) [5]. Nematode strains can be stored indefinitely in liquid nitrogen allowing for generating large mutant collections and public mutant repositories. *C. elegans* is amenable to unbiased forward and reverse genetic screens and particularly susceptible to gene inactivation by RNA interference [6, 7]. Due to its value as a research tool, a large set of methods that include advanced high resolution imaging techniques have been developed.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*C. elegans* has emerged as a powerful experimental system to study the molecular and cellular aspects of human disease *in vivo*. It has been estimated that about 42% of the human disease genes have an orthologue in the genome of *C. elegans* including those genes associated with Alzheimer disease, juvenile Parkinson's disease, spinal muscular atrophy, hereditary non-polyposis colon cancer, and many others age-related disorders [8-10]. Modelling a human disease in a simple invertebrate, such as *C. elegans* allows the dissection of complex molecular pathways into their component parts and thus provides meaningful insight into the pathogenesis of a complex disease phenotype. Here, we survey nematode models of human disease, highlighting recent discoveries that shed light on the molecular mechanisms underlying disease pathogenesis.

## 2 *C. elegans* models of neurodegenerative disorders

Neurodegeneration has a profound effect on human health, yet the mechanisms underlying neuronal injury and death remain unclear. *C. elegans* models of neuronal dysfunction have been established for a number of neurodegenerative diseases, including Alzheimer disease, Parkinson disease and polyglutamin-expansion disorders. These models typically involve the transgenic expression of human disease genes.

### 2.1 Alzheimer's disease (AD)

Alzheimer's disease (AD) is a progressive neurologic disease that results in the irreversible loss of neurons particularly in the neocortex. The brain of AD affected patients is characterized by the accumulation of intracellular neurofibrillary tangles (NFTs) composed of the microtubule-associated protein tau that stabilizes microtubules when phosphorylated [11] and of extracellular senile plaques primarily composed of the  $\beta$ -amyloid peptide (A $\beta$ ) [12]. Compelling evidence supports the causative role of (A $\beta$ )<sub>1-42</sub>,

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

which derives from the proteolysis of Amyloid Precursor Protein (APP) by  $\beta$ - and  $\gamma$ -secretase, in AD. Similarly, mutations in the tau gene lead to familiar tauopathies.

Autosomal dominant mutations in APP are correlated with rare cases of early-onset AD. Determining the *in vivo* functions of APP is difficult because mammals have an APP gene family containing two APP-related genes. *C. elegans* has a single APP-related gene, the *apl-1* mapped on the X chromosome. *Apl-1* is expressed in multiple cell types and is necessary for many developmental processes, including proper molting and morphogenesis. Loss of *apl-1* causes larval lethality which can be rescued by neuronal expression of the extracellular domain of APL-1. Similarly, the overexpression of APL-1 causes defects in movement, brood size and larval viability of transgenic nematodes. *Apl-1* overexpression-induced lethality is partially rescued by the reduced activity of *sel-12*, a *C. elegans* homologue of the human  $\gamma$ -secretase gene component presenilin 1, suggesting that SEL-12, like mammalian PS1, regulates either directly or indirectly the activity of APL-1 [13].

*C. elegans* AD models were developed that express A $\beta$  variants under the control of male-specific, body wall muscle-specific and neuronal-specific promoters. Only transgenic lines carrying the *unc-54/A $\beta$ <sub>1-42</sub>* minigene accumulate intracellular A $\beta$  and show a progressive paralysis beginning in young adulthood. Co-immunoprecipitation studies coupled with mass-spectrometry reveal that chaperone-related proteins (two human HSP70, 3 HSP-16 homologous to  $\alpha$ B-crystallin and a putative negative regulator of HSP70 function) interact with A $\beta$  *in vivo*. This interaction is considered as part of a cellular protective response since the overexpression of either HSP-16.2 or a human HSP70 partially suppresses the A $\beta$ -induced paralysis in worms [14]. Heat shock treatment reduces A $\beta$  oligomeric protein and delays paralysis in transgenic A $\beta$  worms probably through the activation of heat shock proteins regulated by the HSF-1 transcription factor [15]. HSF-1 has been coupled to normal ageing and age-related



1  
2  
3  
4  
5  
6 diseases [16]. Previous studies showed that constituents of *Ginkgo biloba* leaf extract  
7 (EGb 761 and ginkgolides) delay beta amyloid-induced paralysis in transgenic *C. elegans*  
8 [17]. This effect was not additive to protective heat shock suggesting a shared  
9 mechanism of action by two treatments [15]. Microarray analysis in a transgenic worm  
10 strain, that accurately expresses a temperature inducible A $\beta$ <sub>42</sub> in body wall muscle,  
11 reveals that the gene encoding the arsenite-inducible protein (*aip-1*) is upregulated as  
12 part of a cellular protective response. Worms overexpressing AIP-1 show decreased  
13 accumulation of A $\beta$ <sub>42</sub> peptide and attenuated paralysis. The AIP-1 human homologue,  
14 AIRAPL, but not AIRAP when expressed under the control of *myo-3* promoter also  
15 confers protection against A $\beta$  toxicity [18]. Both AIP-1 and AIRAPL contain a predicted  
16 farnesylation site known to be critical for the AIP-1 mediated longevity in *C. elegans* [19].  
17 AIP-1 and AIRAPL enhance at least partially general protein turnover by acting as  
18 positive regulators of proteosomal function and thus protect against A $\beta$ -induced toxicity  
19 [18]. The effects of tetracycline and its analogues doxycycline and minocycline on A $\beta$ <sub>42</sub>-  
20 induced toxicity were assayed [20] using the worms expressing a temperature inducible  
21 A $\beta$ <sub>1-42</sub> transgene [21]. Tetracyclines successfully protected transgenic worms from the A $\beta$   
22 insult by reducing the concentration of oligomers considered to be responsible for the  
23 toxic phenotype. These effects were specific, dose-dependent and not linked to any  
24 antibiotic activity. Furthermore, tetracyclines protect A $\beta$ -expressing nematodes from  
25 oxidative stress by reducing the superoxide production suggesting a potential use of  
26 these drugs in targeting A $\beta$  aggregates.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

## 49 2.2 Tauopathies

50  
51 Abnormal phosphorylation, aggregation and ultimately functional alteration of the main  
52 microtubule-associated protein, tau, is the major cause of neurodegenerative disorders  
53 known as tauopathies [22, 23]. Tau aggregates are found in AD, Pick's disease,  
54 corticobasal degeneration, Down's syndrome and FTDP-17 (frontotemporal dementia  
55 with parkinsonism chromosome 17 type) [24]. The most common tauopathy is  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 Alzheimer's disease. To analyze the role of tau modification during AD progression,  
7  
8 human tau and a pseudohyperphosphorylated (10 serine/threonine residues were  
9  
10 changed to glutamates) PHP tau, that mimics AD- relevant tau modification, were brought  
11  
12 under the control of a pan neuronal (a 3.4 kb upstream of the gene F25B3.3) promoter.  
13  
14 Transgenic worms expressing either tau or PHP tau develop a phenotype of  
15  
16 uncoordinated (*Unc*) locomotion, characteristic of a variety of *C. elegans* nervous system  
17  
18 defects. However, only PHP tau- expressing worms show a defective pattern of motor  
19  
20 neuron development suggesting that PHP tau may better mimic the stable disease-like  
21  
22 tau modification which interferes with mechanisms of axonal outgrowth and path finding  
23  
24 [25]. To identify specific modifiers for the tau-induced pathologic phenotype, a genome-  
25  
26 wide RNAi screen was conducted using transgenic worms expressing mutations in tau  
27  
28 (FTDP-17 tau) driven by the *aex-3* pan neuronal promoter. 60 enhancer genes modified  
29  
30 only the tau-induced *Unc* phenotype. These genes normally function in protein  
31  
32 phosphorylation, protein folding, stress response, nucleic acid function, proteolysis and  
33  
34 neurotransmission [26]. In addition, random mutational screens on these transgenic  
35  
36 animals resulted in the identification of two novel tauopathy-associated genes, *sut-1* and  
37  
38 *sut-2* (suppressor of tauopathy) [27, 28]. SUT-1 requires the activity of UNC-34 which  
39  
40 plays a role in the correct neuronal cell migration and axonal guidance. SUT-2 encodes a  
41  
42 CCCH zinc finger protein, primarily expressed in the nucleus of neuronal cells, which  
43  
44 binds ZYG-12, a protein of the HOOK family. In turn, ZYG-12 acts as a linker connecting  
45  
46 membrane compartments with the microtubule cytoskeleton protein.  
47  
48

### 49 **2.3 Polyglutamine-expansion disorders**

50  
51 Numerous human neurodegenerative disorders including Huntington's disease (HD),  
52  
53 several spinocerebellar ataxias and Kennedy's disease [29, 30] are associated with  
54  
55 misfolded and aggregation-prone proteins that interfere with protein homeostasis leading  
56  
57 to cellular dysfunction. Polyglutamine diseases are caused by a CAG triplet expansion in  
58  
59  
60

1  
2  
3  
4  
5  
6 the coding regions of seemingly unrelated genes resulting in a pathogenic protein with  
7 abnormally expanded track of glutamines (Fig.1) [31].  
8  
9

10  
11 To address the mechanisms underlying the impact of aggregation-prone proteins  
12 on cellular function, transgenic *C. elegans* lines expressing different lengths of polyQ-  
13 YFP (yellow fluorescent protein) or CFP (cyan fluorescent protein) from integrated arrays  
14 in muscle (polyQm) [32] or neuronal (polyQn) [33] cells have been utilized in genetic  
15 screens. These polyglutamine aggregation models show polyQ-length dependent  
16 aggregation and toxicity. *C. elegans* polyQm or polyQn strains were crossed with animals  
17 expressing temperature-sensitive (ts) mutations considered to be sensitive indicators of  
18 a disruption in protein homeostasis. These experiments have shown that chronic  
19 expression of an aggregation-prone protein interferes with the function of multiple  
20 structurally and functionally unrelated proteins in a polyQ-length dependent manner.  
21 PolyQ expansions uncover the folding defect of proteins harbouring mildly destabilizing  
22 temperature-sensitive mutations, which in turn modify polyQ aggregation and toxicity and  
23 enhance the disruption of cellular folding homeostasis. Thus, mild folding mutations in the  
24 genetic background can modify the toxic phenotypes of polyglutamine diseases [34].  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 Protein aggregation and cellular dysfunction can occur at a threshold of  
42 approximately 40 glutamine residues. However, the threshold for polyQ aggregation and  
43 toxicity is dynamic and ageing-dependent in *C. elegans*. At 3 days of age or less, only  
44 worms expressing Q40 or greater exhibit aggregates, while in older animals aggregates  
45 appear even in Q33 and Q35 expressing worms (32).  
46  
47  
48  
49  
50

51  
52 A link between molecular determinants of ageing and aggregation-mediated  
53 proteotoxicity was established in *C. elegans*. Polyglutamine toxicity was dependent on  
54 the activity of *daf-16*, a forkhead transcription factor of the FOXO family functioning  
55 downstream of *age-1* (a phosphoinositide-3 kinase) in the insulin-like signalling pathway  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 (ILS) [35]. Work on polyQ suppression shows that TOR-2, the worm ortholog of the  
7 mammalian torsinA, is an ER-associated protein with chaperone activity which  
8 suppresses polyglutamine aggregation in *C. elegans* [36].  
9  
10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Huntington's disease (HD) is the most frequent autosomal dominant inherited  
polyQ disorder caused by an expansion of a CAG trinucleotide sequence in the huntingtin  
(Htt) protein. Normal Huntington alleles encode up to 37 CAG repeats, whereas HD-  
affected individuals consist of more than 40.

23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
Previous work has shown that ubiquilin is an interactor of presenilin, a protein  
associated with Alzheimer's disease. Ubiquilin proteins contain multiple ubiquitin related  
domains and are linked to the ubiquitin-proteasome system of protein degradation  
because both their ubiquitin-like domain and the ubiquitin-associated domain have been  
shown to bind the proteosomal subunit S5a [37]. The role of ubiquilin in HD pathogenesis  
was investigated using *C. elegans* expressing GFP-Htt-polyQ fusion proteins in body wall  
muscle. Transgenic worms display a polyQ length-dependent motility defect which is  
suppressed by ubiquilin overexpression through a mRFP-tagged ubiquilin fusion protein.  
Conversely, knockdown by RNAi of *C. elegans* ubiquilin gene exacerbated the motility  
defect associated with the expression of GFP-Htt (Q55) fusion protein. These findings  
show that, similar to AD, ubiquilin protects against polyglutamine-induced toxicity and  
they suggest a general role for ubiquilin in regulating a number of different  
neurodegenerative disorders [38]. The motility defect that these *C. elegans* HD models  
display is reduced by the knockdown of the gene encoding the dynamin-related protein 1  
(Drp-1) which controls mitochondrial fission. This result establishes a link between HD  
pathogenesis and mitochondrial dysfunction [39]. A role for insulin-like signaling pathway  
(IIS) in the amelioration of Huntington's-associated proteotoxicity has been reported [16].

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

Two different *C. elegans* models of human disease were used to explore the effects of dietary restriction (DR) on polyglutamine-associated age-related diseases. A HD model in which 35 glutamine residues were fused to YFP (Q35YFP) and expressed in the body wall muscles [32] and a transgenic model expressing a 42-amino acid amyloid-beta peptide (A $\beta$ <sub>42</sub>) under the control of the *unc-54* promoter [40]. The results demonstrate that DR suppresses age-associated paralysis in these worms. To test if DR acts as a general suppressor of proteotoxicity, the effects of food deprivation on transgenic worms expressing an aggregation-prone form of GFP (GFP-degron) were also examined. It was shown that both environmental and genetic (represented by the RNAi inhibition of *eat-2* and the knockdown of *rab-10*, *sams-1* or *drr-1* genes that act downstream of food consumption) models of DR confer protection against proteotoxicity in *C. elegans*. Compelling evidence was presented that the general protective effect of DR is mediated by the activity of the heat shock transcription factor, *hsf-1* [41].

#### 33 **2.4 Parkinson's disease**

34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Parkinson's disease (PD) is the second most common neurodegenerative disorder characterized by involuntary movements including bradykinesia and difficulty in balance and muscle rigidity due to the progressive loss of dopaminergic (DA) neurons from the substantia nigra. At cellular level, PD is characterized by cytoplasmic protein inclusions called Lewy bodies that are the pathological feature of PD and other neurodegenerative disorders known as synucleopathies [42].  $\alpha$ -synuclein ( $\alpha$ -syn), a major constituent of Lewy bodies, is causatively related to PD since missense mutations in the  $\alpha$ -syn gene or duplications and triplications of the  $\alpha$ -syn locus cause familial forms of PD [43-45]. Sporadic PD affecting 1-4% of those over 65 years of age appears to be unrelated to mutations or multiplications of the  $\alpha$ -syn locus and implicates a combination of environmental factors. How  $\alpha$ -syn inclusions are formed and how this links to disease is poorly understood. Synucleopathies have been associated with mitochondria dysfunction,

1  
2  
3  
4  
5  
6 defective autophagy, impairment of the ubiquitin-proteasome system, production of  
7  
8 reactive oxygen species (ROS), stress within the ER [46].  
9

10  
11 The worm genome includes orthologs of all established familiar PD genes  
12 (*Parkin*, *DJ-1*, *PINK1*, *UCHL-1*, *LRRK2*, *PARK9* and *NURR1*) except the  $\alpha$ -syn. A  
13  
14 transgene bearing Rab1 guanosine triphosphate, a GTPase involved in endoplasmic  
15  
16 reticulum to Golgi transport, under the control of the dopamine transporter (*DAT-1*) gene  
17  
18 promoter was used to drive the expression of Rab1 along with  $\alpha$ -syn in DA neurons in a  
19  
20 worm PD model. Human  $\alpha$ -syn overexpression in *C. elegans* leads to the formation of  
21  
22 misfolded  $\alpha$ -syn aggregates that increase with age. Coexpression of Rab1 significantly  
23  
24 rescued the neurodegeneration caused by the expression of  $\alpha$ -syn alone. Considering  
25  
26 that Rab1 plays an essential role in ER-Golgi vesicles trafficking and its enhanced  
27  
28 expression protects against  $\alpha$ -syn pathology, these combined results indicate that the  
29  
30 accumulation of  $\alpha$ -syn inhibits the ER-Golgi trafficking [47]. Once coexpressed, TOR-2  
31  
32 [48] ameliorated the formation of these aggregates by associating with  $\alpha$ -syn in Lewy  
33  
34 bodies [49].  
35  
36  
37  
38

39 Worms expressing both  $\alpha$ -syn::GFP and TOR-2 in body wall muscles were used  
40  
41 in a large-scale RNAi screen for modifiers of  $\alpha$ -syn misfolding and neurodegeneration.  
42  
43 868 candidate genes potentially associated with existing PD genes and pathways were  
44  
45 targeted for knockdown. 20 gene products that enhance misfolding of human  $\alpha$ -syn upon  
46  
47 depletion were isolated. The neuroprotective genes, F32A6.3, the *C. elegans* ortholog of  
48  
49 VPS41, a conserved vesicular protein, essential for lysosomal biogenesis, W08D2.5, the  
50  
51 worm orthologue of PARK9 and F55A4.1 (orthologue of Sec22p), a well characterized  
52  
53 vesicular trafficking protein in yeast, were among those identified. The identification of *C.*  
54  
55 *elegans* ATG7 as a neuroprotective gene indicates the significant role of autophagy and  
56  
57 lysosomal function in preventing the DA neurons loss caused by excess  $\alpha$ -syn [48]. To  
58  
59 further characterize the processes involved in inclusion bodies formation, a genome-wide  
60

1  
2  
3  
4  
5  
6 RNAi screen was performed in transgenic worms expressing the  $\alpha$ -synuclein-YFP  
7  
8 chimaera driven by the *unc-54* promoter. The screen identified 80 suppressors, of which  
9  
10 49 have an established human ortholog. Most of the genes are involved in protein quality  
11  
12 control and vesicle trafficking between the ER-Golgi as well as other vesicular  
13  
14 compartments. Moreover, this screen uncovered several regulators of life span (the  
15  
16 NAD<sup>+</sup> dependent protein deacetylase *sir-2.1*, the sphingolipid synthase *lagr-1*) as  
17  
18 modifiers of inclusion formation suggesting a link between the molecular mechanisms of  
19  
20 ageing and  $\alpha$ -synuclein pathology [50].  
21  
22

23  
24 Mutations in GTPase (R1441C/G) and kinase (G2019S) domains of leucine-rich  
25  
26 repeat kinase (LRRK2) are the most frequent cause of autosomal dominant and  
27  
28 idiopathic PD [51, 52]. Expression of human LRRK2 in *C. elegans* neurons modifies the  
29  
30 response to mitochondrial inhibition, inferred by the increased survival of worms exposed  
31  
32 to rotenone or paraquat, which are agents that impair mitochondrial function. These  
33  
34 findings demonstrate the power of the nematode as a model to study the possible gain-  
35  
36 of-function phenotype acquired by LRRK2 mutations. *lrk-1* is the sole *C. elegans*  
37  
38 orthologue of LRRK2. *lrk-1* and LRRK2 seems to share similar functions since LRRK2  
39  
40 complements the sensitization to rotenone caused by the *lrk-1* knockdown. These results  
41  
42 indicate a role for LRRK2 and its worm orthologue in regulating mitochondrial dysfunction  
43  
44 [53]. R1441C/G and G2019S mutations increase kinase activity of LRRK2 *in vitro* [54,  
45  
46 55]. The role of LRRK2 in PD pathogenesis was investigated *in vivo* using transgenic *C.*  
47  
48 *elegans* overexpressing human LRRK2 wild type, R1441C/G and G2019S mutations and  
49  
50 a LRRK2-inactive mutation K1347A in dopaminergic neurons. All transgenic worms show  
51  
52 an age-dependent degeneration of DA neurons, locomotor dysfunction, behavioural  
53  
54 abnormalities and dopamine deficiency. In contrast, loss of endogenous LRRK-1 and the  
55  
56 GTP binding defective mutation, K1347A, ameliorated LRRK2-associated phenotypes  
57  
58 [56].  
59  
60

## 2.5 Amyotrophic lateral sclerosis (ALS)

Amyotrophic lateral sclerosis (ALS), the most common motor neuron disorder, is characterized by dysfunction or death of motor neurons in the motor cortex, brain stem and spinal cord. Approximately 10% of ALS cases are familial with an autosomal dominant inheritance [57] and ~20% of familial ALS (FALS) are associated with mutations in the gene coding for Cu/Zn superoxide dismutase (SOD1) [58] that catalyzes the conversion of highly reactive superoxide anions into hydrogen peroxide [59]. While it is accepted that toxic gain of function mutations in SOD1 gene evoke the disease phenotype, the mechanisms of toxicity remain unclear.

Transgenic *C. elegans* expressing FALS- related mutant (A4V, G37R, G93A) human SOD1 are more sensitive to paraquat-induced oxidative stress than control worms expressing wild-type human SOD1. Moreover, the transgenic worms that harbour gain of function alleles show an aberrant accumulation and aggregate formation of mutant proteins under oxidative stress. Thus, oxidative damage during ageing might cause FALS by inhibiting degradation and enhancing aggregate formation of mutant SOD1 proteins [60].

G85R, an ALS-associated human mutant SOD1, causes a severe locomotor defect in transgenic *C. elegans* when expressed pan-neuronally under the control of the synaptobrevin (*snb-1*) gene promoter. Paralyzed animals have soluble oligomers and insoluble aggregates in neurons. The movement defect results from the dysfunction of synapses, the connections between neurons and between neurons and muscles. An RNAi screen identified molecular chaperones as critical modifiers of protein aggregation – mediated toxicity [61]. Expression of three distinct SOD1 mutations (G85R, G93A, 127X) in the body muscle cells of wild type (N2) worms cause only mild cellular toxicity. However, when SOD1 mutants are expressed in the background of destabilized protein polymorphisms, their toxicity is enhanced. These results suggest that the genetic



1  
2  
3  
4  
5  
6 background may intensify the specific toxic phenotypes of misfolded and aggregation-  
7  
8 prone proteins that characterize conformational diseases [62].  
9

10  
11 A recent study identified the gene ALS8 as a novel cause of ALS [63]. Mutations  
12  
13 in ALS8, as in SOD, result in a wide range of defects which vary in the age of onset, the  
14  
15 speed of progression and the motor neurons affected. This variation may be due to  
16  
17 genetic modifiers, redundancy or environmental contribution. ALS8 codes for VAMP  
18  
19 (Synaptobrevin) Associated Protein B (VAPB) which is closely related to VAPA, a protein  
20  
21 associated with the cytoplasmic face of the endoplasmic reticulum (ER) and the Golgi  
22  
23 apparatus [64-66]. Many species have VAPs proteins characterized by the major sperm  
24  
25 protein domain (MSP) of about 125 residues in the amino (N)-terminal end [67], the  
26  
27 central region of coiled-form motif and the hydrophobic carboxy (C)-terminus which acts  
28  
29 as a membrane anchor. The MSP domain is named from its similarity to *C. elegans*  
30  
31 MSPs, the most abundant proteins in the worm sperm [68].  
32  
33

34  
35 MSP function in the oocyte maturation and sheath contraction in *C. elegans* [69].  
36  
37 After secretion, extracellular MSP directly binds to the VAB-1 Eph receptor and other yet  
38  
39 to be identified receptors on oocyte and ovarian cell surfaces [70].  
40  
41

42  
43 The biological function of VAPs and the mechanisms underlying their impact in  
44  
45 ALS pathogenesis is not well understood. To define the role of VAPs, *C. elegans* was  
46  
47 used together with other model systems. *C. elegans* has a single VAP homolog, the VPR-  
48  
49 1 and VAP MSP domains share a 25% identity in their primary sequences with MSPs.  
50  
51 Injection of recombinant VPR-1 MSP stimulated oocyte maturation and sheath  
52  
53 contraction in the gonads of worms lacking MSP and sperm, suggesting that *C. elegans*  
54  
55 MSP and VAP MSP domains have evolutionary conserved extracellular signalling activity.  
56  
57 Microinjection of *Drosophila* VAP (dVAP) and hVAP also induced both responses. It is  
58  
59 shown that VAPs regulate Eph receptor signalling *in vivo*. MSP domains of VAP proteins  
60

1  
2  
3  
4  
5  
6 are cleaved and secreted ligands for Eph receptors. Mutations in VAPs may result in a  
7  
8 failure to secrete the MSP domain, accumulation of inclusions in the endoplasmic  
9  
10 reticulum and an unfolded protein response. These findings provide insight into the  
11  
12 disease mechanism [71].

### 13 14 15 **2.6 Spinal muscular atrophy (SMA)**

16  
17 Spinal muscular atrophy (SMA), the most common genetic cause of infant mortality, is an  
18  
19 autosomal recessive neuromuscular disorder. SMA patients suffer from degeneration of  
20  
21 lower motor neurons in the anterior horn of the spinal cord leading to atrophy of the  
22  
23 corresponding muscles [72]. The survival motor neuron gene (SMN) has been  
24  
25 causatively linked with the SMA, but the mechanisms underlying the disease  
26  
27 pathogenesis are poorly understood. Given that SMN is widely expressed both within and  
28  
29 outside the nervous system [73], the specific motor neuron defect that the SMA patients  
30  
31 face may be rather linked to the gene function [74]. *C. elegans smn-1* gene has been  
32  
33 identified as the orthologue of human SMN [75] with a widespread expression in various  
34  
35 tissues including the nervous system and body wall muscles. The mutant *smn-1(ok355)*  
36  
37 harbours a deletion allele which removes most of *smn-1* including the translational start  
38  
39 codon. Having reduced SMN-1 levels throughout the organism, *smn-1(ok355)* is used for  
40  
41 modelling aspects of SMA. Mutant animals proceed through embryogenesis due to  
42  
43 maternally contributed SMN-1, but showed late larval arrest, decreased lifespan, defects  
44  
45 in motility and pharyngeal pumping. Neuronal, but not muscle-directed expression of full  
46  
47 length *smn-1* partially rescues the *smn-1(ok355)* phenotype [76]. A genome-wide RNAi  
48  
49 screen and behavioural studies in the loss of function *smn-1(ok355)* mutant strain provide  
50  
51 preliminary evidence that the TGF $\beta$ /BMP signalling pathway plays a role in the SMN-  
52  
53 mediated neuromuscular pathology in *C. elegans* [77].

### 54 55 56 **3 Stroke – Excitotoxicity**

57  
58  
59  
60

1  
2  
3  
4  
5  
6 Necrotic cell death underlies the pathology of numerous human degenerative conditions  
7 [78]. In *C. elegans*, non-apoptotic cell death (necrosis) is triggered by specific genetic  
8 lesions or environmental conditions such as energy depletion, toxin exposure or extreme  
9 temperatures. Severe deprivation of energy resources can rapidly develop during  
10 ischaemic or hypoglycaemic episodes. Prolonged hypoxia, a condition of oxygen  
11 deprivation that takes place in ischemic episodes and stroke, also induces cell death in *C.*  
12 *elegans* [79]. The most thoroughly characterized necrosis-induced mutations in the  
13 nematode are gain of function mutations in specific ion channel genes, such as degenerin  
14 genes *deg-1* and *mec-4*, the acetylcholine receptor channel subunit gene *deg-3* and the  
15 Gs protein  $\alpha$ -subunit gene *gsa-1*, which result in necrotic cell death of neurons expressing  
16 the mutant proteins [80]. Execution of necrotic cell death requires the activity of calcium-  
17 regulated CLP-1 and TRA-3 calpain proteases and ASP-3 ASP-4 aspartyl proteases.  
18 Perturbation of intracellular concentrations of calcium, either by extracellular calcium  
19 influx or by release of endoplasmic reticulum stores, activates calpain proteases which in  
20 turn engage executioner aspartyl proteases leading to cell destruction [81]. In mammalian  
21 models of excitotoxicity, calpains increase  $\text{Ca}^{2+}$  concentration by mediating the cleavage  
22 of specific glutamate receptors [82]. Aspartyl proteases are either cytoplasmic or  
23 lysosomal that are released into the cytoplasm as a result of lysosomal rupture.  
24 Acidification of the cytoplasm due to the action of hydrolytic enzymes deliberated during  
25 the lysosomal rupture requires the vacuolar  $\text{H}^+$ -V-ATPase (V-ATPases), a proton pump  
26 that acidifies endocytic compartments. Similar mechanisms may underlying the extreme  
27 acidosis that is observed during stroke in humans [83]. Evidence supports that lysosomes  
28 contribute to execution of necrosis. Suppression of lysosome biogenesis and of  
29 lysosome-mediated cytoplasmic acidification, that develops during necrosis and is  
30 required for cell death, affects necrosis in *C. elegans* [84]. A working model for necrotic  
31 cell death and cellular defense pathways such as autophagy that may underlie diverse  
32 aspects of development, tissue homeostasis and disease pathogenesis in *C. elegans* is  
33 outlined in Fig. 2.

## 4 Muscle-associated disorders

Dystrophin is a huge cytosolic protein that links the intracellular F-actin filaments to the members of the dystrophin-glycoprotein complex (DGC) [85, 86]. Genetic defects in the complex components cause a variety of pathological conditions including muscular dystrophy, cardiomyopathy and vasospasm. Determining the molecular function of the complex will increase our understanding of muscle diseases pathogenesis. *C. elegans* has a dystrophin complex (DAPC) and mutations in its components result in head bending and hypercontraction [87, 88]. Several lines of evidence couple defects in the DAPC with calcium homeostasis disruption through a mechanism involving the ISLO-1 which mediates the interaction of SLO-1 (the orthologue of the mammalian BK channel) with DAPC. A defect in either DAPC or ISLO-1 impacts SLO-1 channel localization resulting in muscle hyperactivity [89].

### 4.1 Duchenne muscular dystrophy (DMD)

Duchenne muscular dystrophy (DMD) is one of the most severe X-linked childhood diseases, characterized by progressive muscle wasting and weakness due to mutations in dystrophin gene [90]. A *C. elegans* DMD model carrying mutations in *dys-1* and *h1h-1*, the worm orthologues of the human dystrophin and MyoD respectively, was used to investigate the effects of *chn-1* on muscle degeneration. *Chn-1* encodes the homologue of the human E3/E4 ubiquitylation enzyme CHIP (carboxyl-terminus of Hsc-70 interacting protein) that is strongly expressed in both neurons and muscles [91]. Downregulation of *chn-1* by mutation, as well as by pharmacological treatment with a proteasomal inhibitor (MG132) improves the motility of DMD worms. Muscle-specific decrease of CHN-1 activity is sufficient to control muscle degeneration in dystrophin mutant animals. These data suggest a specific role of CHN-1/CHIP in the control of muscle degeneration through the ubiquitin-proteasome pathway [92].

## 4.2 Oculopharyngeal muscular dystrophy (OPMD)

Oculopharyngeal muscular dystrophy (OPMD) is a late-onset autosomal dominant myopathy characterized by eyelid ptosis, dysphagia and limb-girdle muscle weakness [93]. A short polyalanine (GCG-triplet) stretch from 10 alanines to at least 13 alanines in the gene encoding polyalanine binding protein 1 (PABPN1) on chromosome 14q cause gain of function mutations leading to misfolding, aggregate formation and dysfunction of the ubiquitin proteasome system [94-96]. However, the pathogenesis of disease remains unclear.

In a *C. elegans* OPMD model, loss of function mutations in *sir-2.1/SIRT1* and *daf-16* protect transgenic worms from the muscle decline and abnormal motility induced by expression of mutant (13 alanines) PAPBN1 protein [97].

## 5 Cancer

Cancer is a major cause of human mortality and typically evolves as a multistep process from mutations in key cellular genes. Although *C. elegans* does not form tumours *per se*, other mutant phenotypes are linked to worm homologues of human cancer genes, which often function at distinct steps within evolutionary conserved pathways. Thus, identifying *C. elegans* genes that disrupt processes relevant to cancer could contribute importantly to our understanding of the cancer gene functions.

Several studies show that c-Met, a receptor tyrosine kinase (RTK) is overexpressed and amplified in many human cancers [98-102]. Mutations in the c-Met juxtamembrane domain as well as germline mutations have been identified in small cell lung cancer [103] and in non-small cell lung cancer [104] cases respectively.

Transgenic worms expressing wild type c-Met or the frequent human lung cancer specific c-Met mutants (R988C and T1010I) were generated in order to investigate a

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

cancer gene function in a whole organism context. Mutant transgenic worms show locomotion defects, low fecundity and vulval hyperplasia, phenotypes intensified upon exposure to nicotine [105].

The RTK/Ras/Raf/MAPK signalling pathway is highly conserved and interfaces with many aspects of animal development. Work in *C. elegans* has greatly contributed in our understanding of this pathway and its regulation [106]. In *C. elegans*, Ras controls vulval induction and its hyperactivation interferes with vulval cell differentiation leading to a multivulval phenotype [107]. Vulval development provides a model system to understand cell proliferation and differentiation, deregulation of which is causatively linked to human tumours.

Recently, the role of miRNAs in human cancer is extensively studied. miRNAs are small non-protein coding RNA molecules of ~21-23 nucleotides which control gene expression at the post-transcriptional level acting as developmental timing regulators [108]. Many of the 112 miRNAs identified so far in *C. elegans* are conserved across phylogeny. Compelling evidence suggests that miRNAs are involved in the regulation of ageing and metabolism and provides insight into their function as tumour suppressor genes or oncogenes.

Several miRNAs act in the Notch and Ras signalling pathways [109-111]. miR-84, a *let-7* family miRNA, targets *let-60*, a *C. elegans* orthologue of HRAS, KRAS and NRAS, the human ras oncogenes. All of the human ras genes have complementary sites for human *let-7* family miRNAs in their 3'UTRs, suggesting a similar mode of regulation as the worm *let-60* [110]. Gain of function mutations in *let-60* result in a multivulval phenotype which is partially suppressed by mir-84 overexpression [112]. Accumulating evidence suggests that *let-7* acts as a tumour suppressor gene. Similarly, miR-61 which

1  
2  
3  
4  
5  
6 targets *vav-1*, a homologue of the Vav oncogene family in vertebrates [113] may  
7  
8 influence cancer through the regulation of Notch signalling [111].  
9

10  
11 The role of apoptosis in cancer treatment sensitivity is well established. Apoptosis  
12 is a genetically regulated cell death mechanism that is required for normal development  
13 and homeostasis. Defects in key apoptotic proteins can interfere with tumourigenesis and  
14 chemotherapy [114]. Therefore, understanding at the molecular level the events that  
15 affect DNA damage- induced apoptosis is important for cancer research. The apoptotic  
16 molecular pathway is highly conserved [115].  
17  
18  
19  
20  
21  
22  
23  
24

25 In *C. elegans*, germline apoptosis limits the number of oocytes competing for  
26 nutrients in the gonad. It can also be activated in response to pathogenic microorganisms  
27 invasion or DNA damageing agents and genotoxic stress [116]. The p-53/ CEP-1, CED-9  
28 (Bcl-2), CED-4 (Apaf1) and CED-3 (caspase) proteins function in the checkpoint  
29 machinery of DNA damage-induced germline apoptosis. A recent study reveals that the  
30 hypoxia-inducible transcription factor (HIF-1) antagonizes the function of CEP-1 in DNA-  
31 damage induced apoptosis. HIF-1 inhibits apoptosis by transcriptionally upregulating the  
32 tyrosinase family member TYR-2 in the ASJ neurons. Then, TYR-2 is secreted and once  
33 entering the gonad, it blocks apoptosis. These findings yield insight into a novel pathway  
34 through which two single neurons regulate apoptotic fate in another tissue. Ultimately,  
35 this pathway links hypoxia to apoptosis. Knock down of the TYR-2 homologue TRP2 in  
36 human melanoma cells increases basal and cisplatin-induced p53 apoptosis, indicating  
37 an evolutionary conserved function. TYR-2 would be a candidate target in new  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60 therapeutic strategies [117].

## 6 Metabolic diseases

The O-GlcNAc signalling pathway, one of the most evolutionary ancient pathways,  
couples the nutrient-dependent synthesis of UDP-GlcNAc to O-GlcNAc modification of

1  
2  
3  
4  
5  
6 Ser/Thr residues of a large number of nuclear and cytoplasmic protein targets.

7 Accumulating data suggest the involvement of O-GlcNAc cycling in age-associated  
8 human diseases such as insulin resistance, X-linked Dystonia Parkinsonism, cancer and  
9 disease of the immune system.  
10  
11  
12

13  
14  
15 Two highly conserved enzymes are responsible for the cycling of O-GlcNAc on  
16 Ser/Thr residues of target proteins. The glycosyltransferase, OGT and the O-GlcNAcase  
17 which mediate the addition and the removal of O-GlcNAc respectively. The identification  
18 of the enzymes involved in O-GlcNAc cycling led to the detection of their interacting  
19 partners. Two-hybrid screens reveal diverse proteins associated with vesicle and  
20 organelle movements in neurons, mitochondrial movement or factors involved in the  
21 control of gene expression that interact with OGT. In *C. elegans*, two-hybrid analysis  
22 indicates the interaction of OGT-1 with MAPK, PMK-1. The highly conserved hexosamine  
23 biosynthetic pathway is upstream of O-GlcNAc cycling. Accumulating findings suggested  
24 a link between the hexosamine biosynthetic pathway, O-GlcNAc cycling and mammalian  
25 insulin resistance [118].  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 *C. elegans* strains having putative null alleles of *ogt-1* (ortholog of O-GlcNAc  
40 transferase) or *oga-1* (ortholog of O-GlcNAcase) provide versatile platforms for studying  
41 the role of O-GlcNAc in cellular signaling and metabolism. Nuclear transport of  
42 transcription factors is normal in homozygous *ogt-1(ok430)* [119] and *oga-1(ok1207)*  
43 knockout animals [120], yet the *ogt-1(ok430)* mutant completely lack any detectable O-  
44 GlcNAc on nuclear pore proteins, whereas the *oga-1(ok1207)* accumulated O-GlcNAc.  
45 Both mutant strains show increased glycogen and trehalose levels and decreased lipid  
46 storage. The link of human OGT and OGA to insulin resistance, prompted the  
47 examination of the impact of *ogt-1* and *oga-1* deletions on the insulin- like- signaling  
48 pathway controlling nutrient storage, dauer formation, longevity and stress responses in  
49 *C. elegans*. *oga-1 (ok1207)* knockout enhances dauer formation induced by a  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6 temperature sensitive allele of the insulin-like receptor *daf-2* (*e1370*) [120] under  
7  
8 conditions in which the *ogt-1(ok430)* null allele lowers dauer formation [119]. *C. elegans*  
9  
10 is the first animal with viable and fertile knockout alleles of OGA or OGT, which provide  
11  
12 genetically amenable models of non-insulin dependent diabetes. Use of O-GlcNAc  
13  
14 cycling mutants in high-throughput RNAi screening revealed the interaction of *ogt-1* with  
15  
16 essential components of important signaling pathways such as  $\beta$ -catenin. In mammals,  $\beta$ -  
17  
18 catenin triggers key developmental switches including the formation of T-cells and B-cells  
19  
20 in adaptive immunity [121].  
21

22  
23 Accumulating evidence suggests that the enzymes of O-GlcNAc cycling may play  
24  
25 a role in modulating transcription [122, 123]. The impact of null alleles of O-GlcNAc  
26  
27 cycling genes on chromatin structure and gene expression can be globally examined in  
28  
29 *C. elegans*. Whole-genome chromatin immunoprecipitation (ChIP)-on- chip tiling arrays  
30  
31 revealed over 800 promoters where O-GlcNAc cycling occurs including microRNA genes  
32  
33 and multigene operons. Transcriptional profiling of O-GlcNAc knockout mutants  
34  
35 confirmed the deregulation of numerous genes associated with insulin-like signaling,  
36  
37 ageing, stress response, innate immunity and carbohydrate/lipid metabolism [124]. These  
38  
39 findings suggest that altered O-GlcNAc cycling has important implications in the diseases  
40  
41 associated with ageing, including obesity, type II diabetes mellitus, cardiovascular  
42  
43 diseases, cancer and neurodegenerative diseases.  
44  
45

46  
47 In *C. elegans*, insulin and its signaling pathway is implicated in the regulation of  
48  
49 fat storage and reproduction and in modulating lifespan. *daf-2*, the homologue of the  
50  
51 mammalian insulin receptor, is part of a signal transduction cascade homologous to the  
52  
53 insulin pathway in mammals. DAF-2 knockout induces dauer formation and extends  
54  
55 lifespan. Restoration of DAF-2 function specifically in neurons is sufficient to restore  
56  
57 lifespan and reproduction of *daf-2* knockouts to wild-type levels. Growing evidence  
58  
59 suggests that impaired insulin signaling in the central nervous system (CNS) plays a  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

fundamental role in the pathogenesis of common metabolic diseases, including obesity and type 2 diabetes [125].

*C. elegans* provides a versatile platform for dissecting the molecular mechanisms underlying glucose-mediated impact on cellular and mitochondrial function in diabetes. An intracellular glucose concentration of 14mmol/l, which is within the range observed in poorly controlled diabetes patients, results in methylglyoxal-derived modifications of mitochondrial proteins and increase in ROS formation thus reducing lifespan in *C. elegans*. These effects are independent of osmotic pressure and the pathways activated by caloric restriction in *eat-2* mutants and the insulin-like receptor *daf-2/daf-16* pathway. Conversely, high glucose-mediated lifespan reduction in *C. elegans* is dependent on glyoxalase -1-controlled mitochondrial complexes II and III. Further studies should delineate the relevance of these mechanisms described in *C. elegans* to the situation in diabetic patients [126].

MicroRNAs, have also been implicated in the regulation of glucose and lipid metabolism and hence in the initiation and progression of diabetes and its specific complications. miRNAs which play a critical role in the pathogenesis of the disease may serve as potential biomarkers for prognosis and diagnosis of diabetes. Furthermore, the altered expression profiles of miRNAs in the pancreas and insulin-targeted tissues in diabetic patients could provide insights for treating this complex metabolic disease [127].

## 7 Concluding remarks

Accumulating evidence demonstrates that most of the important pathways such as insulin signalling, Ras/Notch signalling, p53, many miRNAs have remain essentially unchanged during evolution. Many human diseases, including cancer, originate from mutations in genes functioning in pathways conserved in *C. elegans*. Although nematode disease models might not perfectly recapitulate the pathophysiology of human disease, the small,

1  
2  
3  
4  
5 transparent worm enables the use of powerful genetic and molecular approaches to  
6 dissect the toxicity mechanisms and determine the *in vivo* factors that exacerbate or cure  
7 pathological conditions. Besides being an excellent model for drug target identification, *C.*  
8 *elegans* is amenable to high-throughput screens for rapid and inexpensive drug  
9 evaluation before embarking on expensive and elaborate animal models.  
10  
11  
12  
13  
14  
15  
16  
17

18 *N.T. is supported by grants from the European Research Council (ERC), the*  
19 *European Commission 6th and 7th Framework programs and the European*  
20 *Molecular Biology Organization (EMBO).*  
21

22  
23 *The authors have declared no conflict of interest.*  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 8 References

- [1] Brenner, S., The genetics of *Caenorhabditis elegans*. *Genetics* 1974, *77*, 71-94.
- [2] Sulston, J. E., Schierenberg, E., White, J. G., Thomson, J. N., The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Dev Biol* 1983, *100*, 64-119.
- [3] Harris, T. W., Chen, N., Cunningham, F., Tello-Ruiz, M., *et al.*, WormBase: a multi-species resource for nematode biology and genomics. *Nucleic Acids Res* 2004, *32*, D411-417.
- [4] Hariharan, I. K., Haber, D. A., Yeast, flies, worms, and fish in the study of human disease. *N Engl J Med* 2003, *348*, 2457-2463.
- [5] Chen, N., Harris, T. W., Antoshechkin, I., Bastiani, C., *et al.*, WormBase: a comprehensive data resource for *Caenorhabditis* biology and genomics. *Nucleic Acids Res* 2005, *33*, D383-389.
- [6] Kamath, R. S., Ahringer, J., Genome-wide RNAi screening in *Caenorhabditis elegans*. *Methods* 2003, *30*, 313-321.
- [7] Fire, A., Xu, S., Montgomery, M. K., Kostas, S. A., *et al.*, Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 1998, *391*, 806-811.
- [8] Baumeister, R., Ge, L., The worm in us - *Caenorhabditis elegans* as a model of human disease. *Trends Biotechnol* 2002, *20*, 147-148.
- [9] Poulin, G., Nandakumar, R., Ahringer, J., Genome-wide RNAi screens in *Caenorhabditis elegans*: impact on cancer research. *Oncogene* 2004, *23*, 8340-8345.
- [10] Kenyon, C., The plasticity of aging: insights from long-lived mutants. *Cell* 2005, *120*, 449-460.
- [11] Kosik, K. S., Joachim, C. L., Selkoe, D. J., Microtubule-associated protein tau (tau) is a major antigenic component of paired helical filaments in Alzheimer disease. *Proc Natl Acad Sci U S A* 1986, *83*, 4044-4048.
- [12] Glenner, G. G., Wong, C. W., Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem Biophys Res Commun* 1984, *120*, 885-890.
- [13] Hornsten, A., Lieberthal, J., Fadia, S., Malins, R., *et al.*, APL-1, a *Caenorhabditis elegans* protein related to the human beta-amyloid precursor protein, is essential for viability. *Proc Natl Acad Sci U S A* 2007, *104*, 1971-1976.
- [14] Link, C. D., *C. elegans* models of age-associated neurodegenerative diseases: lessons from transgenic worm models of Alzheimer's disease. *Exp Gerontol* 2006, *41*, 1007-1013.
- [15] Wu, Y., Cao, Z., Klein, W. L., Luo, Y., Heat shock treatment reduces beta amyloid toxicity in vivo by diminishing oligomers. *Neurobiol Aging*, *31*, 1055-1058.
- [16] Hsu, A. L., Murphy, C. T., Kenyon, C., Regulation of aging and age-related disease by DAF-16 and heat-shock factor. *Science* 2003, *300*, 1142-1145.
- [17] Wu, Y., Wu, Z., Butko, P., Christen, Y., *et al.*, Amyloid-beta-induced pathological behaviors are suppressed by Ginkgo biloba extract EGb 761 and ginkgolides in transgenic *Caenorhabditis elegans*. *J Neurosci* 2006, *26*, 13102-13113.
- [18] Hassan, W. M., Merin, D. A., Fonte, V., Link, C. D., AIP-1 ameliorates beta-amyloid peptide toxicity in a *Caenorhabditis elegans* Alzheimer's disease model. *Hum Mol Genet* 2009, *18*, 2739-2747.

- 1  
2  
3  
4  
5  
6 [19] Yun, C., Stanhill, A., Yang, Y., Zhang, Y., *et al.*, Proteasomal adaptation to  
7 environmental stress links resistance to proteotoxicity with longevity in  
8 *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* 2008, *105*, 7094-7099.
- 9 [20] Diomede, L., Cassata, G., Fiordaliso, F., Salio, M., *et al.*, Tetracycline and its  
10 analogues protect *Caenorhabditis elegans* from beta amyloid-induced toxicity by  
11 targeting oligomers. *Neurobiol Dis*.
- 12 [21] Link, C. D., Invertebrate models of Alzheimer's disease. *Genes Brain Behav* 2005, *4*,  
13 147-156.
- 14 [22] Goedert, M., Jakes, R., Mutations causing neurodegenerative tauopathies. *Biochim*  
15 *Biophys Acta* 2005, *1739*, 240-250.
- 16 [23] Iqbal, K., Alonso Adel, C., Chen, S., Chohan, M. O., *et al.*, Tau pathology in  
17 Alzheimer disease and other tauopathies. *Biochim Biophys Acta* 2005, *1739*, 198-  
18 210.
- 19 [24] Lee, V. M., Goedert, M., Trojanowski, J. Q., Neurodegenerative tauopathies. *Annu*  
20 *Rev Neurosci* 2001, *24*, 1121-1159.
- 21 [25] Brandt, R., Gergou, A., Wacker, I., Fath, T., Hutter, H., A *Caenorhabditis elegans*  
22 model of tau hyperphosphorylation: induction of developmental defects by  
23 transgenic overexpression of Alzheimer's disease-like modified tau. *Neurobiol*  
24 *Aging* 2009, *30*, 22-33.
- 25 [26] Kraemer, B. C., Burgess, J. K., Chen, J. H., Thomas, J. H., Schellenberg, G. D.,  
26 Molecular pathways that influence human tau-induced pathology in *Caenorhabditis*  
27 *elegans*. *Hum Mol Genet* 2006, *15*, 1483-1496.
- 28 [27] Kraemer, B. C., Schellenberg, G. D., SUT-1 enables tau-induced neurotoxicity in *C.*  
29 *elegans*. *Hum Mol Genet* 2007, *16*, 1959-1971.
- 30 [28] Guthrie, C. R., Schellenberg, G. D., Kraemer, B. C., SUT-2 potentiates tau-induced  
31 neurotoxicity in *Caenorhabditis elegans*. *Hum Mol Genet* 2009, *18*, 1825-1838.
- 32 [29] Orr, H. T., Beyond the Qs in the polyglutamine diseases. *Genes Dev* 2001, *15*, 925-  
33 932.
- 34 [30] Ross, C. A., Polyglutamine pathogenesis: emergence of unifying mechanisms for  
35 Huntington's disease and related disorders. *Neuron* 2002, *35*, 819-822.
- 36 [31] Orr, H. T., Zoghbi, H. Y., Trinucleotide repeat disorders. *Annu Rev Neurosci* 2007,  
37 *30*, 575-621.
- 38 [32] Morley, J. F., Brignull, H. R., Weyers, J. J., Morimoto, R. I., The threshold for  
39 polyglutamine-expansion protein aggregation and cellular toxicity is dynamic and  
40 influenced by aging in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* 2002, *99*,  
41 10417-10422.
- 42 [33] Brignull, H. R., Moore, F. E., Tang, S. J., Morimoto, R. I., Polyglutamine proteins at  
43 the pathogenic threshold display neuron-specific aggregation in a pan-neuronal  
44 *Caenorhabditis elegans* model. *J Neurosci* 2006, *26*, 7597-7606.
- 45 [34] Gidalevitz, T., Ben-Zvi, A., Ho, K. H., Brignull, H. R., Morimoto, R. I., Progressive  
46 disruption of cellular protein folding in models of polyglutamine diseases. *Science*  
47 2006, *311*, 1471-1474.
- 48 [35] Guarente, L., Kenyon, C., Genetic pathways that regulate ageing in model  
49 organisms. *Nature* 2000, *408*, 255-262.
- 50 [36] Caldwell, G. A., Cao, S., Sexton, E. G., Gelwix, C. C., *et al.*, Suppression of  
51 polyglutamine-induced protein aggregation in *Caenorhabditis elegans* by torsin  
52 proteins. *Hum Mol Genet* 2003, *12*, 307-319.
- 53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6 [37] Walters, K. J., Kleijnen, M. F., Goh, A. M., Wagner, G., Howley, P. M., Structural  
7 studies of the interaction between ubiquitin family proteins and proteasome subunit  
8 S5a. *Biochemistry* 2002, *41*, 1767-1777.
- 9 [38] Wang, H., Lim, P. J., Yin, C., Rieckher, M., *et al.*, Suppression of polyglutamine-  
10 induced toxicity in cell and animal models of Huntington's disease by ubiquilin.  
11 *Hum Mol Genet* 2006, *15*, 1025-1041.
- 12 [39] Wang, H., Lim, P. J., Karbowski, M., Monteiro, M. J., Effects of overexpression of  
13 huntingtin proteins on mitochondrial integrity. *Hum Mol Genet* 2009, *18*, 737-752.
- 14 [40] Link, C. D., Expression of human beta-amyloid peptide in transgenic *Caenorhabditis*  
15 *elegans*. *Proc Natl Acad Sci U S A* 1995, *92*, 9368-9372.
- 16 [41] Steinkraus, K. A., Smith, E. D., Davis, C., Carr, D., *et al.*, Dietary restriction  
17 suppresses proteotoxicity and enhances longevity by an hsf-1-dependent  
18 mechanism in *Caenorhabditis elegans*. *Aging Cell* 2008, *7*, 394-404.
- 19 [42] Vila, M., Przedborski, S., Genetic clues to the pathogenesis of Parkinson's disease.  
20 *Nat Med* 2004, *10 Suppl*, S58-62.
- 21 [43] Polymeropoulos, M. H., Lavedan, C., Leroy, E., Ide, S. E., *et al.*, Mutation in the  
22 alpha-synuclein gene identified in families with Parkinson's disease. *Science* 1997,  
23 *276*, 2045-2047.
- 24 [44] Singleton, A. B., Farrer, M., Johnson, J., Singleton, A., *et al.*, alpha-Synuclein locus  
25 triplication causes Parkinson's disease. *Science* 2003, *302*, 841.
- 26 [45] Chartier-Harlin, M. C., Kachergus, J., Roumier, C., Mouroux, V., *et al.*, Alpha-  
27 synuclein locus duplication as a cause of familial Parkinson's disease. *Lancet*  
28 2004, *364*, 1167-1169.
- 29 [46] Dauer, W., Przedborski, S., Parkinson's disease: mechanisms and models. *Neuron*  
30 2003, *39*, 889-909.
- 31 [47] Cooper, A. A., Gitler, A. D., Cashikar, A., Haynes, C. M., *et al.*, Alpha-synuclein  
32 blocks ER-Golgi traffic and Rab1 rescues neuron loss in Parkinson's models.  
33 *Science* 2006, *313*, 324-328.
- 34 [48] Hamamichi, S., Rivas, R. N., Knight, A. L., Cao, S., *et al.*, Hypothesis-based RNAi  
35 screening identifies neuroprotective genes in a Parkinson's disease model. *Proc*  
36 *Natl Acad Sci U S A* 2008, *105*, 728-733.
- 37 [49] Shashidharan, P., Good, P. F., Hsu, A., Perl, D. P., *et al.*, TorsinA accumulation in  
38 Lewy bodies in sporadic Parkinson's disease. *Brain Res* 2000, *877*, 379-381.
- 39 [50] van Ham, T. J., Thijssen, K. L., Breitling, R., Hofstra, R. M., *et al.*, *C. elegans* model  
40 identifies genetic modifiers of alpha-synuclein inclusion formation during aging.  
41 *PLoS Genet* 2008, *4*, e1000027.
- 42 [51] Paisan-Ruiz, C., Jain, S., Evans, E. W., Gilks, W. P., *et al.*, Cloning of the gene  
43 containing mutations that cause PARK8-linked Parkinson's disease. *Neuron* 2004,  
44 *44*, 595-600.
- 45 [52] Zimprich, A., Biskup, S., Leitner, P., Lichtner, P., *et al.*, Mutations in LRRK2 cause  
46 autosomal-dominant parkinsonism with pleomorphic pathology. *Neuron* 2004, *44*,  
47 601-607.
- 48 [53] Saha, S., Guillily, M. D., Ferree, A., Lanceta, J., *et al.*, LRRK2 modulates vulnerability  
49 to mitochondrial dysfunction in *Caenorhabditis elegans*. *J Neurosci* 2009, *29*, 9210-  
50 9218.
- 51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6 [54] West, A. B., Moore, D. J., Biskup, S., Bugayenko, A., *et al.*, Parkinson's disease-  
7 associated mutations in leucine-rich repeat kinase 2 augment kinase activity. *Proc*  
8 *Natl Acad Sci U S A* 2005, *102*, 16842-16847.
- 9 [55] Gloeckner, C. J., Kinkl, N., Schumacher, A., Braun, R. J., *et al.*, The Parkinson  
10 disease causing LRRK2 mutation I2020T is associated with increased kinase  
11 activity. *Hum Mol Genet* 2006, *15*, 223-232.
- 12 [56] Yao, C., El Khoury, R., Wang, W., Byrd, T. A., *et al.*, LRRK2-mediated  
13 neurodegeneration and dysfunction of dopaminergic neurons in a *Caenorhabditis*  
14 *elegans* model of Parkinson's disease. *Neurobiol Dis.*
- 15 [57] Mulder, D. W., Kurland, L. T., Offord, K. P., Beard, C. M., Familial adult motor neuron  
16 disease: amyotrophic lateral sclerosis. *Neurology* 1986, *36*, 511-517.
- 17 [58] Cudkovicz, M. E., McKenna-Yasek, D., Sapp, P. E., Chin, W., *et al.*, Epidemiology of  
18 mutations in superoxide dismutase in amyotrophic lateral sclerosis. *Ann Neurol*  
19 1997, *41*, 210-221.
- 20 [59] Hart, P. J., Pathogenic superoxide dismutase structure, folding, aggregation and  
21 turnover. *Curr Opin Chem Biol* 2006, *10*, 131-138.
- 22 [60] Oeda, T., Shimohama, S., Kitagawa, N., Kohno, R., *et al.*, Oxidative stress causes  
23 abnormal accumulation of familial amyotrophic lateral sclerosis-related mutant  
24 SOD1 in transgenic *Caenorhabditis elegans*. *Hum Mol Genet* 2001, *10*, 2013-2023.
- 25 [61] Wang, J., Farr, G. W., Hall, D. H., Li, F., *et al.*, An ALS-linked mutant SOD1 produces  
26 a locomotor defect associated with aggregation and synaptic dysfunction when  
27 expressed in neurons of *Caenorhabditis elegans*. *PLoS Genet* 2009, *5*, e1000350.
- 28 [62] Gidalevitz, T., Krupinski, T., Garcia, S., Morimoto, R. I., Destabilizing protein  
29 polymorphisms in the genetic background direct phenotypic expression of mutant  
30 SOD1 toxicity. *PLoS Genet* 2009, *5*, e1000399.
- 31 [63] Nishimura, A. L., Mitne-Neto, M., Silva, H. C., Richieri-Costa, A., *et al.*, A mutation in  
32 the vesicle-trafficking protein VAPB causes late-onset spinal muscular atrophy and  
33 amyotrophic lateral sclerosis. *Am J Hum Genet* 2004, *75*, 822-831.
- 34 [64] Soussan, L., Burakov, D., Daniels, M. P., Toister-Achituv, M., *et al.*, ERG30, a VAP-  
35 33-related protein, functions in protein transport mediated by COPI vesicles. *J Cell*  
36 *Biol* 1999, *146*, 301-311.
- 37 [65] Skehel, P. A., Fabian-Fine, R., Kandel, E. R., Mouse VAP33 is associated with the  
38 endoplasmic reticulum and microtubules. *Proc Natl Acad Sci U S A* 2000, *97*,  
39 1101-1106.
- 40 [66] Kaiser, S. E., Brickner, J. H., Reilein, A. R., Fenn, T. D., *et al.*, Structural basis of  
41 FFAT motif-mediated ER targeting. *Structure* 2005, *13*, 1035-1045.
- 42 [67] Nishimura, Y., Hayashi, M., Inada, H., Tanaka, T., Molecular cloning and  
43 characterization of mammalian homologues of vesicle-associated membrane  
44 protein-associated (VAMP-associated) proteins. *Biochem Biophys Res Commun*  
45 1999, *254*, 21-26.
- 46 [68] Bottino, D., Mogilner, A., Roberts, T., Stewart, M., Oster, G., How nematode sperm  
47 crawl. *J Cell Sci* 2002, *115*, 367-384.
- 48 [69] Miller, M. A., Nguyen, V. Q., Lee, M. H., Kosinski, M., *et al.*, A sperm cytoskeletal  
49 protein that signals oocyte meiotic maturation and ovulation. *Science* 2001, *291*,  
50 2144-2147.
- 51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6 [70] Miller, M. A., Ruest, P. J., Kosinski, M., Hanks, S. K., Greenstein, D., An Eph  
7 receptor sperm-sensing control mechanism for oocyte meiotic maturation in  
8 *Caenorhabditis elegans*. *Genes Dev* 2003, *17*, 187-200.
- 9 [71] Tsuda, H., Han, S. M., Yang, Y., Tong, C., *et al.*, The amyotrophic lateral sclerosis 8  
10 protein VAPB is cleaved, secreted, and acts as a ligand for Eph receptors. *Cell*  
11 2008, *133*, 963-977.
- 12 [72] Ogino, S., Wilson, R. B., Genetic testing and risk assessment for spinal muscular  
13 atrophy (SMA). *Hum Genet* 2002, *111*, 477-500.
- 14 [73] Lefebvre, S., Burglen, L., Reboullet, S., Clermont, O., *et al.*, Identification and  
15 characterization of a spinal muscular atrophy-determining gene. *Cell* 1995, *80*,  
16 155-165.
- 17 [74] Briese, M., Esmaili, B., Sattelle, D. B., Is spinal muscular atrophy the result of  
18 defects in motor neuron processes? *Bioessays* 2005, *27*, 946-957.
- 19 [75] Miguel-Aliaga, I., Culetto, E., Walker, D. S., Baylis, H. A., *et al.*, The *Caenorhabditis*  
20 *elegans* orthologue of the human gene responsible for spinal muscular atrophy is a  
21 maternal product critical for germline maturation and embryonic viability. *Hum Mol*  
22 *Genet* 1999, *8*, 2133-2143.
- 23 [76] Briese, M., Esmaili, B., Fraboulet, S., Burt, E. C., *et al.*, Deletion of *smn-1*, the  
24 *Caenorhabditis elegans* ortholog of the spinal muscular atrophy gene, results in  
25 locomotor dysfunction and reduced lifespan. *Hum Mol Genet* 2009, *18*, 97-104.
- 26 [77] Dimitriadi, M., Hart, A. C., Neurodegenerative disorders: Insights from the nematode  
27 *Caenorhabditis elegans*. *Neurobiol Dis*.
- 28 [78] Syntichaki, P., Tavernarakis, N., Death by necrosis. Uncontrollable catastrophe, or is  
29 there order behind the chaos? *EMBO Rep* 2002, *3*, 604-609.
- 30 [79] Scott, B. A., Avidan, M. S., Crowder, C. M., Regulation of hypoxic death in *C.*  
31 *elegans* by the insulin/IGF receptor homolog DAF-2. *Science* 2002, *296*, 2388-  
32 2391.
- 33 [80] Syntichaki, P., Tavernarakis, N., The biochemistry of neuronal necrosis: rogue  
34 biology? *Nat Rev Neurosci* 2003, *4*, 672-684.
- 35 [81] Syntichaki, P., Xu, K., Driscoll, M., Tavernarakis, N., Specific aspartyl and calpain  
36 proteases are required for neurodegeneration in *C. elegans*. *Nature* 2002, *419*,  
37 939-944.
- 38 [82] Xu, W., Wong, T. P., Chery, N., Gaertner, T., *et al.*, Calpain-mediated mGluR1alpha  
39 truncation: a key step in excitotoxicity. *Neuron* 2007, *53*, 399-412.
- 40 [83] Syntichaki, P., Samara, C., Tavernarakis, N., The vacuolar H<sup>+</sup>-ATPase mediates  
41 intracellular acidification required for neurodegeneration in *C. elegans*. *Curr Biol*  
42 2005, *15*, 1249-1254.
- 43 [84] Artal-Sanz, M., Samara, C., Syntichaki, P., Tavernarakis, N., Lysosomal biogenesis  
44 and function is critical for necrotic cell death in *Caenorhabditis elegans*. *J Cell Biol*  
45 2006, *173*, 231-239.
- 46 [85] Ervasti, J. M., Campbell, K. P., Membrane organization of the dystrophin-  
47 glycoprotein complex. *Cell* 1991, *66*, 1121-1131.
- 48 [86] Ervasti, J. M., Campbell, K. P., A role for the dystrophin-glycoprotein complex as a  
49 transmembrane linker between laminin and actin. *J Cell Biol* 1993, *122*, 809-823.
- 50 [87] Giesel, K., Bessou, C., Segalat, L., Dystrobrevin- and dystrophin-like mutants  
51 display similar phenotypes in the nematode *Caenorhabditis elegans*.  
52 *Neurogenetics* 1999, *2*, 87-90.
- 53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3  
4  
5  
6 [88] Kim, H., Rogers, M. J., Richmond, J. E., McIntire, S. L., SNF-6 is an acetylcholine  
7 transporter interacting with the dystrophin complex in *Caenorhabditis elegans*.  
8 *Nature* 2004, *430*, 891-896.
- 9 [89] Kim, H., Pierce-Shimomura, J. T., Oh, H. J., Johnson, B. E., *et al.*, The dystrophin  
10 complex controls bk channel localization and muscle activity in *Caenorhabditis*  
11 *elegans*. *PLoS Genet* 2009, *5*, e1000780.
- 12 [90] Bulfield, G., Siller, W. G., Wight, P. A., Moore, K. J., X chromosome-linked muscular  
13 dystrophy (*mdx*) in the mouse. *Proc Natl Acad Sci U S A* 1984, *81*, 1189-1192.
- 14 [91] Ballinger, C. A., Connell, P., Wu, Y., Hu, Z., *et al.*, Identification of CHIP, a novel  
15 tetratricopeptide repeat-containing protein that interacts with heat shock proteins  
16 and negatively regulates chaperone functions. *Mol Cell Biol* 1999, *19*, 4535-4545.
- 17 [92] Nyamsuren, O., Faggionato, D., Loch, W., Schulze, E., Baumeister, R., A mutation in  
18 CHN-1/CHIP suppresses muscle degeneration in *Caenorhabditis elegans*. *Dev Biol*  
19 2007, *312*, 193-202.
- 20 [93] Brais, B., Oculopharyngeal muscular dystrophy: a late-onset polyalanine disease.  
21 *Cytogenet Genome Res* 2003, *100*, 252-260.
- 22 [94] Wang, Q., Mosser, D. D., Bag, J., Induction of HSP70 expression and recruitment of  
23 HSC70 and HSP70 in the nucleus reduce aggregation of a polyalanine expansion  
24 mutant of PABPN1 in HeLa cells. *Hum Mol Genet* 2005, *14*, 3673-3684.
- 25 [95] Klein, A. F., Ebihara, M., Alexander, C., Dicaire, M. J., *et al.*, PABPN1 polyalanine  
26 tract deletion and long expansions modify its aggregation pattern and expression.  
27 *Exp Cell Res* 2008, *314*, 1652-1666.
- 28 [96] Abu-Baker, A., Messaed, C., Laganier, J., Gaspar, C., *et al.*, Involvement of the  
29 ubiquitin-proteasome pathway and molecular chaperones in oculopharyngeal  
30 muscular dystrophy. *Hum Mol Genet* 2003, *12*, 2609-2623.
- 31 [97] Catoire, H., Pasco, M. Y., Abu-Baker, A., Holbert, S., *et al.*, Sirtuin inhibition protects  
32 from the polyalanine muscular dystrophy protein PABPN1. *Hum Mol Genet* 2008,  
33 *17*, 2108-2117.
- 34 [98] Di Renzo, M. F., Olivero, M., Katsaros, D., Crepaldi, T., *et al.*, Overexpression of the  
35 Met/HGF receptor in ovarian cancer. *Int J Cancer* 1994, *58*, 658-662.
- 36 [99] Di Renzo, M. F., Olivero, M., Giacomini, A., Porte, H., *et al.*, Overexpression and  
37 amplification of the met/HGF receptor gene during the progression of colorectal  
38 cancer. *Clin Cancer Res* 1995, *1*, 147-154.
- 39 [100] Ferracini, R., Di Renzo, M. F., Scotlandi, K., Baldini, N., *et al.*, The Met/HGF  
40 receptor is over-expressed in human osteosarcomas and is activated by either a  
41 paracrine or an autocrine circuit. *Oncogene* 1995, *10*, 739-749.
- 42 [101] Hellman, A., Zlotorynski, E., Scherer, S. W., Cheung, J., *et al.*, A role for common  
43 fragile site induction in amplification of human oncogenes. *Cancer Cell* 2002, *1*, 89-  
44 97.
- 45 [102] Maulik, G., Kijima, T., Ma, P. C., Ghosh, S. K., *et al.*, Modulation of the c-  
46 Met/hepatocyte growth factor pathway in small cell lung cancer. *Clin Cancer Res*  
47 2002, *8*, 620-627.
- 48 [103] Ma, P. C., Kijima, T., Maulik, G., Fox, E. A., *et al.*, c-MET mutational analysis in  
49 small cell lung cancer: novel juxtamembrane domain mutations regulating  
50 cytoskeletal functions. *Cancer Res* 2003, *63*, 6272-6281.
- 51 [104] Ma, P. C., Jagadeeswaran, R., Jagadeesh, S., Tretiakova, M. S., *et al.*, Functional  
52 expression and mutations of c-Met and its therapeutic inhibition with SU11274 and  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6 small interfering RNA in non-small cell lung cancer. *Cancer Res* 2005, *65*, 1479-  
7 1488.
- 8 [105] Siddiqui, S. S., Loganathan, S., Krishnaswamy, S., Faoro, L., *et al.*, *C. elegans* as a  
9 model organism for in vivo screening in cancer: effects of human c-Met in lung  
10 cancer affect *C. elegans* vulva phenotypes. *Cancer Biol Ther* 2008, *7*, 856-863.
- 11 [106] Tan, P. B., Kim, S. K., Signaling specificity: the RTK/RAS/MAP kinase pathway in  
12 metazoans. *Trends Genet* 1999, *15*, 145-149.
- 13 [107] Moghal, N., Sternberg, P. W., The epidermal growth factor system in  
14 *Caenorhabditis elegans*. *Exp Cell Res* 2003, *284*, 150-159.
- 15 [108] Nilsen, T. W., Mechanisms of microRNA-mediated gene regulation in animal cells.  
16 *Trends Genet* 2007, *23*, 243-249.
- 17 [109] Sundaram, M. V., The love-hate relationship between Ras and Notch. *Genes Dev*  
18 2005, *19*, 1825-1839.
- 19 [110] Johnson, S. M., Grosshans, H., Shingara, J., Byrom, M., *et al.*, RAS is regulated by  
20 the let-7 microRNA family. *Cell* 2005, *120*, 635-647.
- 21 [111] Yoo, A. S., Greenwald, I., LIN-12/Notch activation leads to microRNA-mediated  
22 down-regulation of Vav in *C. elegans*. *Science* 2005, *310*, 1330-1333.
- 23 [112] Esquela-Kerscher, A., Johnson, S. M., Bai, L., Saito, K., *et al.*, Post-embryonic  
24 expression of *C. elegans* microRNAs belonging to the lin-4 and let-7 families in the  
25 hypodermis and the reproductive system. *Dev Dyn* 2005, *234*, 868-877.
- 26 [113] Tybulewicz, V. L., Vav-family proteins in T-cell signalling. *Curr Opin Immunol* 2005,  
27 *17*, 267-274.
- 28 [114] Johnstone, R. W., Ruefli, A. A., Lowe, S. W., Apoptosis: a link between cancer  
29 genetics and chemotherapy. *Cell* 2002, *108*, 153-164.
- 30 [115] Ellis, H. M., Horvitz, H. R., Genetic control of programmed cell death in the  
31 nematode *C. elegans*. *Cell* 1986, *44*, 817-829.
- 32 [116] van Haaften, G., Romeijn, R., Pothof, J., Koole, W., *et al.*, Identification of  
33 conserved pathways of DNA-damage response and radiation protection by  
34 genome-wide RNAi. *Curr Biol* 2006, *16*, 1344-1350.
- 35 [117] Sendoel, A., Kohler, I., Fellmann, C., Lowe, S. W., Hengartner, M. O., HIF-1  
36 antagonizes p53-mediated apoptosis through a secreted neuronal tyrosinase.  
37 *Nature*, *465*, 577-583.
- 38 [118] Hanover, J. A., Krause, M. W., Love, D. C., The hexosamine signaling pathway: O-  
39 GlcNAc cycling in feast or famine. *Biochim Biophys Acta*, *1800*, 80-95.
- 40 [119] Hanover, J. A., Forsythe, M. E., Hennessey, P. T., Brodigan, T. M., *et al.*, A  
41 *Caenorhabditis elegans* model of insulin resistance: altered macronutrient storage  
42 and dauer formation in an OGT-1 knockout. *Proc Natl Acad Sci U S A* 2005, *102*,  
43 11266-11271.
- 44 [120] Forsythe, M. E., Love, D. C., Lazarus, B. D., Kim, E. J., *et al.*, *Caenorhabditis*  
45 *elegans* ortholog of a diabetes susceptibility locus: oga-1 (O-GlcNAcase) knockout  
46 impacts O-GlcNAc cycling, metabolism, and dauer. *Proc Natl Acad Sci U S A* 2006,  
47 *103*, 11952-11957.
- 48 [121] Hanover, J. A., Love, D. C., Prinz, W. A., Calmodulin-driven nuclear entry: trigger  
49 for sex determination and terminal differentiation. *J Biol Chem* 2009, *284*, 12593-  
50 12597.
- 51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6 [122] Sinclair, D. A., Syrzycka, M., Macauley, M. S., Rastgardani, T., *et al.*, *Drosophila* O-  
7 GlcNAc transferase (OGT) is encoded by the Polycomb group (PcG) gene, super  
8 sex combs (*sxc*). *Proc Natl Acad Sci U S A* 2009, *106*, 13427-13432.
- 9 [123] Gambetta, M. C., Oktaba, K., Muller, J., Essential role of the glycosyltransferase  
10 *sxc/Ogt* in polycomb repression. *Science* 2009, *325*, 93-96.
- 11 [124] Love, D. C., Ghosh, S., Mondoux, M. A., Fukushige, T., *et al.*, Dynamic O-GlcNAc  
12 cycling at promoters of *Caenorhabditis elegans* genes regulating longevity, stress,  
13 and immunity. *Proc Natl Acad Sci U S A*, *107*, 7413-7418.
- 14 [125] Porte, D., Jr., Baskin, D. G., Schwartz, M. W., Insulin signaling in the central  
15 nervous system: a critical role in metabolic homeostasis and disease from *C.*  
16 *elegans* to humans. *Diabetes* 2005, *54*, 1264-1276.
- 17 [126] Schlotterer, A., Kukudov, G., Bozorgmehr, F., Hutter, H., *et al.*, *C. elegans* as model  
18 for the study of high glucose- mediated life span reduction. *Diabetes* 2009, *58*,  
19 2450-2456.
- 20 [127] Pandey, A. K., Agarwal, P., Kaur, K., Datta, M., MicroRNAs in diabetes: tiny players  
21 in big disease. *Cell Physiol Biochem* 2009, *23*, 221-232.
- 22 [128] Pasco, M. Y., Catoire, H., Parker, J. A., Brais, B., *et al.*, Cross-talk between  
23 canonical Wnt signaling and the sirtuin-FoxO longevity pathway to protect against  
24 muscular pathology induced by mutant PABPN1 expression in *C. elegans*.  
25 *Neurobiol Dis*, *38*, 425-433.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1.** Selected human disease models in *Caenorhabditis elegans*.

Disease	Disease-associated protein	Characteristic pathology in <i>C. elegans</i>	<i>C. elegans</i> findings with potential therapeutic relevance
AD	$\beta$ -amyloid peptide	Muscle-associated human $A\beta_{1-42}$ deposits [14] Neuronal $A\beta$ expression induces defect in chemotaxis [17]	Delay of $A\beta_{1-42}$ -mediated paralysis by heat shock treatment [14] and Ginkgolide A [17]
	APP/APL-1	Severe developmental defects induced by inactivation or overexpression of <i>apl-1</i> [13]	AIRAPL/AIP-1 suppresses $A\beta$ toxicity [18] Tetracycline and its analogues protect from $\beta$ -amyloid- induced toxicity [20]
AD-relevant tau	PHP-tau	Insoluble highly phosphorylated tau aggregates in neurons linked to a defective pattern of motor neuron development [25]	
PG	polyQ aggregates	Muscle (polyQm) [32] or neuronal (polyQn) [33] induced toxicity	TOR-2 mediated suppression of polyQ aggregation [36]
HD	Expression of human Htt (Q>40) in body wall muscle	Motility defects [38]	Rescue of disease phenotype by ubiquitin overexpression [38] Amelioration of the motility defect by Drp-1 knockdown [39] Suppression of proteotoxicity by DR [41]
PD	Human $\alpha$ -syn expression in dopaminergic neurons W08D2.5/PARK9, B0432.2/DJ-1, E0432.9/PINK1 LRRK2/LRK-1	Lewy body formation, selective degeneration of dopaminergic neurons [47] Knockdown by RNAi enhances misfolding of human $\alpha$ -synuclein [48] Enhanced loss of dopaminergic neurons [53]	Co-expression of Rab1 rescue the degeneration [47] Co-expression of TOR-2 ameliorates the aggregate formation [48] <i>sir-2.1/SIRT1</i> , <i>lagr-1/LASS2</i> suppress $\alpha$ -synuclein inclusion formation [50] Autophagy confers neuroprotection [48]
ALS	SOD1 ALS8/VPR-1	Pan-neuronal expression caused severe locomotion defect [61] Mild body muscle toxicity influenced by the genetic background [62]	
SMA	SMN/SMN-1	Late larval arrest, decreased lifespan, defects in motility and pharyngeal pumping [76]	<i>smn-1(ok355)</i> deletion mutant as a useful platform for functional analysis of human SMN protein [77]

Stroke- Excitotoxicity	Specific ion channels (DEG-1, MEC-4, DEG-3, GSA-1 in nematode)  Specific proteases (calpains CLP-1, TRA-3 and aspartyl proteases ASP-3, ASP-4 in nematode)	Neurodegeneration [80, 81]	Inhibition of executioner proteases (ASP-3, ASP-4) suppresses necrosis [81]  Alterations of lysosome biogenesis and function by genetic or pharmacological treatment could affect necrotic cell death [84]
DMD	Dystrofin/dys-1	Muscle degeneration [90, 91]	<i>chn-1</i> /CHIP elimination in the musculature suppresses muscle degeneration [92]
OPMD	PABPN1	Muscle cell decline and abnormal motility [97]	Treatment with sirtuin inhibitors protect against mutant PABPN1[97, 128]
Cancer	c-Met	Locomotion defects, low fecundity and vulval hyperplasia [105] Anti-apoptotic factor, antagonizes p53/CEP-1, interferes with tumor progression and therapy [117]	
	<i>let-60</i> /ras oncogenes <i>vav-1</i> /Vav	Multivulval phenotype [109- 111]	
	<i>cep-1</i> /p53	Apoptotic defects interfere with tumorigenesis and resistance to chemotherapeutic drugs [114]	
	HIFa/HIF-1	Anti-apoptotic factor, antagonizes p53/ <i>cep-1</i> , interferes with tumour progression and therapy [117]	Target TYR-2/TRP2 [117]
Metabolic diseases (insulin resistance, type 2 diabetes)	<i>ogt-1</i> , <i>oga-1</i>	Null mutations cause alterations in carbohydrate and lipid metabolism [119, 120, 124]	Overexpression or inhibition of specific miRNAs may be a future tool for targeting diabetes [127]

## Figure legends

**Figure 1. Pathogenesis of polyglutamine diseases.** Polyglutamine expansion disorders are caused by abnormally expanded tracks of polyglutamine residues (polyQ) in seemingly unrelated proteins. In many cases, abnormal proteolytic cleavage of the affected proteins could result in toxic folding intermediates, oligomers and aggregates which ultimately form intracellular inclusions. Damaged proteins cannot be degraded through the ubiquitin (proteosomal) and lysosomal degradation pathways. Thus, the accumulation of unwanted proteins may lead to neurodegeneration. Mitochondrial dysfunction has also been linked to polyglutamine-expansion proteotoxicity.

**Figure 2. The complex interplay between autophagy, cell survival and cell death.** Autophagy is a catabolic process responsible for the degradation of long-lived proteins, molecules and organelles. Cellular material to be degraded is engulfed in double-membrane vesicles, called autophagosomes, which undergo fusion with lysosomes to form autolysosomes where hydrolases degrade the cytoplasmic material. Free fatty acids and amino acids produced by this process can be reutilized by the cell to maintain ATP levels and protein synthesis, contributing to cell survival. Starvation or growth factor deprivation induces autophagy by inhibiting TOR signalling. Autophagy contributes to life span extension under caloric restriction in many organisms including yeast, nematodes, flies and mammals. Mutations in genes functioning in various steps of the autophagy pathway suppress cell death triggered by diverse stimuli. Pharmacological inhibitors block distinct steps of the autophagic pathway (red blocking lines). Autophagy is implicated in neurodegeneration in *C. elegans*. Diverse genetic or environmental insults lead to elevated cytoplasmic calcium concentration disturbing calcium homeostasis. Increased calcium concentration activates cytoplasmic calpain proteases causing lysosomal membrane rupture and release of lysosomal components into the cytoplasm which becomes acidified. The vacuolar H<sup>+</sup> ATPase (V-ATPase) pump is responsible for lysosomal and

1  
2  
3  
4  
5  
6 subsequent cytoplasmic acidification. Low pH conditions favour the activation of  
7  
8 cathepsin proteases contributing to cell death. Autophagy is up-regulated upon induction  
9  
10 of necrosis directly and / or through calpain activation and synergizes with lysosomal  
11  
12 cathepsin proteases to mediate cell death. Similarly impaired autophagy can be  
13  
14 deleterious by allowing damaged organelles and proteins to accumulate. Expression of  
15  
16 human disease genes in worms disturbs cellular homeostatic mechanisms and induces  
17  
18 stress, which beyond a certain threshold, becomes detrimental. TCA: tricarboxylic acid; 3-  
19  
20 MA: 3-methyladenine;  $[Ca^{2+}]$ : cytoplasmic calcium concentration; V-ATPase: vacuolar H<sup>+</sup>-  
21  
22 ATPase; ER: endoplasmic reticulum; SERCA: sarco-endoplasmic reticulum; InsP3R:  
23  
24 inositol triphosphate receptor; RyR: ryanodine receptor.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 ((Author's CVs))  
7

## 8 **Bios**

9

### 12 **Nektarios Bio**

14 Nektarios Tavernarakis is a Research Director at the Institute of Molecular  
15 Biology and Biotechnology, and Professor of Molecular Systems Biology at the  
16 Medical School of the University of Crete in Heraklion, Greece. He earned his  
17 PhD at the University of Crete, studying gene expression in yeast, and trained in  
18 *Caenorhabditis elegans* biology at Rutgers University, New Jersey, USA. His  
19 research focuses on neuronal function and dysfunction. His main interests are the  
20 molecular mechanisms of necrotic cell death and neurodegeneration, the  
21 molecular mechanisms of sensory transduction and integration by the nervous  
22 system, the interplay between cellular metabolism and ageing, and the  
23 development of novel genetic tools for *C. elegans* research. He is the recipient of  
24 a European Research Council (ERC) Advanced Investigator grant award, a  
25 European Molecular Biology Organisation (EMBO) Young Investigator award,  
26 the Alexander von Humboldt Foundation, Friedrich Wilhelm Bessel research  
27 award, and is member of EMBO.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

### 42 **Maria Bio**

44 Maria Markaki is a staff scientist in the *Caenorhabditis elegans* molecular genetics  
45 laboratory at the Institute of Molecular Biology and Biotechnology, in Heraklion,  
46 Crete, Greece. She earned her Ph.D. degree at the University of Crete studying the  
47 ecophysiology of *Oxalis pes-caprae* L. Her main interests lie in the molecular  
48 mechanisms that underlie the effects of autophagy on ageing.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49




Figure 1 - Markaki & Tavernarakis




Figure 2 - Markaki & Tavernarakis

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



73x85mm (150 x 150 DPI)



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



57x76mm (150 x 150 DPI)

