



HAL
open science

CH₄ production and oxidation processes in a boreal fen ecosystem after long-term water table drawdown

Kim Yrjälä, Tero Tuomivirta, Heli Juottonen, Anuliina Putkinen, Kaisa Lappi, Eeva-Stiina Tuittila, Timo Penttilä, Kari Minkkinen, Jukka Laine, Krista Peltoniemi, et al.

► **To cite this version:**

Kim Yrjälä, Tero Tuomivirta, Heli Juottonen, Anuliina Putkinen, Kaisa Lappi, et al.. CH₄ production and oxidation processes in a boreal fen ecosystem after long-term water table drawdown. *Global Change Biology*, 2010, 17 (3), pp.1311. <10.1111/j.1365-2486.2010.02290.x>. <hal-00599524>

HAL Id: hal-00599524

<https://hal.science/hal-00599524v1>

Submitted on 10 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



CH₄ production and oxidation processes in a boreal fen ecosystem after long-term water table drawdown



| | |
|-------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Journal: | <i>Global Change Biology</i> |
| Manuscript ID: | GCB-10-0362 |
| Wiley - Manuscript type: | Primary Research Articles |
| Date Submitted by the Author: | 04-May-2010 |
| Complete List of Authors: | Yrjälä, Kim; University of Helsinki, Department of Biosciences Tuomivirta, Tero; Finnish Forest Research Institute, Southern Unit Juottonen, Heli; University of Helsinki, Department of Biosciences Putkinen, Anuliina; Finnish Forest Research Institute, Southern Unit; University of Helsinki, Department of Biosciences Lappi, Kaisa; University of Helsinki, Department of Biosciences Tuittila, Eeva-Stiina; University of Helsinki, Department of Forest Sciences Penttilä, Timo; Finnish Forest Research Institute, Southern Unit Minkkinen, Kari; University of Helsinki, Department of Forest Sciences Laine, Jukka; Finnish Forest Research Institute, Western Unit Peltoniemi, Krista; Finnish Forest Research Institute, Southern Unit Fritze, Hannu; Finnish Forest Research Institute, Southern Unit |
| Keywords: | greenhouse gas, peatlands, methanogens, methanotrophs, microbial community, DGGE, T-RFLP |
| Abstract: | Mires, especially sedge dominated fens, are sources of the greenhouse gas CH ₄ . Climate change scenarios predict a lowering water table (WT) in mires. To study the effect of WT drawdown on CH ₄ dynamics in a fen ecosystem, we took advantage of a WT drawdown gradient near a ground water extraction plant. Methane fluxes, CH ₄ production and oxidation potentials, were related to microbial communities responsible for the processes in four mire locations (wet, semi-wet, semi-dry and dry). Principal component analyses (PCA) performed on the vegetation, pH, CH ₄ and WT results clearly separated the four sampling locations in the gradient. Long-term lowering of WT was associated with decreased coverage of Sphagnum and aerenchymatic plants, decreased CH ₄ field emissions and CH ₄ production potential. Based on mcrA T-RF the methanogen community structure correlated best with the methane production and coverage of aerenchymatic plants along the gradient. Methanosarcinaceae and Methanocellales were found at the pristine wet end of the gradient, whereas the Fen cluster |

1
2
3
4 characterized the dry end. The methane-oxidizing bacterial (MOB)
5 community consisted exclusively of Methylocystis bacteria, but
6 interestingly of five different alleles (T, S, R, M, and O) of the
7 particulate methane monooxygenase marker gene pmoA. The M
8 allele was dominant in the wet locations, and the occurrence of
9 alleles O, S and T increased with drainage. The R allele that
10 characterized the upper peat layer correlated with CH₄ oxidation
11 potential. These results advance our understanding of mire
12 dynamics after long-term WT drawdown and of the microbiological
13 bases of methane emissions from mires.
14
15



For Review Only

1
2
3 **CH₄ production and oxidation processes in a boreal fen ecosystem after long-term water**
4 **table drawdown**
5
6
7
8
9

10 **Running title: CH₄ production and oxidation processes**
11

12 Kim Yrjälä¹, Tero Tuomivirta², Heli Juottonen¹, Anuliina Putkinen^{1,2}, Kaisa Lappi¹, Eeva-Stiina
13 Tuittila³, Timo Penttilä², Kari Minkkinen³, Jukka Laine⁴, Krista Peltoniemi², Hannu Fritze²
14
15
16

17
18
19
20 ¹ MEM-group, Department of Biosciences, PO Box 56, 00014 University of Helsinki, Finland
21

22 ² Finnish Forest Research Institute, Southern Unit, PO Box 18, 01301 Vantaa, Finland
23

24 ³ Peatland Ecology Group, Department of Forest Sciences, PO Box 27, 00014 University of
25 Helsinki, Finland
26
27

28
29 ⁴ Finnish Forest Research Institute, Western Unit, Kaironiementie 54, 39700 Parkano, Finland
30
31
32
33
34
35

36 Corresponding author: Kim Yrjälä; Tel. +358 (0)9-19159220; Fax, +358 (0)9-19159262; E-mail:
37

38 kim.yrjala@helsinki.fi
39
40
41
42
43
44
45
46
47

48 Keywords: greenhouse gas, peatlands, methanogens, methanotrophs, microbial communities,
49

50 DGGE, T-RFLP
51
52
53
54
55
56
57
58
59
60

Abstract

Mires, especially sedge dominated fens, are sources of the greenhouse gas CH₄. Climate change scenarios predict a lowering water table (WT) in mires. To study the effect of WT drawdown on CH₄ dynamics in a fen ecosystem, we took advantage of a WT drawdown gradient near a ground water extraction plant. Methane fluxes, CH₄ production and oxidation potentials, were related to microbial communities responsible for the processes in four mire locations (wet, semi-wet, semi-dry and dry). Principal component analyses (PCA) performed on the vegetation, pH, CH₄ and WT results clearly separated the four sampling locations in the gradient. Long-term lowering of WT was associated with decreased coverage of *Sphagnum* and aerenchymatic plants, decreased CH₄ field emissions and CH₄ production potential. Based on *mcrA* T-RF the methanogen community structure correlated best with the methane production and coverage of aerenchymatic plants along the gradient. Methanosarcinaceae and Methanocellales were found at the pristine wet end of the gradient, whereas the Fen cluster characterized the dry end. The methane-oxidizing bacterial (MOB) community consisted exclusively of *Methylocystis* bacteria, but interestingly of five different alleles (T, S, R, M, and O) of the particulate methane monooxygenase marker gene *pmoA*. The M allele was dominant in the wet locations, and the occurrence of alleles O, S and T increased with drainage. The R allele that characterized the upper peat layer correlated with CH₄ oxidation potential. These results advance our understanding of mire dynamics after long-term WT drawdown and of the microbiological bases of methane emissions from mires.

Introduction

In the Northern hemisphere mires extend over vast areas that typically are water-saturated ecosystems where a part of the plant litter production avoids aerobic decomposition and accumulates as peat. The high water table (WT) creates anaerobic microbial habitats where methanogenesis occurs (Rydin *et al.* 2006). Hence boreal mires act as methane (CH₄) sources into the atmosphere. Due to climate change, a growing attention has been devoted to these areas, where over one-third of the global terrestrial carbon is stored (Gorham, 1991). The rise of earth temperatures may have profound effects on the microbial life of these ecosystems and thus on microbially-mediated gas emissions (IPCC 2007).

Methane, which is a 25 times more powerful greenhouse gas than carbon dioxide, is produced microbiologically in the final stage of anaerobic degradation of organic matter (Whalen, 2005). Methane is solely produced by methanogens that are classified into Euryarchaeota. Methanogens are highly diverse constituting of four classes and six orders of archaea. Methane-producing archaeal communities have been described from various mire ecosystems (Edwards *et al.* 1998; Basiliko *et al.* 2003; Galand *et al.* 2003; Galand *et al.* 2005; Galand *et al.* 2005; Juottonen *et al.* 2008; Putkinen *et al.* 2009) including a range of nutrient conditions (Juottonen *et al.* 2005) and mire successional stages (Merilä *et al.* 2006). Methanogens of Methanosarcinales, Methanomicrobiales, Methanobacteriales and Methanocellales (Rice cluster I) have frequently been detected.

The CH₄ emissions are, however, dependent on the activity and abundance of methane-oxidizing bacteria (MOB) that are able to oxidize CH₄ to CO₂ in aerobic conditions (Hanson and Hanson 1996). The aerobic layer of peat is roughly defined by the WT position (Bubier & Moore, 1993). Low emissions in mires can occur when the CH₄ never reaches the atmosphere because it is consumed by MOB in the uppermost aerobic layer (Whalen, 2005). In this way the CH₄ flux to

1
2
3 the atmosphere is the sum of the functionally opposite actions of archaea and bacteria involved in
4
5 the CH₄ cycle. The highest MOB activity in mires is observed just above the WT, where CH₄ and
6
7 oxygen levels are adequate for CH₄ oxidation (Sundh *et al.* 1994). While estimates of the
8
9 oxidation efficiency of the produced CH₄ in different mires vary considerably, estimates of
10
11 between 20% in *Carex* dominated fens (Popp *et al.* 2000) and 78% in *Sphagnum* dominated bogs
12
13 (Yavitt *et al.* 1988) have been given. The latter phenomena is explained by the fact that MOB
14
15 inhabit *Sphagnum* species providing the plant CO₂ through CH₄ oxidation under wet conditions
16
17 (Raghoebarsing *et al.* 2005; Larmola *et al.* 2010).
18
19
20
21

22 MOB are traditionally divided into two taxonomic groups, type I and II, within the
23
24 Proteobacteria. Type I MOB includes genera like *Methylobacter* and *Methylomicrobium*, which
25
26 belong to the Gammaproteobacteria. The type II MOB, *Methylocystis*, *Methylosinus*,
27
28 *Methylocella* and *Methylocapsa* belong to the Alphaproteobacteria (Hanson & Hanson 1996;
29
30 Dedysch, 2009). These types differ in their carbon assimilation pathway, phylogenetic affiliation,
31
32 and intracellular membrane arrangement.
33
34
35

36 The methanogenic and methanotrophic communities in Finnish mires have been studied
37
38 by molecular genetic methods. Methanogens were analyzed by using both functional, *mcrA*
39
40 (Galand *et al.* 2002a), and phylogenetic 16S rRNA marker genes (Galand *et al.* 2003). MOB
41
42 have been studied by functional (*pmoA*) marker gene analysis in a pristine and drained fen and
43
44 bog within a mire complex (Jaatinen *et al.* 2005). The fingerprinting of Finnish fen MOB was
45
46 improved by modifications of the *pmoA* reverse primer for DGGE analysis protocols (Tuomivirta
47
48 *et al.* 2009). Methanogens and methanotrophs in mires have, however, rarely been analyzed
49
50 within the same study (McDonald *et al.* 1999).
51
52
53
54

55 The CH₄ turnover in mires is highly dependent on the ecohydrological conditions. In
56
57 climate change scenarios where the raise of temperature is 3 °C, the summertime WT in northern
58
59
60

1
2
3 mires is expected to drop 10-20 cm (Gorham 1991; Roulet *et al.* 1992; Gitay *et al.* 2001). A WT
4
5 drawdown of that magnitude (15 cm) has occurred in the northern Suonukkasuo fen where a
6
7 groundwater extraction plant has generated a WT gradient that has gradually changed part of the
8
9 wet fen into a forested peatland (Jaatinen *et al.* 2008). Such a persistent change in the WT
10
11 influence the plant community structure (Weltzin *et al.* 2000; Weltzin *et al.* 2003) that may lead
12
13 to complete replacement with species better adapted to the new conditions (Laine *et al.* 1995;
14
15 Strack *et al.* 2006).
16
17

18
19
20 The WT gradient was studied combining methane gas flux measurements, environmental
21
22 data and measurements of microbial communities responsible for CH₄ turnover. This gradient
23
24 study prompted the formulation of hypotheses concerning CH₄ turnover in relation to detected
25
26 microbial communities, namely: (1) as the WT drawdown affects the plant cover and stimulate
27
28 tree growth, the CH₄ emissions are reduced in the resulting dryer conditions; (2) the emissions in
29
30 WT drawdown conditions decrease primarily as a consequence of decrease in potential CH₄
31
32 production and simultaneous decrease in diversity of methanogens; (3) the potential CH₄
33
34 oxidation decreases when there is less CH₄ available; (4) the depth is an important factor
35
36 determining microbial community structure, both that of methanogens and of methanotrophs; (5)
37
38 there are specific types of methane-producing archaea and MOB that are favored by the WT
39
40 drawdown.
41
42
43
44
45
46
47

48 **Materials and methods**

49 50 51 52 **Experimental site and sampling**

53
54
55 The study site, Suonukkasuo, is a mesotrophic pine fen located in Rovaniemi, northern Finland
56
57 (66°28'N, 25°51'E) within the aapa mire zone. Mesotrophic pine fens (RhSR in the Finnish mire
58
59
60

1
2
3 site-type nomenclature of Laine and Vasander (2005)) are typically sites where wet lawns and
4
5 drier hummocks form a mosaic-like vegetation pattern. A groundwater extraction plant on an
6
7 esker bordering the mire downstream has affected the WT at the study site since 1959, resulting
8
9 in a clear hydrological gradient where the pristine wet fen (location S4) becomes semi-wet
10
11 (location S3), semi-dry (location S2), and finally a dry pine dominated mesotrophic peatland
12
13 forest (MtkgII, location S1).
14
15

16
17 The average long term water-table (WT) values for the ground-frost free periods of the
18
19 years 2001–2003 and 2005 expressed as distances from soil surface were 26, 21, 15 and 9 cm for
20
21 the locations S1 to S4, respectively. The differences in WT between locations were statistically
22
23 significant, except between locations S1 and S2 ($p=0.066$). The first- and third-quartile ranges of
24
25 the WTs were 21–30, 13–30, 10–21 and 5–13 cm for the locations S1 to S4, respectively. The
26
27 momentary WT at the time of sampling in 2006 was 32, 50, 47 and 23 cm, respectively. The
28
29 hydrological gradient is reflected in vegetation composition as reported in Jaatinen *et al.* (2008).
30
31
32
33

34 Sampling was conducted in September 2006. Intact triplicate peat cores were taken from
35
36 the four locations using a corer (size 4 x 6 cm) to a depth of 70 cm. Depth samples were prepared
37
38 by dividing peat cores at 10 cm intervals. The samples between 0 - 50 cm depth were used for
39
40 estimation of the methanotroph activity and community analyses and the samples between depths
41
42 20 - 70 cm for the respective analyses for methanogens. For convenience we call the samples
43
44 between 0 to 10 cm as 0 cm, between 10 to 20 cm as 10 cm and so on. To measure the peat dry
45
46 weight subsamples were kept at 105 °C for over 12 h and subsequently weighted.
47
48
49
50
51

52 **Field CH₄ measurements**

53
54
55 Gas efflux measurement plots ($n = 3$) were installed at each location in June 2001 (Dahlin *et al.*,
56
57 2003). Collars made of metal tubes (diameter 31.5 cm) were inserted 15-30 cm deep in peat. An
58
59
60

1
2
3 aluminum groove was attached on the top of each collar. A closed chamber (height 30 cm)
4
5 equipped with a fan was placed on the groove filled with water before sampling to seal the
6
7 chamber. At each sampling occasion, four successive samples of 30 ml were taken from the
8
9 chamber with plastic syringes during a 35-minute measurement period at 10 minute intervals (5,
10
11 15, 25, 35 minutes). The syringes were then taken to laboratory and analyzed for CH₄
12
13 concentrations within 24 hours from sampling by a gas chromatograph equipped with a FI
14
15 detector. CH₄ fluxes were calculated from a linear change of CH₄ concentrations during the
16
17 sampling period of 35 minutes. Obvious ebullition events were deleted from the data.
18
19
20
21
22
23

24 **Measurements of potential CH₄ production and oxidation**

25
26 Measurements of potential CH₄ production and oxidation (4.13 μmol in 125 ml infusion bottle)
27
28 with incubation at 15 °C were carried out as described in Juottonen *et al.* (2008) and Jaatinen *et*
29
30 *al.* (2005), respectively.
31
32
33
34
35

36 **Analysis of methanogen communities by *mcrA*-TRFLP**

37
38 DNA for methanogen community analysis was extracted from 0.25 g of wet mass peat with
39
40 PowerSoil DNA Isolation Kit (MoBio Laboratories). Methyl coenzyme M reductase (*mcrA*) gene
41
42 fragments were amplified with the primers of Luton *et al.* (2002). The reactions (50 μl) contained
43
44 1 × PCR buffer with 2 mM MgCl₂ (Biotools), 200 μM dNTPs, 0.4 μM of both primers, 1U of
45
46 DNA polymerase (Biotools), and 1-2 μl of undiluted DNA extract as template. The reaction
47
48 conditions were initial denaturation (94 °C, 3 min) followed by 31 cycles of 94 °C 45 s, 52 °C 1
49
50 min, 72 °C 1 min 30 s, and a final elongation (72 °C, 7 min). In PCR for T-RFLP, the forward
51
52 primer was 5'-labelled with 6-carboxyfluorescein (FAM).
53
54
55
56
57
58
59
60

1
2
3 For T-RFLP, approximately 25-50 ng of PCR products was digested with 3 U of *MspI*
4 (Fermentas) at 37 °C overnight. The T-RFLP analysis was carried out as described in Juottonen
5
6 (*et al.* (2008)). The range of fragment lengths included in the analysis was 75 to 500 bp. Minimum
7
8 peak height threshold was 100 fluorescence units. Results are presented based on relative peak
9
10 area. For identification of T-RFs, three clone libraries (S4 20 cm; S2 30 cm; S1 50cm) were
11
12 constructed as in Juottonen *et al.* (2008). Inserts from clone colonies were re-amplified with
13
14 *mcrA* primers (33-54 clone per library) and screened by RFLP with *MspI*. From RFLP groups
15
16 with > 5 clones, 2-5 clones from each group were sequenced, and from groups with < 5 clones,
17
18 one member was sequenced as in Juottonen *et al.* (2008). Terminal restriction fragments were
19
20 identified based on *in silico* digestion of the sequences and T-RFLP analysis of clones.
21
22
23
24
25
26
27
28

29 **Investigation of MOB by DGGE analysis of the *pmoA* marker gene**

30
31 DNA for methane-oxidizing bacteria (MOB) community analysis was extracted from ca. 0.5 g
32
33 wet mass peat using the FastDNA kit for soil (MP Biomedicals, Ohio, USA) according to Yeates
34
35 & Gillings (1998) modified as in Tuomivirta *et al.* (2009). A region of subunit α of particulate
36
37 methane monooxygenase (*pmoA*) was PCR targeted with the A189f/A621r with a GC-clamp
38
39 attached to the reverse primer (see Tuomivirta *et al.* 2009 for details). Also broad specificity
40
41 A189f/A682r (Holmes *et al.* 1995) primers were applied for reference (Tuomivirta *et al.* 2009).
42
43
44
45

46 Fingerprinting of the MOB diversity was performed in a denaturing gradient gel
47
48 electrophoresis (DGGE) described in Jaatinen *et al.* (2005). The DGGE gel photographs were
49
50 screened for the presence (1) or absence (0) of *pmoA* bands using the AlphaImager 2.1 program
51
52 of the AlphaDigiDoc gel documentation system (Alpha Innotech Corp., CA. US). A binary
53
54 matrix was generated with this data using only the positions of successfully sequenced bands.
55
56
57
58
59
60

1
2
3 Single DGGE bands of interest were excised from the gel, purified and sequenced as described in
4
5 Tuomivirta *et al.* (2009) and Larmola *et al.* (2010).
6
7
8
9

10 **Sequence analysis and phylogeny**

11
12 *mcrA* and *pmoA* sequences were compared to database sequences with BLAST (Altschul *et al.*
13
14 1997) analysis of the National Center for Biotechnology Information (NCBI). Deduced amino
15
16 acid sequences of *mcrA* together with selected reference sequences were aligned with ClustalW
17
18 (Larkin *et al.* 2007). Suitable evolutionary model was selected with ProtTest (Abascal *et al.*
19
20 2005). Maximum likelihood trees were constructed with PhyML (Guindon & Gascuel 2003) with
21
22 model LG+G+F. Bootstrap values were generated from 100 replicates in PhyML. Pairwise
23
24 distance calculations of nucleotide *pmoA* sequences were analyzed with MEGA 4.0 (Tamura *et*
25
26 *al.* 2007). The *mcrA* and *pmoA* sequences have been submitted to the EMBL database under
27
28 accession nos. FN564006-FN564029 (*mcrA*) and GQ279342-GQ279346 (*pmoA*).
29
30
31
32
33
34
35

36 **Statistical evaluation**

37
38 Methane fluxes between locations were compared by one-way analysis of variance (ANOVA).
39
40 Significance of pairwise differences was assessed by Tukey test. The effect of categorical
41
42 variables depth, location and their interaction on potential CH₄ production and oxidation were
43
44 compared with generalized linear models (GLM). Tukey post hoc test was applied to determine
45
46 which pairs of means differ significantly. Because location and depth had significant interactions,
47
48 ANOVA followed by Tukey test was performed separately for each depth. Level of significance
49
50 in all statistical analyses was P=0.05.
51
52
53
54

55 We applied principal component analysis (PCA) to compare the four locations based on
56
57 their environmental variables, i.e., long-term and momentary WT, CH₄ field emission and
58
59
60

1
2
3 oxidation and production potential, coverage of *Sphagnum* and aerenchymatic plants, and pH. To
4
5 examine variation in methanogen and MOB communities, we first applied detrended
6
7 correspondence analysis (DCA). Based on the DCA showing rather small compositional
8
9 variation, (ter Braak & Prentice 1988) we applied redundancy analysis (RDA) to test which
10
11 environmental variables best explained variation in methanogen and MOB communities. The
12
13 significance of the above listed explanatory factors was assessed using Monte Carlo permutation
14
15 test. The multivariate analyses were carried out with Canoco for Windows (ter Braak & Smilauer
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Methane emissions

In the years (2001—2004) preceding the sampling for microbial community analysis, the CH₄ emissions from location S4, were substantial, averaging 73 mg m⁻² day⁻¹ (Fig. 1). As a result of WT drawdown, the emissions were much smaller in the locations S3 (22 mg m⁻² day⁻¹), S2 (0.9 mg m⁻² day⁻¹) and S1 (1.9 mg m⁻² day⁻¹; Fig. 1). Similarly to long term WT the emissions differed significantly between all pairs of locations (P=0.001—0.004) except for locations S2 and S1.

Methane production and oxidation potential

The wettest location S4 had the highest CH₄ production potential at depths from 20 to 50 cm (P=0.001—0.036, Fig. 2) in correspondence with the flux rates recorded in earlier years. The S4 production was 43.2 nmol g dry weight⁻¹ day⁻¹ compared to 0.4 (S3, p=0.005), 4.0 (S2, p=0.010) and 0.4 (S1, p=0.005) at the depth of 20 cm. The production potential of the three other dryer

1
2
3 locations was much lower and did not differ between the locations at any depth. The laboratory
4
5 measurements showed that location and depth significantly affected the CH₄ production potential
6
7 through the WT gradient (p<0.001).
8
9

10 The highest methane oxidation potential was observed in location S3, 561.2 nmol g dry
11 weight⁻¹ day⁻¹, but the sampling depth influenced the oxidation potential (p=0.004) in addition to
12 location (p=0.033) (Fig. 2). In the top layer the potential was significantly higher in S3 than in
13 S2, 62.2 nmol g dry weight⁻¹ day⁻¹ (p=0.042) and S1, 28.9 nmol g dry weight⁻¹ day⁻¹ (p=0.031)
14 and second highest in S4, 177,7 nmol g dry weight⁻¹ day⁻¹. The dry S1 location had the lowest
15 potential oxidation rate. The oxidation potential did not differ between locations in deeper peat
16 layers.
17
18
19
20
21
22
23
24
25
26
27
28

29 **Environmental factors characterizing the WT gradient**

30
31 Principal component analyses (PCA) performed on the vegetation, potential CH₄ production and
32 oxidation, CH₄ field emissions, pH, and long term and momentary (date of sampling) water-table
33 depth clearly separated the four sampling locations along the first principal axis (PC1 in Fig. 3a).
34
35 With the provided measurements the second principal axis (PC2) separated all samples according
36 to depth. Increase in long term WT, coverage of *Sphagnum* and aerenchymatic plants, high CH₄
37 field emissions together with CH₄ production potential characterized the pristine end of the PC1
38 (Fig. 3b). Decreasing momentary WT and CH₄ oxidation potential together with increasing pH
39 characterized the peat layers with increasing depth along PC2.
40
41
42
43
44
45
46
47
48
49
50
51
52

53 **Microbial communities involved in the CH₄ cycle along the gradient**

54
55 T-RFLP fingerprinting of *mcrA* representing methanogens resulted in twelve T-RFs (Table 1). In
56 the RDA T-RFs 468 and 471 were combined since the large size of T-RFs prohibited reliable
57
58
59
60

1
2
3 distinction between them. According to RDA the methanogen community structure showed
4
5 highest correlation with methane production and coverage of aerenchymatic plants along the
6
7 gradient (RD1 in Fig. 4a-c). The second RDA axis showed that the pH influenced the community
8
9 structure according to depth. The first RDA axis separated the four locations when the
10
11 methanogen communities at different depths were analyzed together with environmental factors
12
13 (Fig. 4b). The top layers were most variable through the gradient, but deeper down they became
14
15 more alike.
16
17

18
19
20 In the wet end of the gradient with higher CH₄ production potential the methanogen
21
22 diversity was highest with 8 to 10 T-RFs. The locations S2 and S1 sampled from three depths
23
24 showed lower diversity with 6 T-RFs.
25

26
27 The occurrence of the T-RFs 115 bp (Methanocellales), 272 bp (putatively
28
29 Methanosarcinaceae) and 298 bp (Methanosarcinaceae) correlated most strongly with the wet
30
31 location S4 (Fig. 4c, Fig. 5). In addition to identified methanogens, T-RF 193 bp affiliated with a
32
33 cluster with no cultured representatives in the phylogenetic tree (Fig. 5). The T-RF 278 bp,
34
35 representing Methanomicrobiales-associated Fen cluster, was one of two T-RFs prone to the dry
36
37 end of the gradient despite of overall decrease in methanogen groups along the gradient.
38
39

40
41 In analysis of MOB communities, the broad specificity *pmoA* primer set A189f/A682r did
42
43 not produce correct amplicons (Tuomivirta *et al.* 2009). Five different *pmoA* alleles (T, S, R, M,
44
45 and O), all *Methylocystis* spp according to sequence comparisons, could, however, be isolated
46
47 from the gradient by PCR-DGGE and sequencing with the A189f/A621r primer set (Table 2).
48
49 RDA (Fig. 6a) of the *pmoA* alleles showed that their occurrence was best correlated with CH₄
50
51 oxidation potential, pH and vegetation. Similarly to methanogen communities (Fig. 4a) the
52
53 locations from wet to dry were separated along the first RDA axis, and the second RDA axis
54
55 characterized the depth of the peat (Fig. 6b). The two wettest locations had similar MOB
56
57
58
59
60

1
2
3 communities in their whole profile (Fig. 6b). The deep layers (30 - 40 cm) of the locations S4, S3
4
5 and S2 were also characterized by a similar MOB. The MOB community of the dry location S1
6
7 differed from the other sites. The M allele, dominating the wet end of the gradient (Fig. 6a) was
8
9 identical to a previously detected Finnish fen sequence FJ930087 (Tuomivirta *et al.* 2009) and
10
11 was most different from the other alleles with 11 to 12 point mutations. The R allele inhabiting
12
13 the wetter part of the gradient was identical to sequences FJ930090 and FJ930091 of a Finnish
14
15 fen (Tuomivirta *et al.* 2009), AY309209, a North-American fen (Nold, personal communication),
16
17 and the sequence GQ121280 (Larmola *et al.* 2010) occurring in methane oxidizing community
18
19 within *Sphagnum* mosses. Drying correlated with increasing abundances of alleles O, S and T.
20
21 The R allele dominated the pristine location and the upper layer of the peat (Fig. 6a).
22
23
24
25
26
27
28

29 Discussion

30
31
32
33
34 Generally the study supports the five hypotheses concerning CH₄ turnover in relation to detected
35
36 microbial communities along WT drawdown gradient. The study of greenhouse gas turnover
37
38 along a peatland WT gradient showed that drying appeared to cause a dramatic reduction in CH₄
39
40 emissions and in potential CH₄ production in agreement with our hypotheses 1 and 2. Despite
41
42 more oxic conditions the methane oxidation also decreased substantially along the gradient. The
43
44 phenomena supported our third hypothesis on the superior role of the availability of CH₄ for the
45
46 potential CH₄ oxidation over the direct WT control. In agreement with the fifth hypothesis drying
47
48 decreased the diversity of the methanogen communities and brought forth an adapted group of
49
50 archaea, the Fen cluster methanogens, which was characteristic of dryer conditions. As
51
52 hypothesized (hypothesis 4) the depth appeared to be an important factor for the community
53
54 structure of methanogens and of methanotrophs. The depth effect in WT drawdown brought
55
56
57
58
59
60

1
2
3 about a peculiar shift in methanotroph populations showing specific *pmoA* allele types of
4
5 *Methylocystis* bacteria distributed to specific locations in the gradient. Previously, at the same
6
7 site, the CO₂ flux increased from location S4 to S1 and fungi and bacteria benefited from the WT
8
9 drawdown whereas actinobacteria did not (Jaatinen *et al.* 2008).
10
11

12
13 Separation of the four locations along the WT gradient and the sampling depths (peat
14
15 layers) was clear and could statistically be explained by eight variables. The WT gradient was
16
17 best described in PCA by coverage of aerenchymatic plants, high CH₄ emissions at the wet end,
18
19 coverage of *Sphagnum* moss and CH₄ production potential in this order. The CH₄ oxidation was
20
21 reduced as a consequence of WT drawdown but not as clearly as the production. This may well
22
23 be indicative of the difference in the two processes of production (anaerobic) and the following
24
25 consumption (aerobic) of CH₄. The CH₄ oxidation potential indeed correlated positively with the
26
27 WT of the time of sampling and the coverage of *Sphagnum*, whereas production correlated with
28
29 other variables like pH and emissions (Fig. 3b). The high oxidation in wet locations seemed to be
30
31 coupled with high CH₄ production potential. When more CH₄ is potentially produced by
32
33 methanogenic archaea, more is simultaneously oxidized by the methanotrophic bacteria.
34
35
36
37
38

39
40 Very recently methane oxidation was reported from 23 different *Sphagnum* species
41
42 originating from a Finnish boreal mire site and the activity was related to a *Methylocystis*
43
44 signature similar to the R allele of this study (Larmola *et al.* 2010). The correlation of the
45
46 oxidation potential with *Sphagnum* coverage might be a direct regulation instead of a covariation
47
48 with more pristine conditions.
49

50
51 The methanogen community structure according to T-RFLP analysis changed with
52
53 decrease in diversity along the gradient. The maximum CH₄ production potential for each vertical
54
55 peat core correlated with the diversity of methanogens so that decreasing production was
56
57 reflected in decreasing diversity. WT gradients have mainly been studied in meso- and
58
59
60

1
2
3 microcosms (see, however, Talbot *et al.* 2010), and as far as we know reports from natural
4
5 gradients combined with microbial community analysis of the CH₄ are lacking. In agreement
6
7 with our results from the gradient, experimental field studies in Canada (Strack *et al.* 2004) and
8
9 in Ireland (Laine *et al.* 2009) have shown decrease in CH₄ emission as a result of lowered WT.
10
11 Short term effects of WT on carbon balance in pore water from acidic, oligotrophic mire have
12
13 been reported from a mesocosm study (Blodau *et al.* 2004) where decreased CH₄ production and
14
15 emission was reported as a result of lower WT. Short term effect of drought (4 week) on the
16
17 methanogen community structure in a bog and a fen mesocosm could not be observed (Kim *et al.*
18
19 2008), but a significant decrease in the *mcrA* abundance was detected in the fen mesocosm.
20
21
22
23

24
25 Sampling depth has been shown to affect the methanogen community structure in mires
26
27 (Galand *et al.* 2002b) where the changing conditions with depth (oxygen and quality of growth
28
29 substrates for the microbes active in methanogenesis) will select for methanogen groups. The
30
31 RDA analysis of the methanogen communities together with environmental factors showed
32
33 nicely how the locations differ at the surface, but become much alike in the deepest sampling
34
35 layer where the conditions in the gradient apparently are more similar (Fig. 4b). The discrepancy
36
37 of location S2 to the general trend may well be explained by larger pH variation at this location
38
39 (results not shown) and lowest emissions telling that this location is the most disturbed one in the
40
41 gradient.
42
43
44

45
46 The methanogen group most strongly associated with the pristine end of the gradient was
47
48 Methanosarcinaceae. In the drier locations, Methanosarcinaceae T-RFs were less abundant or
49
50 absent. This finding was surprising considering that many members Methanosarcinaceae are
51
52 versatile methanogens that are able to use several different substrates for growth (Garcia *et al.*
53
54 2000). They also possess enzymes for detoxification of oxygen, which could favor their
55
56 occurrence in the dry end of the gradient, which is the opposite of what was observed. Additional
57
58
59
60

1
2
3 data, such as availability of methanogenic substrates along the gradient, would be required to
4
5 better explain the disappearance of Methanosarcinaceae at the dry end. A minor group also found
6
7 primarily at the wet end was an uncultured cluster (Mx cluster in Fig. 5) which has been detected
8
9 in other wetland ecosystems (Juottonen *et al.* 2005; Castro *et al.* 2004; Smith *et al.* 2007) and in
10
11 animal feces (Ufnar *et al.* 2007), and it has been proposed to form an entirely new order of
12
13 methanogens (Mihajlovski *et al.* 2008). Only two methanogenic groups were typical to the dry
14
15 end of the gradient, and they were affiliated with the Fen cluster which has commonly been found
16
17 in Finnish mires (Juottonen *et al.* 2008). One member of this cluster, *Methanoregula boonei*, was
18
19 isolated from an acidic bog and has the lowest pH optimum known for a methanogen (Bräuer *et*
20
21 *al.* 2006). It is tempting to say that the Fen cluster/Methanoregula-type hydrogenotrophic
22
23 methanogens are well adapted to the conditions of boreal fens characterized by changing WT and
24
25 low pH.
26
27
28
29
30

31 All detected MOBs were by *pmoA* sequence analysis found to be of the *Methylocystis*
32
33 genus, but differed by minute changes, point mutations, in the gene (Table 2). This suggests a
34
35 phylogenetically very narrow MOB population of this fen. RDA showed that the occurrence of
36
37 the five alleles was clearly dependent on location and peat layer in the gradient. It can be
38
39 assumed that before WT drawdown, when the whole site was botanically a homogenous fen, all
40
41 the locations had a similar MOB population with dominating alleles R and T in surface peat and
42
43 then changing to T and M alleles in the deeper peat. Continual drainage of the fen has a selective
44
45 pressure on MOB. The R allele was lost upon drying since methanotrophy in *Sphagnum* mosses
46
47 is only related to high WT (Larmola *et al.* 2010). For reasons we do not know the most different
48
49 M allele, in respect to other ones, disappears in WT drawdown and is replaced by the O and S
50
51 alleles through out the peat layer. The S allele could originate from the generalist T through a
52
53
54
55
56
57
58
59
60

1
2
3 single silent point mutation in *pmoA*. The O has probably been introduced to the location as it
4
5 contains four silent point mutations in respect with S.
6
7

8 *Methylocystis* was a dominating genus also in Russian and German bogs, which showed
9
10 that 60-95% of MOB belonged to type II with *Methylocystis* being the dominant genus (Dedysh
11
12 *et al.* 2003). A study, in which stable isotope probing (SIP) with PLFA, mRNA and microarray
13
14 methods were used, revealed the dominance of type II genera *Methylocystis* and *Methylosinus* in
15
16 blanket peat samples from England (Chen *et al.* 2008). Bearing in mind that also *Sphagnum*
17
18 species possess *Methylocystis* (Larmola *et al.* 2010), one can say that it most likely dominates in
19
20 northern peatlands.
21
22
23

24 With regard to climate change effect on boreal fens, the average drop in WT between
25
26 location S4 (wet fen) and location S3 (semi-wet) was ca. 6 cm. This was accompanied by a
27
28 statistically significant drop in potential CH₄ production together with a reduction in emissions
29
30 from previous years. The potential oxidation changed in the other direction with increasing
31
32 amounts of CH₄ oxidized at location S3 as compared to location S4. This would indicate that
33
34 already a small, 6 cm, WT drawdown will change the CH₄ turnover. The methanogen community
35
36 structure most clearly shifted between locations S3 and S2 and the distribution of the
37
38 methanotrophic allele types is also changing most clearly in this part of the gradient. It can,
39
40 however, be stated that as the underlying microbial communities are changing with a 12 cm
41
42 drop in WT their activity is already altered after the 6-cm initial drop in WT.
43
44
45
46
47

48 Comparison of behavior of methanogens and methanotrophs in the WT gradient using
49
50 CH₄ production and oxidation measurements showed a difference in sensitivity to changing
51
52 conditions. Much of the methanogenic activity is evidently lost already in the semi-wet location
53
54 S3 and after that decreases drastically, whereas the methanotroph potential activity is high in the
55
56 wet locations and decreases much less in the dryer locations. This can be explained by more oxic
57
58
59
60

1
2
3 conditions that should favor these aerobic bacteria. The drop is best explained by the lack of
4
5 substrate; CH₄.
6

7
8 As a general conclusion, we have been able to show that even a minor WT drawdown,
9
10 mimicking the climate change impact affects the dynamics of CH₄ production and oxidation in a
11
12 mire. This study thus shows how ecohydrology shapes the dynamics of mires changing the plant
13
14 community structures, and has a drastic effect on the microbial communities that are the basis for
15
16 the environmentally important function of greenhouse gas turnover.
17
18
19
20
21

22 **Acknowledgements**

23
24 We thank Sirpa Tiikainen for the valuable laboratory help with MOB. This study was financed by
25
26 the Academy of Finland (project number 109816).
27
28
29
30
31
32
33

34 **References**

35
36 Abascal F, Zardoya R, Posada D (2005) ProtTest: selection of best-fit models of protein
37
38 evolution. *Bioinformatics*, **21**, 2104.
39

40
41 Altschul S, Madden T, Schaeffer A, Zhang J, Zhang Z, Miller W, Lipman D (1997) Gapped
42
43 BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic acids*
44
45 *research*, **25**, 3389-3402.
46
47

48
49 Basiliko N, Yavitt J, Dees P, Merkel S (2003) Methane biogeochemistry and methanogen
50
51 communities in two northern peatland ecosystems, New York State. *Geomicrobiology Journal*,
52
53 **20**, 563-577.
54

55
56 Blodau C, Basiliko N, Moore TR (2004) Carbon turnover in peatland mesocosms exposed to
57
58 different water table levels. *Biogeochemistry*, **67**, 331-351.
59
60

1
2
3 Braak CJF, Smilauer P (2002) CANOCO reference manual and CanoDraw for Windows user's
4
5 guide: software for canonical community ordination (version 4.5).
6
7

8 Bräuer SL, Cadillo-Quiroz H, Yashiro E, Yavitt JB, Zinder SH (2006) Isolation of a novel
9
10 acidiphilic methanogen from an acidic peat bog. *Nature*, **442**, 192-194.
11

12 Castro H, Ogram A, Reddy KR (2004) Phylogenetic characterization of methanogenic
13
14 assemblages in eutrophic and oligotrophic areas of the Florida Everglades. *Applied and*
15
16 *environmental microbiology*, **70**, 6559-6568.
17

18
19
20 Chen Y, Dumont MG, Neufeld JD, et al (2008) Revealing the uncultivated majority: combining
21
22 DNA stable-isotope probing, multiple displacement amplification and metagenomic analyses of
23
24 uncultivated *Methylocystis* in acidic peatlands. *Environmental microbiology*, **10**, 2609-2622.
25

26
27 Dahlin T, Minkkinen K, Penttilä T (2003) Methane emissions from a northern boreal pine fen
28
29 along water table gradient – preliminary results. In Proceedings of Sixth Finnish Conference of
30
31 Environmental Sciences. Joensuu, May 8–9, 2003. Current Perspectives in Environmental
32
33 Science and Technology. Honkanen JO and Koponen PS (eds). University of Joensuu, Joensuu,
34
35 Finland.
36
37

38
39 Dedysh SN (2009) Exploring Methanotroph Diversity in Acidic Northern Wetlands:
40
41 Molecular and Cultivation-Based Studies. *Microbiology*, **78**, 655-669.
42

43
44 Dedysh SN, Dunfield PF, Derakshani M, Stubner S, Heyer J, Liesack W (2003) Differential
45
46 detection of type II methanotrophic bacteria in acidic peatlands using newly developed 16S
47
48 rRNA-targeted fluorescent oligonucleotide probes. *FEMS microbiology ecology*, **43**, 299-308.
49

50
51 Edwards C, Hales B, Hall G, et al (1998) Microbiological processes in the terrestrial carbon
52
53 cycle: methane cycling in peat. *Atmospheric Environment*, **32**, 3247-3255.
54
55
56
57
58
59
60

1
2
3 Galand PE, Fritze H, Conrad R, Yrjälä K (2005) Pathways for methanogenesis and diversity of
4
5 methanogenic archaea in three boreal peatland ecosystems. *Applied and environmental*
6
7 *microbiology*, **71**, 2195-2198.

8
9
10 Galand PE, Fritze H, Yrjälä K (2003) Microsite-dependent changes in methanogenic populations
11
12 in a boreal oligotrophic fen. *Environmental microbiology*, **5**, 1133-1143.

13
14 Galand PE, Juottonen H, Fritze H, Yrjala K (2005) Methanogen communities in a drained bog:
15
16 effect of ash fertilization. *Microbial ecology*, **49**, 209-17.

17
18
19 Galand PE, Saarnio S, Fritze H, Yrjälä K (2002a) Depth related diversity of methanogen Archaea
20
21 in Finnish oligotrophic fen. *FEMS microbiology ecology*, **42**, 441-449.

22
23 Galand PE, Saarnio S, Fritze H, Yrjälä K (2002b) Depth related diversity of methanogen Archaea
24
25 in Finnish oligotrophic fen. *FEMS microbiology ecology*, **42**, 441-449.

26
27
28 Garcia JL, Patel BK, Ollivier B (2000) Taxonomic, phylogenetic, and ecological diversity of
29
30 methanogenic Archaea. *Anaerobe*, **6**, 205-226.

31
32
33 Gitay H, Brown S, Easterling W, et al (2001) Ecosystems and their goods and services. In:
34
35 *Climate change 2001; impacts, adaptation, and vulnerability* (eds McCarthy JJ, Canziani OF,
36
37 Leary NA, Dokken DJ, White KS), Cambridge University Press, Cambridge, United Kingdom
38
39 (GBR), United Kingdom (GBR).

40
41
42
43 Gorham E (1991) Northern peatlands: role in the carbon cycle and probable responses to climatic
44
45 warming. *Ecological Applications*, **1**, 182-195.

46
47
48 Guindon S, Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large
49
50 phylogenies by maximum likelihood. *Systematic Biology*, **52**, 696-704.

51
52
53 Hanson R, Hanson T (1996) Methanotrophic bacteria. *Microbiology and Molecular Biology*
54
55 *Reviews*, **60**, 439.

1
2
3 Holmes A, Costello A, Lidstrom M, Murrell J (1995) Evidence that particulate methane
4 monooxygenase and ammonia monooxygenase may be evolutionarily related. *FEMS*
5 *microbiology letters*, **132**, 203-208.

6
7
8
9
10 Jaatinen K, Laiho R, Vuorenmaa A, del Castillo U, Minkkinen K, Pennanen T, Penttilä T, Fritze
11 H (2008) Responses of aerobic microbial communities and soil respiration to water-level
12 drawdown in a northern boreal fen. *Environmental microbiology*, **10**, 339-353.

13
14
15
16
17 Jaatinen K, Tuittila E-S, Laine J, Yrjala K, Fritze H (2005) Methane-Oxidizing Bacteria in a
18 Finnish Raised Mire Complex: Effects of Site Fertility and Drainage. *Microbial ecology*, **50**, 429-
19 439.

20
21
22
23
24 Juottonen H, Tuittila ES, Juutinen S, Fritze H, Yrjala K (2008) Seasonality of rDNA-and rRNA-
25 derived archaeal communities and methanogenic potential in a boreal mire. *The ISME Journal*, **2**,
26 1157–1168.

27
28
29
30
31 Juottonen H, Galand PE, Tuittila ES, Laine J, Fritze H, Yrjala K (2005) Methanogen
32 communities and Bacteria along an ecohydrological gradient in a northern raised bog complex.
33 *Environmental microbiology*, **7**, 1547-1557.

34
35
36
37
38 Kim SY, Lee SH, Freeman C, Fenner N, Kang H (2008) Comparative analysis of soil microbial
39 communities and their responses to the short-term drought in bog, fen, and riparian wetlands. *Soil*
40 *Biology and Biochemistry*, **40**, 2874-2880.

41
42
43
44
45 Laine A, Byrne KA, Kiely G, Tuittila E-S (2009) The effect of altered water level on carbon
46 fluxes in a blanket bog. *SUO* **60**, 65-83.

47
48
49
50 Laine J, Vasander H (2005) Suotyypit ja niiden tunnistaminen.[Mire site types and their
51 identification.].

52
53
54
55 Laine J, Vasander H, Laiho R (1995) Long-term effects of water level drawdown on the
56 vegetation of drained pine mires in southern Finland. *Journal of Applied Ecology*, **32**, 785-802.

- 1
2
3 Larkin M, Blackshields G, Brown N, et al (2007) Clustal W and Clustal X version 2.0.
4
5 *Bioinformatics*, **23**, 2947.
6
7
8 Larmola T, Tuittila E-S, Tirola M, Nykänen H, Martikainen PJ, Yrjälä K, Tuomivirta T, Fritze H
9
10 (2010) The role of *Sphagnum* mosses in the methane cycling of a boreal mire. *Ecology* (in press).
11
12
13 Luton PE, Wayne JM, Sharp RJ, Riley PW (2002) The *mcrA* gene as an alternative to 16S rRNA
14
15 in the phylogenetic analysis of methanogen populations in landfill. *Microbiology*, **148**, 3521-
16
17 3530.
18
19
20 McDonald I, Upton M, Hall G, Pickup R, Edwards C, Saunders J, Ritchie D, Murrell J (1999)
21
22 Molecular ecological analysis of methanogens and methanotrophs in blanket bog peat. *Microbial*
23
24 *ecology*, **38**, 225-233.
25
26
27 Merilä P, Galand PE, Fritze H, Tuittila E, Kukko-Oja K, Laine J, Yrjälä K (2006) Methanogen
28
29 communities along a primary succession transect of mire ecosystems. *FEMS microbiology*
30
31 *ecology*, **55**, 221-229.
32
33
34 Mihajlovski A, Alric M, Brugère JF (2008) A putative new order of methanogenic Archaea
35
36 inhabiting the human gut, as revealed by molecular analyses of the *mcrA* gene. *Research in*
37
38 *microbiology*, **159**, 516-521.
39
40
41 Popp TJ, Chanton JP, Whiting GJ, Grant N (2000) Evaluation of methane oxidation in the
42
43 rhizosphere of a *Carex* dominated fen in northcentral Alberta, Canada. *Biogeochemistry*, **51**, 259-
44
45 281.
46
47
48 Putkinen A, Juottonen H, Juutinen S, Tuittila ES, Fritze H, Yrjälä K (2009) Archaeal rRNA
49
50 diversity and methane production in deep boreal peat. *FEMS microbiology ecology*, **70**, 87-98.
51
52
53 Raghoebarsing AA, Smolders AJP, Schmid MC, et al (2005) Methanotrophic symbionts provide
54
55 carbon for photosynthesis in peat bogs. *Nature*, **436**, 1153-1156.
56
57
58
59
60

1
2
3 Roulet N, Moore T, Bubier J, Lafleur P (1992) Northern fens: methane flux and climatic change.
4
5 *Tellus*, **44**, 100-105.
6

7
8 Rydin H, Jeglum J, Hooijer A (2006) *The biology of peatlands*. Oxford University Press, USA, .
9

10 Smith JM, Castro H, Ogram A (2007) Structure and function of methanogens along a short-term
11 restoration chronosequence in the Florida Everglades. *Applied and Environmental Microbiology*,
12
13 **73**, 4135.
14

15
16 Strack M, Waddington JM, Rochefort L, Tuittila E-S (2006) Response of vegetation and net
17 ecosystem carbon dioxide exchange at different peatland microforms following water table
18
19 drawdown. *Journal of Geophysical Research*, **111**, 1-10.
20
21

22
23 Strack M, Waddington JM, Tuittila E-S (2004) Effect of water table drawdown on peatland
24 methane dynamics: Implications for climate change. *Global Biogeochemical Cycles*, **18**, 1-7.
25
26

27
28 Sundh I, Nilsson M, Granberg G, Svensson B (1994) Depth distribution of microbial production
29 and oxidation of methane in northern boreal peatlands. *Microbial ecology*, **27**, 253-265.
30
31

32
33 Talbot J, Richard PJH, Roulet NT, Booth R (2010) Assessing long-term hydrological and
34
35 vegetation changes following drainage in a bog using paleoecological reconstructions and a
36
37 hydrosequence. *Vegetation Science* (in press).
38
39

40
41 Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: molecular evolutionary genetics analysis
42 (MEGA) software version 4.0. *Molecular biology and evolution*, **24**, 1596.
43
44

45
46 ter Braak CJF, Prentice IC (1988) A theory of gradient analysis. *Advances in Ecological*
47
48 *Research*, **18**, 1-3.
49

50
51 Tuomivirta TT, Yrjälä K, Fritze H (2009) Quantitative PCR of *pmoA* using a novel reverse
52 primer correlates with potential methane oxidation in Finnish fen. *Research in microbiology*, **160**,
53
54 751-756.
55
56
57
58
59
60

1
2
3 Ufnar JA, Wang SY, Ufnar DF, Ellender R (2007) *Methanobrevibacter ruminantium* as an
4
5 indicator of domesticated-ruminant fecal pollution in surface waters. *Applied and Environmental*
6
7 *Microbiology*, **73**, 7118.

8
9
10 Weltzin JF, Bridgham SD, Pastor J, Chen J, Harth C (2003) Potential effects of warming and
11
12 drying on peatland plant community composition. *Global Change Biology*, **9**, 141-151.

13
14
15 Weltzin JF, Pastor J, Harth C, Bridgham SD, Updegraff K, Chapin CT (2000) Response of bog
16
17 and fen plant communities to warming and water-table manipulations. *Ecology*, **81**, 3464-3478.

18
19
20 Yavitt J, Lang G, Downey D (1988) Potential Methane Production and Methane Oxidation Rates
21
22 in Peatland Ecosystems of the Appalachian Mountains, United States. *Global Biogeochemical*
23
24 *Cycles*, **2**, 253–268.

25
26
27 Yeates C, Gillings MR, Davison AD, Altavilla N, Veal DA (1998) Methods for microbial DNA
28
29 extraction from soil for PCR amplification. *Biological Procedures Online*, **1**, 40-47.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Identification of T-RFs detected in analysis methanogens in the mires forming a water table (WT) gradient. The occurrence of PCR-T-RFLP detected methanogens in the WT gradient formed by the four locations S4-S1 and the relative abundance of T-RFs in each location.

| T-RF (bp) | Methanogen group | Occurrence of T-RF (%) | | | |
|-----------|----------------------------------|------------------------|-----------------|-----------------|----------------|
| | | S4 | S3 | S2 | S1 |
| 105 | no sequences | 3 | 2 | - | - |
| 115 | Methanocellales | 6 | 4 | - | - |
| 193 | Mx cluster | 5 | 1 | 2 | - |
| 243 | Methanosarcinaceae | - | - | - | 1 |
| 251 | Methanocellales | 13 | 18 | 7 | 6 |
| 272 | Methanosarcinaceae ^a | 14 | 6 | 5 | 1 |
| 278 | Fen cluster (Methanomicrobiales) | 31 | 41 | 70 | 69 |
| 293 | Fen cluster (Methanomicrobiales) | 16 | 15 | 3 | 14 |
| 298 | Methanosarcinaceae | 2 | - | - | - |
| 468 | Methanobacteriaceae | 9 ^b | 13 ^b | 13 ^b | 8 ^b |
| 471 | Fen cluster (Methanomicrobiales) | | | | |
| 489 | Methanosarcinaceae | 1 | - | - | - |
| | | wet | semi | semi | dry |
| | | | wet | dry | |

^a No sequences were detected in this study but identification is based on sequences from other mire sites (Juottonen *et al.*, unpublished data).

^b Includes T-RFs 468 and 471 bp.

Table 2 Number of point mutations (integers) between different *pmoA* allele sequences and standard errors (in parenthesis) of pairwise distance calculations. Accession numbers in Genbank are GQ279342 (T), GQ279343 (S), GQ279344 (R), GQ279345 (M) and GQ279346 (O), respectively.

| Sequence of <i>pmoA</i> | | | | | |
|----------------------------|---------|------------|------------|------------|------------|
| O | 5 | 4 | 7 | 11 | |
| M | 12 | 11 | 12 | | |
| R | 2 | 3 | | | |
| S | 1 | | | | |
| T | | | | | |
| | T | S | R | M | O |
| T | | [0.00198] | [0.00286] | [0.00977] | [0.00513] |
| S | 0.00122 | | [0.00376] | [0.00918] | [0.00455] |
| R | 0.00244 | 0.00367 | | [0.01011] | [0.00648] |
| M | 0.01518 | 0.0139 | 0.01522 | | [0.00944] |
| O | 0.00616 | 0.00492 | 0.00867 | 0.01382 | |

1
2
3 Fig. 1 Average methane emission 2001-2004 from the four locations forming a water table (WT)
4
5
6 gradient, from wet fen location S4 to dry forested location S1.
7
8
9

10 Fig. 2 (a) The potential methane production, (b) the potential methane oxidation in the four
11
12 locations S4 - S1. The depths are given as distance from the surface of the mires.
13
14
15
16

17 Fig. 3 (a) Principal component analysis (PCA) of environmental factors, emissions, vegetation,
18
19 water table (WT) for locations S4 to S1 coupled with CH₄ production and oxidation. The
20
21 direction of the arrow describes the correlation between measured factors. The length of the
22
23 arrow gives the degree of explanation of each variable in the ordination. The two first principal
24
25 components together explained 55.3% of the variation. Eigenvalues of the first and second axis
26
27 were 0.366 and 0.186, respectively. (b) Score plot of the PCA.
28
29
30
31
32
33

34 Fig. 4 (a) Redundancy analysis (RDA) of statistically significant environmental factors and
35
36 methanogen communities. The length of an arrow describes the degree of explanation of
37
38 environmental factor. The first two axes explained 28.0% of the compositional variation. (b)
39
40 RDA of methanogen phylotypes (T-RFs) at different depths and environmental factors. (c) RDA
41
42 of T-RFs detected in the water table (WT) gradient. The length of arrow describes how well the
43
44 phylotype correlates with locations in the gradient.
45
46
47
48
49

50 Fig. 5 Maximum likelihood tree of partial *mcrA* sequences (128 aa) from Suonukkasuo fen clones
51
52 (in bold) and reference methanogen sequences. Size of terminal restriction fragment is indicated
53
54 for each group. Scale indicates 10% sequence divergence. Nodes with bootstrap values >95% are
55
56 marked with filled circles and >75% with open circles. The tree was rooted with *Methanopyrus*
57
58
59
60

1
2
3 *kandleri*. In bold sequence names Suon1, Suon2 and Suon4 refer to locations S1, S2 and S4 of
4
5
6 the water table gradient.
7
8
9

10 Fig. 6 (a) RDA of methanotroph *pmoA* alleles showing correlation strength (length of arrow) with
11 statistically significant environmental factors. The first two axes explained 47.1% of the
12
13 compositional variation. (b) The change in methanotroph communities according to depth and
14
15 environmental factors picturing the WT gradient.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



206x103mm (600 x 600 DPI)

Review Only


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



195x133mm (600 x 600 DPI)


www Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



166x167mm (600 x 600 DPI)


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



170x112mm (600 x 600 DPI)

ew Only


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



166x135mm (600 x 600 DPI)

Only


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



196x114mm (600 x 600 DPI)

View Only


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



177x122mm (600 x 600 DPI)


www.Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




202x210mm (600 x 600 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



157x217mm (600 x 600 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



171x138mm (600 x 600 DPI)

Only