



**HAL**  
open science

## Warming Alters the Size Spectrum and Shifts the Distribution of Biomass in Aquatic Ecosystems

Gabriel Yvon-Durocher, Jose Maria Montoya, Mark Trimmer, Guy Woodward

► **To cite this version:**

Gabriel Yvon-Durocher, Jose Maria Montoya, Mark Trimmer, Guy Woodward. Warming Alters the Size Spectrum and Shifts the Distribution of Biomass in Aquatic Ecosystems. *Global Change Biology*, 2010, 17 (4), pp.1681. 10.1111/j.1365-2486.2010.02321.x . hal-00599527

**HAL Id: hal-00599527**

**<https://hal.science/hal-00599527>**

Submitted on 10 Jun 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Warming Alters the Size Spectrum and Shifts the Distribution of Biomass in Aquatic Ecosystems



Journal:	<i>Global Change Biology</i>
Manuscript ID:	GCB-10-0491
Wiley - Manuscript type:	Primary Research Articles
Date Submitted by the Author:	14-Jun-2010
Complete List of Authors:	Yvon-Durocher, Gabriel; Queen Mary University of London, School of Biological and Chemical Sciences Montoya, Jose; 2 Institute of Marine Sciences Trimmer, Mark; Queen Mary University of London, School of Biological and Chemical Sciences Woodward, Guy; Queen Mary University of London, School of Biological and Chemical Sciences
Keywords:	Global Warming, Body Size, Size Spectra, Mass-abundance relationships, Biomass
Abstract:	<p>Body size is one of the key determinants of community structure. The relationship between abundance and body size can explain how community biomass is partitioned among the biota of an ecosystem. We used an aquatic mesocosm experiment to determine how warming of <math>\sim 4^{\circ}\text{C}</math> would affect the body size, biomass and taxonomic structure of planktonic communities. We found that warming increased the steepness of the slope of the community size spectrum, primarily by altering the phytoplankton size spectrum. Warming also reduced the mean and maximum body size of phytoplankton by approximately one order of magnitude. The observed shifts in phytoplankton size structure were reflected in large shifts in phytoplankton community composition, though zooplankton taxonomic composition was unaffected by warming. Furthermore, warming reduced community biomass and total phytoplankton biomass, although zooplankton biomass was unaffected. This resulted in an increase in the zooplankton to phytoplankton biomass ratio in the warmed mesocosms, which could be explained by faster turnover within the phytoplankton assemblages. Overall, warming shifted the distribution of phytoplankton body size towards smaller individuals with rapid turnover and low standing biomass, resulting in a reorganisation of the biomass structure of the food webs. These results indicate future environmental warming may have profound effects on the structure of aquatic communities.</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



SCHOLARONE™  
Manuscripts

For Review Only

1  
2  
3  
4 1 **Warming Alters the Size Spectrum and Shifts the Distribution of**  
5  
6 2 **Biomass in Aquatic Ecosystems**  
7  
8  
9 3

10  
11 4 **Running title:** Warming Alters Community Size Structure  
12

13 5 **Gabriel Yvon-Durocher<sup>1</sup>, José M. Montoya<sup>1,2</sup>, Mark Trimmer<sup>1</sup>, and Guy**  
14 **Woodward<sup>1</sup>**  
15  
16

17  
18  
19 8 <sup>1</sup> *School of Biological & Chemical Sciences, Queen Mary University of London, London*  
20 *E1 4NS. U.K.*  
21

22 10 <sup>2</sup> *Institute of Marine Sciences (ICM-CSIC) Pg. Marítim de la Barceloneta, 37-49*  
23 *E-08003 Barcelona, Spain*  
24

25  
26 12 emails (g.yvon-durocher@qmul.ac.uk; montoya@icm.csic.es; m.trimmer@qmul.ac.uk;  
27 g.woodward@qmul.ac.uk)  
28  
29  
30  
31  
32

33  
34 16 **Keywords:** Global Warming, Body Size, Size Spectra, Mass-abundance relationships,  
35  
36 17 Biomass, Food Webs, Phytoplankton, Zooplankton.  
37  
38  
39  
40  
41  
42  
43

44 21 **Authors for correspondence:** Gabriel Yvon-Durocher (g.yvon-durocher@qmul.ac.uk)  
45 and Guy Woodward (g.woodward@qmul.ac.uk)  
46

47 23 School of Biological & Chemical Sciences,  
48  
49 24 Queen Mary University of London,  
50  
51 25 London E1 4NS. U.K.

52 26 TEL:+44 20 7882 3640  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 29 **ABSTRACT**  
4

5 30 Body size is one of the key determinants of community structure. The relationship  
6  
7 31 between abundance and body size can explain how community biomass is partitioned  
8  
9  
10 32 among the biota of an ecosystem. We used an aquatic mesocosm experiment to determine  
11  
12 33 how warming of ~4°C would affect the body size, biomass and taxonomic structure of  
13  
14 34 planktonic communities. We found that warming increased the steepness of the slope of  
15  
16  
17 35 the community size spectrum, primarily by altering the phytoplankton size spectrum.  
18  
19 36 Warming also reduced the mean and maximum body size of phytoplankton by  
20  
21 37 approximately one order of magnitude. The observed shifts in phytoplankton size  
22  
23  
24 38 structure were reflected in large shifts in phytoplankton community composition, though  
25  
26 39 zooplankton taxonomic composition was unaffected by warming. Furthermore, warming  
27  
28  
29 40 reduced community biomass and total phytoplankton biomass, although zooplankton  
30  
31 41 biomass was unaffected. This resulted in an increase in the zooplankton to phytoplankton  
32  
33 42 biomass ratio in the warmed mesocosms, which could be explained by faster turnover  
34  
35  
36 43 within the phytoplankton assemblages. Overall, warming shifted the distribution of  
37  
38 44 phytoplankton body size towards smaller individuals with rapid turnover and low  
39  
40 45 standing biomass, resulting in a reorganisation of the biomass structure of the food webs.  
41  
42  
43 46 These results indicate future environmental warming may have profound effects on the  
44  
45 47 structure of aquatic communities.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 48 INTRODUCTION

49 Body size can play a key role in determining community structure (Elton 1927, Lindeman  
50 1942, Damuth 1981, Peters 1983, Brown *et al.* 2004, Petchey *et al.* 2008) because it  
51 influences ecological processes across multiple levels of organisation; from individuals  
52 (Peters 1983, Brown *et al.* 2004), and their interactions (Emmerson & Raffaelli 2004,  
53 Berlow *et al.* 2009), to populations (Damuth 1981, Jennings & Mackinson 2003, Reuman  
54 *et al.* 2008), communities and ecosystems (Petchey *et al.* 2008). Understanding how this  
55 “size-structure” might then be altered by human impacts is an important contemporary  
56 challenge for ecology given recent concerns over accelerating rates of biodiversity loss  
57 and climate change (Pimm 2009, Montoya & Raffaelli 2010).

58 The relationship between abundance and body size (equal to body mass; terms are  
59 interchangeable hereafter) is potentially a very powerful descriptor of how energy and  
60 nutrients are partitioned within the biomass of an ecosystem (White *et al.* 2007). It is also  
61 an emergent property of size structure at lower levels of organisation: for example, body  
62 size can be important for determining the presence and strength of trophic interactions  
63 between individuals because it constrains their metabolic requirements (Berlow *et al.*  
64 2009). The trophic architecture of the community determines the amount of energy  
65 available to an organism of a given size, and therefore its population abundance (Damuth  
66 1981). The relationship between abundance and body mass therefore integrates size-  
67 structure over many levels of organisation.

68 Since the pioneering work of Sheldon *et al.* (1972) the relationship between  
69 abundance and body size in pelagic food webs has typically been conceptualised as a  
70 frequency distribution of individual body sizes (Sheldon *et al.* 1972). This relationship

1  
2  
3 71 has been dubbed the “size spectrum” (Kerr & Dickie 2001). The negative slopes of size  
4  
5 72 spectra describe how quickly abundance decreases with size, and have often been used to  
6  
7  
8 73 assess the ecological status of ecosystems impacted by fisheries (Rice & Gislason 1995)  
9  
10 74 and, more recently, agricultural practices in terrestrial ecosystems (Mulder & Elser 2009).  
11  
12 75 For example, steep size spectra with negative slopes in marine ecosystems are indicative  
13  
14 76 of over-fishing because the relative abundance of large organisms is suppressed by size-  
15  
16  
17 77 selective harvesting (Pauly *et al.* 1998).

18  
19  
20 78 Understanding how the distribution of biomass in aquatic ecosystems might  
21  
22 79 respond to warming is crucial for predicting the robustness and functioning of these  
23  
24 80 ecosystems in the future warmer climate. New evidence suggests that “*reduced body size*  
25  
26 81 *is the third universal response to global warming, besides range, and phenological*  
27  
28 82 *shifts*” (Daufresne *et al.* 2009). Changes in the size-structure of communities in response  
29  
30 83 to warming are now being documented across a range of ecosystem types and spatial  
31  
32 84 scales. For instance, experiments on aquatic micro-organisms have found that warmed  
33  
34 85 communities tend to be dominated by smaller bacteria (Daufresne *et al.* 2009).  
35  
36  
37 86 Macroecological studies across latitudinal temperature gradients (Moran *et al.* 2010), and  
38  
39 87 paleoecological studies (Finkel *et al.* 2005) in the open ocean have revealed an increased  
40  
41 88 prevalence of small phytoplankton in warmer oceanic regions. These studies suggest that  
42  
43 89 the underlying size structure of aquatic ecosystems might not be robust to global warming  
44  
45  
46 90 (Finkel *et al.* 2005, Falkowski & Oliver 2007, Daufresne *et al.* 2009, Winder *et al.* 2009,  
47  
48 91 Finkel *et al.* 2010, Moran *et al.* 2010).  
49  
50

51  
52 92 However, these studies have tended to focus on the effects of warming on  
53  
54 93 restricted subsets of species (e.g. diatoms or phytoplankton) within an ecosystem (Finkel  
55  
56  
57  
58  
59  
60

1  
2  
3 94 *et al.* 2005, Winder *et al.* 2009) and documented changes in body size across latitudinal  
4  
5 95 gradients where other factors (i.e. nutrient limitation) are potentially confounded with  
6  
7  
8 96 temperature (Moran *et al.* 2010). At present, we still lack sufficient data documenting the  
9  
10 97 effects of warming *per se* on the size-structure of entire local communities comprising  
11  
12 98 multiple trophic levels to be able to isolate its effects at this level of biological  
13  
14  
15 99 organisation.  
16

17  
18 100 Here we attempt to address this current knowledge gap by measuring for the first  
19  
20 101 time the consequences of experimental warming on the community size structure and  
21  
22 102 distribution of biomass of entire planktonic food webs from 20 replicated freshwater  
23  
24 103 mesocosms. These experimental systems were maintained at either ambient temperature  
25  
26  
27 104 ( $n = 10$ ), or  $\sim 4^{\circ}\text{C}$  above ambient ( $n = 10$ ), in line with warming scenarios predicted for  
28  
29 105 temperate latitudes by the end of the 21<sup>st</sup> century (IPCC 2007). Mesocosm scale  
30  
31  
32 106 experiments such as these afford the opportunity to isolate the effects of temperature from  
33  
34 107 other potentially confounding variables (e.g. spatial gradients in available nutrients) on  
35  
36  
37 108 the structure of entire replicated communities. They also permit direct comparisons to be  
38  
39 109 made between the structure of communities under ambient conditions with that of their  
40  
41 110 “future” warmed counterparts. We used this experiment to test the following hypotheses:  
42

43  
44 111 (i) Warming will shift the distribution of body size by increasing the prevalence  
45  
46 112 of small sized species, resulting in an overall steepening of the slope of the community  
47  
48 113 size spectrum. We expect this effect to be most pronounced in the phytoplankton  
49  
50 114 assemblages because phytoplankton size structure tends to be strongly related to the  
51  
52 115 prevailing physical and chemical environment (Reynolds 1984) and recent observations  
53  
54  
55 116 in aquatic ecosystems suggest that warming tends to favour smaller phytoplankton  
56  
57  
58  
59  
60



1  
2  
3 117 (Finkel *et al.* 2005, Falkowski & Oliver 2007, Daufresne *et al.* 2009, Winder *et al.* 2009,  
4  
5  
6 118 Moran *et al.* 2010).

7  
8 119 (ii) Warming will reduce total standing community biomass. Again, we expect  
9  
10 120 this effect to be most pronounced for phytoplankton for two reasons. First, a shift in the  
11  
12 121 community size spectrum towards smaller species should result in an overall reduction in  
13  
14 122 the standing biomass. Second, theoretical expectations from the metabolic theory of  
15  
16 123 ecology (MTE) suggest total standing biomass should decline with increasing  
17  
18 124 temperature (Allen *et al.* 2002), such that the total standing biomass in a community ( $B_{\text{tot}}$ )  
19  
20 125 is predicted to vary as  $B_{\text{tot}} = r_0 e^{-E/kT} M^{1/4}$  where  $r_0$  is the resource supply rate,  $e^{-E/kT}$  is  
21  
22 126 the Boltzmann factor where  $E$  is the activation energy of metabolism,  $k$  is Boltzmann's  
23  
24 127 constant and  $T$  is absolute temperature. Therefore, holding  $r_0$  constant (i.e. if the supply  
25  
26 128 rate of limiting resources does not vary with  $T$ ),  $B_{\text{tot}}$  should decline with increases in  
27  
28 129 environmental temperature according to  $e^{-E/kT}$ .

29  
30  
31 130 (iii) Warming will alter the relative distribution of biomass between  
32  
33 131 phytoplankton and zooplankton assemblages. We expect that a shift in phytoplankton size  
34  
35 132 structure and a concomitant reduction in standing biomass will result in elevated  
36  
37 133 zooplankton-to-phytoplankton biomass ratios in the warmed mesocosms. We also predict  
38  
39 134 that relatively high zooplankton biomass will be retained in the warmed mesocosms,  
40  
41 135 because phytoplankton turnover rates should increase in response to metabolic  
42  
43 136 stimulation by warming and by a shift towards smaller species with faster generation  
44  
45 137 times. Comparable shifts in the organisation of plankton communities have been  
46  
47 138 observed in the open ocean (Gasol *et al.* 1997) and in lakes (del Giorgio & Gasol 1995)  
48  
49 139 along large scale spatial gradients of nutrient limitation.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 1404  
5 141 **MATERIAL AND METHODS**6  
7 142 *Experimental Design*

8  
9  
10 143 The experiment was carried out between December 2005 and April 2008 at the  
11  
12 144 Freshwater Biological Association, River Laboratory (2°10`W, 50°13`N), East Stoke,  
13  
14 145 Dorset, UK. A detailed description of the experimental set-up has been described  
15  
16 146 elsewhere (Yvon-Durocher *et al.* 2010). The experiment consisted of twenty freshwater  
17  
18 147 mesocosms (~1 m<sup>3</sup>, 0.5 m water depth): ten replicates remained at ambient temperature,  
19  
20 148 whilst the other ten were warmed and maintained at 3-5°C (mean 4°C) above ambient.  
21  
22 149 The mesocosms were seeded in December 2005 with organic substrates and a suite of  
23  
24 150 organisms to include the main components of organismal composition and physical  
25  
26 151 structure of shallow lake ecosystems. The established communities contained  
27  
28 152 representative species from primary producers (phytoplankton, including: *Botryococcus*,  
29  
30 153 *Chlorella*, *Volvox*, *Scenedesmus*) to top predators (Roach, *Rutilus rutilus*. Linnaeus), and  
31  
32 154 a suite of zooplankton consumers (including cyclopoid and calanoid copepods,  
33  
34 155 caldocerans, and rotifers). The biota was left to establish for ten months prior to  
35  
36 156 experimental warming, which commenced in September 2006, thereby allowing time for  
37  
38 157 further, natural colonisation before the onset of the study in April 2007. Populations of  
39  
40 158 the introduced top predator, *R. rutilus* were maintained at a constant densities [two  
41  
42 159 individuals (age 1+) per mesocosm (~12 g C m<sup>-3</sup>)] in all mesocosms and monitored via  
43  
44 160 regular electro-fishing surveys. Because the fish were maintained at predetermined  
45  
46 161 biomass-densities they merely served to “complete” the food webs to mimic natural  
47  
48 162 shallow lakes and were not considered further in the analyses.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 1634  
5 164 *Measuring the Size Spectrum*

6  
7  
8 165 The plankton communities from each of the 20 mesocosms were sampled at the beginning  
9  
10 166 and end of the growing season in April and October 2007 respectively (Yvon-Durocher *et*  
11  
12 167 *al.* 2010). The entire water column (depth 0.5m) from the sediment surface to the water  
13  
14 168 surface was sampled using a 0.8m – long tube sampler (Volume: 2L), which was  
15  
16 169 positioned at random in each mesocosm on each date. Each sample was divided into two  
17  
18 170 size categories for preservation and subsequent analyses, via filtration through a 80µm  
19  
20 171 sieve: organisms that were retained were preserved in 4% Formalin, and of the remaining  
21  
22 172 sample (i.e. organisms <80 µm), a 100ml sub-sample was preserved in 1% Lugol's iodine  
23  
24 173 for microscopy analyses.

25  
26  
27  
28  
29 174 Plankton >80 µm were counted, measured and identified by microscopy (using a  
30  
31 175 Nikon SMZ1500 dissection microscope). Zooplankton > 80 µm were typically assigned  
32  
33 176 to taxonomic orders, though in a number of cases rotifers were identified to species level.  
34  
35 177 Planktonic organisms <80µm were counted, measured and identified by inverted  
36  
37 178 microscopy. Phytoplankton <80µm were typically identified to genus level, which is  
38  
39 179 generally considered to be of sufficient taxonomic resolution to detect the effects of a  
40  
41 180 perturbation (Cottingham & Carpenter 1998). Organisms were settled for 24 h in a 10ml  
42  
43 181 Utermöhl sedimentation chamber before viewing under an inverted light microscope  
44  
45 182 (Leica DMIRE2). An initial scan of the sample, viewed under low magnification (150×),  
46  
47 183 of a fixed area (50 mm<sup>2</sup>) was used to count and measure large, rare organisms. At higher  
48  
49 184 magnification (630×), *n* fields of view were chosen at random and all organisms were  
50  
51 185 counted, sized and identified until a minimum of 400 individuals were measured from  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 186 each sample. This was sufficient to estimate 95% of the variance in the distribution of  
4  
5 187 body size (Fig S1 in supplementary material) given that settlement of organisms followed  
6  
7  
8 188 a Poisson distribution within the sedimentation chamber (Fig S2 in supplementary  
9  
10 189 material).

11  
12  
13 190 Linear body dimensions were determined with an interactive image analysis  
14  
15 191 system (Hamamatsu C4742-95 camera and Openlab software). Body size of all  
16  
17 192 organisms was expressed in units of carbon ( $\mu\text{g C}$ ). For organisms  $>80\mu\text{m}$  (typically  
18  
19 193 zooplankton), biovolumes were determined by assigning organisms to geometric shapes  
20  
21 194 that closely represented the real shape of the organism (Ruttner-Kolisko 1977, Reiss &  
22  
23 195 Schmid-Araya 2008). Body mass was determined by converting biovolume to  
24  
25 196 freshweight using a factor of 1.1, and carbon content was then estimated from a dry/wet  
26  
27 197 weight ratio of 0.25 and a dry carbon content of 40% (Reiss & Schmid-Araya 2008). For  
28  
29 198 organisms  $<80\mu\text{m}$  (typically phytoplankton) biovolumes, were similarly estimated from  
30  
31 199 geometric shapes that were most similar to the shape of the organism (Hillebrand *et al.*  
32  
33 200 1999). Biovolume was then converted into carbon units assuming a multiplication factor  
34  
35 201 of 0.109 (Montagnes *et al.* 1994). In total 47,699 individual organisms of both  
36  
37 202 phytoplankton and zooplankton were measured.  
38  
39  
40  
41  
42  
43  
44  
45

#### 204 ***Phytoplankton turnover***

46  
47 205 Turnover rates of the phytoplankton assemblages ( $\mu\text{g C m}^{-3} \text{ d}^{-1}/\mu\text{g C m}^{-3}$ ) were estimated  
48  
49 206 for each mesocosm on each sampling occasion ( $n = 40$ ). Phytoplankton turnover was  
50  
51 207 calculated as the quotient of primary production and standing phytoplankton biomass  
52  
53 208 after Gasol *et al.* (1997). This gives an estimate of the biomass specific production, or the  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 209 rate at which the carbon in the assemblage turns over. Measurements of primary  
4  
5 210 production were made simultaneously using the dissolved oxygen change technique and  
6  
7  
8 211 are presented in detail in Yvon-Durocher *et al.* (2010). Benthic metabolism measured  
9  
10 212 using *in-situ* benthic chambers contributed, on average, 35 % of whole system respiration  
11  
12 213 (see S7 in supplementary material for further details). From this we infer that our  
13  
14 214 measures of primary production predominantly reflect planktonic metabolism and  
15  
16 215 provide reliable estimates of carbon turnover within the phytoplankton assemblages.  
17  
18  
19  
20 216

### 21 217 ***Constructing the Size Spectrum***

22  
23  
24 218 The community size spectrum ( $n = 40$ ), which included phytoplankton and zooplankton,  
25  
26 219 and the phytoplankton assemblage size spectrum ( $n = 40$ ) were constructed for each  
27  
28 220 mesocosm in April and October 2007. The size spectrum of the zooplankton assemblage  
29  
30 221 alone could not be constructed accurately due to the relatively small body mass range and  
31  
32 222 the low number of individuals present in some samples. Size spectra were constructed by  
33  
34 223 logarithmic binning of the body masses ( $M$ ) of the individuals measured in each  
35  
36 224 mesocosm (either the entire community or just the phytoplankton). The total range of  
37  
38 225  $\log_{10}(M)$  values was divided into 10 bins of equal width and the  $\log_{10}$  of the total  
39  
40 226 population abundance of all organisms with  $\log_{10}(M)$  in each bin was regressed against  
41  
42 227 the bin centres (Reuman *et al.* 2008, White *et al.* 2008). The slope of the linear model  
43  
44 228 describes how quickly the abundance of individuals declines with increasing size in the  
45  
46 229 size spectrum (see Tables S5 and S6 in supplementary material). We also measured two  
47  
48 230 normalisation constants of the linear model. The intercept at  $x = 0$ : its variation between  
49  
50 231 warmed and ambient treatments gives information on the relative abundance of large  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 232 organisms, and the intercept at  $x = -8$ : its variation provides information on the  
4  
5 233 differences among treatments in the relative abundance of the smallest organisms. For  
6  
7  
8 234 both the community and the phytoplankton size spectrum, non-significant coefficients of  
9  
10 235 the linear models (i.e. at  $P > 0.05$ ) were excluded from further analyses ( $n = 5$  out of 40 for  
11  
12 236 the phytoplankton size spectrum).  
13  
14

15 237

16  
17 238 *Statistical Analyses*

18  
19  
20 239 We analysed differences between treatments in the following community properties: size  
21  
22 240 spectrum slopes and intercepts; total community biomass; total phytoplankton biomass;  
23  
24 241 total zooplankton biomass; and mean individual body mass, using ANOVA, with  
25  
26  
27 242 treatment (either warmed or ambient) and sampling occasion (April or October) as fixed  
28  
29 243 factors. The relationships between phytoplankton and zooplankton biomass and the  
30  
31 244 biomass ratio of zooplankton to phytoplankton were determined using ANCOVA, again  
32  
33 245 using treatment and sampling occasion as factors. In all statistical modelling procedures  
34  
35 246 the most parsimonious model was identified using the Akaike Information Criterion  
36  
37 247 (AIC). Statistical analyses were performed using R statistical software (R. Development.  
38  
39 248 Core. 2006).  
40  
41  
42

43 249 Multivariate analysis of phytoplankton taxonomic composition was conducted  
44  
45 250 using the vegan package in R. Redundancy analysis (RDA) was used to test for a  
46  
47 251 significant linear trend in community composition. RDA is a constrained form of  
48  
49 252 principal components analysis and assesses the variation in taxonomic composition that  
50  
51 253 can be explained by specific environmental variables defined as the constraints. Here, the  
52  
53 254 first RDA axis quantified the linear component of the between treatment variation in  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 255 phytoplankton taxonomic composition. Consequently, it was used to assess the strength  
4  
5 256 of the trend and its significance was tested using permutation tests. The  $F$ -ratio of the first  
6  
7  
8 257 RDA axis was compared with those of 999 permutations, to assess the statistical  
9  
10 258 significance of the linear trend. As well as treatment (warming),  $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ,  $\text{NH}_4^+$ ,  $\text{PO}_4^{3-}$ ,  
11  
12 259 and total inorganic N:P (see S3 for details on nutrient measurements) were tested as  
13  
14 260 constraining environmental variables. Phytoplankton taxon biomass was transformed  
15  
16 261 prior to the construction of the RDA by taking the proportional contribution of a given  
17  
18 262 taxa as a fraction of the total biomass in a given mesocosm. Furthermore, rare genera  
19  
20 263 defined as those occurring in less than two mesocosms per sampling date were excluded  
21  
22 264 from the RDA analysis to reduce noise in the data.  
23  
24  
25  
26  
27  
28

## 29 266 **RESULTS**

### 30 267 *Effects of Warming on the Size Spectrum*

31  
32 268 Warming significantly increased the steepness of the slope of the community size  
33  
34 269 spectrum from -0.86 (95% CI -0.83 to -0.89) in the systems at ambient temperature to -  
35  
36 270 0.95 (95% CI -0.92 to -0.98) in the warmed mesocosms (Fig. 1 a, b & c; Table 1), i.e.,  
37  
38 271 smaller organisms were relatively more abundant than large organisms in the warmed  
39  
40 272 communities. Furthermore, the intercept of the community size spectrum at  $x = 0$  (i.e. at  
41  
42 273 large body masses) was significantly reduced, whilst the intercept at  $x = -8$  (i.e. at small  
43  
44 274 body masses) was significantly elevated in the warmed mesocosms (Table 1). Thus, the  
45  
46 275 abundance of larger organisms declined on average, while the abundance of small  
47  
48 276 organisms increased in response to warming.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 277 Comparable patterns were observed for the phytoplankton size spectrum (Fig. 1d,  
4  
5  
6 278 e & f). Warming significantly increased the steepness of the slope of the phytoplankton  
7  
8 279 size spectrum from -0.36 (95% CI -0.32 to -0.40) in the systems at ambient temperature  
9  
10 280 to -0.49 (95% CI -0.43 to -0.55) in the warmed mesocosms (Table 1; Fig. 1d, e & f).  
11  
12 281 Warming also significantly reduced the intercept of the phytoplankton size spectrum  
13  
14 282 (Table 1). Therefore, small organisms were relatively more abundant than large  
15  
16 283 organisms in the warmed mesocosms. Additionally, warming truncated the upper size  
17  
18 284 classes of the phytoplankton size spectrum (Fig. 1 d). The maximum phytoplankton body  
19  
20 285 mass in the ambient mesocosms was  $1.36 \times 10^{-2}$   $\mu\text{g C}$ , while in the heated the maximum  
21  
22 286 body mass was only  $3.88 \times 10^{-3}$   $\mu\text{g C}$ . Furthermore, the average body mass of an individual  
23  
24 287 phytoplankter was almost an order of magnitude smaller in the warmed mesocosms  
25  
26 288 relative to the ambient systems (Fig. 2; Table 1), while the average size of an individual  
27  
28 289 zooplankter was unaffected by warming (Fig. 2; Table 1).  
29  
30  
31  
32  
33

### 34 290

### 35

### 36 291 *Effects of Warming on Community Composition*

### 37

38 292 Redundancy analysis of the phytoplankton taxa revealed that the composition of the  
39  
40 293 phytoplankton assemblages were significantly different between warmed and ambient  
41  
42 294 treatments in both April (Fig. 3a;  $F$ -ratio = 5.72;  $P$  = 0.011; permutation number = 999)  
43  
44 295 and October (Fig. 3b;  $F$ -ratio = 5.87;  $P$  = 0.001; permutation number = 999). RDA1  
45  
46 296 which was constrained by treatment, explained 24.1% and 24.6% of the variation in the  
47  
48 297 taxonomic composition of the phytoplankton assemblages in April and October  
49  
50 298 respectively, which in both cases was greater than the variation explained by PCA1,  
51  
52 299 indicating that treatment effects were the dominant predictor of phytoplankton taxonomic  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 300 composition. We also tested for significant relationships between phytoplankton  
4  
5 301 taxonomic composition and other environmental variables ( $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ,  $\text{NH}_4^+$ ,  $\text{PO}_4^{3-}$ , total  
6  
7 302 inorganic N:P) using permutation tests, though none of these variables significantly  
8  
9 303 predicted taxonomic composition. Certain taxa were strongly associated with either  
10  
11 304 warmed or ambient treatments. For example, in both April and October, the large  
12  
13 305 chlorophyte, *Botryococcus* clustered towards the ambient treatment centroid, while the  
14  
15 306 small cyanophyte *Synechocystis*, and the small chlorophyte *Monoraphidium*, typically  
16  
17 307 clustered towards the heated centroid. The phytoplankton assemblages consisted of many  
18  
19 308 rare, generalist taxa that were present in both treatments; however, in most of the  
20  
21 309 mesocosms the biomass was dominated by a few indicator taxa (named above) that were  
22  
23 310 associated with either the heated or the ambient treatments. Furthermore, figures 3a and  
24  
25 311 3b show that a large core contingent of the phytoplankton assemblages were present in  
26  
27 312 both April and October and that only a few taxa were present in only one month,  
28  
29 313 suggesting that temporal succession was less important than treatment effects in  
30  
31 314 determining phytoplankton community composition.

32  
33  
34  
35  
36  
37  
38  
39 315 In contrast to the phytoplankton assemblages the taxonomic composition of the  
40  
41 316 zooplankton assemblages differed very little between treatments in both April and  
42  
43 317 October (Fig. 4a & b). In heated and ambient treatments calanoid and cyclopoid copepods  
44  
45 318 dominated zooplankton biomass with cladocerans and rotifers forming a smaller  
46  
47 319 secondary contingent of the assemblages. These patterns were consistent between April  
48  
49 320 and October, though ostracods, oligochates and the rotifer *Asplanchna* were absent from  
50  
51 321 the zooplankton assemblage in October.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 323 *Effects of Warming on the Distribution of Biomass*  
4

5  
6 324 Total planktonic community biomass differed between April and October in the ambient  
7

8 325 but not in the warmed mesocosms (Fig. 5). Overall, warming significantly reduced total  
9

10 326 community biomass (Fig. 5; Table 1). This was principally driven by a considerable  
11

12 327 reduction in total phytoplankton biomass in the warmed mesocosms (Fig. 5; Table 1).  
13

14  
15 328 Overall, warming shifted the distribution of biomass and body size of phytoplankton from  
16

17 329 assemblages comprised of large individuals with high standing biomass to assemblages  
18

19 330 with low standing biomass and many small individuals. In contrast, warming appeared to  
20

21 331 have no effect on the distribution of size and biomass within the zooplankton  
22

23 332 assemblages (Fig. 5; Table 1).  
24  
25  
26

27 333 Zooplankton and phytoplankton biomass were not correlated within the  
28

29 334 mesocosms (Fig. 6a; Table 2). The former varied by about two orders of magnitude and  
30

31 335 the latter by three orders of magnitude among mesocosms (Fig. 6a). The ratio of  
32

33 336 zooplankton to phytoplankton biomass (Z:P) was significantly and negatively correlated  
34

35 337 with phytoplankton biomass (Fig. 6b; Table 1). Therefore, zooplankton biomass exceeded  
36

37 338 phytoplankton biomass (i.e. Z:P >1) when phytoplankton biomass was low and vice versa  
38

39 339 (i.e. Z:P <1) when phytoplankton biomass was high. Warming significantly increased the  
40

41 340 ratio of Z:P biomass (Table 2). Furthermore, the ratio of Z:P biomass was strongly and  
42

43 341 positively correlated with the turnover rates of the phytoplankton assemblages, which  
44

45 342 exhibited distinct variation between warmed and ambient mesocosms (Fig. 6c; Table 2).  
46

47 343 In summary, the warmed mesocosms were characterised by phytoplankton assemblages  
48

49 344 comprised of small individuals with low standing stocks of biomass and rapid turnover  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 345 rates which supported relatively high standing stocks of zooplankton, exemplified by  
4  
5 346 high Z:P biomass ratios.  
6  
7

8 347

9  
10 348 **DISCUSSION**

11  
12 349 There is ample evidence that ecological responses to recent climate change are already  
13  
14 350 occurring at the species (and therefore the population) level (Walther *et al.* 2002), but  
15  
16 351 scaling from populations to communities and ecosystems is challenging because of the  
17  
18 352 perceived indeterminacy of ecological interactions (Yodzis & Innes 1992). Therefore,  
19  
20 353 there is an increasingly urgent need to explore the effects of the principal components of  
21  
22 354 climate change (e.g., warming) on community structure and ecosystem functioning  
23  
24 355 (Tylianakis *et al.* 2008, Montoya & Raffaelli 2010). Our results broadly supported our  
25  
26 356 experimental hypotheses: i.e., that warming would increase the steepness of the size  
27  
28 357 spectrum slope, reduce total community biomass, and increase the zooplankton to  
29  
30 358 phytoplankton biomass ratio. These findings could provide some novel insights into how  
31  
32 359 future warming might change the distribution of body size and biomass in aquatic  
33  
34 360 ecosystems. The size structure of plankton communities in aquatic ecosystems is a key  
35  
36 361 driver of rates of carbon sequestration and nutrient cycling (Laws *et al.* 2000), and  
37  
38 362 therefore changes in the distribution of planktonic body size and biomass could alter the  
39  
40 363 regulation of biotic feedbacks with warming on a potentially global scale (Falkowski *et*  
41  
42 364 *al.* 1998).  
43  
44  
45  
46  
47  
48  
49

50 365 The general increase in the prevalence of small organisms with increases in  
51  
52 366 environmental temperature that we observed experimentally agrees well with recent  
53  
54 367 studies that have either focused on specific taxa, or subsets of taxa (Atkinson *et al.* 2003,  
55  
56  
57  
58  
59  
60

1  
2  
3 368 Finkel *et al.* 2005, Daufresne *et al.* 2009, Winder *et al.* 2009), or analysed correlational  
4  
5 369 trends in community structure across latitudinal gradients in temperature (Moran *et al.*  
6  
7  
8 370 2010). However, here we have developed this understanding further by documenting the  
9  
10 371 effects of warming on the body size, biomass and taxonomic structure of entire  
11  
12 372 planktonic food webs subjected to experimental warming. Experimental mesocosm  
13  
14  
15 373 studies, although inevitably an abstraction of natural ecosystems, afford us the  
16  
17 374 opportunity to isolate the effects of temperature from other potentially confounding  
18  
19 375 variables (e.g. latitudinal and biogeographical effects) whilst studying entire replicated  
20  
21 376 plankton communities.  
22  
23

24 377 The increase in the dominance of small phytoplankton and the truncation of the  
25  
26 378 larger size classes in their size spectrum resulted in a general increase in the steepness of  
27  
28 379 the slope of the community size spectrum in the warmed mesocosms. Changes in the  
29  
30 380 distribution of organism size might arise from at least two broad mechanisms, which are  
31  
32 381 not necessarily mutually exclusive. Firstly, organisms might exhibit a degree of  
33  
34 382 phenotypic plasticity to changes in temperature. This hypothesis has been termed the  
35  
36 383 temperature-size rule (Atkinson *et al.* 2003) and posits that reduced organism size at  
37  
38 384 higher temperatures is an adaptive plastic response that results from selection for earlier  
39  
40 385 reproduction as population growth rate increases. The accelerated completion of the life  
41  
42 386 cycle occurs at the expense of maturation size (Atkinson *et al.* 2003). In the second  
43  
44 387 mechanism, changes in the physicochemical environment created by warming select for  
45  
46 388 smaller sized species. In this case, changes in community size structure occur as an  
47  
48 389 indirect effect of warming, mediated for example, by concomitant nutrient limitation,  
49  
50  
51 390 resulting in the competitive exclusion of larger species (Finkel *et al.* 2005, Irwin *et al.*  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 391 2006, Falkowski & Oliver 2007, Winder *et al.* 2009, Finkel *et al.* 2010). Here, small cell  
4  
5 392 size increases the efficiency of the acquisition of limiting nutrients because of a higher  
6  
7  
8 393 surface area to volume ratio and is therefore competitively advantageous under conditions  
9  
10 394 of nutrient limitation (Litchman *et al.* 2009).

11  
12 395 Our results support the second mechanism. Redundancy analysis revealed that  
13  
14 396 warming dramatically shifted the taxonomic composition of the phytoplankton  
15  
16  
17 397 assemblages. Moreover, warming favoured smaller phytoplankton genera, resulting in a  
18  
19 398 reduction in mean and maximum body size by almost an order of magnitude. For  
20  
21 399 example, the large chlorophyte *Botryococcus* dominated the biomass of the ambient  
22  
23 400 mesocosms in both April and October, but was almost entirely absent from the warmed  
24  
25 401 mesocosms. Similarly the small cyanophyte *Synechocystis* and the small chlorophyte  
26  
27 402 *Monoraphidium* were strongly associated with the warmed mesocosms but were only  
28  
29 403 peripheral members of the assemblages in the ambient mesocosms. Warming therefore  
30  
31 404 resulted in the establishment of phytoplankton assemblages dominated by small species,  
32  
33 405 rather than reducing the body size of the same species composition present in the ambient  
34  
35 406 mesocosms.

36  
37  
38  
39 407 Our experimental design adopted a space-for-time substitution approach to  
40  
41 408 attempt to understand the consequences of warming on the structure of plankton  
42  
43 409 communities. The relatively infrequent but highly replicated sampling regime adopted in  
44  
45 410 our study was a necessary compromise. For example, we documented the size, biomass  
46  
47 411 and taxonomic structure of 20 replicated experimental ecosystems on two separate  
48  
49 412 sampling occasion at the beginning and end of the growing season (identified from  
50  
51 413 measures of primary production; see Yvon-Durocher *et al.*, (2010) for details) rather than  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 414 focusing on the complex temporal dynamics of the plankton assemblages of one or two  
4  
5 415 systems, as would typically be logistically feasible within such a study. As a result, our  
6  
7 416 results come with an associated caveat: we are unable to discern the effects of warming  
8  
9 417 on the temporal succession of the plankton communities. However, analysis of the  
10  
11 418 phytoplankton taxonomic composition suggests that a large, core contingent of these  
12  
13 419 assemblages are present in both April and October but which differ markedly between  
14  
15 420 treatments in both months. These results suggest that temporal succession in the plankton  
16  
17 421 communities was less important than the effect of treatment (i.e. warming) in determining  
18  
19 422 the taxonomic and therefore the body size and biomass structure of these assemblages.  
20  
21  
22  
23

24  
25 423 Inorganic nitrogen was limiting in our experiment (N:P ratios were  $\approx 11:1$ , and  
26  
27 424 were below the 16N:1P expected at Redfield; see S3 & S4 in supplementary material for  
28  
29 425 further details) but to the same extent in both warmed and ambient treatments: i.e.,  
30  
31 426 warming did not exacerbate nutrient limitation. Therefore, it is unlikely that a direct  
32  
33 427 effect of nutrient limitation induced by warming caused the observed shifts in  
34  
35 428 phytoplankton size structure that have been frequently documented in the open ocean and  
36  
37 429 in lake ecosystems (Finkel *et al.* 2005, Falkowski & Oliver 2007, Winder *et al.* 2009,  
38  
39 430 Finkel *et al.* 2010). Furthermore, redundancy analysis revealed that inorganic nutrient  
40  
41 431 concentrations ( $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ,  $\text{NH}_4^+$ ,  $\text{PO}_4^{3-}$ , and total inorganic N:P) were not significantly  
42  
43 432 correlated with phytoplankton taxonomic composition. Therefore, the shift in  
44  
45 433 phytoplankton size and taxonomic structure in the warmed treatments might simply  
46  
47 434 reflect the fact that smaller phytoplankton have lower specific nitrogen requirements than  
48  
49 435 large phytoplankton (Litchman *et al.* 2007). Litchman *et al.* (2007) found that the  
50  
51 436 minimum nitrogen quota required to support growth,  $Q_{\min}$ , across a wide range of  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 437 phytoplankton taxa increases allometrically, resulting in a disproportionate increase in  
4  
5 438 cellular nitrogen quota with size. Because metabolic rates and nutrient uptake rates  
6  
7  
8 439 increase with temperature and size (Gillooly *et al.* 2001, Allen & Gillooly 2009), under  
9  
10 440 conditions of nutrient limitation, small cell size should provide a competitive advantage  
11  
12 441 as environmental temperatures rise. This is because species with lower  $Q_{\min}$  will be better  
13  
14 442 able to balance the increased demand for limiting nutrients imposed by temperature  
15  
16 443 driven elevated metabolic rates.  
17  
18

19  
20 444 An alternative mechanism for the shifts in phytoplankton size and taxonomic  
21  
22 445 structure in the warmed mesocosms is that warming served to increase “top down”  
23  
24 446 control of the phytoplankton community by increasing zooplankton grazing rates. We  
25  
26 447 have previously demonstrated that heterotrophic metabolism increased more rapidly than  
27  
28 448 autotrophic metabolism with increasing temperature in the same experimental system  
29  
30 449 (measurements made simultaneously; see Yvon-Durocher *et al.*, (2010) for details).  
31  
32 450 Therefore, because ingestion rates increase in proportion with metabolic rates (Berlow *et*  
33  
34 451 *al.* 2009), warming might have increased the strength of top down control of  
35  
36 452 phytoplankton populations by zooplankton grazing. Moreover, zooplankton are often size  
37  
38 453 selective when feeding on phytoplankton, typically consuming the largest size classes  
39  
40 454 possible (Porter 1973, Hall *et al.* 1976, Katechakis *et al.* 2002). Warming might therefore  
41  
42 455 have increased the prevalence of small sized phytoplankton indirectly, by elevating  
43  
44 456 grazing pressure on the larger size classes of the phytoplankton community due to the  
45  
46 457 elevated metabolic demands of zooplankton at higher temperature. Importantly, both the  
47  
48 458 “top down” and “bottom up” hypotheses stated here are not mutually exclusive: both  
49  
50 459 bottom up regulation of phytoplankton competitive ability for limiting nutrients, and top  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 460 down control of large phytoplankters by zooplankton grazing could occur simultaneously,  
4  
5 461 and combine with the direct effects of warming on metabolism to produce the observed  
6  
7  
8 462 shifts in size, biomass taxonomic structure.  
9

10 463 Warming reduced total standing community biomass, largely via a reduction in  
11  
12 464 phytoplankton biomass. These results confirmed our qualitative theoretical predictions.  
13  
14 465 For example, because the potential resource supply rate (i.e. the concentrations of  
15  
16  
17 466 limiting inorganic nutrients) remained constant, we predicted that elevated metabolic  
18  
19  
20 467 demands at higher temperatures should have resulted in a decline in standing community  
21  
22 468 biomass in the warmed mesocosms. Assuming  $B_{\text{tot}} = r_0 e^{-E/kT} M^{1/4}$  and that  $r_0$  (i.e. the  
23  
24 469 resource supply rate) and  $M^{1/4}$  (i.e. the allometric scaling of biomass with body mass) are  
25  
26  
27 470 constant with temperature we can predict that for  $\approx 4^\circ\text{C}$  warming (i.e. the average annual  
28  
29 471 temperature increase in our experiment) standing community biomass should decline  
30  
31 472 approximately 1.54 fold according to:  $e^{-E/kT_h} / e^{-E/kT_a}$  where  $T_h$  and  $T_a$  are the mean annual  
32  
33 473 temperatures of the heated and ambient mesocosms (290.9 and 286.1 K, respectively) and  
34  
35 474  $E$  is the activation energy of metabolism  $\approx 0.65\text{eV}$  (Gillooly *et al.* 2001). In our  
36  
37 475 experiment average total community biomass declined 2.53 fold (i.e. the ratio of mean  
38  
39 476 biomass in the heated and ambient mesocosms), almost double that predicted by  
40  
41 477 metabolic costs alone, suggesting that additional factors may be operating.  
42  
43  
44  
45

46 478 The large shift in the distribution of body size from large to small phytoplankton  
47  
48 479 might further reduce standing biomass. For example, the above prediction assumes that  
49  
50 480 the allometric scaling of biomass with body mass (i.e.  $B_{\text{tot}} = M^{1/4}$ ) remains constant with  
51  
52 481 warming. However, we have demonstrated that the slope of the community size spectrum  
53  
54  
55 482 (i.e. the log-log relationship) which is equivalent to the exponent (i.e.  $-\alpha$ ) of  $N = M^\alpha$ ,  
56  
57  
58  
59  
60



1  
2  
3 483 where  $N$  is abundance (White *et al.* 2007, Reuman *et al.* 2008, White *et al.* 2008),  
4  
5 484 changes from -0.86 to -0.95 in response to warming. Therefore, because  $B_{\text{tot}} = N \times M$  the  
6  
7  
8 485 allometric scaling of  $B_{\text{tot}}$  declined from  $B_{\text{tot}} = M^{0.14}$  in the ambient mesocosms to  
9  
10 486  $B_{\text{tot}} = M^{0.05}$  in the warmed mesocosm: i.e., more standing biomass was retained in larger  
11  
12 487 body size classes in the ambient relative to the warmed mesocosms. We suggest that the  
13  
14  
15 488 effects of increased metabolic costs, associated with warmer temperatures and the shift in  
16  
17 489 the distribution of body size and taxonomic composition of the phytoplankton  
18  
19  
20 490 assemblage, could have acted synergistically to reduce total community biomass in the  
21  
22 491 warmed mesocosms.

23  
24  
25 492 The ratio of zooplankton to phytoplankton biomass, Z:P, declined as a function of  
26  
27 493 phytoplankton biomass, in line with our third experimental hypothesis. Our results are  
28  
29 494 qualitatively similar to the findings of Gasol *et al.* (1997) who also demonstrated that the  
30  
31 495 ratio of heterotroph to autotroph biomass (H:A) was a declining function of autotroph  
32  
33 496 biomass in the open ocean and coastal seas, although they attributed the relationship to a  
34  
35  
36 497 nutrient gradient rather than temperature. In our case, the large shifts in community size  
37  
38 498 structure and the distribution of biomass between zooplankton and phytoplankton were  
39  
40 499 independent of the inorganic nutrient status of the mesocosms and appear to have been  
41  
42  
43 500 driven largely by the effects of temperature on metabolism and the relative competitive  
44  
45 501 abilities of large and small phytoplankton.

46  
47  
48 502 We found a strong, positive correlation between the Z:P biomass ratio and the  
49  
50 503 turnover rate of the phytoplankton assemblages, which differed profoundly between  
51  
52 504 warmed and ambient treatments. These results offer insight into how these communities  
53  
54  
55 505 might function: the inverted pyramid or squared biomass distributions (i.e. Z>P or Z=P)

1  
2  
3 506 in the warmed mesocosms contrasted markedly with the pyramidal biomass structure (i.e.  
4  
5 507  $Z < P$ ) of the mesocosms at ambient temperature. This suggests that warming of  $\sim 4^\circ\text{C}$   
6  
7  
8 508 fundamentally altered the structure and functioning (i.e. energy transfer) of our  
9  
10 509 experimental ecosystems. For instance, in the heated mesocosms the high relative  
11  
12 510 biomass of zooplankton may have been supported by a fast turnover rate of the  
13  
14 511 phytoplankton assemblage. For example, for the low standing stocks of phytoplankton  
15  
16 512 biomass in the warmed mesocosms ( $2.93 \times 10^5 \mu\text{g C m}^{-3}$  in heated;  $1.12 \times 10^6 \mu\text{g C m}^{-3}$  in  
17  
18 513 ambient) to sustain the equivalent biomass of zooplankton as the mesocosms at ambient  
19  
20 514 temperature ( $1.71 \times 10^5 \mu\text{g C m}^{-3}$  in ambient;  $1.36 \times 10^5 \mu\text{g C m}^{-3}$  heated), the turnover rate  
21  
22 515 of the phytoplankton community would need to be elevated by a factor of  $\sim 4$ . The  
23  
24 516 average turnover rates of the phytoplankton community in the warmed treatments was  
25  
26 517 actually elevated by a factor of  $\sim 5$  (i.e.  $40.9 \mu\text{g C m}^{-3} \text{ d}^{-1} / \mu\text{g C m}^{-3}$  in heated;  $8.25 \mu\text{g C}$   
27  
28 518  $\text{m}^{-3} \text{ d}^{-1} / \mu\text{g C m}^{-3}$  in the ambient) and was therefore sufficient to support the biomass of  
29  
30 519 zooplankton in these systems. Taken together, these results suggest that warming  
31  
32 520 increases the rate of carbon flux between autotrophs and heterotrophs. This effect appears  
33  
34 521 to be driven by the relative increase in small phytoplankton, which have faster turnover  
35  
36 522 times due to the  $-1/4$  allometry of mass specific metabolic rate and generation time  
37  
38 523 (Gillooly *et al.* 2002, Brown *et al.* 2004), and also the direct stimulation of metabolism  
39  
40 524 and generation time by temperature (Gillooly *et al.* 2002).  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

## 51 **Conclusion**

52  
53 527 In general, the results of our experiment reflect patterns in empirical surveys that  
54  
55 528 have analysed phytoplankton communities over macroevolutionary time (Finkel *et al.*  
56  
57  
58  
59  
60

1  
2  
3 529 2005, Finkel *et al.* 2007), across latitudinal gradients in temperature (Moran *et al.* 2010),  
4  
5 530 and across gradients of nutrient regime and productivity (del Giorgio & Gasol 1995,  
6  
7  
8 531 Gasol *et al.* 1997). Our results, however, represent the first experimental evidence for a  
9  
10 532 shift in the distribution of body size and biomass of whole plankton communities that can  
11  
12 533 be attributed directly to the effects of warming via a controlled and replicated whole  
13  
14 534 ecosystem manipulation. Although we now have some tantalising hints, the precise  
15  
16 535 mechanism behind the size shifts we observed requires further research. Also, the  
17  
18 536 consequences of such shifts in community size structure for the functioning (e.g. carbon  
19  
20 537 sequestration capacity) of aquatic ecosystems remains an unexplored avenue in  
21  
22 538 ecological research, though with no doubt one that will prove fundamental in addressing  
23  
24 539 the future challenges posed by environmental change.  
25  
26  
27  
28  
29  
30  
31  
32

540

541

#### 542 **Acknowledgements**

543 We thank Brian Godfrey, Dan Perkins, and the Freshwater Biological Association for  
544 their help with the experiment. Sarah Tuffin and Rebecca Stewart contributed to the  
545 counting of phytoplankton G. Yvon-Durocher was supported by a Natural Environment  
546 Research Council studentship (NER/S/A2006/14029). J. Montoya was funded by the  
547 NERC Fellowship Scheme (NE/C002105/1), and a Ramon y Cajal Fellowship (RYC-  
548 2008-03664).

549

#### 550 **References**

551 Allen AP, Brown JH, Gillooly JF (2002) Global biodiversity, biochemical kinetics, and  
552 the energetic-equivalence rule. *Science*, **297**, 1545-1548.

552

553

554

555

556

557

558

559

560

- 1  
2  
3 553 Allen AP, Gillooly JF (2009) Towards an integration of ecological stoichiometry and the  
4  
5 554 metabolic theory of ecology to better understand nutrient cycling. *Ecology*  
6  
7 555 *Letters*, **12**, 369-384.
- 8  
9  
10 556 Atkinson D, Ciotti BJ, Montagnes DJS (2003) Protists decrease in size linearly with  
11  
12 557 temperature: ca. 2.5% degrees C<sup>-1</sup>. *Proceedings of the Royal Society of London*  
13  
14 558 *Series B-Biological Sciences*, **270**, 2605-2611.
- 15  
16  
17 559 Berlow EL, Dunne JA, Martinez ND, Stark PB, Williams RJ, Brose U (2009) Simple  
18  
19 560 prediction of interaction strengths in complex food webs. *Proceedings of the*  
20  
21 561 *National Academy of Sciences of the United States of America*, **106**, 187-191.
- 22  
23  
24 562 Brown JH, Gillooly JF, Allen AP, Savage VM, West GB (2004) Toward a metabolic  
25  
26 563 theory of ecology. *Ecology*, **85**, 1771-1789.
- 27  
28  
29 564 Cottingham KL, Carpenter SR (1998) Population, community, and ecosystem variates as  
30  
31 565 ecological indicators: Phytoplankton responses to whole-lake enrichment.  
32  
33 566 *Ecological Applications*, **8**, 508-530.
- 34  
35  
36 567 Damuth J (1981) Population density and body size in mammals. *Nature*, **290**, 699-700.
- 37  
38  
39 568 Daufresne M, Lengfellner K, Sommer U (2009) Global warming benefits the small in  
40  
41 569 aquatic ecosystems. *Proceedings of the National Academy of Sciences of the*  
42  
43 570 *United States of America*, **106**, 12788-12793.
- 44  
45  
46 571 del Giorgio PA, Gasol JM (1995) Biomass distribution in freshwater plankton  
47  
48 572 communities. *American Naturalist*, **146**, 135-152.
- 49  
50 573 Elton C (1927) *Animal Ecology*. Kluwer Academics Publishers, B.V.
- 51  
52  
53 574 Emmerson MC, Raffaelli D (2004) Predator-prey body size, interaction strength and the  
54  
55 575 stability of a real food web. *Journal of Animal Ecology*, **73**, 399-409.  
56  
57  
58  
59  
60

- 1  
2  
3 576 Falkowski PG, Barber RT, Smetacek V (1998) Biogeochemical controls and feedbacks  
4  
5 577 on ocean primary production. *Science*, **281**, 200-206.  
6  
7  
8 578 Falkowski PG, Oliver MJ (2007) Mix and match: how climate selects phytoplankton.  
9  
10 579 *Nature Reviews Microbiology*, **5**, 813-819.  
11  
12 580 Finkel ZV, Beardall J, Flynn KJ, Quigg A, Rees TAV, Raven JA (2010) Phytoplankton in  
13  
14 581 a changing world: cell size and elemental stoichiometry. *Journal of Plankton*  
15  
16 582 *Research*, **32**, 119-137.  
17  
18  
19 583 Finkel ZV, Katz ME, Wright JD, Schofield OME, Falkowski PG (2005) Climatically  
20  
21 584 driven macroevolutionary patterns in the size of marine diatoms over the  
22  
23 585 cenozoic. *Proceedings of the National Academy of Sciences of the United States*  
24  
25 586 *of America*, **102**, 8927-8932.  
26  
27  
28 587 Finkel ZV, Sebbo J, Feist-Burkhardt S, *et al.* (2007) A universal driver of  
29  
30 588 macroevolutionary change in the size of marine phytoplankton over the Cenozoic.  
31  
32 589 *Proceedings of the National Academy of Sciences of the United States of America*,  
33  
34 590 **104**, 20416-20420.  
35  
36  
37 591 Gasol JM, del Giorgio PA, Duarte CM (1997) Biomass distribution in marine planktonic  
38  
39 592 communities. *Limnology and Oceanography*, **42**, 1353-1363.  
40  
41  
42 593 Gillooly JF, Brown JH, West GB, Savage VM, Charnov EL (2001) Effects of size and  
43  
44 594 temperature on metabolic rate. *Science*, **293**, 2248-2251.  
45  
46  
47 595 Gillooly JF, Charnov EL, West GB, Savage VM, Brown JH (2002) Effects of size and  
48  
49 596 temperature on developmental time. *Nature*, **417**, 70-73.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 597 Hall DJ, Threlkeld ST, Burns CW, Crowley PH (1976) Size efficiency hypothesis and the  
4  
5 598 size structure of zooplankton communities. *Annual Review of Ecology and*  
6  
7 599 *Systematics*, **7**, 177-208.
- 10 600 Hillebrand H, Durselen CD, Kirschtel D, Pollinger U, Zohary T (1999) Biovolume  
11  
12 601 calculation for pelagic and benthic microalgae. *Journal of Phycology*, **35**, 403-  
13  
14 602 424.
- 17 603 IPCC (2007) Climate Change 2007: The Physical Sciences Basis. . In: *Contribution of*  
18  
19 604 *Working Group I to the Fourth Assessment Report of the Intergovernmental Panel*  
20  
21 605 *on Climate Change*. (ed Parry ML, Canziani, O.F., Palutikof, J. P., van der  
22  
23 606 Linden, P.J. & Hanson, C. E.), pp. 7-22. Cambridge University Press, Cambridge.
- 26 607 Irwin AJ, Finkel ZV, Schofield OME, Falkowski PG (2006) Scaling-up from nutrient  
27  
28 608 physiology to the size-structure of phytoplankton communities. *Journal of*  
29  
30 609 *Plankton Research*, **28**, 459-471.
- 33 610 Jennings S, Mackinson S (2003) Abundance-body mass relationships in size-structured  
34  
35 611 food webs. *Ecology Letters*, **6**, 971-974.
- 38 612 Katechakis A, Stibor H, Sommer U, Hansen T (2002) Changes in the phytoplankton  
39  
40 613 community and microbial food web of Blanes Bay (Catalan Sea, NW  
41  
42 614 Mediterranean) under prolonged grazing pressure by doliolids (Tunicata),  
43  
44 615 cladocerans or copepods (Crustacea). *Marine Ecology-Progress Series*, **234**, 55-  
45  
46 616 69.
- 49 617 Kerr SR, Dickie LM (2001) *The Biomass Spectrum: A predator Prey Theory of Aquatic*  
50  
51 618 *Production*. Columbia University Press, New York.
- 52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 619 Laws EA, Falkowski PG, Smith WO, Ducklow H, McCarthy JJ (2000) Temperature  
4  
5 620 effects on export production in the open ocean. *Global Biogeochemical Cycles*,  
6  
7 621 **14**, 1231-1246.  
8  
9  
10 622 Lindeman RL (1942) The trophic-dynamic aspect of ecology. . *Ecology*, **23**, 399-418.  
11  
12 623 Litchman E, Klausmeier CA, Schofield OM, Falkowski PG (2007) The role of functional  
13  
14 624 traits and trade-offs in structuring phytoplankton communities: scaling from  
15  
16 625 cellular to ecosystem level. *Ecology Letters*, **10**, 1170-1181.  
17  
18  
19 626 Litchman E, Klausmeier CA, Yoshiyama K (2009) Contrasting size evolution in marine  
20  
21 627 and freshwater diatoms. *Proceedings of the National Academy of Sciences of the*  
22  
23 628 *United States of America*, **106**, 2665-2670.  
24  
25  
26 629 Montagnes DJS, Berges JA, Harrison PJ, Taylor FJR (1994) Estimating carbon, nitrogen,  
27  
28 630 protein and chlorophyll-a from volume in marine phytoplankton. *Limnology and*  
29  
30 631 *Oceanography*, **39**, 1044-1060.  
31  
32  
33 632 Montoya JM, Raffaelli D (2010) Climate change, biotic interactions and ecosystem  
34  
35 633 services. *Philosophical Transactions of the Royal Society of London Series B-*  
36  
37 634 *Biological Sciences*, **365**, 2013-2018.  
38  
39  
40 635 Moran XAG, Lopez-Urrutia A, Calvo-Diaz A, Li WKW (2010) Increasing importance of  
41  
42 636 small phytoplankton in a warmer ocean. *Global Change Biology*, **16**, 1137-1144.  
43  
44  
45 637 Mulder C, Elser JJ (2009) Soil acidity, ecological stoichiometry and allometric scaling in  
46  
47 638 grassland food webs. *Global Change Biology*, **15**, 1-9.  
48  
49  
50 639 Pauly D, Christensen V, Dalsgaard J, Froese R, Torres F (1998) Fishing down marine  
51  
52 640 food webs. *Science*, **279**, 860-863.  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 641 Petchey OL, Beckerman AP, Riede JO, Warren PH (2008) Size, foraging, and food web  
4  
5  
6 642 structure. *Proceedings of the National Academy of Sciences of the United States of*  
7  
8 643 *America*, **105**, 4191-4196.
- 9  
10 644 Peters RH (1983) *The ecological implications of body size*. Cambridge University Press,  
11  
12 645 Cambridge.
- 13  
14  
15 646 Pimm SL (2009) Climate Disruption and Biodiversity. *Current Biology*, **19**, 595-601.
- 16  
17 647 Porter KG (1973) Selective grazing and differential digestion of algae by zooplankton.  
18  
19 648 *Nature*, **244**, 179-180.
- 20  
21  
22 649 R. Development. Core. T (2006). R Foundation for Statistical Computing,  
23  
24 650 , Vienna, Austria.
- 25  
26  
27 651 Reiss J, Schmid-Araya JM (2008) Existing in plenty: abundance, biomass and diversity  
28  
29 652 of ciliates and meiofauna in small streams. *Freshwater Biology*, **53**, 652-668.
- 30  
31  
32 653 Reuman DC, Mulder C, Raffaelli D, Cohen JE (2008) Three allometric relations of  
33  
34 654 population density to body mass: theoretical integration and empirical tests in 149  
35  
36 655 food webs. *Ecology Letters*, **11**, 1216-1228.
- 37  
38  
39 656 Reynolds CS (1984) *The Ecology of Freshwater Phytoplankton* Cambridge University  
40  
41 657 Press Cambridge.
- 42  
43  
44 658 Rice J, Gislason H (1995) In: *ICES International Symposium on changes in the North Sea*  
45  
46 659 *Ecosystem and Their Causes - Arhus 1975 Revisited*, pp. 1214-1225. Academic  
47  
48 660 Press Ltd, Aarhus, Denmark.
- 49  
50  
51 661 Ruttner-Kolisko A (1977) Suggestions for biomass calculations of plankton rotifers.  
52  
53 662 *Archiv für Hydrobiologie, Beiheft Ergebnisse Limnologie*, **8**, 71-76.  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3 663 Sheldon RW, Prakash A, Sutcliffe WH (1972) The size distribution of particles in the  
4  
5 664 ocean. *Limnology and Oceanography*, **17**, 327-340.  
6  
7  
8 665 Tylianakis JM, Didham RK, Bascompte J, Wardle DA (2008) Global change and species  
9  
10 666 interactions in terrestrial ecosystems. *Ecology Letters*, **11**, 1351-1363.  
11  
12 667 Walther GR, Post E, Convey P, *et al.* (2002) Ecological responses to recent climate  
13  
14 668 change. *Nature*, **416**, 389-395.  
15  
16  
17 669 White EP, Enquist BJ, Green JL (2008) On estimating the exponent of power-law  
18  
19 670 frequency distributions. *Ecology*, **89**, 905-912.  
20  
21  
22 671 White EP, Ernest SKM, Kerkhoff AJ, Enquist BJ (2007) Relationships between body size  
23  
24 672 and abundance in ecology. *Trends in Ecology & Evolution*, **22**, 323-330.  
25  
26  
27 673 Winder M, Reuter JE, Schladow SG (2009) Lake warming favours small-sized planktonic  
28  
29 674 diatom species. *Proceedings of the Royal Society B-Biological Sciences*, **276**, 427-  
30  
31 675 435.  
32  
33  
34 676 Yodzis P, Innes S (1992) Body size and consumer-resource dynamics. *American*  
35  
36 677 *Naturalist*, **139**, 1151-1175.  
37  
38  
39 678 Yvon-Durocher G, Jones JI, Woodward G, Trimmer M, Montoya JM (2010) Warming  
40  
41 679 alters the metabolic balance of ecosystems. *Philosophical Transactions of the*  
42  
43 680 *Royal Society of London Series B-Biological Sciences*, **365**, 2117-2126.  
44  
45  
46  
47

48 **Figure Legends**

49  
50 683 **Figure. 1.** The size spectrum. (a) The community size spectrum of a heated (red circles)  
51  
52 684 and ambient (black circles) mesocosm, highlighting the increase in the steepness of the  
53  
54  
55 685 slope in the warmed mesocosm. (b) Frequency distribution of the slope of the community  
56  
57  
58  
59  
60

1  
2  
3 686 size spectrum in the ambient mesocosms ( $n=20$ ), (c) frequency distribution of the slope  
4  
5 687 of the community size spectrum in the warmed mesocosms ( $n=20$ ). On average the  
6  
7  
8 688 slope of the community size spectrum in the warmed mesocosms was significantly  
9  
10 689 steeper than the ambient mesocosms (Table 1). (d) The phytoplankton size spectrum of a  
11  
12 690 heated (red circles) and ambient (black circles) mesocosm, highlighting the increase in  
13  
14 691 the steepness of the slope and the truncation of large sized individuals in the warmed  
15  
16 692 mesocosm. (e) Frequency distribution of the slope of the phytoplankton size spectrum in  
17  
18 693 the ambient mesocosms ( $n=17$ ), (f) frequency distribution of the slope of the community  
19  
20 694 size spectrum in the warmed mesocosms ( $n=18$ ).  
21  
22  
23  
24  
25  
26

27 696 **Figure. 2.** Effects of warming on mean body mass ( $\pm 1$  s.e.m) of phytoplankton (a) and  
28  
29 697 zooplankton (b) individuals. Data are presented as the overall average of the mean body  
30  
31 698 mass of phytoplankton and zooplankton individuals over 20 mesocosms for each  
32  
33 699 treatment. The mean cell mass of phytoplankton is significantly reduced in response to  
34  
35 700 warming while there is no significant difference in the mean body mass of zooplankton  
36  
37 701 between heated and unheated treatments (table 1).  
38  
39  
40  
41  
42

43 703 **Figure. 3.** Redundancy analysis (RDA) biplot for sites (i.e. mesocosms) and species  
44  
45 704 scores for phytoplankton taxa recorded in the mesocosm experiment in April (a) and  
46  
47 705 October (b). In both cases RDA 1 was constrained by treatment and accounted for 24.1%  
48  
49 706 and 24.6% of the variation in the taxonomic composition of the mesocosms in April and  
50  
51 707 October respectively. In the plot the dotted lines denote the 95% confidence ellipses  
52  
53 708 around the centroids for both treatments. In both April and October these ellipses do not  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 709 overlap indicating that the community composition was significantly different between  
4  
5 710 warmed and ambient treatments. The solid lines enclose all mesocosms that belong to a  
6  
7  
8 711 particular treatment; in both cases heated treatments (1, 4, 6, 8, 9, 12, 14, 15, 17, 19)  
9  
10 712 cluster to the left, while ambient treatments (2, 3, 5, 7, 10, 11, 13, 16, 18, 20) cluster to  
11  
12 713 the right. Genus abbreviations are as follows: *Aphanothece* (Aph), *Asterococcus* (Ast),  
13  
14 714 *Botryococcus* (Bty), *Bumilleriopsis* (Bum), *C.dinobryonis* (C.d), *Chlorella* (Chl),  
15  
16 715 *Chlorococcum* (Coc), *Chroococcus* (Chr), *Chroomonas* (Cho), *Coencococcus* (Coe),  
17  
18 716 *Cosmarium* (Cos), *Cryptomonas* (Cry), *Goniochloris* (Gon), *Kirchneriella* (Kri),  
19  
20 717 *Monoraphidium* (Mon), *Navicula* (Nav), *Nephrocytium* (Nep), *Rhodomonas* (Rho),  
21  
22 718 *Scenedesmus* (Sce), *Synechococcus* (Syn), *Synechocystis* (Syc), *Spermatozopsis* (Spe).  
23  
24  
25  
26  
27  
28

719

29 720 **Figure. 4.** Mean biomass of the major zooplankton taxonomic groups documented in the  
30  
31 721 mesocosms in (a) April and (b) October. Note that there is very little difference in the  
32  
33 722 biomass contribution of the different zooplankton taxa between treatments suggesting  
34  
35 723 that the zooplankton community composition was unaffected by warming.  
36  
37  
38

724

39  
40  
41 725 **Figure.5.** Effects of warming on mean total planktonic biomass ( $\pm 1$  s.e.m). Data are  
42  
43 726 presented as the averages of the total biomass of either phytoplankton and/or zooplankton  
44  
45 727 across the mesocosms for each treatment (n=20 per treatment for the overall mean; n=10  
46  
47 728 per treatment for each sampling occasion). Total biomass is significantly reduced by  
48  
49 729 warming. This is mainly driven by a reduction in phytoplankton biomass, while there is  
50  
51 730 no significant difference in the biomass of zooplankton in response to warming (table 1).  
52  
53  
54

731

1  
2  
3 732 **Figure. 6.** (a) Relationship between zooplankton and phytoplankton biomass. (b)  
4  
5 733 Relationship between the ratio of zooplankton to phytoplankton biomass (Z:P) and total  
6  
7  
8 734 phytoplankton biomass. (c) The relationship between Z:P and the turnover rate of the  
9  
10 735 phytoplankton communities. Each data point corresponds to either the total zooplankton  
11  
12 736 or phytoplankton biomass or the Z:P in either a heated (red circles) or ambient mesocosm  
13  
14  
15 737 (black circles).  
16  
17  
18 738

19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Review Only

739 **Table 1.** The effect of treatment (heated or ambient) on community-level properties. CSS  
 740 is the community size spectrum and PSS is the phytoplankton size spectrum. ANOVAs  
 741 were used to isolate treatment effects on individual community-level properties. In each  
 742 ANOVA month (either April or October) was added as a factor. For each community-  
 743 level property there was no significant effect of month, which was removed from the  
 744 model using the AIC score.

745

<b>Community Property</b>	<b>DF</b>	<b>F</b>	<b>P</b>
CSS slope	1, 38	11.1	<b>0.002</b>
CSS intercept ( $x = 0$ )	1, 38	8.2	<b>0.007</b>
CSS intercept ( $x = -8$ )	1, 38	4.2	<b>0.047</b>
PSS slope	1, 33	11.8	<b>0.002</b>
PSS intercept	1, 33	8.27	<b>0.007</b>
Total community biomass	1, 38	10.8	<b>0.002</b>
Total phytoplankton biomass	1, 38	13.1	<b>&lt;0.001</b>
Total zooplankton biomass	1, 38	0.47	0.5 (NS)
Mean phytoplankton body mass	1, 38	18.9	<b>&lt;0.001</b>
Mean zooplankton body mass	1, 38	1.4	0.2 (NS)
Z:P Biomass ratio	1, 38	4.82	<b>0.034</b>

746

747

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

748 **Table 2.** Analysis of covariance for the relationships between zooplankton and  
 749 phytoplankton biomass, the Z:P biomass ratio and phytoplankton biomass, and the Z:P  
 750 biomass ratio and phytoplankton turnover time.

Relationship	DF	F	P	r <sup>2</sup>
<b>Log<sub>10</sub>(Zoo biomass) vs Log<sub>10</sub> (Phyto biomass)</b>	1, 38	0.062	0.805 (NS)	0.002
<i>Difference in slope</i>	1, 38	3.021	0.073(NS)	N/A
<i>Difference in intercept</i>	1, 38	0.195	0.661 (NS)	N/A
<b>Log<sub>10</sub>(Z:P) vs Log<sub>10</sub> (Phytoplankton biomass)</b>	1, 38	32.65	<b>&lt;0.0001</b>	0.58
<i>Difference in slope</i>	1, 38	1.806	0.187 (NS)	N/A
<i>Difference in intercept</i>	1, 38	0.002	0.956 (NS)	N/A
<b>Log<sub>10</sub>(Z:P) vs Log<sub>10</sub> (Phytoplankton turnover)</b>	1, 38	52.51	<b>&lt;0.0001</b>	0.58
<i>Difference in slope</i>	1, 38	2.171	0.147 (NS)	N/A
<i>Difference in intercept</i>	1, 38	0.538	0.468 (NS)	N/A

751

For Review Only




Figure. 1. The size spectrum. (a) The community size spectrum of a heated (red circles) and ambient (black circles) mesocosm, highlighting the increase in the steepness of the slope in the warmed mesocosm. (b) Frequency distribution of the slope of the community size spectrum in the ambient mesocosms (n=20), (c) frequency distribution of the slope of the community size spectrum in the warmed mesocosms (n=20). On average the slope of the community size spectrum in the warmed mesocosms was significantly steeper than the ambient mesocosms (Table 1). (d) The phytoplankton size spectrum of a heated (red circles) and ambient (black circles) mesocosm, highlighting the increase in the steepness of the slope and the truncation of large sized individuals in the warmed mesocosm. (e) Frequency distribution of the slope of the phytoplankton size spectrum in the ambient mesocosms (n=17), (f) frequency distribution of the slope of the community size spectrum in the warmed mesocosms (n=18).

198x328mm (72 x 72 DPI)




Figure 3. Redundancy analysis (RDA) biplot for sites (i.e. mesocosms) and species scores for phytoplankton taxa recorded in the mesocosm experiment in April (a) and October (b). In both cases RDA 1 was constrained by treatment and accounted for 24.1% and 24.6% of the variation in the taxonomic composition of the mesocosms in April and October respectively. In the plot the dotted lines denote the 95% confidence ellipses around the centroids for both treatments. In both April and October these ellipses do not overlap indicating that the community composition was significantly different between warmed and ambient treatments. The solid lines enclose all mesocosms that belong to a particular treatment; in both cases heated treatments (1, 4, 6, 8, 9, 12, 14, 15, 17, 19) cluster to the left, while ambient treatments (2, 3, 5, 7, 10, 11, 13, 16, 18, 20) cluster to the right. Genus abbreviations are as follows: Aphanothece (Aph), Asterococcus (Ast), Botryococcus (Bty), Bumilleriopsis (Bum), C.dinobryonis (C.d), Chlorella (Chl), Chlorococcum (Coc), Chroococcus (Chr), Chroomonas (Cho), Coencococcus (Coe), Cosmarium (Cos), Cryptomonas (Cry), Goniochloris (Gon), Kirchneriella (Kri), Monoraphidium (Mon), Navicula (Nav), Nephrocystium (Nep), Rhodomonas (Rho), Scenedesmus (Sce), Synechococcus (Syn), Synechocystis (Syc), Spermatozopsis (Spe).

391x198mm (72 x 72 DPI)






Figure 4. Mean biomass of the major zooplankton taxonomic groups documented in the mesocosms in (a) April and (b) October. Note that there is very little difference in the biomass contribution of the different zooplankton taxa between treatments suggesting that the zooplankton community composition was unaffected by warming.  
 119x220mm (72 x 72 DPI)




Figure.5. Effects of warming on mean total planktonic biomass ( $\pm 1$  s.e.m). Data are presented as the averages of the total biomass of either phytoplankton and/or zooplankton across the mesocosms for each treatment ( $n=20$  per treatment for the overall mean;  $n=10$  per treatment for each sampling occasion). Total biomass is significantly reduced by warming. This is mainly driven by a reduction in phytoplankton biomass, while there is no significant difference in the biomass of zooplankton in response to warming (table 1).

144x252mm (72 x 72 DPI)




Figure 6. (a) Relationship between zooplankton and phytoplankton biomass. (b) Relationship between the ratio of zooplankton to phytoplankton biomass (Z:P) and total phytoplankton biomass. (c) The relationship between Z:P and the turnover rate of the phytoplankton communities. Each data point corresponds to either the total zooplankton or phytoplankton biomass or the Z:P in either a heated (red circles) or ambient mesocosm (black circles).  
180x328mm (72 x 72 DPI)

1  
2  
3 1 **Warming Alters the Size Spectrum and Shifts the Distribution of Biomass in**  
4  
5  
6 2 **Aquatic Ecosystems**  
7  
8  
9 3

10  
11 4 **SUPPLEMENTARY MATERIAL**  
12  
13  
14 5

15  
16 6 **Gabriel Yvon-Durocher<sup>1</sup>, José M. Montoya<sup>1,2</sup>, Mark Trimmer<sup>1</sup>, and Guy Woodward<sup>1</sup>**  
17  
18 7


19 8 <sup>1</sup> *School of Biological & Chemical Sciences, Queen Mary University of London, London E1 4NS.*  
20  
21 9 *U.K.*  
22

23 10 <sup>2</sup> *Institute of Marine Sciences (ICM-CSIC) Pg. Marítim de la Barceloneta, 37-49*  
24  
25 11 *E-08003 Barcelona, Spain*  
26  
27 12

28  
29 13 **Authors for correspondence:** Gabriel Yvon-Durocher (g.yvon-durocher@qmul.ac.uk) and Guy  
30  
31 14 Woodward (g.woodward@qmul.ac.uk)  
32  
33 15 School of Biological & Chemical Sciences,  
34  
35 16 Queen Mary University of London,  
36  
37 17 London E1 4NS. U.K.  
38  
39 18 TEL:+44 20 7882 3640  
40  
41 19


For Review Only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




20  
21  
22 **Figure S1.** Frequency distributions of individual body mass for (a) all individuals measured, (b)  
23 a random sample of 400 (i.e. the number of individuals actually measured in a sample) from a,  
24 (c) a random sample of 2000 from a, (d) a random sample of 100 from a. Data highlight that a  
25 sample of 400 individuals is sufficient to estimate the variance in the distribution of body size  
26 comparable to the whole community. When measuring the phytoplankton a minimum of 400  
27 individuals from any given pond were measured over the number of fields of view required to  
28 count 400 from the sample in the sedimentation chamber. It is also clear that a sample of 100 is  
29 not sufficient to accurately reproduce the variance in the body mass distribution of the whole  
30 community. Assuming that organisms of a given body mass are Poisson distributed (figure S2,  
31 table S3) on the surface of the sedimentation chamber, the measurement of 400 individuals  
32 should be sufficient to attain an error of 5% [if error =  $1/\sqrt{n}$ ].

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Figure S2.** Size-frequency distribution for phytoplankton in pond 14 from April 2007. Panels show the size-frequency distribution after analysing all fields of view (FOV) taken to measure ~400 individuals in the sedimentation chamber, 1 FOV, 2 FOVs, 3 FOVs and 4 FOVs. Data highlight the equitable distribution of body size among fields of view which reflects the random settlement of phytoplankton cells in the sedimentation chamber. Tests for dispersion were carried for all samples and settlement conformed to Poisson statistics in every case (data not shown).



43  
44 **Figure S3.** Seasonality of inorganic nutrients in the warmed (red lines) and ambient (black lines)  
45 mesocosms. (a) Nitrite, (b) Nitrate, (c) Ammonium, (d) Silicate, (e) Phosphate, (f) the  
46 stoichiometry of the inorganic nutrient pool, N:P. Water samples for measuring dissolved  
47 inorganic nutrient concentrations were collected from mid depth in the mesocosm at 9am on each  
48 sampling occasion. Samples were filtered (Whatmann GF/F) and stored frozen ( $-20^\circ\text{C}$ ) for  
49 subsequent determination of  $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ,  $\text{NH}_4^+$ ,  $\text{PO}_4^{3-}$  and dissolved silica (Si) using a segmented  
50 flow auto-analyser (Skalar, San++, Breda, Netherlands), according to (Kirkwood 1996).  
51 Inorganic nutrients ( $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ,  $\text{NH}_4^+$ ,  $\text{PO}_4^{3-}$  & Si) exhibited strong seasonal trends. For  
52 example,  $\text{NO}_3^-$  concentrations peaked in spring and declined progressively throughout the  
53 summer, when rates of primary production were maximal (Yvon-Durocher *et al.* 2010), and were  
54 depleted to  $\approx 0.005 \mu\text{mol l}^{-1}$  by October, before regeneration in the winter. Concentrations of  
55  $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ,  $\text{NH}_4^+$  and  $\text{PO}_4^{3-}$  showed identical seasonal patterns in the warmed and ambient

1  
2  
3 56 treatments, with no significant differences in the overall mean annual concentrations of these  
4  
5  
6 57 nutrients (table S4). Furthermore, the stoichiometry of the inorganic nutrient pool exhibited  
7  
8 58 remarkable similarity between treatments, with a mean annual ratio of total inorganic N to P of  
9  
10 59  $\approx 11:1$  in both heated and ambient mesocosms.  
11  
12  
13 60

14  
15 61 **Table S4.** Results of the linear mixed effects model testing for differences in the concentration of  
16  
17 62 inorganic nutrients between heated and ambient mesocosms. A linear mixed effects model was  
18  
19 63 conducted with restricted maximum likelihood methods using the *lme* (linear mixed-effects  
20  
21 64 model) function in R, treatment (heated or unheated) was the fixed effect, and temporal pseudo-  
22  
23 65 replication from repeated sampling of the mesocosms over the year was accounted for by  
24  
25 66 including mesocosm identity nested with sampling occasion as random effects.  
26  
27  
28  
29

Inorganic Nutrient	DF	F	P
NO <sub>2</sub> <sup>-</sup>	1, 120	0.06	0.812 (NS)
NO <sub>3</sub> <sup>-</sup>	1, 120	0.65	0.420 (NS)
NH <sub>4</sub> <sup>+</sup>	1, 120	0.23	0.632 (NS)
Si	1, 120	6.08	<b>0.015</b>
PO <sub>4</sub> <sup>3-</sup>	1, 120	0.68	0.412 (NS)
Total inorganic N to P	1, 120	0.009	0.922 (NS)

30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40 67  
41 68  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



69 **Table S5.** Regression statistics for the community size spectrum of each mesocosm for the  
 70 relationship:  $\log(N_i) = b * \log(M_i) + a$ . Where  $N_i$  is the abundance of the size class  $i$  and is the  
 71 mass at the centre of the  $i^{\text{th}}$  size bin,  $b$  and  $a$  are the slope and the intercept respectively. These  
 72 data highlight that the size spectrum was linear for each of the mesocosms and that the individual  
 73 size distribution was a power law.


Pond	Treatment	Month	Slope	Intercept	$r^2$	$P$ -value
1	Heated	April	-0.92	4.64	0.91	0.00020
2	Ambient	April	-0.94	4.58	0.81	0.00040
3	Ambient	April	-0.93	4.80	0.86	0.00030
4	Heated	April	-0.93	4.50	0.90	0.00003
5	Ambient	April	-0.79	5.28	0.84	0.00020
6	Heated	April	-1.12	3.30	0.78	0.00060
7	Ambient	April	-0.83	5.09	0.84	0.00020
8	Heated	April	-0.90	4.39	0.80	0.00040
9	Heated	April	-1.03	3.94	0.85	0.00040
10	Ambient	April	-0.92	3.89	0.80	0.00100
11	Ambient	April	-0.86	4.71	0.78	0.00200
12	Heated	April	-0.90	4.52	0.97	0.00004
13	Ambient	April	-0.98	4.21	0.74	0.00100
14	Heated	April	-0.88	4.25	0.91	0.00080
15	Heated	April	-0.94	4.64	0.80	0.00100
16	Ambient	April	-0.91	4.90	0.93	0.00001
17	Heated	April	-1.05	4.05	0.88	0.00050
18	Ambient	April	-0.71	5.58	0.81	0.00030
19	Heated	April	-0.92	5.09	0.97	0.00000
20	Ambient	April	-0.75	5.49	0.90	0.00010
1	Heated	October	-0.87	4.58	0.94	0.00001
2	Ambient	October	-0.94	4.15	0.95	0.00001
3	Ambient	October	-0.72	5.46	0.70	0.00200
4	Heated	October	-0.94	4.02	0.90	0.00001
5	Ambient	October	-0.94	4.10	0.81	0.00040
6	Heated	October	-1.06	3.68	0.85	0.00100
7	Ambient	October	-0.97	4.12	0.83	0.00020
8	Heated	October	-0.92	4.50	0.89	0.00040
9	Heated	October	-0.90	4.50	0.84	0.00040
10	Ambient	October	-0.84	5.09	0.87	0.00070
11	Ambient	October	-0.80	5.27	0.83	0.00020
12	Heated	October	-0.93	4.42	0.88	0.00006

13	Ambient	October	-0.78	5.30	0.88	0.00010
14	Heated	October	-0.94	3.75	0.82	0.00080
15	Heated	October	-0.95	4.17	0.93	0.00010
16	Ambient	October	-0.79	5.00	0.87	0.00020
17	Heated	October	-1.01	4.25	0.93	0.00010
18	Ambient	October	-0.95	3.77	0.93	0.00009
19	Heated	October	-0.85	4.70	0.89	0.00010
20	Ambient	October	-0.94	4.24	0.92	0.00004

**Table S6.** Regression statistics for the phytoplankton size spectrum of each mesocosm for the relationship:  $\log(N_i) = b * \log(M_i) + a$ . Where  $N_i$  is the abundance of the size class  $i$  and is the mass at the centre of the  $i^{\text{th}}$  size bin,  $b$  and  $a$  are the slope and the intercept respectively.

Pond	Treatment	Month	Slope	Intercept	$r^2$	$P$ -Value
1	Heated	April	-0.41	7.50	0.42	0.040000
2	Ambient	April	-0.31	8.20	0.70	0.005000
4	Heated	April	-0.50	6.89	0.90	0.000092
6	Heated	April	-0.35	7.70	0.71	0.002200
7	Ambient	April	-0.22	8.39	0.57	0.012000
8	Heated	April	-0.27	8.03	0.55	0.014000
9	Heated	April	-0.55	6.78	0.88	0.000160
10	Ambient	April	-0.47	6.72	0.91	0.000020
11	Ambient	April	-0.30	7.82	0.53	0.018000
12	Heated	April	-0.76	5.08	0.62	0.011000
13	Ambient	April	-0.34	7.77	0.56	0.013000
14	Heated	April	-0.65	5.52	0.70	0.004600
15	Heated	April	-0.57	6.86	0.87	0.000200
16	Ambient	April	-0.45	7.28	0.75	0.001200
17	Heated	April	-0.52	7.20	0.67	0.004000
18	Ambient	April	-0.27	7.90	0.48	0.030000
19	Heated	April	-0.65	6.36	0.93	0.000006
20	Ambient	April	-0.34	7.60	0.77	0.000900
2	Ambient	October	-0.48	6.65	0.82	0.000300
4	Heated	October	-0.50	6.41	0.78	0.000680
5	Ambient	October	-0.34	7.70	0.85	0.000300
7	Ambient	October	-0.34	7.70	0.62	0.007000
8	Heated	October	-0.30	8.14	0.68	0.004000
9	Heated	October	-0.42	7.16	0.94	0.000005
10	Ambient	October	-0.43	7.50	0.79	0.001400

11	Ambient	October	-0.32	7.90	0.77	0.000880
12	Heated	October	-0.45	7.09	0.87	0.000240
13	Ambient	October	-0.40	7.43	0.90	0.000099
14	Heated	October	-0.35	7.41	0.58	0.010000
15	Heated	October	-0.57	6.25	0.80	0.000400
16	Ambient	October	-0.25	8.20	0.60	0.023000
17	Heated	October	-0.60	6.70	0.91	0.000200
18	Ambient	October	-0.46	6.70	0.82	0.000300
19	Heated	October	-0.36	7.49	0.69	0.003000
20	Ambient	October	-0.38	7.38	0.86	0.000300




**Figure S7.** Quotient of benthic to ecosystem metabolism. On average over the course of the year benthic metabolism represented ~35% of whole ecosystem metabolism measured using the dissolved oxygen change technique (see Yvon-Durocher *et al.*, (2010) for details). Benthic metabolism was measured using dark in-situ benthic chambers which enclosed a sample of 500 mL at the sediment-water interface. A magnetic stirrer in the chamber ensured that the sample was evenly mixed. Benthic respiration was measured by the removal of 25mL samples at the beginning and the end of the 6 hour incubations. The samples were gently discharged into gas-

1  
2  
3 91 tight vials (12ml, Exetainers, Labco Ltd, High Wycombe, UK) and allowed to overflow twice (to  
4  
5 92 minimize atmospheric gas exchange), and fixed for Winkler analysis. The samples were  
6  
7  
8 93 immediately fixed and stored in a fridge at 5 °C to minimize light and temperature fluctuations  
9  
10 94 until they could be titrated in the laboratory (< 5 d). To ensure linearity of oxygen uptake a timed  
11  
12 95 series of samples were taken, subsequently only T = 0 and T = final samples were taken to limit  
13  
14 96 sample extraction from the chambers.  
15  
16  
17  
18  
19 97

## 20 98 **References**

- 21  
22 99 Kirkwood D.S. (1996). *Nutrients: practical notes on their determination in seawater*. ICES,  
23  
24  
25 100 Copenhagen.  
26  
27 101 Reynolds C.S. (1984). *The Ecology of Freshwater Phytoplankton* Cambridge University Press  
28  
29 102 Cambridge.  
30  
31 103 Yvon-Durocher G., Jones J.I., Woodward G., Trimmer M. & Montoya J.M. (2010). Warming  
32  
33 104 alters the metabolic balance of ecosystems. *Philosophical Transactions of the Royal*  
34  
35 105 *Society of London Series B-Biological Sciences*, 365, 2117-2126.  
36  
37  
38  
39 106  
40  
41 107  
42  
43  
44 108  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



114x225mm (72 x 72 DPI)