



HAL
open science

A different way of looking at depression

Evelyn Van Weel-Baumgarten, Peter Lucassen, Lieke Hassink-Franke, Henk Schers

► **To cite this version:**

Evelyn Van Weel-Baumgarten, Peter Lucassen, Lieke Hassink-Franke, Henk Schers. A different way of looking at depression. *International Journal of Clinical Practice*, 2010, 64 (11), pp.1493. 10.1111/j.1742-1241.2010.02405.x . hal-00599538

HAL Id: hal-00599538

<https://hal.science/hal-00599538>

Submitted on 10 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



A different way of looking at depression

Journal:	<i>International Journal of Clinical Practice</i>
Manuscript ID:	IJCP-01-10-0042.R1
Manuscript Type:	Perspective
Date Submitted by the Author:	29-Jan-2010
Complete List of Authors:	Weel-Baumgarten, Evelyn; Radboud University Nijmegen Medical Centre, Department of Primary and Community Care Lucassen, Peter; Radboud University Nijmegen Medical Centre, Department of Primary and Community Care Hassink-Franke, Lieke; Radboud University Nijmegen Medical Centre, Department of Primary and Community Care Schers, Henk; Radboud University Nijmegen Medical Centre, Department of Primary and Community Care
Specialty area:	



A different way of looking at depression

E. van Weel-Baumgarten, P Lucassen, L. Hassink-Franke and H Schers.

The value of the diagnosis of depression in primary care has been under debate for many years now, even though the focus of this discussion has changed from time to time, shifting from under to over-diagnosis. In this contribution we suggest a different way of looking at depression in primary care, and we will substantiate this opinion with arguments why it might be more appropriate using a stepped model for diagnosis as well as for treatment.

Why do we need a different way of looking at depression?

In a recent systematic review Mitchell et al. concluded that general practitioners over-diagnose depression more than they miss cases: for every 100 unselected cases seen in primary care, there are more false positives (n=15) than either missed (n=10) or identified cases (n=10); general practitioners (GP's) diagnose depression too frequently and not too in-frequently(1). This is a relatively new sound in scientific literature. For a long time the importance of a diagnosis and therefore the need to screen has been stressed. The focus was on under-diagnosis of depression in primary care, supposing that detection will lead to a better outcome. But evidence showed that, unfortunately, this is not always the case in primary care(2). Both conclusions, that of over-diagnosis as well as under-diagnosis, suggest that there is something seriously wrong with the diagnosis of depression in primary care settings. This is important because only a correct diagnosis can be followed by adequate and effective treatment as set out in various guidelines(3;4). A further problem is that results of treatment for depression in primary care are disappointing. Antidepressants have a strong placebo effect and in general practice only one in about every 5 or 6 patients benefit specifically from medication(5). This result is often attributed to inadequate treatment by general practitioners or to a low compliance by patients for instance because of side effects of medication. But over-diagnosis and over-treatment might also explain the disappointing effectiveness of antidepressants in primary care, using these medications in patients who do not need them. Furthermore the effectiveness of psychotherapy is also is not very large as it is roughly equal to the effectiveness of medication.

Field Code Changed

Field Code Changed

Field Code Changed

Deleted: About

Deleted:

Deleted: T

Different views on depression

We would like to discuss the diagnosis of depression in primary care from another perspective, believing that the main problem is not compliance with diagnostic criteria, but having a different view on the reality of patients consulting their GP with depressed feelings. General practitioners' guidelines advocate using strict criteria for the diagnosis of depression. These criteria are provided by the DSM-IV classification system. Since the overall acceptance of this classification, all depression is looked upon as one and the same. According to the DSM, the diagnosis is symptom-based without reference to a cause. The classification is said to be value-neutral. However, it denies the fact that the concept of

1
2 depression is relative; it is a diagnosis by agreement and not by essence. It has been argued 'that
3 depression has reached a status which goes far beyond its true status and that the concept has been
4 widened gradually to a concept containing many heterogeneous categories'(6). This might also be one
5 of the reasons why general practitioners currently prescribe antidepressants too often. Due to this
6 model many patients receive antidepressants even when consulting with depressive symptoms for the
7 first time(7). Even though antidepressants are sometimes combined with psychotherapy, the question
8 remains how many of these patients really need these depression specific medications and treatments.
9 Perhaps prescribing of antidepressants to too many patients is also causing a dilution of their
10 effectiveness, which is much lower in primary care where the whole spectrum of depressive symptoms
11 is seen, than in secondary care; with on average more severe and more chronic cases in which
12 antidepressants are really needed(8:9).

Deleted: (5)

Field Code Changed

Deleted: (6)

Field Code Changed

Field Code Changed

Deleted: (7:8)

13 The value-neutral, symptom-based DSM-IV approach leaves out the existence of situations where
14 patients experience intense sadness (and other symptoms of the list of items for depression) as a
15 completely adequate response to what has happened in their life. Many of these patients fulfil the
16 criteria for major depression. However, the intense sadness has a function and as such distinguishes
17 between normality and disorder(10). We think that treating these patients with antidepressants or
18 psychological treatments might be one of the causes of the low effectiveness of treatments for
19 depression in primary care as a whole, medicalising too many cases of human suffering. Many of
20 these people might be more in need of talking through what has happened, thus experiencing support
21 and stimulating the restoration of hope. We think that such approaches and generic interventions are
22 often already used in primary care.

Deleted: (9)

Field Code Changed

31 A new approach

32 Underpinning the last statement and the relativity of the classification system, we return to the
33 systematic review by Mitchell et al. They mention many possible reasons for their findings of more
34 false positive depression diagnoses than true positives. They point out the strikingly high specificity
35 and sensitivity of the diagnosis in a study that we performed in the practices of the Continuous
36 Morbidity Registration (CMR)(11). This is the registry of the department of Primary and Community
37 Care of the Radboud University Nijmegen Medical Centre. It is a network of four practices in the
38 Nijmegen region that records all morbidity on an ongoing basis since 1971 with a total practice list of
39 about 12.000 patients. The authors contributed this high specificity and sensitivity of the diagnosis to
40 the training these GPs received in using diagnostic criteria of the ICPC(12). Of course training in
41 diagnostic criteria does contribute to a correct diagnosis, but in this case there was no specific training
42 for the diagnosis of depression, no training in DSM diagnosis, nor had the training been given
43 recently. The real reason for the adequate diagnoses might be that in these practices most patients have
44 been registered in the same practice for many years. They receive a person-centred longitudinal

Field Code Changed

Deleted: (10)

Field Code Changed

Deleted: (11)

1 approach in which the GP knows most patients well (as is often the case in The Netherlands), and
2 takes into account symptoms as well as context.

3 Having mentioned all these arguments we propose a stepped care model, not just for treatment but also
4 for diagnosis(13). We believe this will decrease the number of over-diagnoses of depression, make it
5 easier to distinguish between 'normal' human suffering and the illness of depression, and act
6 accordingly. The main arguments for our proposal are the insufficient distinction between depression
7 and 'normal' sadness and the disappointing effectiveness of treatments for depression in primary care.

8 With reference to the diagnosis it means that symptoms, when first presented, should be seen in the
9 light of what is going on in that person's life, taking into account context matters as well as symptoms.

10 We should try to avoid labelling symptoms to illness immediately, giving the patient's own
11 explanation and story an important role. Checking symptoms may be helpful, and many patients even
12 like discussing their symptoms in a structured, detailed way using checklists. It is probably good to
13 discuss symptoms in detail as long as patients have the opportunity to tell 'the story' as well as the
14 symptoms(14;15). Using checklists also helps to choose between watchful waiting and active
15 treatment as it sheds light on the symptoms and their severity, as much as the story tells about
16 circumstances and possible causes. Watchful waiting is safe as in primary care many cases of
17 depression are mild and patients might recover without medical interventions, looking at the 'natural'
18 history of depression in primary care(16). Watchful waiting is not the same as doing nothing. We think
19 that it is often justified to follow-up and see if the symptoms improve with a few supportive
20 consultations aiming at restoration of hope, or with a generic intervention or a specific treatment for
21 relevant co-morbidity which is likely to contribute to depressive feelings. Patients for whom the
22 depressive symptoms do not improve as a consequence of our activities will eventually be subject to a
23 formal diagnostic procedure, and if necessary receive depression specific treatment. In this person
24 centred stepped approach it is appropriate to wait a little longer before making a diagnosis. It serves to
25 see if the symptoms are a 'normal' response to circumstances the patient experiences, and if this
26 develops in a positive direction or whether an intervention is needed.

27
28
29
30
31
32
33
34
35
36
37
38
39

Conclusion

40 We do not deny the value of using classification systems for patients, but we think that in primary care
41 it might be suitable to use it as a second step believing that most patients with 'normal sadness' are
42 better off without an immediate diagnosis.

43 Having made this case, we believe that this new approach deserves a try-out. We would then like to
44 study what GP's discuss with their patients, how they diagnose and treat depressive symptoms over
45 several consultations, and the influence of this approach on prognosis and well-being, rather than
46 trying to mould stories from patients into the categories of the DSM classification immediately in first
47 consultations. It will also be interesting to see if a new classification system for mental health in
48
49
50

Field Code Changed

Deleted: (12)

Deleted: (13;14)

Field Code Changed

Field Code Changed

Deleted: (15)

Deleted: a

Formatted: Font: (Default) Times
New Roman, 11 pt, English (U.S.)

primary care, as suggested by Gask et al. will help in distinguishing between distress and disorder, and lead to more appropriate subsequent interventions(17).

- Formatted: Font: (Default) Times New Roman, 11 pt
- Formatted: Font: (Default) Times New Roman, 11 pt, English (U.S.)
- Formatted: Font: (Default) Times New Roman, 11 pt, English (U.S.)
- Formatted: Font: (Default) Times New Roman, 11 pt

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Reference List

- (1) Mitchell AJ, Vaze A, Rao S. Clinical diagnosis of depression in primary care: a meta-analysis. Lancet 2009 August 22;374(9690):609-19.
- (2) Dowrick C, Buchan I. Twelve month outcome of depression in general practice: does detection or disclosure make a difference? [see comments]. BMJ 1995 November 11;311(7015):1274-6.
- (3) NHG. Guideline on depression. Huisarts en Wetenschap 1994;11:482-90.
- (4) Van Marwijk HWJ et al. NHG- Standaard Depressieve stoornis (depressie). (eerste herziening). Huisarts en Wetenschap 2003;46 (11):614-23.
- (5) Walsh BT, Seidman SN, Sysko R, Gould M. Placebo response in studies of major depression: variable, substantial, and growing. JAMA 2002 April 10;287(14):1840-7.
- (6) Parker G. Is depression overdiagnosed? Yes. BMJ 2007 August 18;335(7615):328.
- (7) Volkers A, De Jong A, de Bakker D, van Dijk L. Doelmatig voorschrijven van antidepressiva in de huisartspraktijk. Utrecht: Nivel; 2005.
- (8) Fournier JC, DeRubeis RJ, Hollon SD, Dimidjian S, Amsterdam JD, Shelton RC, Fawcett J. Antidepressant drug effects and depression severity: a patient-level meta-analysis. JAMA 2010 January 6;303(1):47-53.
- (9) Arroll B, Macgillivray S, Ogston S, Reid I, Sullivan F, Williams B, Crombie I. Efficacy and tolerability of tricyclic antidepressants and SSRIs compared with placebo for treatment of depression in primary care: a meta-analysis. Ann Fam Med 2005 September;3(5):449-56.
- (10) Horwitz AV, Wakefield JC. The loss of sadness. Oxford: Oxford University Press; 2007.
- (11) Weel-Baumgarten EM, van den Bosch WJ, van den Hoogen HJ, Zitman FG. The validity of the diagnosis of depression in general practice: is using criteria for diagnosis as a routine the answer? Br J Gen Pract 2000 April;50(453):284-7.
- (12) Van Weel C. Validating long term morbidity recording. J Epidemiol Community Health 1995 August;49 Suppl 1:29-32.
- (13) Lucassen P vREvW-BEDC. Making less depression diagnoses: beneficial for patients? Mental Health in Family Medicine 2008;5:161-5.
- (14) Dowrick C, Leydon GM, McBride A, Howe A, Burgess H, Clarke P, Maisey S, Kendrick T. Patients' and doctors' views on depression severity questionnaires incentivised in UK quality and outcomes framework: qualitative study. BMJ 2009;338:b663.
- (15) Kendrick T, Dowrick C, McBride A, Howe A, Clarke P, Maisey S, Moore M, Smith PW. Management of depression in UK general practice in relation to scores on depression severity questionnaires: analysis of medical record data. BMJ 2009;338:b750.

Formatted: Left

Formatted: Space After: 12 pt

Formatted: Dutch (Netherlands)

Formatted: Dutch (Netherlands)

Formatted: Dutch (Netherlands)

(16) Weel-Baumgarten EM, Schers HJ, van den Bosch WJ, van den Hoogen HJ, Zitman FG. Long-term follow-up of depression among patients in the community and in family practice settings - A systematic review. J Fam Pract 2000 December;49(12):1113-20.

(17) Gask L, Klinkman M, Fortes S, Dowrick C. Capturing complexity: the case for a new classification system for mental disorders in primary care. Eur Psychiatry 2008 October;23(7):469-76.

Deleted: Reference List¶

- ¶
- . (1) . Mitchell AJ, Vaze A, Rao S. Clinical diagnosis of depression in primary care: a meta-analysis. *Lancet* 2009 August 22;374(9690):609-19.¶
- . (2) . Dowrick C, Buchan I. Twelve month outcome of depression in general practice: does detection or disclosure make a difference? [see comments]. *BMJ* 1995 November 11;311(7015):1274-6.¶
- . (3) . NHG. Guideline on depression. *Huisarts en Wetenschap* 1994;11:482-90.¶
- . (4) . Van Marwijk HWJ et al. NHG-Standaard Depressieve stoornis (depressie). (eerste herziening). *Huisarts en Wetenschap* 2003;46 (11):614-23.¶
- . (5) . Parker G. Is depression overdiagnosed? Yes. *BMJ* 2007 August 18;335(7615):328.¶
- . (6) . Volkers A, De Jong A, de Bakker D, van Dijk L. Doelmatig voorschrijven van antidepressiva in de huisartspraktijk. *Utrecht: Nivel*; 2005. ¶
- . (7) . Fournier JC, DeRubeis RJ, Hollon SD, Dimidjian S, Amsterdam JD, Shelton RC, Fawcett J. Antidepressant drug effects and depression severity: a patient-level meta-analysis. *JAMA* 2010 January 6;303(1):47-53.¶
- . (8) . Arroll B, Macgillivray S, Ogston S, Reid I, Sullivan F, Williams B, Crombie I. Efficacy and tolerability of tricyclic antidepressants and SSRIs compared with placebo for treatment of depression in primary care: a meta-analysis. *Ann Fam Med* 2005 September;3(5):449-56.¶
- . (9) . Horwitz AV, Wakefield JC. *The loss of sadness*. Oxford: Oxford University Press; 2007.¶
- . (10) . Weel-Baumgarten EM, van den Bosch WJ, van den Hoogen HJ, Zitman FG. The validity of the diagnosis of depression in general practice: is using criteria for diagnosis as a routine the answer? *Br J Gen Pract* 2000 April;50(453):284-7.¶
- . (11) . Van Weel C. Validating long term morbidity recording. *J Epidemiol Community Health* 1995 August;49 Suppl 1:29-32.¶
- . (12) . Lucassen P vREvW-BEDC. Making less depression diagnoses: beneficial for patients? *Mental Health in Family Medicine* 2008;5:161-5.¶
- . (13) . Dowrick C, Leydon GM, McBride A, Howe A, Burgess H, Clarke P, Maisey S, Kendrick T. Patients' and doctors' views on depression severity questionnaires incentivised in UK quality and outcomes framework: qualitative study. *BMJ* 2009;338:b663.¶
- . (14) . Kendrick T, Dowrick C, McBride A, Howe A, Clarke P, Maisey S, Moore M, Smith PW. Management of depression in UK general practice in relation to scores on depression severity questionnaires: analysis of medical record data. *BMJ* 2009;338:b750.¶
- . (15) . Weel-Baumgarten EM, Schers HJ, van den Bosch WJ, van den Hoogen HJ, Zitman FG. Long-term follow-up... [1]

Formatted: Indent: Left: 0 pt,
Hanging: 36 pt

Reference List

- (1) Mitchell AJ, Vaze A, Rao S. Clinical diagnosis of depression in primary care: a meta-analysis. *Lancet* 2009 August 22;374(9690):609-19.
- (2) Dowrick C, Buchan I. Twelve month outcome of depression in general practice: does detection or disclosure make a difference? [see comments]. *BMJ* 1995 November 11;311(7015):1274-6.
- (3) NHG. Guideline on depression. *Huisarts en Wetenschap* 1994;11:482-90.
- (4) Van Marwijk HWJ et al. NHG- Standaard Depressieve stoornis (depressie). (eerste herziening). *Huisarts en Wetenschap* 2003;46 (11):614-23.
- (5) Parker G. Is depression overdiagnosed? Yes. *BMJ* 2007 August 18;335(7615):328.
- (6) Volkers A, De Jong A, de Bakker D, van Dijk L. Doelmatig voorschrijven van antidepressiva in de huisartspraktijk. Utrecht: Nivel; 2005.
- (7) Fournier JC, DeRubeis RJ, Hollon SD, Dimidjian S, Amsterdam JD, Shelton RC, Fawcett J. Antidepressant drug effects and depression severity: a patient-level meta-analysis. *JAMA* 2010 January 6;303(1):47-53.
- (8) Arroll B, Macgillivray S, Ogston S, Reid I, Sullivan F, Williams B, Crombie I. Efficacy and tolerability of tricyclic antidepressants and SSRIs compared with placebo for treatment of depression in primary care: a meta-analysis. *Ann Fam Med* 2005 September;3(5):449-56.
- (9) Horwitz AV, Wakefield JC. *The loss of sadness*. Oxford: Oxford University Press; 2007.
- (10) Weel-Baumgarten EM, van den Bosch WJ, van den Hoogen HJ, Zitman FG. The validity of the diagnosis of depression in general practice: is using criteria for diagnosis as a routine the answer? *Br J Gen Pract* 2000 April;50(453):284-7.
- (11) Van Weel C. Validating long term morbidity recording. *J Epidemiol Community Health* 1995 August;49 Suppl 1:29-32.
- (12) Lucassen P vREvW-BEDC. Making less depression diagnoses: beneficial for patients? *Mental Health in Family Medicine* 2008;5:161-5.
- (13) Dowrick C, Leydon GM, McBride A, Howe A, Burgess H, Clarke P, Maisey S, Kendrick T. Patients' and doctors' views on depression severity questionnaires incentivised in UK quality and outcomes framework: qualitative study. *BMJ* 2009;338:b663.
- (14) Kendrick T, Dowrick C, McBride A, Howe A, Clarke P, Maisey S, Moore M, Smith PW. Management of depression in UK general practice in relation to scores on depression severity questionnaires: analysis of medical record data. *BMJ* 2009;338:b750.
- (15) Weel-Baumgarten EM, Schers HJ, van den Bosch WJ, van den Hoogen HJ, Zitman FG. Long-term follow-up of depression among patients in the community and in family practice settings - A systematic review. *J Fam Pract* 2000 December;49(12):1113-20.