



**HAL**  
open science

## Low prevalence of hepatitis E infection among pregnant women in Madrid, Spain

Maria Luisa Mateos, Gustavo Gabilondo, Beatriz Romero, Oihane S. de La Maza, Teresa Perez-Gracia

► **To cite this version:**

Maria Luisa Mateos, Gustavo Gabilondo, Beatriz Romero, Oihane S. de La Maza, Teresa Perez-Gracia. Low prevalence of hepatitis E infection among pregnant women in Madrid, Spain. *Journal of Medical Virology*, 2010, 82 (10), pp.1666. 10.1002/jmv.21840 . hal-00599782

**HAL Id: hal-00599782**

**<https://hal.science/hal-00599782>**

Submitted on 11 Jun 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Low prevalence of hepatitis E infection among pregnant women in Madrid, Spain

Journal:	<i>Journal of Medical Virology</i>
Manuscript ID:	JMV-10-1762.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	31-Mar-2010
Complete List of Authors:	Mateos, Maria; Hospital Ramon y Cajal, Microbiología Gabilondo, Gustavo; Hospital Ramon y Cajal, Microbiología Romero, Beatriz; Hospital Ramon y Cajal, Microbiología S. de la Maza, Oihane; Hospital Ramon y Cajal, Microbiología Perez-Gracia, teresa; Facultad de Ciencias de la Salud. Universidad CEU Cardenal Herrera, Microbiología
Keywords:	hepatitis E, pregnant women, viral hepatitis, imported diseases



## Low prevalence of hepatitis E infection among pregnant women in Madrid, Spain

Maria Luisa Mateos Lindemann<sup>1</sup>, Gustavo Gabilondo<sup>1</sup>, Beatriz Romero<sup>1</sup>, Oihane Martín S. de la Maza<sup>1</sup>, Maria Teresa Pérez-Gracia<sup>2</sup>

1. Servicio de Microbiología y Parasitología. Hospital Ramón y Cajal. Ctra. Colmenar Km 9.1 Madrid 28029

2. Dep. Microbiología. Facultad de Ciencias de la Salud. Universidad CEU Cardenal Herrera. Moncada, Valencia.

Shortened title: HEV infection in pregnant women. Subclinical infection in Spain

Correspondence to: Maria Luisa Mateos, Servicio de Microbiología y Parasitología. Hospital Ramón y Cajal. Ctra. Colmenar Km 9.1 Madrid 28029. Fax 34 91 3368809  
E-mail: mmateos.hrc@salud.madrid.org

Key words: hepatitis E, HEV, pregnant women, imported diseases, viral hepatitis

### SUMMARY:

Hepatitis E virus, a cause of sporadic hepatitis in developed countries is present in Spain. The aims of this study were to determine whether the prevalence of IgG anti-hepatitis E virus antibodies has increased in recent years in Spain and the prevalence in a study population at risk of developing the disease following the infection. Serum samples from 1040 women in the first trimester of pregnancy were studied. The prevalence rate of 3.6% has not increased significantly since 1999 in spite of the large number of immigrants from endemic regions. IgM anti-HEV antibodies were detected in 0.67% of the patients suggesting that although subclinical infection exists the prevalence is very low. The reasons for the differences in the morbidity of HEV in pregnant women in different geographical areas remain unclear. Good hygienic measures and public health conditions are important factors for ensuring low prevalence of hepatitis E in Spain.

### INTRODUCTION

Hepatitis E virus (HEV) is a major cause of acute viral hepatitis in developing countries. In Europe many cases of autochthonous hepatitis E have been recognized recently in

1  
2  
3 patients without any evidence of travel to endemic areas; this has been suggested as a  
4 distinct clinico-epidemiological entity and the name of "indigenous to developed  
5 countries hepatitis E" (IDC HE) has been proposed (Teo,2006). Acute infection by HEV  
6 generally causes mild and self-limiting hepatitis but it can also result severe clinical  
7 fulminant hepatic failure in pregnant women and is associated with very high mortality  
8 particularly during the third trimester in endemic areas (Mushahwar, 2008).

9  
10 Interestingly, this feature has never been reported in industrialized countries where  
11 genotype 3 is the prevalent genotype in contrast to genotypes 1 and 2 which are found  
12 most frequently in developing countries. The reasons for the differences in the  
13 morbidity and outcome of HEV in pregnant women in individual countries remain  
14 unclear. Prevalence studies conducted in Spain have reported a wide range of  
15 prevalence rates from 0.6% in pregnant women in the north of Spain (Suarez Gonzalez  
16 et al, 2004) to 7.3 % in the general population of Catalonia (Buti et al, 2006).

17  
18 There has been a large influx of immigrants to Madrid in recent years with figures in  
19 June, 2009 as high as 17.1% of the total population and 27.3 % in some of the poorer  
20 neighbourhoods (Consejeria de Inmigración, 2009). Since a large number of pregnant  
21 women in Spain come from endemic countries and hepatitis E is more severe in this  
22 group of patients, it is important to determine the characteristics of HEV infection in  
23 this risk group.

24  
25 The aim of this study was to investigate the seroprevalence of HEV infection in a  
26 representative sample of pregnant women and determine the clinical features in relation  
27 to the presence of IgM anti-HEV antibodies in the blood of the patients.

## 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 **PATIENTS AND METHODS**

45  
46  
47 Serum samples from 1040 pregnant women, who attended the Obstetric Department of  
48 the Ramon and Cajal Hospital (Madrid, Spain) for routine screening were included in  
49 this study. Serum was taken from the patient during the first trimester of pregnancy  
50 during routine antenatal check-ups. One hundred and eighty one patients were studied in  
51 2007, 588 patients in 2009 and 271 patients in 2010. Habitat (rural or urban), age, ALT  
52 and AST and clinical symptoms were assessed in all patients. The sample size for an  
53 expected prevalence of IgG HEV antibodies of 2.9% in Madrid (Mateos, 1999), an  
54 alpha error of 5% and a precision of + 0.001 was 1,046 pregnant women.  
55  
56  
57  
58  
59  
60

1  
2  
3 IgG anti-HEV antibodies were detected in serum by a commercial enzyme  
4 immunoassay (EIA) (HEV Ab, DiaPro Diagnostic Bioprobes, Milan, Italy) following  
5 the manufacturer's instructions. All positive samples were studied further for the  
6 presence of IgM anti-HEV antibodies (HEV IgM, Dia.Pro Diagnostic Bioprobes, Milan,  
7 Italy). A result was considered positive by both tests when the ratio of the sample  
8 optical density and the cut off value was higher than 2. Positive results by EIA were  
9 confirmed by Western blot analysis (RecomBlot HEV IgG/IgM, Mikrogen, Martinsried,  
10 Germany). In addition, HEV RNA was amplified by reverse transcriptase (RT)-nested  
11 PCR (Fernández Barredo et al, 2006) in all serum samples with IgM anti-HEV.  
12  
13 The Institutional Ethic Committee approved the study and written informed consent was  
14 obtained from all the patients.  
15  
16  
17  
18  
19  
20  
21  
22  
23

## 24 RESULTS

25  
26  
27  
28 All the patients included in this study were living in urban areas in Madrid. The age  
29 range was 14 to 45 years (media: 30.6 years). None of the patients presented clinical  
30 symptoms associated with hepatitis when the blood sample was drawn. IgG anti-HEV  
31 antibodies were found in 2.1% (5/181) of the 2007 group, in 4.08 % (24/588) of the  
32 2009 group and in 3.3% (9/271) of the 2010 group. The overall prevalence was 3.6%  
33 (38/1040). IgM anti-HEV antibodies were detected in 0.67 % (7/1040) suggesting acute  
34 or recent infection. ARN HEV was negative in all 7 samples IgM anti-HEV positive  
35 tested. None of them presented clinical symptoms related to viral acute hepatitis. ALT  
36 and AST had normal values in all the patients who presented IgG and IgM anti-HEV in  
37 serum. Six oh those patients were Spanish and 1 was from Egypt.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

## 49 DISCUSSION

50  
51  
52  
53 HEV circulates in Spain. The virus has been isolated from sewage (Pina S et al, 2000)  
54 and it is considered a zoonotic disease and the most probable reservoir for the human  
55 population is swine (Fernández Barredo et al, 2007). HEV has been the cause of  
56 sporadic acute hepatitis E in Spain (Perez-Gracia et al, 2004) and is considered as an  
57 occupational disease (Perez Gracia et al, 2007). However in industrialized countries,  
58 the transmission routes for autochthonous HEV infections remain unknown although  
59  
60

1  
2  
3 food and environmental routes are considered the most important (Teo, 2010). Since all  
4 the patients included in this study were living in urban settings, direct animal contact  
5 was not a probable transmission route. Obviously, exposure to food in the kitchen to  
6 raw pork or poultry cannot be excluded neither can shellfish which is consumed  
7 frequently in Spain. The prevalence of IgG anti-HEV antibodies was 2.1 %, 4.08% and  
8 3.3 % in the group of patients studied in 2007, 2009 and 2010 respectively. The overall  
9 prevalence was 3.6%. In studies conducted in 1999 in the same area, a prevalence of  
10 2.9% was found (Mateos et al, 1999). The odds ratio was 0.8 and the 95% CI was 0.67-  
11 1.73. This difference was not statistically significant ( $p < 0.810$ ). Interestingly, seven  
12 (0.67 %) patients presented IgM anti-HEV antibodies in blood. All of them were  
13 asymptomatic cases of hepatitis E occurring during the first trimester of pregnancy.  
14 Unfortunately it was not possible to test any further blood samples since these patients  
15 were healthy during pregnancy. No risk factors for transmission were found by  
16 interviews held with these infected patients. In contrast, pregnant women in endemic  
17 areas suffer very frequently severe hepatitis and even death. The reasons for the  
18 difference in the outcome of HEV in different geographical areas remain unclear but it  
19 could be the result of early childhood exposure to HEV, producing long-lasting  
20 immunity and/or modified subsequent responses of exposure to the virus. Alternatively,  
21 the predominant HEV genotypes in Spain could be less virulent than those in  
22 developing countries. The HEV genotypes circulating in endemic areas, mainly  
23 genotype 1 and 2, are very uncommon in industrialized countries. In conclusion, the  
24 data obtained during this study demonstrate that asymptomatic hepatitis E occurs in  
25 pregnant women in non-endemic countries at a very low rate (0.67 %). The HEV  
26 genotypes and subtypes circulating in different countries can be responsible for the  
27 different geographical morbidity in pregnant women (Navaneethan et al, 2008);  
28 seroprevalence of IgG HEV antibodies has not increased in Spain in these recent years  
29 (3.6 %), considering the seroprevalence studies conducted in the same area in 1999  
30 (2.9%) (Mateos et al, 1999). Discrepancies between the prevalence rates found in Spain  
31 by other authors (0.6%-7.3%) (Suarez Gonzalez et al, 2004; Buti et al, 2006) may be  
32 due to epidemiological characteristics of the patients studied (rural or urban areas,  
33 gender, age, profession), methodology and lack of confirmation with other tests such as  
34 Western blot. The geographical situation inside the country can also have a role since a  
35 very high prevalence has been reported in south west France (Mansuy et al, 2008). The  
36 good hygienic measures and public health conditions in Spain may be the most  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 important factors in maintaining the low prevalence rate of hepatitis E which is not  
4 considered a major health problem at the moment. Nonetheless if the number of  
5 clandestine immigrants from endemic countries continues to increase in Spain and the  
6 route of transmission of HEV is not established completely in order to prevent the  
7 spread of HEV, this favourable epidemiological situation may change in future years.  
8  
9  
10  
11  
12  
13  
14  
15

### 16 **Conflict of interests**

17  
18 The authors declare that there is not any conflict of interest in the present report.  
19  
20  
21  
22  
23  
24

### 25 **REFERENCES**

- 26  
27  
28 Buti M, Domínguez A, Plans P, Jardí R, chaper M, Espuñes J, Cardeñosa N,  
29 Rodríguez Frias F, Esteban R, Plasencia A, Salleras L. 2006. Prevalence of hepatitis  
30 E Virus infection in children in the northeast of Spain. *Clin Vacc Immunol* ;13:1328-  
31 1333  
32  
33  
34 Consejería de Inmigración. Población empadronada junio. 2009. Diario ABC,  
35 26/12/2009. [www.abc.es/madrid](http://www.abc.es/madrid).  
36  
37  
38 Fernández Barredo S, Galiana C, Garcia A, Vega S, Gómez MT, Perez Gracia  
39 MT.2007. Detection of hepatitis E virus shedding in feces of pigs at different stages  
40 of production using reverse transcription-polymerase chain reaction. *J Vet Diag*  
41 *Invest*;18:462-465  
42  
43  
44 Fernández-Barredo S, Galiana C, Garcia A, Vega S, Gómez MT, Perez-Gracia MT.  
45 2007. Prevalence and genetic characterization of hepatitis E virus (HEV) in paired  
46 samples of feces and serum from naturally infected pigs. *Can J Vet Res* ; 71 (3):  
47 236-242  
48  
49  
50 Mateos M L, Camarero C, Lasa E, Teruel JL, Mir N, Baquero F. 1999. Hepatitis E  
51 virus: relevance in blood donors and risk groups. *Vox Sanguinis* ;76:78-80  
52  
53  
54 Mansuy JM, Legrand-Abrevanel F, Calot JP, Peron JM, Alric L, Agudo S, Rech Henri,  
55 Destruel F, Izopet J.2008. High prevalence of anti-hepatitis E virus antibodies in  
56 flood donors from south west France. *J Med Virol*; 80:289-293  
57  
58  
59 Mushahwar I. Hepatitis E Virus: Molecular virology, clinical features, diagnosis,  
60 transmission, epidemiology and prevention. 2008. *J Med Virol* ; 80:646-658

- 1  
2  
3 Navaneethan U, Al Mohajer M, Shata MT. Hepatitis E in pregnancy: understanding the  
4 pathogenesis. *Liver Int* 2008 Nov;28(9):1190-9  
5  
6  
7 Perez-Gracia M.T, Garcia-Valdivia MS, Galan F, and Rodriguez-Iglesias MA, 2004.  
8 Detection of hepatitis E virus in patients sera in southern Spain. *Acta Virol* 48: 197-  
9 200  
10  
11 Perez-Gracia MT, Mateos ML, Galiana C, Fernández-Barredo S, García A, Gómez  
12 MT, Moreira V. 2007. Autochthonous hepatitis E infection in a slaughterhouse  
13 worker. *Am J Trop Med Hyg* 77(5):893-896  
14  
15  
16 Pina S, Buti M, Cotrina M, Piella J, Girones R.2000. HEV identified in serum from  
17 humans with acute hepatitis and in sewage of animal origin in Spain. *J Hepatol* 33:  
18 826-833  
19  
20  
21 Suarez Gonzalez A, Solís Sánchez G, Otero Guerra L, Viejo de la Guerra G, Álvarez  
22 Navascues C, García López R.2004. Prevalencia de inmunidad frente a los virus de  
23 la hepatitis en gestantes del Área Sanitaria de Gijón. *Gastroenterol Hepatol* 27  
24 (6):347-352  
25  
26  
27 Teo C.2006. Hepatitis E indigenous to economically developed countries: to what  
28 extent a zoonosis? *Curr Opin Infect Dis*;19:460-466  
29  
30  
31 Teo CG. 2010. Much meat, much malady: changing perceptions of the epidemiology of  
32 hepatitis E. *Clin Microbiol Infect* 16:24-32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60