



Co-Circulation of Genetically Distinct Human Metapneumovirus and Human Bocavirus Strains in Young Children with Respiratory Tract Infections in Italy

Alessandra Zappa, Marta Canuti, Elena Frati, Elena Pariani, Silvana Perin, Lorena Maria Ruzza, Claudio Farina, Alberto Podestà, Alessandro Zanetti, Antonella Amendola, et al.

► To cite this version:

Alessandra Zappa, Marta Canuti, Elena Frati, Elena Pariani, Silvana Perin, et al.. Co-Circulation of Genetically Distinct Human Metapneumovirus and Human Bocavirus Strains in Young Children with Respiratory Tract Infections in Italy. *Journal of Medical Virology*, 2010, 83 (1), pp.156. 10.1002/jmv.21940 . hal-00599787

HAL Id: hal-00599787

<https://hal.science/hal-00599787>

Submitted on 11 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Co-Circulation of Genetically Distinct Human Metapneumovirus and Human Bocavirus Strains in Young Children with Respiratory Tract Infections in Italy

Journal:	<i>Journal of Medical Virology</i>
Manuscript ID:	JMV-10-1827.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	21-Jul-2010
Complete List of Authors:	<p>Zappa, Alessandra; Università degli Studi di Milano, Dipartimento di Sanità Pubblica-Microbiologia-Virologia</p> <p>Canuti, Marta; Università degli Studi di Milano, Dipartimento di Sanità Pubblica-Microbiologia-Virologia</p> <p>Fрати, Elena; Università degli Studi di Milano, Dipartimento di Sanità Pubblica-Microbiologia-Virologia</p> <p>Pariani, Elena; Università degli Studi di Milano, Dipartimento di Sanità Pubblica-Microbiologia-Virologia</p> <p>Perin, Silvana; AO "Ospedale San Carlo Borromeo", Unità di Microbiologia</p> <p>Ruzza, Lorena; AO "Ospedale San Carlo Borromeo", Unità di Pediatria</p> <p>Farina, Claudio; AO "Ospedale San Carlo Borromeo", Unità di Microbiologia</p> <p>Podestà, Alberto; AO "Ospedale San Carlo Borromeo", Unità di Pediatria</p> <p>Zanetti, Alessandro; Università degli Studi di Milano, Dipartimento di Sanità Pubblica-Microbiologia-Virologia; Università degli Studi di Genova, CIRI-IV, Dipartimento di Scienze della Salute</p> <p>Amendola, Antonella; Università degli Studi di Milano, Dipartimento di Sanità Pubblica-Microbiologia-Virologia</p> <p>Tanzi, Elisabetta; Università degli Studi di Milano, Dipartimento di Sanità Pubblica-Microbiologia-Virologia</p>
Keywords:	Respiratory infections, Phylogenetic analysis, human Metapneumovirus, human Bocavirus, Recombination events

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCHOLARONE™
Manuscripts

For Peer Review

Table I. Primers Used in This Study.**A. Primers Used for Respiratory Infection Screening.^a**

	Primer	Sequence 5'→3'	Gene	Amplicon size (bp)	Reference
hAdV nested-PCR					
I step	ADHEX1f ADHEX1r	AACACCTAYGASTACATGAAC ATGGGGGTARAGCATGTTRGC	Hexon	473	Modified from [Avellón et al., 2001]
II step	ADHEX2f ADHEX2r	AAAYCCMTTYAACCACCACC ACATCCTTYCKGAAGTTCCA		170	
hBoV nested-PCR					
I step	162f 561r	CCAGCAAGTCCTCCAAACTCACCTGC GGAGCTTCAGGATTGGAAGCTCTGTG	NP-1	399	[Manning et al., 2006]
II step	188f 542r	GAGCTCTGTAAGTACTATTAC CTCTGTGTTGACTGAATACAG		354	Modified from [Allander et al., 2005]
hCoV multiplex nested-PCR					
I step					
hCoV-229E	CORO9 CORO10	GCACAGGACCCCATAAAGATGC GAGAACGAGCAAGACTCTTGGCA	N	444	Modified from [Dessau et al., 2001]
hCoV-OC43	CORO11 CORO12	GCAATCCAGTAGTAGAGCGTCC TTGACATCAGCCTGGTTRCTAGCG		445	
II step					
hCoV-229E	CORO1 CORO2	AGGCGCAAGAATTCAGAACCAGAG AGCAGGACTCTGATTACGAGAAGG		308	
hCoV-OC43	CORO5 CORO7	CCCAAGCAAACCTGCTACYTCTCAG GCAGCARTTGACGCTGGTTG		228	
Flu A/B multiplex Real Time PCR					
FluA					
	FLUAV_FOR FLUAV_REV FLUAV_TM	ACAAGACCAATCCTGTCACCTCT GGCATTTTGGACAAAGCGTCTAC FAM-CAGTCCTCGCTCACTGGGCACGGT(p)-BHQ1	M	108	[Valle et al. 2001]
FluB					
	FLUBV_FOR FLUBV_REV FLUBV_TM	CCAGTGGGACAACCAGA TGCTCTTTCCGGGGATG JOE-ATCATCAGACCAGCAACCCTTGCC(p)-BHQ1	NP	89	
hMPV nested-PCR					

I step	HMPV-M1f	GAGTCCTAYCTRGTAGACACC	M	247	This study
	HMPV-M1r	AGTACAGACATDGCWGCACC			
II step	HMPV-M2f	GACCWGCTGTTCAAGTTG		151	
	HMPV-M2r	YTGTGATGYAGCATACAGAG			
hPIV1-4 multiplex nested-PCR					
I step					
hPIV-1	PIP1+	CYTAAATTCAGATATGTAT	HN	480	Modified from [Echevarría et al., 1998]
	PIP1-	GATAAATARTWATTGATACG			
hPIV-2	PIP2+	AACAATCTGCTGCAGSRTTT	HN	507	
	PIP2-	ATGTCAGAYAATGGRCAAAT			
hPIV-3	PIP3+	CTGTAAACTCAGACTTGGTA	HN	477	
	PIP3-	TTTARGCCYTTGTCAACAAC			
hPIV-4	PI4P+	CTGAACGGTTGCATTCAGGT	P	441	[Aguilar et al., 2000]
	PI4P-	TTGCATCAAGAATGAGTCCT			
II step					
hPIV-1	PIS1+	CCGGHAATYTCTCATACCTATG	HN	316	Modified from [Echevarría et al., 1998]
	PIS1-	CYTTGGAGCGGAGTTGTTAWG			
hPIV-2	PIS2+	CCATTTACCTAAGTGATGGAAT	HN	203	
	PIS2-	GCCCTGTTGTATTTGGAAGAGA			
hPIV-3	PIS3+	ACTCCCAARGTTGATGAAAGAT	HN	102	
	PIS3-	TAAATCTTGTTGTTGAGATTG			
hPIV-4	PI4S+	AAAGAATTAGGTGCAACCAGTC	P	244	[Aguilar et al., 2000]
	PI4S-	GTGTCTGATCCCATAAGCAGC			
hRV eminested-PCR					
I step	PR1f	CGGACACCCAAAGTAG	5'UTR	380	[Ireland et al., 1993]
	PR2r	GCACTTCTGTTTCCCC			
II step	PR1f	CGGACACCCAAAGTAG		202	
	PR3r	GGCAGCCACGCAGGCT			
RSV multiplex nested-PCR					
I step					
RSV-A/B	RSVAB1	ATGGAGYTGCRYATCCWCARRRCAARTGCAAT	F	737	[Coiras et al., 2003]
	RSVAB2	AGGTGTWGTTACACCTGCATTRACACTRAATTC			
II step					
RSV-A	RSVA3	TTATACACTCAACAATRCCAAAAAWACC		363	
	RSVA4	AAATTCCTGGTAATCTCTAGTAGTCTGT			

RSV-B	RSVB3 RSVB4	ATCTTCCTAACTCTTGCTRTTAATGCATTG GATGCGACAGCTCTGTTGATTACTATG		611	
B. Primers Used for the Molecular Characterization of hBoV and hMPV Strains.^a					
hBoV sequencing nested-PCR					
I step	VPs1 VPas1	GCACTTCTGTATCAGATGCCTT CGTGGTATGTAGGCGTGTAG	VP1/2	903	[Smuts and Hardie, 2006]
II step	VPs2 VPas1	CTTAGAACTGGTGAGAGCACTG CGTGGTATGTAGGCGTGTAG		850	
hMPV sequencing nested-PCR					
I step	F-f MPVF1r	GTYAGCTTCAGTCAATTCAAC GTCTTCCTGTGCTAACTTTG	F	532	Modified from [Huck et al., 2006] [Peret et al., 2002]
II step	MPVF1f F-r	CTTTGGACTTAATGACAGATG CCTGTGCTGACTTTGCATG		445	[Peret et al., 2002] Modified from [Huck et al., 2006]


^aPrimers used for the molecular surveillance for acute respiratory tract infections (A) and for the molecular characterization of hBoV and hMPV strains (B).

Table II. Molecular detection of respiratory viruses in pharyngeal swabs collected from hospitalized children with acute respiratory tract infections and children with viral respiratory infections.

A. Number (95% CI) of Infections and Age Distribution Children (N = 240) Hospitalized with Acute Respiratory Tract Infections.					
	Total 240	0–6 months 144 (60; 53.7–66.1)	7–12 months 36 (15; 10.9–19.9)	13–24 months 45 (18.8; 14.2–24.1)	25–36 months 15 (6.3; 3.7–9.9)
Viral agents					
RSV	46 (19.2; 14.6–24.5)	35 (24.3; 17.8–31.8)	4 (11.1; 3.6–24.7)	7 (15.5; 7.1–28.4)	0
hAdV	28 (11.7; 8.1–16.2)	13 (9; 5.1–14.6)	7 (19.4; 8.9–34.7)	7 (15.5; 7.1–28.4)	1 (6.7; 0.3–28.7)
hRV	24 (10; 6.7–14.3)	14 (9.7; 5.6–15.4)	5 (13.9; 5.3–28.1)	4 (8.9; 2.9–20.1)	1 (6.7; 0.3–28.7)
FLU	11 (4.6; 2.4–7.8)	5 (3.5; 1.3–7.5)	2 (5.5; 0.9–17.2)	4 (8.9; 2.9–20.1)	0
CoV	7 (2.9; 1.3–5.7)	3 (2.1; 0.5–5.6)	0	3 (6.7; 1.7–17.1)	1 (6.7; 0.3–28.7)
hPIV	1 (0.4; 0.02–2.0)	0	0	1 (2.2; 0.1–10.5)	0
hMPV	20 (8.3; 5.3–12.4)	13 (9; 5.1–14.6)	3 (8.3; 2.2–21)	4 (8.9; 2.9–20.1)	0
hBoV	29 (12; 18.4–16.7)	14 (9.7; 5.6–15.4)	4 (11.1; 3.6–24.7)	11 (24.4;13.6–38.5) ^a	0
B. Number (95% CI) of Single and Co-infections in Patients (N = 131) in the Context of Age.					
Total	131 (54.6; 48.3–60.8)	76 (52.8; 44.6–60.8) ^b	22 (61.1; 44.6–75.9) ^b	30 (66.7; 52–79.2) ^b	3 (20; 5.4–45.4)
Single infections	102 (77.9; 70.2–84.4)	58 (76.3; 65.8–84.9)	19 (86.4; 67.2–96.4)	22 (73.3; 55.6–86.8)	3
Co-infections	29 (22.1; 15.6–29.9)	18 (23.7; 15.2–34.2)	3 (13.6; 3.6–32.8)	8 (26.7; 13.2–44.4)	0


^ap < 0.05, prevalence of hBoV infections vs. that observed in children 0–6 months and 7–12 months of age.

^bp < 0.05, N (%) viral infections vs. N (%) of children 25–36 months of age.




Genetic analysis of hMPV. A) Phylogenetic tree of the Italian I-MI hMPV sequences (boldface) demonstrating the phylogenetic relationship of the I-MI sequences based on partial F gene nucleotide sequences. Bootstrap values (in addition to A1, A2a, A2b, B1 and B2 group/sublineages) are indicated. Reference sequences correspond to the following GenBank accession numbers: A1) AF371337, AY145294, DQ453017, AJ867506, AJ867520, EF612444; A2a) AY304360, AY145296, DQ453014, EF612451; A2b) AY530095, DQ453021, AJ867528, AJ867538, EF612457; B1) AY304361, AY145294, DQ453015, AJ867513, EF612440; B2) AY304362, AY145289, DQ452995, AJ867503, EF612441. B) Evolutionary mean distances over all sequence pairs within and between groups.

258x362mm (150 x 150 DPI)



Genetic analysis of hBoV sequences. Phylogenetic tree of the Italian I-MI human Bocavirus sequences (boldface) based on partial VP1/VP2 region nucleotide sequences. Bootstrap values (in addition to St1 and St2 genotype clusters) are indicated. Reference sequences correspond to the following GenBank accession numbers: St-1) DQ000497, AM160609, AM160611-AM160613, AM160615, AM689298, AM689299, AM689306, AM689307, EU069434, EU069436, EU069437, EF203921; St-2) DQ000495, AM160610, AM160614, AM689297, AM689301-AM689303, AM689305, EF450717, EF450718, EF450722, EF450723, EF450736, EF450740.
178x302mm (150 x 150 DPI)



Recombination recognition of the Italian I-MI/04-08 sequence: bootscan analysis and identities values between the recombinant sequence and the two reference sequence (MPT8 and HK7) before and after the breakpoint (inset).
254x190mm (96 x 96 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Co-Circulation of Genetically Distinct Human Metapneumovirus and Human Bocavirus Strains in Young Children with Respiratory Tract Infections in Italy

Alessandra Zappa¹, Marta Canuti¹, Elena Frati¹, Elena Pariani¹, Silvana Perin², Maria Lorena Ruzza³, Claudio Farina², Alberto Podestà³, Alessandro Zanetti^{1,4}, Antonella Amendola^{1,4}, Elisabetta Tanzi^{1,4,*}

Affiliations:

- ¹ Dipartimento di Sanità Pubblica-Microbiologia-Virologia, Università degli Studi di Milano, Via Pascal 36/38, 20133 Milano, Italy;
- ² Unità di Microbiologia, AO “Ospedale San Carlo Borromeo”, Via Pio II 3, 20153 Milano;
- ³ Unità di Pediatria, AO “Ospedale San Carlo Borromeo”, Via Pio II 3, 20153 Milano;
- ⁴ CIRI-IV, Dipartimento di Scienze della Salute, Università degli Studi di Genova, Via Pastore 1, 16132 Genova.

***Corresponding author:** Elisabetta Tanzi, Ph.D.

Dipartimento di Sanità Pubblica-Microbiologia-Virologia, Università degli Studi di Milano, Via Pascal 36/38, 20133 Milano, Italy;
Tel: +390250315139 - Fax: +390250315120

E-mail address: elisabetta.tanzi@unimi.it

Running Title: hMPV and hBoV infections in Italian children

ABSTRACT

The discovery of human Metapneumovirus (hMPV) and human Bocavirus (hBoV) identified the etiological causes of several cases of acute respiratory tract infections in children. This report describes the molecular epidemiology of hMPV and hBoV infections observed following viral surveillance of children hospitalized for acute respiratory tract infections in Milan, Italy. Pharyngeal swabs were collected from 240 children ≤ 3 years of age (130 males, 110 females; median age, 5.0 months; IQR, 2.0–12.5 months) and tested for respiratory viruses, including hMPV and hBoV, by molecular methods. hMPV-RNA and hBoV-DNA positive samples were characterized molecularly and a phylogenetical analysis was performed. PCR analysis identified 131/240 (54.6%) samples positive for at least one virus. The frequency of hMPV and hBoV infections was similar (8.3% and 12.1%, respectively). Both infections were associated with lower respiratory tract infections: hMPV was present as a single infectious agent in 7.2% of children with bronchiolitis, hBoV was associated with 18.5% of pediatric pneumonias and identified frequently as a single etiological agent. Genetically distinct hMPV and hBoV strains were identified in children examined with respiratory tract infections. Phylogenetic analysis showed an increased prevalence of hMPV genotype A (A2b sublineage) compared to genotype B (80% vs. 20%, respectively) and of the hBoV genotype St2 compared to genotype St1 (71.4% vs. 28.6%, respectively). Interestingly, a shift in hMPV infections resulting from A2 strains has been observed in recent years. In addition, the occurrence of recombination events between two hBoV strains with a breakpoint located in the VP1/VP2 region was identified.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

43 **Key words:** Pediatric respiratory infections, phylogenetic analysis, human Metapneumovirus,
44 human Bocavirus, pediatric hospitalization, viral recombination.

For Peer Review

INTRODUCTION

Acute respiratory tract infections are an important cause of morbidity and mortality worldwide. These infections are ubiquitous and contagious, affecting repeatedly individuals of all ages, particularly children younger than five years of age [World Health Organization, 2009] and many infections have unknown etiologies. The recent discovery of the human Metapneumovirus (hMPV) in 2001 [van den Hoogen et al., 2001] and of the human Bocavirus (hBoV) in 2005 [Allander et al., 2005] identified the etiological causes of several respiratory infections. hMPV is a common cause of respiratory disease in children under five years old and often accounts for severe clinical manifestations, including bronchiolitis or pneumonia [Foulongne et al., 2006]. hMPV is a single-stranded negative RNA virus of the *Paramyxoviridae* family, *Pneumovirinae* subfamily [van den Hoogen et al., 2001; Broor et al., 2008]. Molecular analysis identified two major genotypes (A and B) with genetic lineages A1, A2 and B1, B2, respectively [van den Hoogen et al., 2004]. Lineage A2 has been recently divided into A2a and A2b sublineages [Huck et al., 2006].

hBoV is detected frequently in the respiratory and gastrointestinal tracts, suggesting its pathologic involvement at both sites [Lindner et al., 2008; Schildgen et al., 2008]. hBoV belongs to the *Bocavirus* genus of the *Parvoviridae* family. It is a non-enveloped virus and contains a linear single-stranded DNA molecule. The two major hBoV genotypes correspond to the original St1 (Stockholm 1) and St2 (Stockholm 2) isolates [Allander et al., 2005].

This report describes the molecular epidemiology of hMPV and hBoV identified following surveillance for respiratory viruses in children hospitalized for acute respiratory tract infections in Milan, Italy. A detailed genetic analysis was carried out using the hMPV and hBoV sequences identified.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
92

67 **PATIENTS AND METHODS**

68 **Study Population**

69 Pediatric patients (N=240; 130 males and 110 females; median age, 5.0 months; interquartile
70 range, 2.0–12.5 months) admitted to the San Carlo Borromeo Hospital in Milan between 2004–
71 2008 with a diagnosis of acute respiratory infection were enrolled in the study. Lower respiratory
72 tract infections (*i.e.*, bronchitis, bronchiolitis, pneumonia or asthma) were observed in 66.7% (95%
73 CI: 60.5–72.4) of children. Bronchiolitis and bronchitis were the most frequent outcomes (60.6%,
74 95% CI: 52.9–68.0 and 37.5%, 95% CI: 30.3–45.2, respectively), pneumonia and asthma being less
75 frequent (16.9%, 95% CI: 11.7–23.3 and 9.4%, 95% CI: 5.5–14.7).

76 At the time of hospital admission pharyngeal swabs were collected from each patient for the
77 identification and characterization of respiratory viruses using Plain Swabs and ITM-RT (Copan,
78 Italy). Informed consent was obtained from parents of eligible children and children were divided
79 into four age groups: ≤6 months (60%, 95% CI: 53.7–66.1), 7–12 months (15%, 95% CI: 10.9–
80 19.9), 13–24 months (18.8%, 95% CI: 14.2–24.1) and 25–36 months (6.2%, 95% CI: 3.7–9.9).

81 The seasonal distribution of the samples collected is as follows: 35.4% (95% CI: 29.6–41.6)
82 of samples were collected during autumn (October–December), 52.5% (95% CI: 46.2–55.8) during
83 winter (January–March) and 12.1% (95% CI: 8.4–16.7) during spring (April–June).

85 **Nucleic Acid Extraction and Amplification**

86 Nucleic acids were extracted using the RNeasy Mini Kit and QIAmp DNA Mini Kit (Qiagen
87 GmbH, Germany). For RNA virus detection cDNA was synthesized with pd(N)₆ random hexamer
88 primers (Amersham Biosciences, UK) using a M-MLV reverse transcriptase (Invitrogen Tech-Line,
89 USA). Viral detection was performed by PCR for the identification and typing of both classic viral
90 respiratory pathogens *i.e.*, influenza virus A and B (FluA/B), parainfluenza viruses 1-4 (hPIV1-4),
91 respiratory syncytial virus A and B (RSVA/B), coronavirus (229E and OC43 hCoV229E/OC43),
92 rhinovirus (hRV), adenovirus (hAdV) and hMPV and hBoV (Table IA).

Sequencing of Identified hMPV and hBoV Strain Sequences

Molecular characterization was performed by sequence analysis of a 450 bp amplicons of the fusion (F) gene (spanning nucleotides 653–1,102) for hMPV and an 850 bp amplicon of the VP1/VP2 region (spanning nucleotides 1,084–1,933) for hBoV (Table IB).

Following PCR amplification, amplicons were purified with NucleoSpin® Extract II (Macherey-Nagel GmbH, Germany). Nucleotide sequences were obtained by automated DNA sequencing on the ABI PRISM 3100 genetic analyzer (Applied Biosystems, Foster City, CA). The hMPV-F gene and hBoV-VP1/VP2 region nucleotide sequences obtained were deposited into GenBank.

Molecular Characterization and Phylogenetic Analysis

Multiple nucleotide sequences were aligned using ClustalX version 2.0 [Thompson et al., 1997]. Phylogenetic trees were constructed by means of the Neighbor-Joining method and Kimura 2-Parameter model using the MEGA package, version 4.0 [Kimura, 1980; Saitou and Nei, 1987; Tamura et al., 2007]. A bootstrap resampling analysis was performed (1,000 replicates) to test tree robustness [Felsenstein, 1985].

Mean evolutionary distances over all sequence pairs between and within groups were calculated using the Kimura 2-Parameter model in MEGA software [Kimura, 1980; Tamura et al., 2007]. To detect recombination events, a bootscanning analysis was performed using the Simplot software version 3.5.1 (Kimura 2-Parameter model; window size: 200 bp; step size: 20 bp; 1,000 bootstrap replicates, Neighbour-Joining tree analysis), [Lole et al., 1999]. Predicted protein sequences were obtained by means of BioEdit software [Hall, 1999].

Statistical Analysis

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

118 Data were expressed as median (interquartile range, IQR) and percentages (95% confidence
119 intervals, 95% CI) as appropriate. Comparisons between groups were performed using the Chi-
120 squared test or Fisher’s exact test. A P-value <0.05 was considered statistically significant (two-
121 tailed test). All statistical analyses were performed using the OpenEPI software, version 2.2.1 [Dean
122 et al., 2009].

For Peer Review

RESULTS

Viral Surveillance in Children with Acute Respiratory Tract Infections

One hundred and thirty-one out of 240 (54.6%, 95% CI: 48.3–60.8) samples collected from hospitalized children were PCR-positive for at least one respiratory virus. Forty-six samples (19.2%) were positive for RSVA/B, 11.7% for hAdV, 10% for hRV, 4.6% for Flu A/B, 2.9% for hCoV229E/OC43 and 0.4% for hPIV3. hMPV was detected in 8.3% and hBoV in 12.1% of samples examined (Table IIA). Mixed infections were detected in 22.1% of acute viral respiratory tract infections (Table IIB). Respiratory viruses were detected more frequently in children younger than two years of age (52.8%, 61.1% and 66.7% for children ≤ 6 , 7–12 and 13–24 months of age, respectively) in comparison to children 25–36 months of age (20%, $P < 0.05$, Table IIB).

Viral pathogens were associated with 60.6% (95% CI: 52.9–68.0) of lower respiratory-tract infections, particularly in bronchiolitis cases (60.8%, 95% CI: 50.9–70.2). The most frequently detected viruses were RSV (50.8%, 95% CI: 38.2–63.4), hBoV (20.3%, 95% CI: 11.5–32.0), hRV (16.9%, 95% CI: 8.9–28.1), hAdV (16.9%, 95% CI: 8.9–28.1) and hMPV (11.9%, 95% CI: 5.3–22.1).

Epidemiologic and Molecular Features of hMPV and hBoV Infections

hMPV or hBoV were detected in 37.4% (95% CI: 29.4–45.9) of respiratory virus-positive samples but only in children ≤ 24 months old. hMPV prevalence was similar in the first three age groups (range: 8.3–9.0%) while hBoV was identified more frequently in the 13–24 month old group compared to the other groups (24.4% vs. 9.7% and 11.1%, $P < 0.05$) (Table IIA).

hMPV was identified more often as a single pathogen and only 30% of cases were associated with mixed infections ($P < 0.05$). hBoV was detected as a mixed infection in 58.6% (17/29) of cases, mainly in association with RSV ($N = 10$), hRV ($N = 5$) and hAdV ($N = 4$). Among these, two triple-infection (hBoV-RSV-hRV) cases were detected. The frequency of co-infections was similar in all age groups but different viruses were involved. The median age of hBoV co-

1
2
3 149 infected children was 2.0 months (IQR: 1.5–50.0 months), for hBoV-RSV 8.5 months (IQR: 3.5–
4
5
6 150 13.0 months) and 14 months for both hBoV-hRV and hBoV-hAdV (IQR: 11.5–17.0 months). No
7
8 151 hMPV-hBoV co-infections were identified.

9
10 152 hMPV was found as the single source of infection in 7.2% (95% CI: 3.2–13.8) of
11
12
13 153 bronchiolitis cases and part of co-infections in 11.1% of pneumonia (95% CI: 2.9–27.3) and 13.3%
14
15 154 of asthma cases (95% CI: 2.3–37.5). hBoV was identified in 12.4% of bronchiolitis cases (95% CI:
16
17 155 6.9–20.1), mainly as a hBoV-RSV co-infection. In addition, a high prevalence of hBoV infections
18
19
20 156 (18.5%, 95% CI: 7.1–36.4) were seen in association with pneumonia, most often as a single viral
21
22 157 agent.

23
24
25 158
26
27 159 **Phylogenetic Analysis of hMPV and hBoV Sequences**

28
29 160 Ten out of 20 hMPV-positive samples were characterized at the molecular level and a
30
31
32 161 phylogenetic tree of the hMPV F gene was generated (Fig. 1A). The identity range between tree
33
34 162 sequences was 81.1–100%. Mean evolutionary distances for all sequence pairs are reported in Fig.
35
36 163 1B. Phylogenetic analysis showed a prevalence of genotype A strains (8/10) compared to genotype
37
38
39 164 B strains (2/10). In the context of genotype A, six hMPV-F sequences shared similarity with the
40
41 165 A2b sublineage, one with A2a and one with the A1 group that presented with a single amino acid
42
43 166 change (N358T). Genotype B sequences belonged exclusively to lineage B1.

44
45
46 167 Sixteen out of 29 hBoV-positive samples were analyzed and the hBoV VP1/VP2 region
47
48 168 described (Fig. 2). All sequences shared a high degree of similarity with reference strains and other
49
50
51 169 defined sequences (identity range: 97.3–100%). Most sequences (14/16) were genetically similar to
52
53 170 genotypes St1 (4/14) and St2 (10/14). A third cluster (bootstrap value: 92), including two Italian
54
55 171 sequences (I-MI/04-08 and I-MI/02-07, identity 100%) and the Asian HK7 isolate, were identified
56
57
58 172 between the two main branches. The boot scanning analysis revealed a recombination event with a
59
60 173 breakpoint located at nucleotide position 1,302 of the VP1/VP2 region (Fig. 3). The 5' region of
174
these two Italian sequences was more similar to the MPT-8 strain (identity 99% vs. 97.6%) and the

1
2
3 175 3' portion more similar to the HK7 strain (identity 97.8% vs. 99%) (Fig. 3 inset).
4

5
6 176 Six amino acid substitutions were found: G415S, N474S, F540Y, N546H, A555T, and
7
8 177 G566R. Mutations G415S, F540Y and N546H were identified in the I-MI/04-08 and I-MI/02-07
9
10 178 sequences.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION

A molecular-epidemiologic survey of acute respiratory infections was conducted in children younger than three years old hospitalized between 2004 and 2008 in Milan, Italy. Both hospitalization and positive viral detection were more frequent in children <2 years of age compared to older patients.

The prevalence of both hMPV and hBoV infections (8.3% and 12.1%, respectively) were similar to rates reported previously [Maggi et al., 2003; Bastien et al., 2006; Foulongne et al., 2006; Allander et al., 2007]. Both hMPV and hBoV infections were associated with lower respiratory tract infections. hMPV was present as a single infectious agent in 7.2% of children aged ≤12 months with bronchiolitis. The involvement of hBoV at the onset of bronchiolitis was difficult to assess due to the frequent association with other viruses, particularly RSV. In this study, hBoV was associated with 18.5% of pediatric pneumonias and frequently identified as a single agent.

Twenty percent of hospitalized children had dual or triple co-infections and hBoV was involved in approximately 60% of mixed infections. The extended persistence of hBoV in airways (up to 7 months) increased patient's chances of acquiring a co-infection, explaining the high frequency of co-infections with this virus [Brieu et al., 2008; Lindner et al., 2008; von Linstow et al., 2008]. The seasonal distribution of hBoV, with a peak incidence between December and January, resembled that observed for RSV, hRV and hAdV. This suggested that the overlapping distribution of these viruses during the seasons could have facilitated hBoV co-infections. The epidemiologic model of hMPV infections was distinct from many other respiratory viruses, with peak incidence rates in late spring, mostly between March and April. This observation may account for the absence of hMPV-hBoV co-infections.

Co-circulation of both hMPV genotypes was observed, with predominance of genotype A (particularly the A2b sublineage), supporting previous reports demonstrating an increase in A2 strain infections in recent years [Boivin et al., 2004; Wang et al., 2008; Herbert et al., 2005]. Guant et al. [Guant et al.; 2009] suggested a further potential back-shift to genotype B in Scotland during

the winter season of 2007/2008. Unfortunately, no samples from this study were available for sequencing to confirm this hypothesis. As suggested by Herbert *et al.* [2005], the complex circulation pattern of hMPV may be the reason this strain can escape pre-existing host immune responses, suggesting that additional molecular surveillance studies need to be carried out to confirm genotype B back shift circulation and to clarify the existence of a seasonal distribution of hMPV strains. Phylogenetic analyses demonstrated that both hBoV genotypes co-circulated during the study period and that genotype St2 predominated. Interestingly, evidence of genetic recombination between the two hBoV strains (with a breakpoint site in the VP1/VP2 region) was found. The two mosaic sequences (identity: 100%) were identified in two different years (2007 and 2008), indicating that this variant had been circulating in the population since at least 2007. Genetic recombination has been reported to occur in Parvoviruses [Shackelton *et al.*, 2007; Ohshima *et al.*, 2009] and in different hBoV species [Arthur *et al.*, 2009; Kapoor *et al.*, 2009; Kapoor *et al.*, 2010]. To date, only Lin *et al.* [2009] have reported an intra-species recombination event involving hBoV1, with a breakpoint located in the NS1 gene. Taken together, these data suggested that intra-species recombination could occur in hBoV1 at different positions of its genome, playing a major role in the evolutionary history of this virus. Further studies will be necessary to identify breakpoint patterns in these viruses and to identify their potential association with viral spread among populations.

In conclusion, this study showed that both hMPV and hBoV circulated in Italy and were involved in the pathogenesis of lower respiratory tract diseases in children. Data presented in this report provided evidence for both a shift in the incidence of hMPV A2 strain infections in recent years and the occurrence of recombination events in hBoV strains. Therefore, continuous molecular surveillance is important for the detection of new viral strains as a means of detecting their spread and assessing their epidemiological impact.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

The Authors thank Dr. D. Colzani for skilful technical assistance.

This work was supported by the Italian Ministry of University and Research (MIUR, PRIN-2005, protocol 2005067255_003; MIUR, PRIN-2007, protocol 2007LPAF42_003)

Conflict of interest

None of the authors have financial or other conflicting interests with regard to the information described in this manuscript.

For Peer Review

REFERENCES

- Aguilar JC, Pérez-Breña MP, García ML, Cruz N, Erdman DD, Echevarría JE. 2000. Detection and identification of human parainfluenza viruses 1, 2, 3, and 4 in clinical samples of pediatric patients by multiplex reverse transcription-PCR. *J Clin Microbiol* 38:1191-1195.
- Allander T, Tammi MT, Eriksson M, Bjerkner A, Tiveljung-Lindell A, Andersson B. 2005. Cloning of a human parvovirus by molecular screening of respiratory tract samples. *Proc Natl Acad Sci USA* 102:12891-12896.
- Allander T, Jartti T, Gupta S, Niesters HGM, Lehtinen P, Österback R, Vuorinen T, Waris M, Bjerkner A, Tiveljung-Lindell A, van den Hoogen BG, Hyypiä T, Ruuskanen O. 2007. Human Bocavirus and acute wheezing in children. *Clin Infect Dis* 44:904-910.
- Arthur JL, Higgins GD, Davidson GP, Givney RC, Ratcliff RM. 2009. A novel bocavirus associated with acute gastroenteritis in Australian children. *PLoS Pathog* 5:e1000391.
- Avellón A, Pérez P, Aguilar JC, Lejarazu R, Echevarría JE. 2001. Rapid and sensitive diagnosis of human adenovirus infections by a generic polymerase chain reaction. *J Virol Met* 92:113-120.
- Bastien N, Brandt K, Dust K, Ward D, Li Y. 2006. Human Bocavirus infection, Canada. *Emerg Infect Dis* 12:848-850.
- Boivin G, Mackay I, Sloots TP, Madhi S, Freymuth F, Wolf D, Shemer-Avni Y, Ludewick H, Gray GC, LeBlanc E. 2004. Global genetic diversity of human metapneumovirus fusion gene. *Emerg Infect Dis* 10:1154-1157.
- Brieu N, Guyon G, Rodière M, Segondy M, Foulongne V. 2008. Human bocavirus infection in children with respiratory tract disease. *Pediatr Infect Dis J* 27:969-973.
- Broor S, Bharaj P, Chahar HS. 2008. Human metapneumovirus: a new respiratory pathogen. *J Biosci* 33:483-493.
- Coiras MT, Pérez-Breña P, García ML, Casas I. 2003. Simultaneous detection of influenza A, B, and C viruses, respiratory syncytial virus, and adenoviruses in clinical samples by multiplex reverse transcription nested-PCR assay. *J Med Virol* 69:132-144.

- 1
2
3 263 Dean AG, Sullivan KM, Soe MM. 2009. OpenEpi: Open Source Epidemiologic Statistics for Public
4
5
6 264 Health, Version 2.3. Update 2009/20/05. Available at: <http://www.openepi.com>
7
8 265 Dessau RB, Lisby G, Frederiksen JL. 2001. Coronaviruses in brain tissue from patients with
9
10 266 multiple sclerosis. *Acta Neuropathol* 101:601-604.
11
12
13 267 Echevarría JE, Erdman DD, Swierkosz EM, Holloway BP, Anderson LJ. 1998. Simultaneous
14
15 268 detection and identification of human parainfluenza viruses 1, 2, and 3 from clinical samples by
16
17 269 multiplex PCR. *J Clin Microbiol* 36:1388-1391.
18
19
20 270 Felsenstein J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*
21
22 271 39:783-791.
23
24
25 272 Foulongne V, Guyon G, Rodière M, Segondy M. 2006. Human metapneumovirus infection in
26
27 273 young children hospitalized with respiratory tract disease. *Pediatr Infect Dis J*. 25:354-359.
28
29 274 Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program
30
31
32 275 for windows95/98/nt. *Nucl Acids Symp* 41:95-98.
33
34 276 Huck B, Scharf G, Neumann-Haefelin D, Puppe W, Weigl J, Falcone V. 2006. Novel human
35
36 277 metapneumovirus sublineage. *Emerg Infect Dis* 12:147-150.
37
38
39 278 Ireland DC, Kent J, Nicholson KG. 1993. Improved detection of rhinoviruses in nasal and throat
40
41 279 swabs by seminested RT-PCR. *J Med Virol* 40:96-101.
42
43
44 280 Kapoor A, Slikas E, Simmonds P, Chieochansin T, Naeem A, Shaukat S, Alam MM, Sharif S,
45
46 281 Angez M, Zaidi S, Delwart E. 2009. A newly identified bocavirus species in human stool. *J*
47
48 282 *Infect Dis* 199:196-200.
49
50
51 283 Kapoor A, Simmonds P, Slikas E, Li L, Bodhidatta L, Sethabutr O, Triki H, Bahri O, Oderinde BS,
52
53 284 Baba MM, Bukbuk DN, Besser J, Bartkus J, Delwart E. 2010. Human Bocaviruses are highly
54
55 285 diverse, dispersed, recombination prone, and prevalent in enteric infections. *J Infect Dis*
56
57 286 201:1633-1643.
58
59
60 287 Kimura M. 1980. A simple method for estimating evolutionary rate of base substitutions through
288 comparative studies of nucleotide sequences. *J Mol Evol* 16:111-120.

- Lin JH, Chiu SC, Lin YC, Chen HL, Lin KH, Shan KH, Wu HS, Liu HF. 2009. Clinical and genetic analysis of Human Bocavirus in children with lower respiratory tract infection in Taiwan. *J Clin Virol* 44:219-224.
- Lindner J, Karalar L, Schimanski S, Pfister H, Struff W, Modrow S. 2008. Clinical and epidemiological aspects of human bocavirus infection. *J Clin Virol* 43:391-395.
- Lole KS, Bollinger RC, Paranjape RS, Gadkari D, Kulkarni SS, Novak NG, Ingersoll R, Sheppard HW, Ray SC. 1999. Full-length human immunodeficiency virus type 1 genome from subtype C-infected seroconverters in India, with evidence of intersubtype recombination. *J Virol* 73:152-160.
- Maggi F, Pifferi M, Vatteroni M, Fornai C, Tempestini E, Anzilotti S, Lanini L, Andreoli E, Ragazzo V, Pistello M, Specter S, Bendinelli M. 2003. Human Metapneumovirus associated with respiratory tract infections in a 3-year study of nasal swabs from infants in Italy. *J Clin Microbiol* 4:2987-2991.
- Manning A, Russell V, Eastick K, Leadbetter GH, Hallam N, Templeton K, Simmonds P. 2006. Epidemiological profile and clinical associations of human Bocavirus and other human Parvoviruses. *J Infect Dis* 194:1283-1290.
- Ohshima T, Mochizuki M. 2009. Evidence for recombination between feline panleukopenia virus and canine parvovirus type 2. *J Vet Med Sci* 71:403-408.
- Peret TC, Boivin G, Li Y, Couillard M, Humphrey C, Osterhaus AD, Erdman DD, Anderson LJ. 2002. Characterization of human metapneumoviruses isolated from patients in North America. *J Infect Dis* 185:1660-1663.
- Saitou N and Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406-425.
- Schildgen O, Müller A, Allander T, Mackay IM, Völz S, Kupfer B, Simon A. 2008. Human Bocavirus: passenger or pathogen in acute respiratory tract infections? *Clin Microbiol Rev* 21:291-304.

1
2
3 315 Shackelton LA, Hoelzer K, Parrish CR, Holmes EC. 2007. Comparative analysis reveals frequent
4
5 316 recombination in the parvoviruses. *J Gen Virol* 88:3294-3301.
6
7
8 317 Smuts H, Hardie D. 2006. Human Bocavirus in hospitalized children, South Africa. *Emerg Infect*
9
10 318 *Dis* 12:1457-1458.
11
12
13 319 Tamura K, Dudley J, Nei M, Kumar S. 2007. MEGA4: Molecular Evolutionary Genetics Analysis
14
15 320 (MEGA) software version 4.0. *Mol Biol Evol* 24:1596-1599.
16
17 321 Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. 1997. The CLUSTALX
18
19 windows interface: flexible strategies for multiple sequence alignment aided by quality analysis
20 322 tools. *Nucleic Acids Res* 25:4876-4882.
21
22 323
23
24 324 Valle L, Amicizia D, Bacilieri S, Banfi F, Riente R, Durando P, Sticchi L, Gasparini R, Esposito C,
25
26 Icardi G, Ansaldi F. 2006. Performance testing of two new one-step real time PCR assays for
27 325
28 detection of human influenza and avian influenza viruses isolated in humans and respiratory
29 326
30 syncytial virus. *J Prev Med Hyg* 47:127-133.
31
32 327
33
34 328 van den Hoogen BG, de Jong JC, Groen J, Kuiken T, de Groot R, Fouchier RAM, Osterhaus
35
36 329 ADME. 2001. A newly discovered human pneumovirus isolated from young children with
37
38 respiratory tract disease. *Nat Med* 7:719-724.
39 330
40
41 331 van den Hoogen BG, Herfst S, Sprong L, Cane PA, Forleo-Neto E, de Swart RL, Osterhaus ADME,
42
43 332 Fouchier RAM. 2004. Antigenic and genetic variability of human Metapneumoviruses. *Emerg*
44
45 *Infect Dis* 10:658-666.
46 333
47
48 334 von Linstow ML, Høgh M, Høgh B. 2008. Clinical and epidemiologic characteristics of human
49
50 bocavirus in Danish infants: results from a prospective birth cohort study. *Pediatr Infect Dis J*
51 335
52 27:897-902.
53 336
54
55 337 Wang HC, Huang SW, Wang SW, Tsai HP, Kiang D, Wang SM, Liu CC, Su IJ, Wang JR. 2008.
56
57 Co-circulating genetically divergent A2 human metapneumovirus strains among children in
58 338
59 southern Taiwan. *Arch Virol* 153:2207-2213.
60 339

1
2
3 340 World Health Organization. 2009. Acute Respiratory Infections (Update February 2009). Available
4
5 341 at: http://www.who.int/vaccine_research/diseases/ari/en/index.html
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

1
2
3 342 **FIGURE LEGENDS**

4
5 343 Fig. 1. Genetic analysis of hMPV. A) Phylogenetic tree of the Italian I-MI hMPV sequences
6
7
8 344 (**boldface**) demonstrating the phylogenetic relationship of the I-MI sequences based on partial F
9
10 345 gene nucleotide sequences. Bootstrap values (in addition to A1, A2a, A2b, B1 and B2
11
12 346 group/sublineages) are indicated. Reference sequences correspond to the following GenBank
13
14
15 347 accession numbers: A1) AF371337, AY145294, DQ453017, AJ867506, AJ867520, EF612444;
16
17 348 A2a) AY304360, AY145296, DQ453014, EF612451; A2b) AY530095, DQ453021, AJ867528,
18
19 349 AJ867538, EF612457; B1) AY304361, AY145294, DQ453015, AJ867513, EF612440; B2)
20
21 350 AY304362, AY145289, DQ452995, AJ867503, EF612441. B) Evolutionary mean distances over
22
23
24 351 all sequence pairs within and between groups.
25
26
27 352

28
29 353 Fig. 2. Genetic analysis of hBoV sequences. Phylogenetic tree of the Italian I-MI human Bocavirus
30
31 354 sequences (**boldface**) based on partial VP1/VP2 region nucleotide sequences. Bootstrap values (in
32
33
34 355 addition to St1 and St2 genotype clusters) are indicated. Reference sequences correspond to the
35
36 356 following GenBank accession numbers: St-1) DQ000497, AM160609, AM160611-AM160613,
37
38 357 AM160615, AM689298, AM689299, AM689306, AM689307, EU069434, EU069436, EU069437,
39
40
41 358 EF203921; St-2) DQ000495, AM160610, AM160614, AM689297, AM689301-AM689303,
42
43 359 AM689305, EF450717, EF450718, EF450722, EF450723, EF450736, EF450740.
44
45
46 360

47
48 361 Fig. 3. Recombination recognition of the Italian I-MI/04-08 sequence: bootscan analysis and
49
50 362 identities values between the recombinant sequence and the two reference sequence (MPT8 and
51
52
53 363 HK7) before and after the breakpoint (inset).
54
55
56
57
58
59
60