



Development of helical vortex theory

Valery Okulov

► To cite this version:

Valery Okulov. Development of helical vortex theory. EUROMECH Colloquium 525 - Instabilities and transition in three-dimensional flows with rotation, Jun 2011, Ecully, France. hal-00600414

HAL Id: hal-00600414

<https://hal.science/hal-00600414>

Submitted on 14 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



DEVELOPMENT OF HELICAL VORTEX THEORY

Valery OKULOV

Department of Mechanical Engineering and Center for Fluid Dynamics,
 Technical University of Denmark, 2800 Kgs. Lyngby, Denmark; E-mail: vaok@mek.dtu.dk

The main goal of the current communication is to attract attention to the thriving theory of helical vortices which are 2-D from Euler's consideration and looks from this point of view like "plane" flows (fig. 1a) but with additional uniform rotation of a plane pattern of the flow along fixed axis (fig. 1b). However, from Lagrange's point of view the 2-D helical vortices induce strong 3-D motion of fluid particles (fig. 1c) [1].




Figure 1. Examples of 2-D flows with rectilinear (a) and helical vortices (b); 3-D paths of fluid particles induced by single helical vortex filament in a cylinder (c).

At present the theory of 2-D helical vortex dynamics are based on various analytical components, which holds true for all values of the helix pitch, such as (i) the 2-D Biot-Savart law for helical filaments represented by Kapteyn series [2] or rewriting in a form with singularity separation [3] (fig. 2); (ii) solutions of helical vortex tubes with finite core, governed by series expansion of helical multipoles [4]; (iii) relations between the induction of vortex filaments and the self-induced velocity of helical vortex tubes [5] resulting in a closed analytical solution of the helix motion [6]; (iv) analytical representation of Goldstein's solution for the circulation of a helical vortex sheet in equilibrium [7]; (v) Kelvin's N-gon stability problem of point vortices generalized to multiple helical vortices [6, 8].




Figure 2. Isolines of the azimuthal velocity field induced in cross-section by helical vortex filament:
 (a) inaccurate simulation by finite part of the Kapteyn series with unreal singularities on cylinder supporting the helix;
 (b) accurate simulation after separation of the singularity along the helix filament in the Kapteyn series [3].




Figure 3. Two possibilities for equilibrium of helical vortex pairs and its prototypes from point vortex pairs: (a) centre or axis of rotation lies on the same hand from both point vortices; (b) centre of or axis rotation lies between ones




Figure 4. The circulation ratio β of the two helical vortices in equilibrium rotation as function of their relative proximity for different values of helical pitch h : 1.5 (dash-dotted lines); 2 (dashed lines); 5 (dotted lines); infinity (solid lines). Circle and point indicate equilibrium of the helical pairs with different levers.

An important result of the 2-D helical vortex theory is that the vorticity submitted to the helical symmetry does not vary along a trajectory of fluid particles. It coincides with conclusion of point vortex dynamics but in contrast of the plane one, where the total vortex motion consists of the mutual induction of other point vortices, for helix case in addition to an induced flow field due to induction from the other helical vortices on the surroundings a self-induction of the helical vortex should be included too. In result of this difference unexpressed applications of the 2-D helical vortex dynamics were established. For example figs. 3 and 4 demonstrate how the two helical vortices (vortex pair) with the same or opposite circulations but different levers can be in equilibrium rotation which is impossible for two point vortices. In the presentation it was also shown how the assumption of helical symmetry in the context of 2-D helical vortices can be exploited to analyze and model various cases of rotating flows.

References

- [1] Alekseenko S.V., Kuibin P.A., Okulov V.L. Theory of Concentrated Vortices: An Introduction. Springer (2007)
- [2] Hardin, J.C.: The velocity field induced by a helical vortex filament. Phys. Fluids 25, 1949–1952 (1982)
- [3] Okulov, V.L.: Determination of the velocity field induced by vortex filaments of cylindric and weak conic shapes. Russ. J. Eng. Thermophys. 5(2), 63–75 (1995)
- [4] Fukumoto, Y., Okulov, V.L.: The velocity field induced by a helical vortex tube. Phys. Fluids 17(10), 107101 (2005)
- [5] Boersma, J., Wood, D.H.: On the self-induced motion of a helical vortex. J. Fluid Mech. 384, 263–280 (1999)
- [6] Okulov, V.L.: On the stability of multiple helical vortices. J. Fluid Mech. 521, 319–342 (2004)
- [7] Goldstein, S.: On the vortex theory of screw propellers. Proc. R. Soc. Lond. A 123(792), 440–465 (1929)
- [8] Okulov, V.L., Sørensen, J.N.: Stability of helical tip vortices in a rotor far wake. J. Fluid Mech. 576, 1–25 (2007)