



HAL
open science

A see-saw S4 model for fermion masses and mixings

Davide Meloni

► **To cite this version:**

Davide Meloni. A see-saw S4 model for fermion masses and mixings. *Journal of Physics G: Nuclear and Particle Physics*, 2010, 37 (5), pp.55201. <10.1088/0954-3899/37/5/055201>. <hal-00600796>

HAL Id: hal-00600796

<https://hal.science/hal-00600796v1>

Submitted on 16 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

A See-Saw S_4 model for fermion masses and mixings

Davide Meloni ¹

*Institut für Theoretische Physik und Astrophysik ,
Universität Würzburg, D-97074 Würzburg, Germany*

Abstract

We present a supersymmetric see-saw S_4 model giving rise to the most general neutrino mass matrix compatible with Tri-Bimaximal mixing. We adopt the $S_4 \times Z_5$ flavour symmetry, broken by suitable vacuum expectation values of a small number of flavon fields. We show that the vacuum alignment is a natural solution of the most general superpotential allowed by the flavour symmetry, without introducing any soft breaking terms. In the charged lepton sector, mass hierarchies are controlled by the spontaneous breaking of the flavour symmetry caused by the vevs of one doublet and one triplet flavon fields instead of using the Froggatt-Nielsen $U(1)$ mechanism. The next to leading order corrections to both charged lepton mass matrix and flavon vevs generate corrections to the mixing angles as large as $\mathcal{O}(\lambda_C^2)$. Applied to the quark sector, the symmetry group $S_4 \times Z_5$ can give a leading order V_{CKM} proportional to the identity as well as a matrix with $\mathcal{O}(1)$ coefficients in the Cabibbo 2×2 submatrix. Higher order corrections produce non vanishing entries in the other V_{CKM} entries which are generically of $\mathcal{O}(\lambda_C^2)$.

¹e-mail address: davide.meloni@physik.uni-wuerzburg.de

1 Introduction

The Tri-Bimaximal structure (TBM) [1] of the neutrino mixing matrix is remarkably in agreement with the experimental results in the neutrino sector [2]. Within 1- σ error, the values of the mixing angles can be approximated by their TBM values [3]:

$$\tan^2 \theta_{23} = 1 \quad \tan^2 \theta_{12} = \frac{1}{2} \quad \sin \theta_{13} = 0. \quad (1)$$

The simplified structure of the mixing matrix suggested the possibility to be explained using some discrete non-abelian groups, added to the Standard Model, and containing a triplet representations to fit the number of lepton families observed in Nature. The symmetry A_4 [4, 5] (also in the context of Grand Unified theories [6]) emerged as a natural candidate because it is the smallest discrete group with triplet representation and it is sufficiently manageable to be broken differently in the charged and neutral lepton sectors, a necessary condition if we want to get a mixing matrix different from the identity. In the context of see-saw models, it is interesting to observe that the TBM structure in A_4 is generally associated with a well defined relations among the complex eigenvalues of the light neutrino mass matrix:

$$\frac{1}{m_3} = \frac{1}{m_1} - \frac{2}{m_2}, \quad (2)$$

implying that most A_4 models are quite predictive because of the reduced number of independent parameters. It should be stressed, however, that the realisation of the TBM strongly relies on the choice of symmetry breaking pattern. In fact, in the neutrino sector the group A_4 is usually broken into a subgroup generated by the matrices

$$U_{\mu-\tau} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

and

$$G = \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix},$$

which, in the basis where charged leptons are diagonal, leave invariant the most general neutrino mass matrix diagonalized by TBM:

$$m_{light} = \begin{pmatrix} x & y & y \\ y & x+v & y-v \\ y & y-v & x+v \end{pmatrix}. \quad (3)$$

It turns out that the representations of A_4 only contain G and the invariance under $U_{\mu-\tau}$ arises accidentally, as a consequence of the specific field content of the model. From this

point of view, the group S_4 [7]-[9] arises as a natural candidate as a flavour group for neutrino mixing because one can find a suitable representation of it containing simultaneously the previous elements. It should also be noted that the extension of the A_4 symmetry from lepton to quarks seems to be complicated by the absence of doublet representations, whose use is suggested by the heaviness of the top quark, which S_4 possesses instead. However, the problem of reproducing small mixing angles and strong mass hierarchy in the up-sector at the same time can only be partially alleviated by the bidimensional representations of S_4 because one or more fine tunings between the relevant Yukawas are invoked to correctly reproduce some of the m_{up}/m_{down} quark mass ratios. In this paper, we build a constrained see-saw S_4 -based model for fermion masses and mixing which, compared to models already existent in the literature, realises the most general neutrino mass matrix diagonalized by TBM at leading order (LO). This is obtained allowing the right-handed neutrinos to couple to singlet, doublet and triplet flavon fields. The light neutrino masses depend on six complex Yukawa parameters and the typical A_4 sum rule of eq.(2) does not hold, leaving the model less predictive but more manageable. The mass hierarchy among charged leptons is obtained breaking the S_4 symmetry by the vevs of a doublet and triplet flavon fields, without invoking any Froggatt-Nielsen $U(1)$ symmetry. The unwanted couplings are forbidden imposing an additional Z_5 symmetry to the model. The resulting $S_4 \times Z_5$ symmetry is minimal from the point of view of the flavour symmetry and field content. We extend the $S_4 \times Z_5$ symmetry to the quark sector including the left-handed components into triplet representations (and not into doublets, as usually done); we show that, even using a rigid structure like that proposed in this paper, some of the relevant features of the quark sector, like a good leading order V_{CKM} and quark mass ratios, can still be accounted for. The paper is organized as follows: in Sect.2, we discuss the relevant feature of the S_4 symmetry and the structure of the model, presenting leading order results on neutrino as well as charged lepton mass matrices; in Sect.3 we compute the next to leading order corrections (NLO) to the vacuum alignment and the relevant higher order operators, both responsible for deviations from TBM mixing; in Sect.4 we discuss some phenomenological results obtained from our model with all Yukawas constrained to be $\mathcal{O}(1)$ and we also show that the whole model allows for acceptable leptogenesis parameters. Sect.5 is devoted to the quark sector whereas in Sect.6 we draw our conclusions.

2 The structure of the model

We introduce here the structure of the model which leads to TBM in first approximation. We recall that S_4 , the permutation group of 4 objects, can be generated by the two elements S and T obeying the relations (a "presentation" of the group):

$$S^4 = T^3 = 1, \quad ST^2S = T. \quad (4)$$

The action of the generators S and T can be assigned as follows:

$$(1234) \xrightarrow{S} (2341)$$

$$(1234) \rightarrow^T (2314)$$

and the 24 elements of the group, belonging to 5 conjugate classes, are:

$$\mathcal{C}_1 : 1$$

$$\mathcal{C}_2 : S^2 = (3412), TS^2T^2 = (4321), S^2TS^2T^2 = (2143)$$

$$\mathcal{C}_3 : T, T^2 = (3124), S^2T = (1423), S^2T^2 = (2431), STST^2 = (4132)$$

$$STS = (4213), TS^2 = (4132), T^2S^2 = (1342)$$

$$\mathcal{C}_4 : ST^2 = (1243), T^2S = (4231), TST = (1432)$$

$$TST^2 = (3214), STS^2 = (1324), S^2TS = (2134)$$

$$\mathcal{C}_5 : S, TST^2 = (2413), ST = (3142), TS = (3421), S^3 = (4123), S^3T^2 = (4312)$$

The inequivalent irreducible representations of S_4 are 1_1 , 1_2 , 2 and 3 . It is immediate to see that one-dimensional unitary representations are given by:

$$\begin{aligned} 1_1 : S = 1 \quad T = 1 \\ 1_2 : S = -1 \quad T = 1 \end{aligned} \quad (5)$$

while the two-dimensional unitary representation, in a basis where the element T is diagonal, is given by:

$$T = \begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix}, \quad S = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (6)$$

Finally, the three-dimensional unitary representation is as follows:

$$3_1 : T = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega \end{pmatrix}, \quad S = \frac{1}{3} \begin{pmatrix} -1 & 2\omega & 2\omega^2 \\ 2\omega & 2\omega^2 & -1 \\ 2\omega^2 & -1 & 2\omega \end{pmatrix}, \quad (7)$$

where $\omega = e^{2\pi i/3} = (-1 + \sqrt{3})/2$, whereas in the 3_2 representation the generator T is the same but S is the opposite. It is useful to remind the product rules between the group representations:

$$\begin{aligned} 1_1 \otimes \xi &= \xi \\ 1_2 \otimes 1_2 &= 1_1, \quad 1_2 \otimes 2 = 2, \quad 1_2 \otimes 3_i = 3_j \\ 2 \otimes 2 &= 1_1 \oplus 1_2 \oplus 2, \quad 2 \otimes 3_i = 3_1 \oplus 3_2, \quad 3_i \otimes 3_i = 1_1 \oplus 2 \oplus 3_1 \oplus 3_2, \\ 3_1 \otimes 3_2 &= 1_2 \oplus 2 \oplus 3_2 \oplus 3_2, \end{aligned} \quad (8)$$

where the indices $i, j = 1, 2$, with $i \neq j$ and ξ indicates any other representation. The Clebsch-Gordan coefficients in the basis presented above are reported in Appendix A.

The general neutrino mass matrix of eq.(3) can be obtained in the framework of the see-saw mechanism,

$$m_{light} = -m_D^T m_M^{-1} m_D, \quad (9)$$

where both Majorana (m_M) and Dirac (m_D) mass matrices are needed. They are derived from the most general lagrangian invariant under $S_4 \times Z_5$ and containing fields in any of the S_4 representations, which are singlets, doublets and triplets. The group S_4 is broken by means of suitable vev's of Standard Model singlet fields (flavons), whose alignments have to guarantee the correct entries of the Majorana and Dirac mass matrices. Group theoretical considerations help in understanding the pattern of symmetry breaking needed to generate the wanted matrix; in the representation of App.A, the elements S^2 and $(TSTS^2)$ leave invariant the m_{light} of eq.(3), that is

$$\begin{aligned} S^2 m_{light} S^2 &= m_{light} \\ (TSTS^2) m_{light} (TSTS^2) &= m_{light}. \end{aligned}$$

This means that the vevs of the flavon fields in the neutrino sector have to be invariant under the subgroup generated by them. This strong condition for the flavon alignment is realized, for the triplet representation, by the field configuration

$$\langle \varphi_S \rangle = v_S (1, 1, 1). \quad (10)$$

For the bidimensional representation, the same matrices should leave the vev of a doublet field invariant; if we choose

$$\langle \Delta \rangle = v_\Delta (1, 1) \quad (11)$$

these correspond to the first matrix of the groups $\mathcal{C}_{4,5}$ in the doublet representation. Therefore, we have found a scalar field configuration which remains invariant under the action of the same matrices that leave m_{light} invariant. In addition to the previous fields, we also include in the model a singlet flavon field ξ , with a non-vanishing vev. To realize the classical see-saw mechanism, we need to introduce right-handed neutrinos, which we assume to transform as a triplet representation of S_4 . To avoid large fine-tuning from terms of the form $M\nu^c\nu^c$, like those discussed in [5], we properly tune the ν^c charge under Z_5 ¹; whatever this Z_5 charge is, if we want the singlet, doublet and triplet flavons to contribute to the neutrino mass matrix their charges have to be the same. We also attribute a Z_5 charge to the h_u higgs boson in order to avoid the Weinberg operator $O_5 = \ell h_u \ell h_u$ at the leading order [11], where ℓ is a triplet field of the $SU(2)$ Standard Model lepton doublets. This also forbids a leading order Dirac mass term of the form $(\nu^c \ell) h_u$, which is then generated at the $\mathcal{O}(1/\Lambda)$ through couplings with the same flavon fields ξ , Δ and φ_S . The Z_5 charges for the matter fields as well as for flavons and driving fields (discussed later) are reported in Tab.1, where we used the symbol $\omega = e^{2\pi i/5}$ to indicate the Z_5 unit-charge.

The lagrangian in the neutrino sector is then as follows:

$$W_\nu = \frac{1}{\Lambda} \nu^c \ell h_u (y_{\nu_1} \varphi_S + y_{\nu_2} \Delta + y_{\nu_3} \xi) + \nu^c \nu^c (a \xi + b \varphi_S + c \Delta), \quad (12)$$

¹Similar considerations, but using two discrete groups, have been discussed in [10].

Field	ν^c	ℓ	e^c	μ^c	τ^c	h_d	h_u	φ_T	η	Δ	φ_S	ξ	φ_0^T	φ_0^S	Δ_0	ρ_0
S_4	3	3_1	1_2	1_1	1_1	1_1	1_1	3_1	2	2	3_1	1_1	3_1	3_1	2	1_1
Z_5	ω^2	1	ω^3	ω^2	ω	1	ω^2	ω^4	ω^4	ω	ω	ω	ω^2	ω^3	ω^3	ω^2
$U(1)_R$	1	1	1	1	1	0	0	0	0	0	0	0	2	2	2	2

Table 1: Transformation properties of leptons, electroweak Higgs doublets and flavons under $S_4 \times Z_5$ and $U(1)_R$.

where a , b , c and y_{ν_i} are complex Yukawa couplings. The Dirac mass matrix is obtained from the first term in eq.(12) and it is given by:

$$m_D = \frac{v_u}{\Lambda} \begin{pmatrix} 2y_{\nu_1}v_S + y_{\nu_3}u & y_{\nu_2}v_\Delta - y_{\nu_1}v_S & y_{\nu_2}v_\Delta - y_{\nu_1}v_S \\ y_{\nu_2}v_\Delta - y_{\nu_1}v_S & 2y_{\nu_1}v_S + y_{\nu_2}v_\Delta & y_{\nu_3}u - y_{\nu_1}v_S \\ y_{\nu_2}v_\Delta - y_{\nu_1}v_S & y_{\nu_3}u - y_{\nu_1}v_S & 2y_{\nu_1}v_S + y_{\nu_2}v_\Delta \end{pmatrix} \quad (13)$$

where v_u is the vacuum expectation value of the higgs field h_u . The other terms give the Majorana mass matrix:

$$m_M = \begin{pmatrix} au + 2bv_S & -bv_S + cv_\Delta & -bv_S + cv_\Delta \\ -bv_S + cv_\Delta & 2bv_S + cv_\Delta & au - bv_S \\ -bv_S + cv_\Delta & au - bv_S & 2bv_S + cv_\Delta \end{pmatrix} \quad (14)$$

whose eigenvalues are:

$$\begin{aligned} M_1 &= au + 3bv_S - cv_\Delta \\ M_2 &= au + 2cv_\Delta \\ M_3 &= -au + 3bv_S + cv_\Delta. \end{aligned} \quad (15)$$

Using eq.(9), we can derive the light neutrino mass matrix, diagonalized by tri-bimaximal mixing, whose eigenvalues are:

$$\begin{aligned} m_1 &= -\left(\frac{v_u}{\Lambda}\right)^2 \frac{(3y_{\nu_1}v_S - y_{\nu_2}v_\Delta + y_{\nu_3}u)^2}{au + 3bv_S - cv_\Delta} \\ m_2 &= -\left(\frac{v_u}{\Lambda}\right)^2 \frac{(2y_{\nu_2}v_\Delta + y_{\nu_3}u)^2}{au + 2cv_\Delta} \\ m_3 &= \left(\frac{v_u}{\Lambda}\right)^2 \frac{(3y_{\nu_1}v_S + y_{\nu_2}v_\Delta - y_{\nu_3}u)^2}{au - 3bv_S - cv_\Delta}. \end{aligned} \quad (16)$$

We see that the neutrino masses depend on six unrelated complex Yukawa parameters, which offer more freedom to tune mass differences and then recover the phenomenology associated to neutrino oscillation. Notice also that no sum rules can be found in this case among complex eigenvalues.

2.1 Charged leptons

It could be easier to work in a basis where the charged lepton mass matrix is diagonal. In order to understand how this naturally arises in an S_4 -based model, we observe that a generic diagonal matrix m_l^D (with diagonal entries different to each other) is left invariant under the action of an element A of S_4 only if such an element is itself diagonal, with different phase factors at each diagonal entry, as it can be understood requiring that the relation

$$A^\dagger m_l^{D\dagger} m_l^D A = m_l^{D\dagger} m_l^D \quad (17)$$

is satisfied. The generator T , for example, is such an appropriate matrix. Since the charged lepton matrix is generated after spontaneous symmetry breaking, one could choose flavon fields with vevs invariant under the action of T . Instead, we prefer the choice

$$\langle \varphi_T \rangle = v_T (0, 1, 0) \quad (18)$$

which, even breaking completely the group S_4 , not only guarantees the diagonal form of the mass matrix but also generates the hierarchy among lepton families without introducing any additional $U(1)_{FN}$ symmetry². In the same way, we choose the vev for the doublet flavon η as:

$$\langle \eta \rangle = v_\eta (0, 1); \quad (19)$$

the corresponding lagrangian in the charged lepton sector is as follows:

$$\begin{aligned} \mathcal{L} = & \frac{y_\tau}{\Lambda} \tau^c (\ell \varphi_T) h_d + \\ & \frac{y_{\mu_1}}{\Lambda^2} \mu^c (\ell \varphi_T \varphi_T) h_d + \frac{y_{\mu_2}}{\Lambda^2} \mu^c \ell (\eta \varphi_T) h_d + \\ & \frac{y_{e_1}}{\Lambda^3} e^c \ell [\varphi_T (\varphi_T \varphi_T)_2]_{3_2} h_d + \frac{y_{e_2}}{\Lambda^3} e^c \ell [\varphi_T (\varphi_T \varphi_T)_{3_1}]_{3_2} h_d + \\ & \frac{y_{e_3}}{\Lambda^3} e^c \ell [\eta (\varphi_T \varphi_T)_{3_1}]_{3_2} h_d + \frac{y_{e_4}}{\Lambda^3} e^c \ell [\varphi_T (\eta \eta)_2]_{3_2} h_d + \\ & \frac{y_{e_5}}{\Lambda^2} e^c \ell (\Delta \varphi_S)_{3_2} h_d. \end{aligned} \quad (20)$$

It is interesting to observe that the last term in the lagrangian would be the dominant one and would drive the electron mass to a too large value³. However, the leading order structures of the vacua in eqs.(10)-(11) prevent this term to appear. We will show later that this is not the case when the next to leading order corrections to the flavon alignment

²This works in the same way as described in [5] noticing that, like in the case of A_4 , $(0, 1, 0)^2 = (0, 0, 1)$ and $(0, 1, 0)^3 = (1, 0, 0)$.

³This is because this term explicitly breaks the residual symmetries needed to generate the correct hierarchies between the charged lepton masses.

are taken into account. After symmetry breaking, the mass matrix has the form:

$$m_\ell = \frac{v_d v_T}{\Lambda} \begin{pmatrix} \frac{1}{\Lambda^2} [v_T^2 (y_{e1} + 2 y_{e2}) - 2 v_T v_\eta y_{e3} + v_\eta^2 y_{e4}] & 0 & 0 \\ 0 & \frac{1}{\Lambda} (2 y_{\mu 1} v_T + y_{\mu 2} v_\eta) & 0 \\ 0 & 0 & y_\tau \end{pmatrix}, \quad (21)$$

where $v_d = \langle h_d \rangle$. To estimate the order of magnitude of v_T and v_η , we can use the experimental informations on the ratios of lepton masses. Assuming that the combinations of the y coefficients are of all of $\mathcal{O}(1)$, one obtains:

$$\begin{aligned} \left(\frac{m_\mu}{m_\tau} \right) &\sim 2 \varepsilon_T + \varepsilon_\eta \simeq 0.06 \\ \left(\frac{m_e}{m_\tau} \right) &\sim 3 \varepsilon_T^2 - 2 \varepsilon_T \varepsilon_\eta + \varepsilon_\eta^2 \simeq 0.0003 \end{aligned}$$

where we introduced the small quantities

$$\varepsilon_T = v_T/\Lambda \quad \varepsilon_\eta = v_\eta/\Lambda.$$

These relations are satisfied for:

$$(|\varepsilon_T|, |\varepsilon_\eta|) \sim (0.017, 0.029). \quad (22)$$

so that we can roughly assume that both ε_T and ε_η are of the same order of magnitude, $\varepsilon \sim \mathcal{O}(\lambda_C^2)$.

2.2 Superpotential and vacuum alignment

The most general driving superpotential w_d invariant under $S_4 \times Z_5$ with $R = 2$ is given by

$$\begin{aligned} w_d &= g_1 (\varphi_0^S \varphi_S \varphi_S) + g_2 (\varphi_0^S \varphi_S) \xi + g_3 \varphi_0^S (\varphi_S \Delta) + \\ &g_4 \Delta_0 (\Delta \Delta) + g_5 \Delta_0 (\varphi_S \varphi_S) + g_6 \Delta_0 \Delta \xi + \\ &h_1 \varphi_0^T (\varphi_T \varphi_T) + h_2 \varphi_0^T (\eta \varphi_T) + r_1 \rho_0 (\varphi_T \varphi_T) + r_2 \rho_0 (\eta \eta) \\ &= w_d^{LO}(\varphi_0^S, \Delta_0) + w_d^{LO}(\varphi_0^T, \rho_0) \end{aligned} \quad (23)$$

where all possible contractions among flavon fields are understood. The equations which fix the components of the vevs of the various flavon fields are obtained solving a system of equations obtained deriving w_d with respect to the component of the driving fields. The charge assignment reported in Tab.1 allows to separate the equations into two sets of independent relations among the fields appearing in the neutrino sector and in the charged one, respectively. For the latter we have:

$$\begin{aligned}
\frac{\partial w_d}{\partial \varphi_{01}^T} &= 2h_1(\varphi_{T1}^2 - \varphi_{T2}\varphi_{T3}) + h_2(\eta_1\varphi_{T2} + \eta_2\varphi_{T3}) = 0 \\
\frac{\partial w_d}{\partial \varphi_{02}^T} &= 2h_1(\varphi_{T2}^2 - \varphi_{T1}\varphi_{T3}) + h_2(\eta_1\varphi_{T1} + \eta_2\varphi_{T2}) = 0 \\
\frac{\partial w_d}{\partial \varphi_{03}^T} &= 2h_1(\varphi_{T3}^2 - \varphi_{T1}\varphi_{T2}) + h_2(\eta_1\varphi_{T3} + \eta_2\varphi_{T1}) = 0 \\
\frac{\partial w_d}{\partial \rho_0} &= r_1(\varphi_{T1}^2 + 2\varphi_{T2}\varphi_{T3}) + 2r_2\eta_1\eta_2,
\end{aligned} \tag{24}$$

whose solutions are:

$$\langle \eta \rangle = v_\eta(0, 1), \quad \langle \varphi_T \rangle = v_T(0, 1, 0), \quad v_\eta = -2 \left(\frac{h_1}{h_2} \right) v_T. \tag{25}$$

It is important to observe that the last equation is crucial to avoid another solution of the form $\langle \eta \rangle = (1, -1)$, $\langle \varphi_T \rangle = (1, 1, 1)$. We also recover the relation between the vevs v_T and v_η of eq.(22). In the neutrino sector, the set of equations read as follows:

$$\begin{aligned}
\frac{\partial w_d}{\partial \varphi_{01}^S} &= 2g_1(\varphi_{S1}^2 - \varphi_{S2}\varphi_{S3}) + g_2\xi\varphi_{S1} + g_3(\Delta_1\varphi_{S2} + \Delta_2\varphi_{S3}) = 0 \\
\frac{\partial w_d}{\partial \varphi_{02}^S} &= 2g_1(\varphi_{S2}^2 - \varphi_{S1}\varphi_{S3}) + g_2\xi\varphi_{S3} + g_3(\Delta_1\varphi_{S1} + \Delta_2\varphi_{S2}) = 0 \\
\frac{\partial w_d}{\partial \varphi_{03}^S} &= 2g_1(\varphi_{S3}^2 - \varphi_{S1}\varphi_{S2}) + g_2\xi\varphi_{S2} + g_3(\Delta_1\varphi_{S3} + \Delta_2\varphi_{S1}) = 0 \\
\frac{\partial w_d}{\partial \Delta_{01}} &= g_4\Delta_1^2 + g_5(\varphi_{S3}^2 + 2\varphi_{S1}\varphi_{S2}) + g_6\Delta_2\xi = 0 \\
\frac{\partial w_d}{\partial \Delta_{02}} &= g_4\Delta_2^2 + g_5(\varphi_{S2}^2 + 2\varphi_{S1}\varphi_{S3}) + g_6\Delta_1\xi = 0.
\end{aligned} \tag{26}$$

The system is solved by:

$$\langle \xi \rangle = u, \quad \langle \Delta \rangle = v_\Delta(1, 1), \quad \langle \varphi_S \rangle = v_S(1, 1, 1), \tag{27}$$

with the additional relations

$$v_\Delta = -\frac{g_2 u}{2g_3} \tag{28}$$

$$v_S^2 = \left(\frac{2g_2 g_3 g_6 - g_2^2 g_4}{12g_5 g_3^2} \right) u^2 \tag{29}$$

and u undetermined. We have explicitly checked that the solutions of the vacuum alignment equations are unique; in fact, requiring for the generic field Φ^{LO} to be shifted as $\Phi^{LO} + \delta\Phi$, we found that the components $\delta\Phi$ are all in the same directions of the corresponding Φ^{LO} ; thus, we do not need to introduce any soft term to drive the superpotential into the wanted minimum.

3 Next to leading order

The discussion of the corrections to the previous results starts with a study of the next to leading order structure of the vacuum alignments of the flavon fields. It will turn out that such corrections will be enough to guarantee deviation from TBM at a level compatible with the recent experimental results.

3.1 Corrections to the vacuum alignment

The next to leading order terms mix the charged lepton and neutrino sectors in a non-trivial way. We want to find perturbations of (25,27) of the form

$$\begin{aligned}
\langle \varphi_S \rangle &= (v_S + \delta v_{S_1}, v_S + \delta v_{S_2}, v_S + \delta v_{S_3}) \\
\langle \varphi_T \rangle &= (\delta v_{T_1}, v_T + \delta v_{T_2}, v_T + \delta v_{T_3}) \\
\langle \Delta \rangle &= (v_\Delta + \delta v_{\Delta_1}, v_\Delta + \delta v_{\Delta_2}) \\
\langle \eta \rangle &= (\delta v_{\eta_1}, v_\eta + \delta v_{\eta_2}) \\
\langle \xi \rangle &= u + \delta u.
\end{aligned} \tag{30}$$

On a general ground, we have eleven unknowns but only nine equations (six from the two triplets, two from the doublet and one from the singlet); then, we expect two of the previous shifts to remain unconstrained. To better understand the output of such an analysis, we study the (φ_0^S, Δ_0) and (φ_0^T, ρ_0) sectors separately.

3.1.1 The (φ_0^S, Δ_0) sector

This sector is responsible for the alignment of the fields φ_S and Δ . At order $\mathcal{O}(1/\Lambda)$, the three-field terms entering the superpotential are combination of φ_T and η ; collecting these terms, we have:

$$\delta w_d(\varphi_0^S) = \sum_{i=1}^3 \frac{S_i}{\Lambda} \varphi_0^S (\varphi_T \varphi_T \varphi_T)_i + \sum_{i=4}^5 \frac{S_i}{\Lambda} \varphi_0^S (\varphi_T \eta \eta)_i + \frac{S_6}{\Lambda} \varphi_0^S (\varphi_T \varphi_T \eta)_i \tag{31}$$

$$\delta w_d(\Delta_0) = \sum_{i=1}^2 \frac{\delta_i}{\Lambda} \Delta_0 (\eta \eta \eta)_i + \sum_{i=3}^4 \frac{\delta_i}{\Lambda} \Delta_0 (\varphi_T \varphi_T \eta)_i + \frac{\delta_5}{\Lambda} \Delta_0 (\varphi_T \varphi_T \varphi_T)_i \tag{32}$$

where, in both cases, the index i of the trilinear terms represents different S_4 contractions. These NLO corrections have to be evaluated with the leading order vev's in eq.(25), so that they do not contain any of the unknowns under investigation. In particular, the structure of the vacua (25) produces

$$\delta w_d(\Delta^0) = 0;$$

and the total LO+NLO part of the superpotential responsible for the alignment of φ_S and Δ is given by:

$$w_d(\varphi_0^S, \Delta_0) = w_d^{LO}(\varphi_0^S, \Delta_0) + \delta w_d(\varphi_0^S).$$

Symmetry arguments allow to understand the structure of the solutions, whose detailed expressions can be found by explicitly solving the system of equations according to the procedure of Sect.(2.2). In fact, we see that, after symmetry breaking, the non-vanishing terms in $\delta w_d(\varphi_0^S)$ are proportional to the vector $(1, 0, 0)$, which is left invariant by the S_4 element $(TSTS^2)$; this has a $2 \leftrightarrow 3$ symmetry which forces the new vev of φ_S to have the form $(v_S + \delta v_{S_1}, v_S + \delta v_{S_2}, v_S + \delta v_{S_2})$. At the same time, the shifts δv_{Δ_i} should all be equal because of the vanishing NLO terms $\delta w_d(\Delta^0)$. An explicit computation confirms these speculations and we can write the new vacua in the following form:

$$\varphi_S = v_S \begin{pmatrix} 1 + A_S \left(\frac{\varepsilon}{\varepsilon'}\right)^2 \varepsilon \\ 1 + B_S \left(\frac{\varepsilon}{\varepsilon'}\right)^2 \varepsilon \\ 1 + B_S \left(\frac{\varepsilon}{\varepsilon'}\right)^2 \varepsilon \end{pmatrix} \quad \Delta = v_\Delta \begin{pmatrix} 1 + A_\Delta \left(\frac{\varepsilon}{\varepsilon'}\right)^2 \varepsilon \\ 1 + A_\Delta \left(\frac{\varepsilon}{\varepsilon'}\right)^2 \varepsilon \end{pmatrix} \quad (33)$$

where the coefficients A_S , B_S and A_Δ are linear combinations of leading and next to leading order coefficients and, for the sake of simplicity, we have introduced the small parameter $\varepsilon' = \langle \varphi_S \rangle / \Lambda \sim \langle \Delta \rangle / \Lambda \sim \langle \xi \rangle / \Lambda$. Notice that the three δv_{S_i} also depend on the undetermined parameter δu with identical coefficients, so that they can be reabsorbed in the leading order result.

3.1.2 The (φ_0^T, ρ_0) sector

This sector is responsible for the alignment of the fields φ_T and η . At order $\mathcal{O}(1/\Lambda)$, the new terms in the superpotential are combinations of φ_S , Δ and ξ ; we have:

$$\begin{aligned} \delta w_d(\varphi_0^T) &= \sum_{i=1}^3 \frac{t_i}{\Lambda} \varphi_0^T (\varphi_S \varphi_S \varphi_S)_i + \sum_{i=4}^5 \frac{t_i}{\Lambda} \varphi_0^T (\varphi_S \Delta \Delta)_i + \\ &\frac{t_6}{\Lambda} \varphi_0^T (\varphi_S \varphi_S \Delta) + \frac{t_7}{\Lambda} \varphi_0^T (\varphi_S \varphi_S) \xi + \frac{t_8}{\Lambda} \varphi_0^T \varphi_S \xi^2 + \frac{t_9}{\Lambda} \varphi_0^T (\varphi_S \Delta) \xi \end{aligned} \quad (34)$$

and

$$\begin{aligned} \delta w_d(\rho_0) &= \frac{\rho_1}{\Lambda} \rho_0 (\varphi_S \varphi_S \varphi_S) + \frac{\rho_2}{\Lambda} \rho_0 (\Delta \varphi_S \varphi_S) + \frac{\rho_3}{\Lambda} \rho_0 (\varphi_S \varphi_S) \xi + \\ &\frac{\rho_4}{\Lambda} \rho_0 (\Delta \Delta \Delta) + \frac{\rho_5}{\Lambda} \rho_0 (\Delta \Delta) \xi + \frac{\rho_5}{\Lambda} \rho_0 \xi^3. \end{aligned} \quad (35)$$

In this case, the corrections to φ_0^T do not conserve any of the S_4 basis elements and we expect that the components of $\langle \varphi_T \rangle$ will point to different directions. Also the $\langle \eta \rangle$'s vev is not preserved at the next to leading order and, consequently, the shifts δv_{η_i} are different from zero and different to each other. We choose to treat δv_{η_2} as the second undetermined parameters, so that the new vacua can be cast in the following form:

$$\varphi_T = v_T \begin{pmatrix} A_T \left(\frac{\varepsilon'}{\varepsilon}\right)^2 \varepsilon' \\ 1 + B_T \left(\frac{\varepsilon'}{\varepsilon}\right)^2 \varepsilon' \\ C_T \left(\frac{\varepsilon'}{\varepsilon}\right)^2 \varepsilon' \end{pmatrix} \quad \eta = v_\eta \begin{pmatrix} A_\eta \left(\frac{\varepsilon'}{\varepsilon}\right)^2 \varepsilon' \\ 1 + B_\eta \delta v_{\eta_2} \end{pmatrix}. \quad (36)$$

Notice that a correction proportional to δv_{η_2} also appears in the second component of φ_T in the form $\varphi_{T_2} = v_T - (h_2/2h_1) \delta v_{\eta_2} + \mathcal{O}(\varepsilon')$ which, using the last relation in eq.(25), gives $\varphi_{T_2} = -(h_2/2h_1) (v_T + \delta v_{\eta_2}) + \mathcal{O}(\varepsilon')$, so that the shift δv_{η_2} can be readorbed into a redefinition of v_T .

3.2 Charged lepton mass matrix

The next to leading order corrections to the mass matrix of eq.(21) come from the corrections to the vacuum alignments in eqs.(33) and (36) and from higher dimensional operators, suppressed by a relative $\mathcal{O}(1/\Lambda)$ with respect to each of the terms quoted in eq.(20). In the following we will study them separately.

3.2.1 Corrections from vacuum alignment

The main features of these corrections are related to the fact that all the vanishing entries in the matrix of eq.(21) are suppressed by one additional $\mathcal{O}(1/\Lambda)$ factor compared to the diagonal entries, except for the first line, where the $\mathcal{O}(1/\Lambda^2)$ operator in eq.(20) is non vanishing. As a result, we get the following charged lepton mass matrix at the NLO:

$$m_\ell = v_d \begin{pmatrix} a_1 \varepsilon^3 & a_2 \varepsilon \varepsilon'^2 & -a_2 \varepsilon \varepsilon'^2 \\ b_1 \varepsilon'^3 & b_2 \varepsilon^2 & b_3 \varepsilon'^3 \\ c_1 \frac{\varepsilon'^3}{\varepsilon} & c_2 \frac{\varepsilon'^3}{\varepsilon} & c_3 \varepsilon \end{pmatrix} \quad (37)$$

where the coefficients a_i, b_i and c_i can be easily reconstructed from eqs.(20) and (36). The matrix $m_\ell^\dagger m_\ell$ can be diagonalized by the unitary transformation

$$U_\ell = \begin{pmatrix} 1 & \left(\frac{b_1}{b_2} \frac{\varepsilon'^3}{\varepsilon^2}\right)^* & \left(\frac{c_1}{c_3} \frac{\varepsilon'^3}{\varepsilon^2}\right)^* \\ -\frac{b_1}{b_2} \frac{\varepsilon'^3}{\varepsilon^2} & 1 & \left(\frac{c_2}{c_3} \frac{\varepsilon'^3}{\varepsilon^2}\right)^* \\ -\frac{c_1}{c_3} \frac{\varepsilon'^3}{\varepsilon^2} & -\frac{c_2}{c_3} \frac{\varepsilon'^3}{\varepsilon^2} & 1 \end{pmatrix}. \quad (38)$$

so that, at the lowest order in ε' , the coefficients of the electron row in eq.(37) do not contribute. From the matrix in eq.(38), we can compute the U_{PMNS} mixing matrix from the lepton sector only:

$$U_{\text{PMNS}} = U_\ell^\dagger U_{\text{TBM}} = \begin{pmatrix} \sqrt{\frac{2}{3}} + \frac{1}{\sqrt{6}} (a^* + b^*) \left(\frac{\varepsilon'^3}{\varepsilon^2}\right)^* & \frac{1}{\sqrt{3}} - \frac{1}{\sqrt{3}} (a^* + b^*) \left(\frac{\varepsilon'^3}{\varepsilon^2}\right)^* & \frac{1}{\sqrt{2}} (a^* - b^*) \left(\frac{\varepsilon'^3}{\varepsilon^2}\right)^* \\ -\sqrt{\frac{1}{6}} + \sqrt{\frac{2}{3}} a \left(\frac{\varepsilon'^3}{\varepsilon^2}\right) + \frac{1}{\sqrt{6}} c^* \left(\frac{\varepsilon'^3}{\varepsilon^2}\right)^* & \sqrt{\frac{1}{3}} + \frac{1}{\sqrt{3}} a \left(\frac{\varepsilon'^3}{\varepsilon^2}\right) - \frac{1}{\sqrt{3}} c^* \left(\frac{\varepsilon'^3}{\varepsilon^2}\right)^* & -\sqrt{\frac{1}{2}} - c^* \left(\frac{\varepsilon'^3}{\varepsilon^2}\right)^* \\ -\sqrt{\frac{1}{6}} + \frac{1}{\sqrt{6}} (2b - c) \left(\frac{\varepsilon'^3}{\varepsilon^2}\right) & \sqrt{\frac{1}{3}} + (b + c) \left(\frac{\varepsilon'^3}{\varepsilon^2}\right) & \sqrt{\frac{1}{2}} - \sqrt{\frac{1}{2}} c \left(\frac{\varepsilon'^3}{\varepsilon^2}\right) \end{pmatrix} \quad (39)$$

where $a = b_1/b_2$, $b = c_1/c_3$ and $c = c_2/c_3$. Then, any entries of U_{TBM} get corrected by terms of $\mathcal{O}(\varepsilon'^3/\varepsilon^2)$ which can be as large as $\mathcal{O}(\lambda_C^2)$ to fit the experimental data.

3.2.2 Corrections from higher dimensional operators

For the electron case, there are many of such operators, coming from the following contractions:

$$\begin{aligned}
& (\Delta^3 \varphi_T) (\Delta^3 \eta) (\varphi_S^3 \varphi_T) (\varphi_S^3 \eta) (\Delta^2 \varphi_T \varphi_S) \\
& (\Delta^2 \varphi_T \xi) (\Delta^2 \varphi_S \eta) (\Delta^2 \xi \eta) (\varphi_S^2 \varphi_T \Delta) (\varphi_S^2 \varphi_T \xi) \\
& (\varphi_S^2 \Delta \eta) (\xi^2 \varphi_T \Delta) (\xi^2 \varphi_S \eta) (\xi^2 \varphi_T \varphi_S) (\xi^2 \Delta \varphi_S) .
\end{aligned} \tag{40}$$

It is easy to understand that they contribute to the electron row by terms of order $\mathcal{O}(1/\Lambda^4)$, which are unimportant because they do not sizebly modify the diagonalizing matrix U_ℓ . Also, the corrections to the τ row are of $\mathcal{O}(1/\Lambda^3)$, that is the coefficients b and c appearing in the U_{PMNS} are modified at the next to next to leading order (NNLO) and can be safely neglected. For the muon case, the following higher dimensional operators modify the coefficient a appearing in eq.(39) at the NLO :

$$\begin{aligned}
\mathcal{L}^{NLO} = & \sum_{i=1}^2 \frac{y_{\mu_i}^{NLO}}{\Lambda^3} \mu^c \ell (\varphi_S \varphi_S \varphi_S) h_d + \sum_{i=3}^4 \frac{y_{\mu_i}^{NLO}}{\Lambda^3} \mu^c \ell (\varphi_S \Delta \Delta) h_d + \\
& \frac{y_{\mu_5}^{NLO}}{\Lambda^3} \mu^c \ell \varphi_S \xi^2 + \frac{y_{\mu_5}^{NLO}}{\Lambda^3} \mu^c \ell (\varphi_S \Delta) \xi .
\end{aligned} \tag{41}$$

In conclusion, both types of corrections (from vacuum alignment and from higher dimensional operators) to the charged lepton mass matrix contribute to generate deviation from TBM at $\mathcal{O}(\lambda_C^2)$.

3.3 Neutrinos

Also for neutrinos, we have to take into account corrections from both vacuum alignment and higher dimensional operators. In particular, the NLO corrections of the vevs of the flavon fields affect both Majorana and Dirac masses with the same pattern, as we can see from eq.(12). Different higher order corrections are generated by higher order operators. These corrections turn out to be negligible compared with the NLO and of the same order of magnitude as those induced by the Weinberg operator (see later).

3.3.1 Corrections to the Majorana and Dirac mass matrices

The relevant contributions come from the NLO vacuum alignments. In particular, eq.(14) is modified only by $\langle \varphi_S \rangle$ at the NLO because the other two vevs, namely those of Δ and ξ entering in the lagrangian in eq.(12), are aligned along the LO direction. However, the shifts in $\langle \varphi_S \rangle$ are not enough to introduce independent corrections to all the (six) elements of the Majorana and Dirac mass matrices; in fact, the explicit structure of the

NLO corrections are as follows:

$$\delta m_M = b \left(\frac{\varepsilon}{\varepsilon'} \right)^2 \varepsilon v_S \hat{S} \quad (42)$$

$$\delta m_D = v_u \left(\frac{\varepsilon}{\varepsilon'} \right) \varepsilon^2 y_{\nu_1} \hat{S} \quad (43)$$

where the common matrix \hat{S} is:

$$\hat{S} = \begin{pmatrix} 2A_S & -B_S & -B_S \\ -B_S & 2B_S & -A_S \\ -B_S & -A_S & 2B_S \end{pmatrix}. \quad (44)$$

With these results, we can build the light neutrino mass matrix; since the NLO matrix elements have now complicated expressions in terms of the leading order Yukawas of eq.(12) and of the correction coefficients A_S and B_S , we prefer to summarize its structure as follows:

$$m_{light} = \varepsilon'^2 \begin{pmatrix} x & y & y \\ y & x+v & y-v \\ y & y-v & x+v \end{pmatrix} + \varepsilon^3 \begin{pmatrix} x' & y' & y' \\ y' & z & z' \\ y' & z' & z \end{pmatrix}, \quad (45)$$

where the relation $z+z' \neq x'+y'$ implies that m_{light} is not diagonalized by U_{TBM} . However, it is easy to show that the almost symmetric structure of the ε^3 contribution prevents to generate corrections to the TBM values of θ_{13} and θ_{23} because the diagonalization of m_{light} is achieved by a matrix $U_\nu = U_{TBM} + \delta U$, where δU has a vanishing last column. If we want to obtain a non-vanishing θ_{13} , we need to distinguish the elements (12) from (13) and/or (22) from (33) in eq.(44). Without taking into account the corrections from the charged lepton sector, this can be accomplished only at the NNLO. In fact, a Z_5 singlet built with the $\nu^c \nu^c$ bilinear requires a charge ω for the accompanying fields, and this can be achieved with at least three fields, which induce terms of $\mathcal{O}(1/\Lambda^2)$. There are more than twenty of these terms, generated by the following contractions:

$$\begin{aligned} & (\Delta^2 \varphi_T) (\Delta^2 \eta) (\varphi_S^2 \varphi_T) (\varphi_S^2 \eta) (\xi^2 \varphi_T) (\xi^2 \eta) \\ & (\varphi_T \Delta \varphi_S) (\varphi_T \Delta \xi) (\varphi_T \varphi_S \xi) (\Delta \eta \varphi_S) (\Delta \eta \xi) (\varphi_S \eta \xi). \end{aligned}$$

Similarly, the Dirac mass matrix receives corrections at the same NNLO relative to the leading terms due to the fact that the term $(\nu^c \ell) h_u$ has a total $Z_5 = \omega^4$.

Beside the previous operators, we also have to consider effective terms of the form $llh_u h_u$; given its charge assignment ($Z_5 = \omega^4$), the Weinberg operator arises at $\mathcal{O}(1/\Lambda^2)$ with the insertion of one flavon field, as for the Dirac and Majorana terms:

$$W_\nu^{\text{eff}} = \frac{1}{\Lambda^2} (\alpha_1 \varphi_S + \alpha_2 \Delta + \alpha_3 \xi) (\ell h_u \ell h_u). \quad (46)$$

After spontaneous symmetry breaking, W_ν^{eff} generates terms which are of order:

$$m_W \sim \frac{v_u^2 \langle \Phi \rangle}{\Lambda^2} \sim \varepsilon' \left(\frac{v_u}{\Lambda} \right) v_u. \quad (47)$$

Compared to the NLO corrections of the Dirac mass term, eq.(43), we see that:

$$m_W / \delta m_D \sim \frac{v_u / \Lambda}{\varepsilon} \ll 1;$$

then this type of operators is more suppressed with respect to the Dirac mass corrections (and even more if compared with the Majorana mass terms) and can be safely neglected.

In conclusion, from the neutrino sector only, the following mixing matrix arises:

$$U_\nu = \begin{pmatrix} \sqrt{\frac{2}{3}} - \sqrt{\frac{2}{3}} \frac{(x'+y'-z-z')^*}{9y^*} \left(\frac{\varepsilon^3}{\varepsilon'^2} \right)^* & \frac{1}{\sqrt{3}} + 2 \frac{(x'+y'-z-z')}{9\sqrt{3}y} \left(\frac{\varepsilon^3}{\varepsilon'^2} \right) & 0 \\ -\sqrt{\frac{1}{6}} - \sqrt{\frac{2}{3}} \frac{(x'+y'-z-z')^*}{9y^*} \left(\frac{\varepsilon^3}{\varepsilon'^2} \right)^* & \sqrt{\frac{1}{3}} - \frac{(x'+y'-z-z')}{9\sqrt{3}y} \left(\frac{\varepsilon^3}{\varepsilon'^2} \right) & -\sqrt{\frac{1}{2}} \\ -\sqrt{\frac{1}{6}} - \sqrt{\frac{2}{3}} \frac{(x'+y'-z-z')^*}{9y^*} \left(\frac{\varepsilon^3}{\varepsilon'^2} \right)^* & \sqrt{\frac{1}{3}} - \frac{(x'+y'-z-z')}{9\sqrt{3}y} \left(\frac{\varepsilon^3}{\varepsilon'^2} \right) & \sqrt{\frac{1}{2}} \end{pmatrix} \quad (48)$$

3.3.2 Mixing angles at the NLO

From the previous discussions, it clearly appears that the NLO corrections to θ_{13} and θ_{23} come only from U_ℓ whereas θ_{12} is modified by both U_ℓ and U_ν . They are as follows:

$$\begin{aligned} s_{13} &= |U_{e3}| = \left| \frac{1}{\sqrt{2}} (A - B)^* \left(\frac{\varepsilon'^3}{\varepsilon^2} \right)^* \right| \\ s_{12} &= \frac{|U_{e2}|}{\sqrt{1 - |U_{e3}|^2}} = \left| \frac{1}{\sqrt{3}} - \frac{1}{\sqrt{3}} (A + B)^* \left(\frac{\varepsilon'^3}{\varepsilon^2} \right)^* - \sqrt{2}\alpha \left(\frac{\varepsilon^3}{\varepsilon'^2} \right) \right| \\ s_{23} &= \frac{|U_{\mu 3}|}{\sqrt{1 - |U_{e3}|^2}} = \left| \frac{1}{\sqrt{2}} + C^* \left(\frac{\varepsilon'^3}{\varepsilon^2} \right)^* \right| \end{aligned} \quad (49)$$

where we used the short-hand notation

$$\alpha = -\sqrt{\frac{2}{3}} \frac{(x' + y' - z - z')}{9y}.$$

In particular, the first relation can be used to put a bound on $|\varepsilon'|$; in fact, given the maximum allowed value for θ_{13} and assuming that $|A - B| \sim \mathcal{O}(1)$ we get:

$$|\varepsilon'| \lesssim \left[\sqrt{2} \lambda_C^4 \theta_{13}^{\text{max}} \right]^{1/3} \quad (50)$$

which is close to $(2 \lambda_C^2)$ for $\theta_{13}^{\text{max}} \sim 10^\circ$.

4 A bit of phenomenology

The model we have presented has a huge parameter space, made by the six complex Yukawa couplings y_{ν_i} and a, b and c . It is clear, and we have checked this numerically, that the various phenomenological constraints, coming for example from the smallness of the parameter $r = \Delta m_{sol}^2 / |\Delta m_{atm}^2|$, are easily satisfied, for both type on neutrino mass hierarchies. It seems then more interesting to ask whether the model can still give an acceptable phenomenology in some particular cases, like for example allowing all the Yukawa couplings to be of $\mathcal{O}(1)$ and the flavon vevs $\langle \varphi_S \rangle, \langle \Delta \rangle$ and $\langle \xi \rangle$ to be of the same order of magnitude⁴. In this case, we do not expect any huge hierarchies among the heavy neutrinos, see eq.(15). Our numerical study aims to predict some interesting physical quantities only imposing the following 3- σ experimental constraints [13]:

$$\begin{aligned} \Delta m_{sol}^2 &> 0 \\ |\Delta m_{atm}^2| &= 2.41 \pm 0.34 \times 10^{-3} \text{ eV}^2 \\ r^{\text{exp}} &= 0.032 \pm 0.006 , \end{aligned} \quad (51)$$

also taking $|m_i| \lesssim 0.5 \text{ eV}$. The resulting spectrum of the light neutrino masses and their sum is shown in Fig.(1). In the left panel, we chose to plot the ratio $|m_3|/|m_2|$




Figure 1: *Neutrino mass spectrum and sum of neutrino masses, as predicted from the model with all Yukawa couplings of $\mathcal{O}(1)$ and the flavon vevs of the same order of magnitude. Left panel: behaviour of the ratio $|m_3|/|m_2|$ as a function of the lightest neutrino mass $|m_{\text{lightest}}|$, for both neutrino hierarchies. Right panel: sum of the light neutrino masses as a function of $|m_{\text{lightest}}|$. Also shown the bound from [16] (upper solid lines) and from [16] + [17] (lower solid lines). The vertical line in both panels is the future sensitivity of 0.2 eV on m_{lightest} from the KATRIN experiment [14].*

as a function of the lightest neutrino mass $|m_{\text{lightest}}|$, which is $|m_1|$ ($|m_3|$) for the normal (inverted) hierarchy. We clearly see that the largest hierarchies, obtained at the smallest

⁴In the numerical simulation, this means that we allow the absolute values of the Yukawas to be in the interval $[1/2, 3/2]$, while no restriction whatsoever has been imposed on the flavon vevs a part from being all equal within a factor of 10; also the large scale Λ is left free.

allowed $|m_{lightest}|$, are at the level of $\mathcal{O}(10)$ for the normal ordering and $\mathcal{O}(10^{-2})$ for the inverted one. The ratio tends to a degenerate spectrum in both cases for $|m_{lightest}| \sim 10^{-1}$ eV which, however, is disfavoured in our model. In the right panel we show the sum of the light neutrino masses as a function of the lightest neutrino mass $m_{lightest}$. The vertical line denotes the future sensitivity of 0.2 eV on $|m_{lightest}|$ from the KATRIN experiment [14], and the horizontal lines are the cosmological bounds [15] at 0.60 eV, obtained combining the data from ref.[16], and at 0.19 eV, corresponding to all the previous data combined to the small scale primordial spectrum from Lyman-alpha ($\text{Ly}\alpha$) forest clouds [17]. We see that our model predicts Σm_i too similar for both hierarchies to be distinguished using the current cosmological information on the sum of the neutrino masses; however, such a discrimination could be possible if some improvements on these bounds would be achieved in the near future.

Finally, we present in Fig.(2) the predictions for the values of the effective mass $|m_{ee}|$ as a function of the lightest neutrino mass, for both normal and inverted hierarchy. We also show the future sensitivity of the KATRIN (vertical solid line) and of CUORE [18] (horizontal solid line at 15 meV) experiments. The main feature of the analysis is that




Figure 2: $|m_{ee}|$ as a function of the lightest neutrino mass, for both normal and inverted hierarchy. The filled regions correspond to the possible values of $|m_{ee}|$ in the limit of exact tri-bimaximal mixing, with mass differences computed at the central values in eq.(51). The horizontal and vertical lines are the future expected bounds on $|m_{ee}|$ and $|m_{lightest}|$ from the CUORE and KATRIN experiments, respectively.

a large set of points falls into the region of $|m_{lightest}|$ around 10^{-2} eV but, given the still large number of parameters of the model, many values of $|m_{ee}|$ can be obtained for both hierarchies among the experimental allowed ranges. On the other hand, the region above this value (the regime of degenerate spectra) is strongly disfavoured.

4.1 Leptogenesis

The formal description of the asymmetry parameters can be done in the general context of arbitrary Dirac mass matrix because the final expressions are quite compact and transparent. The asymmetry parameters are defined as follows:

$$\epsilon_i = \frac{1}{8\pi(\hat{Y}\hat{Y}^\dagger)_{ii}} \sum_{j \neq i} \text{Im} \left\{ \left[(\hat{Y}\hat{Y}^\dagger)_{ij} \right]^2 \right\} f \left(\frac{|M_j|^2}{|M_i|^2} \right) \quad (52)$$

where the *hat* matrices are Yukawa matrices evaluated in the basis in which the Majorana mass matrix is diagonal and M_i are the Majorana masses. For supersymmetric theories, the f -function is given by:

$$f(x) = -\sqrt{x} \left[\frac{2}{x-1} + \log \left(\frac{1+x}{x} \right) \right]. \quad (53)$$

Defining Ω as the unitary matrix which diagonalizes the Majorana mass matrix, the LO Yukawa matrix in this basis is given by

$$v_u \hat{Y} = v_u \Omega^T Y_\nu = \Omega^T m_D$$

and the product $\hat{Y}\hat{Y}^\dagger$ reads:

$$\hat{Y}\hat{Y}^\dagger = \Omega^T Y_\nu Y_\nu^\dagger \Omega^* . \quad (54)$$

At LO, $\Omega = U_{TBM}$ and the product $\hat{Y}\hat{Y}^\dagger$ is a diagonal matrix: the ϵ_i parameters are all vanishing [12]. At the next to leading order, one has to take into account the corrections to the Yukawa matrix as well as to the Majorana mass matrix, which reflects in a different structure of both the Ω and Y_ν matrices in such a way that:

$$\begin{aligned} v_u Y_\nu &= \left(m_D + v_u \begin{pmatrix} \varepsilon \\ \varepsilon' \end{pmatrix} \varepsilon^2 y_{\nu_1} \hat{S} \right) = v_u (Y_{LO} + \delta Y) \\ \Omega &= U_{TBM} U_\phi + \delta \Omega \end{aligned} \quad (55)$$

where $\delta\Omega$ has a structure similar to the corrections δU computed in Sect.(3.3.1) and δY is of $\mathcal{O}(\lambda_C^4)$ compared to its leading order result. This means that the correction to the matrix product $\hat{Y}\hat{Y}^\dagger$ is given by:

$$\delta(\hat{Y}\hat{Y}^\dagger) = (\delta\Omega)^T Y_{LO} Y_{LO}^\dagger \Omega^* + \Omega^T Y_{LO} Y_{LO}^\dagger (\delta\Omega)^* + \Omega^T \delta(Y_\nu Y_\nu^\dagger) \Omega^* .$$

The first two terms do not contribute to the ϵ_i parameters because they are complex conjugate of each other and do not contain any imaginary part; then the only contribution arises from the last term. In the basis in which the charged leptons are diagonal and considering ε as a real variable for simplicity, one easily obtains:

$$\delta(\hat{Y}\hat{Y}^\dagger) = \begin{pmatrix} \sigma_1 \varepsilon^3 & \sqrt{2} e^{i(\phi_1 - \phi_2)} \sigma_2 \varepsilon^3 & 0 \\ \sqrt{2} e^{-i(\phi_1 - \phi_2)} \sigma_3 \varepsilon^3 & 0 & 0 \\ 0 & 0 & \sigma_4 \varepsilon^3 \end{pmatrix} \quad (56)$$

where the σ_i coefficients are complicated functions of y_{ν_i} , A_S and B_S . Then, at leading order, the ϵ parameters are given by:

$$\begin{aligned}\epsilon_1 &= \left(\frac{\epsilon}{\epsilon'}\right)^2 \epsilon^4 \frac{(A_S - B_S)^2}{4\pi} \frac{[y_{\nu_1}(y_{\nu_2} + 2y_{\nu_3}) + 3|y_{\nu_1}|^2]^2 f\left(\frac{|M_2|^2}{|M_1|^2}\right) \sin[2(\phi_1 - \phi_2)]}{9|y_{\nu_1}|^2 + |y_{\nu_2}|^2 + |y_{\nu_3}|^2 - 2(y_{\nu_2}y_{\nu_3}) + 3y_{\nu_1}y_{\nu_2} - 3y_{\nu_1}y_{\nu_3}} \\ \epsilon_2 &= \left(\frac{\epsilon}{\epsilon'}\right)^2 \epsilon^4 \frac{(A_S - B_S)^2}{4\pi} \frac{[y_{\nu_1}(y_{\nu_2} + 2y_{\nu_3}) + 3|y_{\nu_1}|^2]^2 f\left(\frac{|M_1|^2}{|M_2|^2}\right) \sin[2(\phi_1 - \phi_2)]}{4|y_{\nu_2}|^2 + |y_{\nu_3}|^2 + 4(y_{\nu_2}y_{\nu_3})} \\ \epsilon_3 &= 0.\end{aligned}\tag{57}$$

A relevant feature of the model is that ϵ_3 always vanishes; however, we can see that, barring possible fine-tunings in the parameters and/or suppressions or enhancements due to $f(|x|^2) \sin[2\Delta\phi]$, $\epsilon_{1,2}$ can be of the right order of magnitude to fulfill the experimental requirements for a successful leptogenesis because

$$\epsilon_{1,2} \sim \lambda_C^8 \sim 6 \times 10^{-6}.\tag{58}$$

5 The quark sector

It is well known that the extension of a flavour symmetry from the neutrino sector to the quark one is highly non trivial due to the “non-trigonometric” structure of the quark mixing matrix V_{CKM} . The common way to introduce quarks in these kind of models is to put the two $SU(2)$ doublets of left-handed quarks into a doublet representation of S_4 and assign the heaviest ones (top and bottom quarks) into singlets, in such a way to easily maintain the hierarchy among mass eigenstates. Better results are obtained if other extra symmetries, i.e. Froggatt-Nielsen or extra Z_N , are introduced in order to further suppress the unwanted couplings; this is particularly true for the m_{up}/m_{top} mass ratio, which otherwise tends to be larger than the experimental counterpart. It is clear that such scenarios are more flexible than the simple $S_4 \times Z_5$ illustrated in this paper; however, it is important to stress that it is still possible to get a satisfactory description of the quark sector without invoking any other extra symmetries and using the triplet representation of S_4 instead of the doublet ones. In the down-quark sector the simplest choice is to *copy* the coupling allowed in the charged lepton sector because the hierarchy between the d , s and b quark masses is quite similar to that existing for e , μ and τ leptons. On the other hand, the mass hierarchy in the up-quark sector does not follow such a simple prescription and we can arrange the $S_4 \times Z_5$ charge assignment in several ways. One possibility is summarized in Tab.(2), where Q is a triplet of $SU(2)$ left-handed doublets. The corresponding leading order lagrangian is the following:

Field	Q	d^c	s^c	b^c	u^c	c^c	t^c
S_4	3_1	1_2	1_1	1_1	1_2	1_1	1_1
Z_5	1	ω^3	ω^2	ω	ω^4	1	ω^4
$U(1)_R$	1	1	1	1	1	1	1

Table 2: Transformation properties of quarks under $S_4 \times Z_5$ and $U(1)_R$.

$$\begin{aligned}
\mathcal{L} = & \frac{y_b}{\Lambda} b^c (Q \varphi_T) h_d + \\
& \frac{y_{s_1}}{\Lambda^2} s^c (Q \varphi_T \varphi_T) h_d + \frac{y_{s_2}}{\Lambda^2} s^c Q (\eta \varphi_T) h_d + \\
& \frac{y_{d_1}}{\Lambda^3} d^c Q [\varphi_T (\varphi_T \varphi_T)_{2}]_{3_2} h_d + \frac{y_{d_2}}{\Lambda^3} d^c Q [\varphi_T (\varphi_T \varphi_T)_{3_1}]_{3_2} h_d + \\
& \frac{y_{d_3}}{\Lambda^3} d^c Q [\eta (\varphi_T \varphi_T)_{3_1}]_{3_2} h_d + \frac{y_{d_4}}{\Lambda^3} d^c Q [\varphi_T (\eta \eta)_{2}]_{3_2} h_d + \\
& \frac{y_{d_5}}{\Lambda^2} d^c Q (\Delta \varphi_S)_{3_2} h_d + \frac{y_t}{\Lambda} t^c (Q \varphi_T) h_u + \\
& \frac{y_{c_1}}{\Lambda^2} c^c (Q \varphi_T \varphi_T) h_u + \frac{y_{c_2}}{\Lambda^2} c^c Q (\eta \varphi_T) h_u + \\
& \frac{y_{u_1}}{\Lambda^3} u^c Q [\Delta (\varphi_T \varphi_T)_{3_1}]_{3_2} h_u + \frac{y_{u_2}}{\Lambda^3} u^c Q [\varphi_S (\varphi_T \varphi_T)_{2}]_{3_2} h_u + \\
& \frac{y_{u_3}}{\Lambda^3} u^c Q [\varphi_S (\varphi_T \varphi_T)_{3_1}]_{3_2} h_u + \frac{y_{u_4}}{\Lambda^3} u^c Q [\varphi_S (\eta \eta)_{2}]_{3_2} h_u + \\
& \frac{y_{u_5}}{\Lambda^3} u^c Q [\eta (\varphi_T \varphi_S)_{3_1}]_{3_2} h_u + \frac{y_{u_6}}{\Lambda^3} u^c Q [\eta (\varphi_T \varphi_S)_{3_2}]_{3_2} h_u.
\end{aligned} \tag{59}$$

This lagrangian gives rise to a diagonal mass matrix for the down-type quarks and an almost diagonal mass matrix for the up-type quarks (all the up entries are filled with terms of the same order $\varepsilon^2 \varepsilon'$). This picture slightly changes when including the next to leading order effects. As for the case of charged leptons, in the down sector the relevant NLO corrections come from the corrections to the vacuum alignment of the flavon fields, which are all of relative $\mathcal{O}(1/\Lambda)$ with respect to their leading order counterparts, whereas the higher order operators give corrections of the same size only for the s -quark, via couplings like

$$(\varphi_S^3), (\varphi_S \Delta^2), (\varphi_S \xi^2), (\varphi_S \Delta \xi), \tag{60}$$

and are much more suppressed for the down and bottom quarks. Similarly, for the up-type quarks the mass matrix is modified by the NLO structure of vacuum alignment of the flavon fields and, for the c -quark, also by the same higher order operators modifying the s -quark entries in eq.(60). All in all, the mass matrices for both type of quarks are given by:

$$m_{down} = v_d \begin{pmatrix} a_1^d \varepsilon^3 & a_2^d \varepsilon \varepsilon'^2 & -a_2^d \varepsilon \varepsilon'^2 \\ b_1^d \varepsilon'^3 & b_2^d \varepsilon^2 & b_3^d \varepsilon'^3 \\ c_1^d \frac{\varepsilon'^3}{\varepsilon} & c_2^d \frac{\varepsilon'^3}{\varepsilon} & c_3^d \varepsilon \end{pmatrix} \quad m_{up} = v_u \begin{pmatrix} a_1^u \varepsilon^2 \varepsilon' & a_2^u \varepsilon^2 \varepsilon' & a_3^u \varepsilon^2 \varepsilon' \\ b_1^u \varepsilon'^3 & b_2^u \varepsilon^2 & b_3^u \varepsilon'^3 \\ c_1^u \frac{\varepsilon'^3}{\varepsilon} & c_2^u \frac{\varepsilon'^3}{\varepsilon} & c_3^u \varepsilon \end{pmatrix} \tag{61}$$

with the following mass eigenvalues:

$$\begin{aligned}
m_d &= v_d \left[a_1^d \varepsilon^3 + \left(\frac{c_1^d}{c_3^d} - \frac{b_1^d}{b_3^d} \right) a_2^d \varepsilon \varepsilon'^3 \right] \\
m_s &= v_d b_2^d \varepsilon^2 + \mathcal{O}(\varepsilon^4) \\
m_b &= v_d c_3^d \varepsilon + \mathcal{O}(\varepsilon^4)
\end{aligned} \tag{62}$$

and

$$\begin{aligned}
m_u &= v_u \left[a_1^u \varepsilon^2 \varepsilon' + \left(\frac{b_1^u a_2^u}{b_2^u} - \frac{a_3^u c_1^u}{c_3^u} \right) \varepsilon'^4 \right] \\
m_c &= v_u b_2^u \varepsilon^2 + \mathcal{O}(\varepsilon'^4) \\
m_t &= v_u c_3^u \varepsilon + \mathcal{O}(\varepsilon'^4).
\end{aligned} \tag{63}$$

The previous mass matrices can be diagonalized by unitary matrices in such a way that:

$$\begin{aligned}
(m_u^\dagger m_u)_{\text{diag}} &= U_{uL}^\dagger (m_u^\dagger m_u) U_{uL} \\
(m_d^\dagger m_d)_{\text{diag}} &= U_{dL}^\dagger (m_d^\dagger m_d) U_{dL},
\end{aligned}$$

where:

$$U_{dL} = \begin{pmatrix} 1 & \left(\frac{b_1^d}{b_2^d} \frac{\varepsilon'^3}{\varepsilon^2} \right)^* & \left(\frac{c_1^d}{c_3^d} \frac{\varepsilon'^3}{\varepsilon^2} \right)^* \\ -\frac{b_1^d}{b_2^d} \frac{\varepsilon'^3}{\varepsilon^2} & 1 & \left(\frac{c_2^d}{c_3^d} \frac{\varepsilon'^3}{\varepsilon^2} \right)^* \\ -\frac{c_1^d}{c_3^d} \frac{\varepsilon'^3}{\varepsilon^2} & -\frac{c_2^d}{c_3^d} \frac{\varepsilon'^3}{\varepsilon^2} & 1 \end{pmatrix}, \quad U_{uL} = \begin{pmatrix} 1 & \left(\frac{b_1^u}{b_2^u} \frac{\varepsilon'^3}{\varepsilon^2} \right)^* & \left(\frac{c_1^u}{c_3^u} \frac{\varepsilon'^3}{\varepsilon^2} \right)^* \\ -\frac{b_1^u}{b_2^u} \frac{\varepsilon'^3}{\varepsilon^2} & 1 & \left(\frac{c_2^u}{c_3^u} \frac{\varepsilon'^3}{\varepsilon^2} \right)^* \\ -\frac{c_1^u}{c_3^u} \frac{\varepsilon'^3}{\varepsilon^2} & -\frac{c_2^u}{c_3^u} \frac{\varepsilon'^3}{\varepsilon^2} & 1 \end{pmatrix} \tag{64}$$

The resulting V_{CKM} is given by:

$$V_{CKM} = U_{uL}^\dagger U_{dL} = \begin{pmatrix} 1 & \left[\left(\frac{b_1^d}{b_2^d} - \frac{b_1^u}{b_2^u} \right) \frac{\varepsilon'^3}{\varepsilon^2} \right]^* & \left[\left(\frac{c_1^d}{c_3^d} - \frac{c_1^u}{c_3^u} \right) \frac{\varepsilon'^3}{\varepsilon^2} \right]^* \\ -\left(\frac{b_1^d}{b_2^d} - \frac{b_1^u}{b_2^u} \right) \frac{\varepsilon'^3}{\varepsilon^2} & 1 & \left[\left(\frac{c_2^d}{c_3^d} - \frac{c_2^u}{c_3^u} \right) \frac{\varepsilon'^3}{\varepsilon^2} \right]^* \\ -\left(\frac{c_1^d}{c_3^d} - \frac{c_1^u}{c_3^u} \right) \frac{\varepsilon'^3}{\varepsilon^2} & -\left(\frac{c_2^d}{c_3^d} - \frac{c_2^u}{c_3^u} \right) \frac{\varepsilon'^3}{\varepsilon^2} & 1 \end{pmatrix}. \tag{65}$$

As for other flavour models, the matching of the V_{CKM} and the quark mass ratios to their experimental values requires some fine-tunings between the Yukawas. As anticipated, all the experimental mass ratios in the down sector are easily reproduced for the natural values $a_1^d, b_2^d, c_3^d \sim \mathcal{O}(1)$ because $m_d/m_s \sim m_s/m_b \sim \varepsilon \sim \lambda_C^2$. A moderate hierarchy is also present in the up sector and it mainly depends on the parameter ε' , which we estimated in Sect.(3.3.2) to be not larger than some units of λ_C^2 . This same parameter appears in the off-diagonal entries of V_{CKM} with the same power in each entries. This is essentially the reason why it is difficult to explain at the same time the mass hierarchy in the up-type quark sector and the off-diagonal values of the quark mixing matrix. For example, one can treat ε' and $\tan \beta$ as free parameters of the model. In that case, one can use the ratio

m_u/m_c to fix the value of ε' and m_u/m_d for that of $\tan\beta$; assuming for all Yukawas the natural $\mathcal{O}(1)$ value, we get:

$$\frac{m_u}{m_c} = \varepsilon' \quad \frac{m_u}{m_d} = \left(\frac{\varepsilon'}{\varepsilon}\right) \tan\beta \quad (66)$$

and then

$$\varepsilon' = \left(\frac{m_u}{m_c}\right)_{\text{exp}} \quad \tan\beta = \lambda_C^2 \left(\frac{m_u}{m_d}\right)_{\text{exp}} / \left(\frac{m_u}{m_c}\right)_{\text{exp}} \sim 12. \quad (67)$$

It is easy to verify that the previous results are enough to accomodate all the independent mass ratios that can be built from six different quarks. However, ε' turns out to be very small and, having assumed all Yukawas of $\mathcal{O}(1)$, it is really difficult to enhance the off-diagonal elements of V_{CKM} to their experimental values. The other possibility is to preserve $\varepsilon' \sim \mathcal{O}(\lambda_C^2)$, requiring that

$$\left[\left(\frac{b_1^d}{b_2^d} - \frac{b_1^u}{b_2^u}\right)\right] \sim \mathcal{O}(1/\lambda_C) \quad \left[\left(\frac{c_1^d}{c_3^d} - \frac{c_1^u}{c_3^u}\right)\right] \sim \mathcal{O}(\lambda_C) \quad \left[\left(\frac{c_2^d}{c_3^d} - \frac{c_2^u}{c_3^u}\right)\right] \sim \mathcal{O}(1). \quad (68)$$

This fine-tuning is a condition on the individual Yukawas, which can be chosen to be:

$$\begin{aligned} c_2^d &\sim b_1^u \sim \mathcal{O}(1) \\ c_1^d &\sim \mathcal{O}(\lambda_C) \\ b_1^d &\sim \mathcal{O}(1/\lambda_C), \end{aligned} \quad (69)$$

providing that $b_2^u, c_3^u \gg 1$ to fit the charm and top quark masses. Although this situation is not completely satisfactory, it illustrates a way to account for many experimental informations in the quark sector using a simple and unconstrained $S_4 \times Z_5$ model, the prize to be paid being a fine-tuning in four of the Yukawa couplings appearing in the lagrangian.

Notice that it is not more difficult to modify the charge assignment proposed in Tab.(2) in such a way to obtain at LO a V_{CKM} matrix with entries of $\mathcal{O}(1)$ in the Cabibbo 2×2 submatrix. This can be accomplished slightly changing only the Z_5 charge of the quark c . In fact, giving the assignment $c \sim (1_1, \omega^3)$ under (S_4, Z_5) , the c -quark part of lagrangian is now:

$$\mathcal{L}_c = \frac{y_{c1}}{\Lambda^2} c^c (Q \varphi_T \Delta) h_u + \frac{y_{c2}}{\Lambda^2} c^c Q (\varphi_T \varphi_S) h_u + \frac{y_{c3}}{\Lambda^2} c^c Q \varphi_T \xi h_u + \frac{y_{c4}}{\Lambda^2} c^c Q (\varphi_S \eta) h_u \quad (70)$$

and the c -quark entries in the mass matrix are modified according to

$$(c^c, u) : b_1^u \varepsilon \varepsilon' \quad (c^c, c) : b_2^u \varepsilon \varepsilon' \quad (c^c, t) : b_3^u \varepsilon \varepsilon'. \quad (71)$$

Correspondingly, the u and c -quark masses are:

$$m_u = v_u \left(a_1^u - \frac{b_1^u a_2^u}{b_2^u} \right) \varepsilon^2 \varepsilon' \quad m_c = v_u b_2^u \varepsilon \varepsilon' \quad (72)$$

and the V_{CKM} has the following structure:

$$V_{CKM} = \begin{pmatrix} -\frac{b_2^u}{K} - \frac{b_1^u b_1^d}{b_2^d K} \frac{\varepsilon'^3}{\varepsilon^2} & \frac{b_1^u}{K} - \frac{b_2^u b_1^d}{b_2^d K} \frac{\varepsilon'^3}{\varepsilon^2} & \frac{c_3^d (b_2^u c_1^u - b_1^u c_2^u) - c_3^u (b_2^d c_1^d - b_1^d c_2^d)}{c_3^d c_3^u K} \frac{\varepsilon'^3}{\varepsilon^2} \\ -\frac{b_1^u}{K} + \frac{b_2^u b_1^d}{b_2^d K} \frac{\varepsilon'^3}{\varepsilon^2} & -\frac{b_2^u}{K} - \frac{b_1^u b_1^d}{b_2^d K} \frac{\varepsilon'^3}{\varepsilon^2} & \frac{c_3^d (b_1^u c_1^u + b_2^u c_2^u) - c_3^u (b_1^d c_1^d + b_2^d c_2^d)}{c_3^d c_3^u K} \frac{\varepsilon'^3}{\varepsilon^2} \\ \left(-\frac{c_1^d}{c_3^d} + \frac{c_1^u}{c_3^u}\right) \frac{\varepsilon'^3}{\varepsilon^2} & \left(-\frac{c_2^d}{c_3^d} + \frac{c_2^u}{c_3^u}\right) \frac{\varepsilon'^3}{\varepsilon^2} & 1 \end{pmatrix} \quad (73)$$

where, for simplicity, we introduced the short-hand notation $K = \sqrt{(b_1^u)^2 + (b_2^u)^2}$ and considered real vevs and Yukawa couplings. In this case, one can reproduce at the same time the correct values for the (11) and (22) entries of V_{CKM} as well as the mass ratio m_u/m_c imposing $b_2^u \sim \mathcal{O}(\lambda_C^2)$; playing with ε' and the other Yukawas one can also reproduce the off-diagonal entries, but one or more fine-tunings in the coefficients in front of $\varepsilon'^3/\varepsilon^2$ ratio are obviously needed.

6 Conclusions

We have presented and discussed an S_4 model for TB mixing (of the see-saw type) and quark mixing which, in spite of being based on a most economical flavour symmetry and field content, it is still phenomenologically viable. In the neutrino sector, we realized the most general neutrino mass matrix diagonalized at LO by TB mixing. At the NLO, all mixing angles receive corrections at higher orders at the level of $\mathcal{O}(\lambda_C^2)$ and, in particular, the elusive θ_{13} is predicted to be within the sensitivity of the experiments which are now in preparation and will take data in the near future. Also, we have shown that the leptogenesis parameters $\varepsilon_{1,2}$ can be of the right order of magnitude to satisfy the requirements needed to reproduce the observed asymmetry and that an acceptable phenomenology is still obtainable even with all the Yukawa couplings at $\mathcal{O}(1)$.

In the charged lepton sector, the mass hierarchy is determined by the $S_4 \times Z_5$ flavour symmetry itself without invoking a Froggatt-Nielsen $U(1)$ symmetry. The NLO corrections to the mass matrix turn out to be relevant for a $\theta_{13} \neq 0$.

In the quark sector, we have discussed in detail to which extent the model can reproduce the data on masses and mixing, emphasizing where it is successful and the reasons for its failures. Two different LO examples for V_{CKM} have been given, one proportional to the identity matrix and the other with $\mathcal{O}(1)$ elements in the 2×2 sector; both realizations need almost the same amount of fine-tuning to reproduce the off-diagonal entries at the NLO.

7 Acknowledgements

I am grateful to Guido Altarelli for very stimulating discussions. I also wish to thank Luca Merlo for some interesting comments. This work was supported by the Deutsche Forschungs-gemeinschaft, contract WI 2639/2-1.

A The Group S_4

We adopt the following convention for the generators S and T , according to [9]

$$S^4 = T^3 = (ST^2)^2 = \mathbb{1} \quad (74)$$

In the different representations, they can be written as reported in Tab.(3):

rep	1_1	1_2	2	3_1	3_2
S	1	-1	$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$	$\frac{1}{3} \begin{pmatrix} -1 & 2\omega & 2\omega^2 \\ 2\omega & 2\omega^2 & -1 \\ 2\omega^2 & -1 & 2\omega \end{pmatrix}$	$\frac{1}{3} \begin{pmatrix} 1 & -2\omega & -2\omega^2 \\ -2\omega & -2\omega^2 & 1 \\ -2\omega^2 & 1 & -2\omega \end{pmatrix}$
T	1	1	$\begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega \end{pmatrix}$	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega \end{pmatrix}$

Table 3: Generators S and T in different representations.

The 24 elements of the group belong to five conjugacy classes

$$\begin{aligned}
\mathcal{C}_1 &: 1 \\
\mathcal{C}_2 &: S^2, TS^2T^2, S^2TS^2T^2 \\
\mathcal{C}_3 &: T, T^2, S^2T, S^2T^2, STST^2 \\
&\quad STS, TS^2, T^2S^2 \\
\mathcal{C}_4 &: ST^2, T^2S, TST \\
&\quad TSTS^2, STS^2, S^2TS \\
\mathcal{C}_5 &: S, TST^2, ST, TS, S^3, S^3T^2.
\end{aligned}$$

The explicit expression of the elements in the 2-dimensional representation is:

$$\begin{aligned}
\bullet \mathcal{C}_{1,2} &: \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\
\bullet \mathcal{C}_3 &: \begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix}, \begin{pmatrix} \omega^2 & 0 \\ 0 & \omega \end{pmatrix} \\
\bullet \mathcal{C}_{4,5} &: \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & \omega \\ \omega^2 & 0 \end{pmatrix}, \begin{pmatrix} 0 & \omega^2 \\ \omega & 0 \end{pmatrix},
\end{aligned}$$

while for the 3-dimensional representation 3_1 the elements are

$$\bullet \mathcal{C}_1 : I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\bullet \mathcal{C}_2: \quad S^2 = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix} \quad TS^2T^2 = \frac{1}{3} \begin{pmatrix} -1 & 2\omega & 2\omega^2 \\ 2\omega^2 & -1 & 2\omega \\ 2\omega & 2\omega^2 & -1 \end{pmatrix}$$

$$S^2TS^2T^2 = \frac{1}{3} \begin{pmatrix} -1 & 2\omega^2 & 2\omega \\ 2\omega & -1 & 2\omega^2 \\ 2\omega^2 & 2\omega & -1 \end{pmatrix}$$

$$\bullet \mathcal{C}_3: \quad T = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega \end{pmatrix} \quad T^2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix} \quad S^2T = \frac{1}{3} \begin{pmatrix} -1 & 2\omega^2 & 2\omega \\ 2 & -\omega^2 & 2\omega \\ 2 & 2\omega^2 & -\omega \end{pmatrix}$$

$$S^2T^2 = \frac{1}{3} \begin{pmatrix} -1 & 2\omega & 2\omega^2 \\ 2 & -\omega & 2\omega^2 \\ 2 & 2\omega & -\omega^2 \end{pmatrix} \quad STST^2 = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2\omega^2 & -\omega^2 & 2\omega^2 \\ 2\omega & 2\omega & -\omega \end{pmatrix}$$

$$STS = \frac{1}{3} \begin{pmatrix} -1 & 2\omega^2 & 2\omega \\ 2\omega^2 & -\omega & 2 \\ 2\omega & 2 & -\omega^2 \end{pmatrix} \quad TS^2 = \frac{1}{3} \begin{pmatrix} -1 & 2\omega & 2\omega^2 \\ 2\omega & -\omega^2 & 2 \\ 2\omega^2 & 2 & -\omega \end{pmatrix}$$

$$T^2S^2 = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2\omega & -\omega & 2\omega \\ 2\omega^2 & 2\omega^2 & -\omega^2 \end{pmatrix}$$

$$\bullet \mathcal{C}_4: \quad ST^2 = \frac{1}{3} \begin{pmatrix} -1 & 2\omega^2 & 2\omega \\ 2\omega & 2 & -\omega^2 \\ 2\omega^2 & -\omega & 2 \end{pmatrix} \quad T^2S = \frac{1}{3} \begin{pmatrix} -1 & 2\omega & 2\omega^2 \\ 2\omega^2 & 2 & -\omega \\ 2\omega & -\omega^2 & 2 \end{pmatrix}$$

$$TST = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & 2 & -1 \\ 2 & -1 & 2 \end{pmatrix} \quad TSTS^2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$STS^2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & \omega \\ 0 & \omega^2 & 0 \end{pmatrix} \quad S^2TS = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & \omega^2 \\ 0 & \omega & 0 \end{pmatrix}$$

$$\bullet \mathcal{C}_5: \quad S = \frac{1}{3} \begin{pmatrix} -1 & 2\omega & 2\omega^2 \\ 2\omega & 2\omega^2 & -1 \\ 2\omega^2 & -1 & 2\omega \end{pmatrix} \quad TST^2 = \frac{1}{3} \begin{pmatrix} -1 & 2\omega^2 & 2\omega \\ 2 & 2\omega^2 & -\omega \\ 2 & -\omega^2 & 2\omega \end{pmatrix}$$

$$ST = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2\omega & 2\omega & -\omega \\ 2\omega^2 & -\omega^2 & 2\omega^2 \end{pmatrix} \quad TS = \frac{1}{3} \begin{pmatrix} -1 & 2\omega & 2\omega^2 \\ 2 & 2\omega & -\omega^2 \\ 2 & -\omega & 2\omega^2 \end{pmatrix}$$

$$S^3 = \frac{1}{3} \begin{pmatrix} -1 & 2\omega^2 & 2\omega \\ 2\omega^2 & 2\omega & -1 \\ 2\omega & -1 & 2\omega^2 \end{pmatrix} \quad S^3T^3 = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2\omega^2 & 2\omega^2 & -\omega^2 \\ 2\omega & -\omega & 2\omega \end{pmatrix}$$

For the 3-dimensional representation 3_2 , the matrices representing the elements of the group can be obtained from the list for the representation 3_1 in the following way: for $\mathcal{C}_{1,2,3}$ are the same, while for $\mathcal{C}_{4,5}$ are the opposite.

In the previous basis, the Clebsch-Gordan coefficients are as follows (α_i indicates the elements of the first representation of the product and β_i the second one):

$$1_1 \otimes \eta = \eta \otimes 1_1 = \eta \quad \text{with } \eta \text{ any representation}$$

$$1_2 \otimes 1_2 = 1_1 \sim \alpha\beta$$

$$1_2 \otimes 2 = 2 \sim \begin{pmatrix} \alpha\beta_1 \\ -\alpha\beta_2 \end{pmatrix}$$

$$1_2 \otimes 3_1 = 3_2 \sim \begin{pmatrix} \alpha\beta_1 \\ \alpha\beta_2 \\ \alpha\beta_3 \end{pmatrix}$$

$$1_2 \otimes 3_2 = 3_1 \sim \begin{pmatrix} \alpha\beta_1 \\ \alpha\beta_2 \\ \alpha\beta_3 \end{pmatrix}.$$

The multiplication rules with the 2-dimensional representation are the following:

$$2 \otimes 2 = 1_1 \oplus 1_2 \oplus 2 \quad \text{with} \quad \left\{ \begin{array}{l} 1_1 \sim \alpha_1\beta_2 + \alpha_2\beta_1 \\ 1_2 \sim \alpha_1\beta_2 - \alpha_2\beta_1 \\ 2 \sim \begin{pmatrix} \alpha_2\beta_2 \\ \alpha_1\beta_1 \end{pmatrix} \end{array} \right.$$

$$2 \otimes 3_1 = 3_1 \oplus 3_2 \quad \text{with} \quad \left\{ \begin{array}{l} 3_1 \sim \begin{pmatrix} \alpha_1\beta_2 + \alpha_2\beta_3 \\ \alpha_1\beta_3 + \alpha_2\beta_1 \\ \alpha_1\beta_1 + \alpha_2\beta_2 \end{pmatrix} \\ 3_2 \sim \begin{pmatrix} \alpha_1\beta_2 - \alpha_2\beta_3 \\ \alpha_1\beta_3 - \alpha_2\beta_1 \\ \alpha_1\beta_1 - \alpha_2\beta_2 \end{pmatrix} \end{array} \right.$$

$$2 \otimes 3_2 = 3_1 \oplus 3_2 \quad \text{with} \quad \left\{ \begin{array}{l} 3_1 \sim \begin{pmatrix} \alpha_1\beta_2 - \alpha_2\beta_3 \\ \alpha_1\beta_3 - \alpha_2\beta_1 \\ \alpha_1\beta_1 - \alpha_2\beta_2 \end{pmatrix} \\ 3_2 \sim \begin{pmatrix} \alpha_1\beta_2 + \alpha_2\beta_3 \\ \alpha_1\beta_3 + \alpha_2\beta_1 \\ \alpha_1\beta_1 + \alpha_2\beta_2 \end{pmatrix} \end{array} \right.$$

The multiplication rules with the 3-dimensional representations are the following:

$$3_1 \otimes 3_1 = 3_2 \otimes 3_2 = 1_1 \oplus 2 \oplus 3_1 \oplus 3_2 \quad \text{with} \quad \left\{ \begin{array}{l} 1_1 \sim \alpha_1\beta_1 + \alpha_2\beta_3 + \alpha_3\beta_2 \\ 2 \sim \begin{pmatrix} \alpha_2\beta_2 + \alpha_1\beta_3 + \alpha_3\beta_1 \\ \alpha_3\beta_3 + \alpha_1\beta_2 + \alpha_2\beta_1 \end{pmatrix} \\ 3_1 \sim \begin{pmatrix} 2\alpha_1\beta_1 - \alpha_2\beta_3 - \alpha_3\beta_2 \\ 2\alpha_3\beta_3 - \alpha_1\beta_2 - \alpha_2\beta_1 \\ 2\alpha_2\beta_2 - \alpha_1\beta_3 - \alpha_3\beta_1 \end{pmatrix} \\ 3_2 \sim \begin{pmatrix} \alpha_2\beta_3 - \alpha_3\beta_2 \\ \alpha_1\beta_2 - \alpha_2\beta_1 \\ \alpha_3\beta_1 - \alpha_1\beta_3 \end{pmatrix} \end{array} \right.$$

$$3_1 \otimes 3_2 = 1_2 \oplus 2 \oplus 3_1 \oplus 3_2 \quad \text{with} \quad \left\{ \begin{array}{l} 1_2 \sim \alpha_1\beta_1 + \alpha_2\beta_3 + \alpha_3\beta_2 \\ 2 \sim \begin{pmatrix} \alpha_2\beta_2 + \alpha_1\beta_3 + \alpha_3\beta_1 \\ -\alpha_3\beta_3 - \alpha_1\beta_2 - \alpha_2\beta_1 \end{pmatrix} \\ 3_1 \sim \begin{pmatrix} \alpha_2\beta_3 - \alpha_3\beta_2 \\ \alpha_1\beta_2 - \alpha_2\beta_1 \\ \alpha_3\beta_1 - \alpha_1\beta_3 \end{pmatrix} \\ 3_2 \sim \begin{pmatrix} 2\alpha_1\beta_1 - \alpha_2\beta_3 - \alpha_3\beta_2 \\ 2\alpha_3\beta_3 - \alpha_1\beta_2 - \alpha_2\beta_1 \\ 2\alpha_2\beta_2 - \alpha_1\beta_3 - \alpha_3\beta_1 \end{pmatrix} \end{array} \right.$$

References

- [1] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. B **530** (2002) 167 [arXiv:hep-ph/0202074]; P. F. Harrison and W. G. Scott, Phys. Lett. B **535** (2002) 163 [arXiv:hep-ph/0203209]; Z. z. Xing, Phys. Lett. B **533** (2002) 85 [arXiv:hep-ph/0204049]; P. F. Harrison and W. G. Scott, Phys. Lett. B **547** (2002) 219 [arXiv:hep-ph/0210197]; P. F. Harrison and W. G. Scott, Phys. Lett. B **557** (2003) 76 [arXiv:hep-ph/0302025]; P. F. Harrison and W. G. Scott, arXiv:hep-ph/0402006; P. F. Harrison and W. G. Scott, Phys. Lett. B **594**, 324 (2004) [arXiv:hep-ph/0403278].
- [2] B. T. Cleveland *et al.*, Astrophys. J. **496** (1998) 505; J. N. Abdurashitov *et al.* [SAGE Collaboration], Phys. Rev. C **60** (1999) 055801 [arXiv:astro-ph/9907113]; W. Hampel *et al.* [GALLEX Collaboration], Phys. Lett. B **447** (1999) 127; S. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **86** (2001) 5651 [arXiv:hep-ex/0103032]; J. P. Cravens *et al.* [Super-Kamiokande Collaboration], Phys. Rev. D

78 (2008) 032002 [arXiv:0803.4312 [hep-ex]]; Q. R. Ahmad *et al.* [SNO Collaboration], Phys. Rev. Lett. **87** (2001) 071301 [arXiv:nucl-ex/0106015]; S. N. Ahmed *et al.* [SNO Collaboration], Phys. Rev. Lett. **92** (2004) 181301 [arXiv:nucl-ex/0309004]; B. Aharmim *et al.* [SNO Collaboration], Phys. Rev. Lett. **101** (2008) 111301 [arXiv:0806.0989 [nucl-ex]]; Y. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **81** (1998) 1562 [arXiv:hep-ex/9807003]; M. Ambrosio *et al.* [MACRO Collaboration], Phys. Lett. B **517** (2001) 59 [arXiv:hep-ex/0106049]; M. Apollonio *et al.* [CHOOZ Collaboration], Phys. Lett. B **466** (1999) 415 [arXiv:hep-ex/9907037]; M. Apollonio *et al.* [CHOOZ Collaboration], Eur. Phys. J. C **27** (2003) 331 [arXiv:hep-ex/0301017]; F. Boehm *et al.*, Phys. Rev. D **64** (2001) 112001 [arXiv:hep-ex/0107009]; K. Eguchi *et al.* [KamLAND Collaboration], Phys. Rev. Lett. **90** (2003) 021802 [arXiv:hep-ex/0212021]; M. H. Ahn *et al.* [K2K Collaboration] Phys. Rev. Lett. **90** (2003) 041801 [arXiv:hep-ex/0212007]; E. Aliu *et al.* [K2K Collaboration], Phys. Rev. Lett. **94** (2005) 081802 [arXiv:hep-ex/0411038]; D. G. Michael *et al.* [MINOS Collaboration], Phys. Rev. Lett. **97** (2006) 191801 [arXiv:hep-ex/0607088]; P. Adamson *et al.* [MINOS Collaboration], Phys. Rev. Lett. **101** (2008) 131802 [arXiv:0806.2237 [hep-ex]].

[3] A. Strumia and F. Vissani, arXiv:hep-ph/0606054; G. L. Fogli *et al.*, Nucl. Phys. Proc. Suppl. **168** (2007) 341; M. C. Gonzalez-Garcia and M. Maltoni, Phys. Rept. **460** (2008) 1 [arXiv:0704.1800 [hep-ph]]; T. Schwetz, AIP Conf. Proc. **981** (2008) 8 [arXiv:0710.5027 [hep-ph]]; M. C. Gonzalez-Garcia and M. Maltoni, Phys. Lett. B **663** (2008) 405 [arXiv:0802.3699 [hep-ph]]; A. Bandyopadhyay, S. Choubey, S. Goswami, S. T. Petcov and D. P. Roy, arXiv:0804.4857 [hep-ph].

[4] E. Ma and G. Rajasekaran, Phys. Rev. D **64** (2001) 113012 [arXiv:hep-ph/0106291]; E. Ma, Mod. Phys. Lett. A **17** (2002) 627 [arXiv:hep-ph/0203238]; K. S. Babu, E. Ma and J. W. F. Valle, Phys. Lett. B **552** (2003) 207 [arXiv:hep-ph/0206292]; M. Hirsch, J. C. Romao, S. Skadhauge, J. W. F. Valle and A. Villanova del Moral, arXiv:hep-ph/0312244; Phys. Rev. D **69** (2004) 093006 [arXiv:hep-ph/0312265]; E. Ma, Phys. Rev. D **70** (2004) 031901; Phys. Rev. D **70** (2004) 031901 [arXiv:hep-ph/0404199]; New J. Phys. **6** (2004) 104 [arXiv:hep-ph/0405152]; arXiv:hep-ph/0409075; S. L. Chen, M. Frigerio and E. Ma, Nucl. Phys. B **724** (2005) 423 [arXiv:hep-ph/0504181]; E. Ma, Phys. Rev. D **72** (2005) 037301 [arXiv:hep-ph/0505209]; M. Hirsch, A. Villanova del Moral, J. W. F. Valle and E. Ma, Phys. Rev. D **72** (2005) 091301 [Erratum-ibid. D **72** (2005) 119904] [arXiv:hep-ph/0507148]; K. S. Babu and X. G. He, arXiv:hep-ph/0507217; E. Ma, Mod. Phys. Lett. A **20** (2005) 2601 [arXiv:hep-ph/0508099]; A. Zee, Phys. Lett. B **630** (2005) 58 [arXiv:hep-ph/0508278]; E. Ma, Phys. Rev. D **73** (2006) 057304 [arXiv:hep-ph/0511133]; X. G. He, Y. Y. Keum and R. R. Volkas, JHEP **0604** (2006) 039 [arXiv:hep-ph/0601001]; B. Adhikary, B. Brahmachari, A. Ghosal, E. Ma and M. K. Parida, Phys. Lett. B **638** (2006) 345 [arXiv:hep-ph/0603059]; Mod. Phys. Lett. A **22** (2007) 101 [arXiv:hep-ph/0610342]; L. Lavoura and H. Kuhbock, Mod. Phys. Lett. A **22** (2007) 181 [arXiv:hep-ph/0610050]; Eur. Phys. J. C **55**, 303 (2008) [arXiv:0711.0670

[hep-ph]]; M. Hirsch, A. S. Joshipura, S. Kaneko and J. W. F. Valle, Phys. Rev. Lett. **99**, 151802 (2007) [arXiv:hep-ph/0703046]; F. Yin, Phys. Rev. D **75** (2007) 073010 [arXiv:0704.3827 [hep-ph]]; F. Bazzocchi, S. Kaneko and S. Morisi, JHEP **0803** (2008) 063 [arXiv:0707.3032 [hep-ph]]; M. Honda and M. Tanimoto, Prog. Theor. Phys. **119** (2008) 583 [arXiv:0801.0181 [hep-ph]]; B. Brahmachari, S. Choubey and M. Mitra, Phys. Rev. D **77** (2008) 073008 [Erratum-ibid. D **77** (2008) 119901] [arXiv:0801.3554 [hep-ph]]; B. Adhikary and A. Ghosal, Phys. Rev. D **78** (2008) 073007 [arXiv:0803.3582 [hep-ph]]; M. Hirsch, S. Morisi and J. W. F. Valle, Phys. Rev. D **78** (2008) 093007 [arXiv:0804.1521 [hep-ph]]; P. H. Frampton and S. Matsuzaki, arXiv:0806.4592 [hep-ph]; F. Feruglio, C. Hagedorn, Y. Lin and L. Merlo, Nucl. Phys. B **809**, 218 (2009) [arXiv:0807.3160 [hep-ph]]; W. Grimus and L. Lavoura, JHEP **0904**, 013 (2009) [arXiv:0811.4766 [hep-ph]]; S. Morisi, Phys. Rev. D **79**, 033008 (2009) [arXiv:0901.1080 [hep-ph]]; M. C. Chen and S. F. King, JHEP **0906**, 072 (2009) [arXiv:0903.0125 [hep-ph]]; G. C. Branco, R. Gonzalez Felipe, M. N. Rebelo and H. Serodio, Phys. Rev. D **79**, 093008 (2009) [arXiv:0904.3076 [hep-ph]]; G. Altarelli and F. Feruglio, Nucl. Phys. B **741** (2006) 215 [arXiv:hep-ph/0512103]; D. Ibanez, S. Morisi and J. W. F. Valle, Phys. Rev. D **80**, 053015 (2009) [arXiv:0907.3109 [hep-ph]]; M. Hirsch, S. Morisi and J. W. F. Valle, Phys. Lett. B **679**, 454 (2009) [arXiv:0905.3056 [hep-ph]]; M. Hirsch, S. Morisi and J. W. F. Valle, Phys. Rev. D **79**, 016001 (2009) [arXiv:0810.0121 [hep-ph]]; E. Ma, arXiv:0908.3165 [hep-ph]; F. Feruglio, C. Hagedorn and L. Merlo, arXiv:0910.4058 [hep-ph]; S. Morisi and E. Peinado, arXiv:0910.4389 [hep-ph]; Y. Lin, Nucl. Phys. B **824**, 95 (2010) [arXiv:0905.3534 [hep-ph]]; C. Hagedorn, E. Molinaro and S. T. Petcov, arXiv:0908.0240 [hep-ph]; S. Morisi and E. Peinado, arXiv:0910.4389 [hep-ph].

[5] Y. Lin, Nucl. Phys. B **813**, 91 (2009) [arXiv:0804.2867 [hep-ph]]; G. Altarelli and D. Meloni, J. Phys. G **36**, 085005 (2009) [arXiv:0905.0620 [hep-ph]].

[6] E. Ma, Mod. Phys. Lett. A **21** (2006) 2931 [arXiv:hep-ph/0607190]; S. F. King and M. Malinsky, Phys. Lett. B **645** (2007) 351 [arXiv:hep-ph/0610250]; S. Morisi, M. Picariello and E. Torrente-Lujan, Phys. Rev. D **75** (2007) 075015 [arXiv:hep-ph/0702034]; F. Bazzocchi, S. Morisi and M. Picariello, Phys. Lett. B **659** (2008) 628 [arXiv:0710.2928 [hep-ph]]; F. Bazzocchi, S. Morisi, M. Picariello and E. Torrente-Lujan, C. Csaki, C. Delaunay, C. Grojean and Y. Grossman, JHEP **0810**, 055 (2008) [arXiv:0806.0356 [hep-ph]]; F. Bazzocchi, M. Frigerio and S. Morisi, Phys. Rev. D **78**, 116018 (2008) [arXiv:0809.3573 [hep-ph]]; P. Ciafaloni, M. Picariello, E. Torrente-Lujan and A. Urbano, Phys. Rev. D **79**, 116010 (2009) [arXiv:0901.2236 [hep-ph]]; G. Altarelli and F. Feruglio, Nucl. Phys. B **720** (2005) 64 [arXiv:hep-ph/0504165]; Nucl. Phys. B **775** (2007) 31 [arXiv:hep-ph/0610165]; G. Altarelli, F. Feruglio and C. Hagedorn, JHEP **0803** (2008) 052 [arXiv:0802.0090 [hep-ph]]; W. Grimus and H. Kuhbock, Phys. Rev. D **77**, 055008 (2008) [arXiv:0710.1585 [hep-ph]]; G. Seidl, arXiv:0811.3775 [hep-ph]; C. Hagedorn, E. Molinaro and S. T. Petcov, arXiv:0908.0240 [hep-ph];

- [7] C. S. Lam, Phys. Rev. D **78**, 073015 (2008) [arXiv:0809.1185 [hep-ph]].
- [8] G. J. Ding, arXiv:0909.2210 [hep-ph]; W. Grimus, L. Lavoura and P. O. Ludl, J. Phys. G **36**, 115007 (2009) [arXiv:0906.2689 [hep-ph]]; G. Altarelli, F. Feruglio and L. Merlo, JHEP **0905**, 020 (2009) [arXiv:0903.1940 [hep-ph]]; F. Bazzocchi, L. Merlo and S. Morisi, Phys. Rev. D **80**, 053003 (2009) [arXiv:0902.2849 [hep-ph]]; H. Ishimori, Y. Shimizu and M. Tanimoto, Prog. Theor. Phys. **121**, 769 (2009) [arXiv:0812.5031 [hep-ph]]; Y. Koide, JHEP **0708**, 086 (2007) [arXiv:0705.2275 [hep-ph]]; H. Zhang, Phys. Lett. B **655**, 132 (2007) [arXiv:hep-ph/0612214]; Y. Cai and H. B. Yu, Phys. Rev. D **74**, 115005 (2006) [arXiv:hep-ph/0608022]; C. Hagedorn, M. Lindner and R. N. Mohapatra, JHEP **0606**, 042 (2006) [arXiv:hep-ph/0602244]; E. Ma, Phys. Lett. B **632**, 352 (2006) [arXiv:hep-ph/0508231]; D. G. Lee and R. N. Mohapatra, Phys. Lett. B **329**, 463 (1994) [arXiv:hep-ph/9403201].
- [9] F. Bazzocchi, L. Merlo and S. Morisi, Nucl. Phys. B **816**, 204 (2009) [arXiv:0901.2086 [hep-ph]].
- [10] F. Bazzocchi and S. Morisi, arXiv:0811.0345 [hep-ph].
- [11] S. Weinberg, Phys. Rev. Lett. **43**, 1566 (1979).
- [12] E. E. Jenkins and A. V. Manohar, Phys. Lett. B **668**, 210 (2008) [arXiv:0807.4176 [hep-ph]].
- [13] T. Schwetz, M. Tortola and J. W. F. Valle, New J. Phys. **10** (2008) 113011 [arXiv:0808.2016 [hep-ph]]; M. Maltoni and T. Schwetz, arXiv:0812.3161 [hep-ph].
- [14] A. Osipowicz *et al.* [KATRIN Collaboration], arXiv:hep-ex/0109033.
- [15] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo and A. M. Rotunno, arXiv:0809.2936 [hep-ph].
- [16] WMAP Collaboration, E. Komatsu *et al.*, arXiv:0803.0547 [astro-ph]; ACBAR Collaboration, C. L. Reichardt *et al.*, arXiv:0801.1491 [astro-ph]; VSA Collaboration, C. Dickinson *et al.*, Mon. Not. Roy. Astron. Soc. **353**, 732 (2004) [arXiv:astro-ph/0402498]; CBI Collaboration, A. C. S. Readhead *et al.*, Astrophys. J. **609**, 498 (2004) [arXiv:astro-ph/0402359]; BOOMERANG Collaboration, C. J. MacTavish *et al.*, Astrophys. J. **647**, 799 (2006) [arXiv:astro-ph/0507503]; SDSS Collaboration, M. Tegmark *et al.*, Phys. Rev. D **74** (2006) 123507 [arXiv:astro-ph/0608632]; SNLS Collaboration, P. Astier *et al.* Astron. Astrophys. **447**, 31 (2006) [arXiv:astro-ph/0510447]; SDSS Collaboration, D. J. Eisenstein *et al.*, Astrophys. J. **633**, 560 (2005) [arXiv:astro-ph/0501171].
- [17] P. McDonald *et al.*, Astrophys. J. Suppl. **163**, 80 (2006); P. McDonald *et al.*, Astrophys. J. **635**, 761 (2005).
- [18] A. Giuliani [CUORE Collaboration], J. Phys. Conf. Ser. **120**, 052051 (2008).