



HAL
open science

Strangeness production at (sub)threshold energies

N Herrmann

► **To cite this version:**

N Herrmann. Strangeness production at (sub)threshold energies. Journal of Physics G: Nuclear and Particle Physics, 2010, 37 (9), pp.94036. 10.1088/0954-3899/37/9/094036 . hal-00600847

HAL Id: hal-00600847

<https://hal.science/hal-00600847>

Submitted on 16 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Strangeness production at (sub)threshold energies

N. Herrmann (FOPI collaboration)

Physikalisches Institut der Universität Heidelberg,
Philosophenweg 12, D-69120 Heidelberg, Germany

E-mail: herrmann@physi.uni-heidelberg.de

Abstract. Strange particles allow to probe the properties of hot and dense baryonic matter. At incident energies close to the production threshold medium effects on the strange hadrons are magnified. New data obtained with the recently completed FOPI upgrade are presented for the system Ni+Ni at 1.91 AGeV and discussed in comparison to earlier findings. Special emphasis is put to the search for multi-baryonic strange bound states, of which the Λp - channel is analyzed in detail.

1. Introduction

Modifications of hadron properties in dense baryonic matter are a current subject of intensive research in hadron physics. Various theoretical approaches [1, 2, 3] agree qualitatively on predicting, for example, modifications of masses and coupling constants for kaons and anti-kaons. Due to the density dependence of the $KN(\bar{K}N)$ potential, the K^- effective mass is expected to drop, whereas the mass of K^+ mesons is predicted to rise with increasing density of nuclear matter. Already two decades ago, Kaplan and Nelson [4] pointed out that due to additional attractive interactions with the surrounding nucleons a condensation of anti-kaons (K^-) may take place in a dense baryonic environment as encountered in the interior of neutron stars.

The strength of the K^- - nucleon potential might give rise to a new class of bound states [5, 6] with quite exotic properties. These bound states are called kaonic clusters and are being searched for in stopped K^- - beams [7, 8] and in pp - reactions [9]. Due to their predicted large central densities and the large compression reached in heavy ion reaction their production might be favored in heavy ion reactions [10] and therefore a complementary program was started to look for bound multibaryonic strange states in heavy ion reactions [11].

In the following new data obtained with the FOPI detector in 2008 at the SIS18 accelerator of GSI are shown for the reaction Ni+Ni at an incident energy of 1.91 AGeV elucidating some key aspects of strangeness physics in the threshold energy range.




Figure 1. Reconstructed masses for a momentum range up to 1.0 GeV/c making use of CDC - RPC matches. Positively charged particles are plotted in red, negative ones are shown in blue. Mass peaks for pions, kaons, proton, deuterons and triton are clearly visible. Positively charged kaons exhibit a signal-to-background ratio of better than 10.

2. Experiment: FOPI - Phase III

In order to allow the detection of strange particles the FOPI detector was upgraded with the installation of a new Time-of-Flight barrel covering the polar angle range $35^\circ < \Theta < 60^\circ$ based on Multigap - Resistive - Plate - Chambers (MRPC) [12] in 2006. Since then this detector was used in major data taking runs for which first result become available now. The performance in identifying the emitted particles is depicted for the reaction Ni+Ni at an incident energy of 1.91 AGeV in Fig. 1 where the mass values are shown that are calculated from the momentum reconstructed in the Central Drift Chamber (CDC) and the velocity measured in the MRPC - barrel. The time resolution and the granularity is sufficient to separate positively (negatively) charged kaons with a signal-to-background ratio in excess of 10 when the maximum momentum is limited to 1.0 (0.8) GeV/c, respectively.

3. Differential kaon flow

The superb kaon identification of the MRPC barrel allows to perform differential flow analysis of these rare particles and thus to continue the effort to determine the in-medium potential for kaons and antikaons [13]. The analysis has progressed to an extend that first surprising data on the differential sideflow of kaons can be shown.

In Fig. 2 the first fourier coefficients v_1 [14] are plotted as function of the normalized rapidity for kaon and protons for two different centralities. The direction of the reaction plane was determined by the transverse momentum method [15] making use of particles detected in the forward wall of FOPI only in order to avoid autocorrelations. The data are corrected for the finite reaction plane resolution making use of the Ollitrault formalism [16]. For better comparability with different reaction plane reconstruction methods the v_1 -data were shifted such that the mean v_1 -value at midrapidity is




Figure 2. Sideflow of kaons and protons in the reaction Ni+Ni at an incident energy of 1.91 AGeV as function of normalized rapidity $y^{(0)} = y/y_{CM} - 1$. The first Fourier coefficients v_1 are plotted for two different centrality bins for kaons (left panel) and protons (right panel). Black data points are obtained for peripheral collisions corresponding to a mean impact parameter of 6 ± 1 fm, the red ones for central collisions with a geometrical impact parameter $b < 3$ fm. The v_1 -values are obtained within the acceptance of the CDC, for details see text.

vanishing as required by symmetry. The centrality selection was chosen such that for both centrality bins the proton flow is approximately the same. Thus the event samples differ by the amount of stopping and the maximum maximum density reached within the collision leading to the same sideways deflection. While the kaon data in the central event sample confirm quantitatively our earlier findings [17, 18], the large sideflow in the most peripheral data sample is rather unexpected: at a normalized rapidity of $y^{(0)} \approx 0.5$ the antiflow of the kaons is as large in magnitude as the flow of the protons. Tentatively this effect can be attributed to the strong momentum dependence of the kaon potential. This hypothesis awaits its confirmation/rejection by transport model calculations.

In order to compare to the data shown in Fig. 2 model calculations have to take into account the acceptance of the detector, that is limited for this detailed analysis to the acceptance of the CDC covering a polar angle range of $26.0^\circ < \Theta_{lab} < 120^\circ$. For kaons additional constraints have to be put on the momenta necessary to obtain sufficient particle identification in the data. In the polar angular range from $30.5^\circ < \Theta_{lab} < 52^\circ$ (MRPC acceptance) kaons are included up to a momentum of 1.0 GeV, in the range $53^\circ < \Theta_{lab} < 120^\circ$ (plastic barrel acceptance) the laboratory momentum is limited to 0.5 GeV. The angular limits correspond to the so called 'shifted target' geometry, where the target was placed 40 cm upstream of its nominal position in the FOPI apparatus.

The available data sample also allow the reconstruction of the K^- -flow. However, this analysis is much more difficult due to the very low yield of K^- -that is at the 3% level of K^+ -production. The analysis is in progress.

4. Λ - baryon reconstruction

Strangeness physics in the threshold energy range is dominated by associate production underlining the importance to investigate hyperons in the same event samples as kaons. Phase space distributions of Λ - baryons are already published from earlier experiments [19]. With the current setup high statistics data samples can be obtained with very high purity as shown in Fig. 3 where a Λ - baryon signal is shown with a signal-to-background ratio of 13. The mean value of the reconstructed invariant mass peak with the current calibration amounts to $M = 1.1156 \pm 0.0002$ GeV in very good agreement with the PDG values. Due to the consistency checks allowed by the presence of the new MRPC barrel the calibration of the CDC could be improved so that a Gaussian width of $\sigma_M = 3.7$ MeV is now achieved for the Ni+Ni data of the 2008 run.




Figure 3. Reconstructed invariant mass of off-vertex pion - proton pairs. A minimal distances of 1.9 (0.6) cm was required for pions (protons) in combination with a pointing angle cut of $|\Delta\phi| < 2^\circ$ and a transverse flight path of at least 4 cm. In the upper panel 29.000 candidates are reconstructed in $5.6 \cdot 10^7$ events in the reaction Ni+Ni at 1.91 AGeV subtracting the mixed event background (blue histogram) from the correlated pairs (red histogram). The signal distribution (lower panel) exhibits a mass resolution of $\sigma_M = 3.7$ MeV. Within a $2.5 \times \sigma_M$ - interval around the nominal mass (hatched area in upper panel) a signal-to-background ratio of 13 is obtained.

The purities shown in Fig. 3 are necessary for a clean study of Λ - baryon flow and for the correlation analysis presented in the next section.

5. Proton - Λ - correlations

In our previous high statistics data samples, besides Λ -deuteron [11] we had analyzed Λ -proton correlations [21, 22] showing a clear signal at an invariant mass of $M = 2.13 \pm 0.02$ GeV for the reactions studied so far: Ni+Ni at 1.93 AGeV and Al+Al at 1.91 AGeV. Note that in these analyses the combinatorial background is reconstructed by using the same techniques that were used for the Σ^* measurement [20] that could be clearly identified in Λ - charged pion correlations. The existence of a peak structure in the Λ -proton correlation is confirmed by the latest data taken with the FOPI apparatus and shown in Fig. 4. The parameters of the peak structure (mean, width and yield) are consistent within statistical errors with the previous findings [21, 22]. The reconstructed yield amounts within the acceptance to 3% of the Λ -yield and is not distributed




Figure 4. Reconstructed invariant mass of Λ - proton pairs. The solid histogram and crosses denote the data (red) and the scaled mixed-events background (blue), respectively (upper panel). Both distributions were normalized over the range indicated by the horizontal arrow. The lower panel shows the difference and an excess at a mean value $M(p\Lambda) = 2.134 \pm 0.004$ GeV with a width of $\Gamma = 26 \pm 14$ MeV (statistical errors only). The statistical information (signal strength, significance and signal-over-background ratio) is extracted in the mass range limited by the vertical dashed lines shown in the upper panel.

according to thermal phase space. In Fig. 4 only proton- Λ -pairs with a combined rapidity of $y_{\Lambda p} < 0.65$ are shown. No significant structure was seen at midrapidity.

The location of the peak at $M(p\Lambda) = 2.134$ GeV agrees within the resolution with the proton- Σ^0 -threshold at 2.1309 GeV and thus drives the hypothesis that the structure is caused by a deuteron-like configuration of a proton and a hyperon. Similar structures have been reported in elementary hadron reactions [23, 24, 25]. However, the interpretation of the peak structure is still controversial since strong dependencies on the kinematics were observed. With our accumulated high statistics FOPI data we now confirm the existence of the structure in heavy-ion collisions where essentially no kinematic constraints exist due to the strong thermalization of the reaction systems.

As indicated by the arrow markers in Fig.4 the structure observed in the heavy ion data is clearly different from the one seen in K^- -absorption in the FINUDA experiment that is considered a candidate for a deeply bound kaonic state. However, the datasets are not necessarily contradicting each other since a) FINUDA has almost no acceptance for an invariant mass of 2.13 GeV and b) the production yield of deeply bound states could be much lower than the one for the observed deuteron-like strange dibaryon.

6. Conclusion

New data have been presented making use of the FOPI phase III setup at GSI elucidating the properties of strange particle in dense baryonic matter. For the first time a strong antiflow component for K^+ is seen for peripheral Ni+Ni reactions at 1.91 AGeV, pointing to a strong momentum dependence of the Kaon-Nucleus potential. The special role of the strange quark is visible in the formation of strange dibaryonic systems. Evidence is presented for the existence of a quasi bound state at an invariant Mass of 2.134 ± 0.004 GeV. Further investigations are necessary to clarify the nature of this state and/or the existence of even heavier multibaryonic strange states. More data are available from the 2009 measurement campaign of FOPI where more than 10^8 events were recorded for the reactions Ni+Pb at an incident energies of 1.91 AGeV and Ru+Ru at 1.65 AGeV opening new possibilities to study hadronic states in the hadronic environment of a heavy-ion collision.

Acknowledgments

This work was supported by the German BMBF under Contract No. 06HD190I, by the Korea Science and Engineering Foundation (KOSEF) under Grant No. F01-2006-000-10035-0, by the mutual agreement between GSI and IN2P3/CEA, by the Hungarian OTKA under Grant No. 47168, within the Framework of the WTZ program (Project RUS 02/021), by DAAD (PPP D/03/44611), by DFG (Projekt 446-KOR-113/76/04) and by the EU, 6th Framework Program, Integrated Infrastructure: Strongly Interacting Matter (Hadron Physics), Contract No. RII3-CT-2004-506078.

References

- [1] G. E. Brown *et al.*, *Nucl. Phys. A* 567 (1994) 937
- [2] W. Weise, *Nucl. Phys. A* 610 (1996) 35c
- [3] G.Q. Li *et al.*, *Nucl. Phys. A* 625 (1997) 372
- [4] D. B. Kaplan and A. E. Nelson, *Phys. Lett. B* 175 (1986) 57
- [5] Y. Akaishi and T. Yamazaki, *Phys. Rev. C* 65 (2002) 044005
- [6] T. Yamazaki, A. Doté, Y. Akaishi, *Phys. Lett. B* 587 (2004) 167
- [7] M. Agnello *et al.*, *Phys. Rev. Lett.* 94 (2005) 212303
- [8] T. Suzuki *et al.* [E549 collaboration], *Mod.Phys.Lett.A* 23 (2008) 2520
- [9] T. Yamazaki *et al.*, *arXiv:0810.5182*
- [10] T. Yamazaki *et al.*, *Nucl. Phys. A* 738 (2004) 168
- [11] N. Herrmann [FOPI collaboration], Proceedings of EXA05 conference, Österreichische Akademie der Wissenschaften, 73 (2005), ISBN 3-7001-3616-1
- [12] A. Schüttauf *et al.*, *Nucl. Instr. Meth. A* 602 (2009) 679
- [13] P. Crochet *et al.* [FOPI collaboration], *Phys. Lett. B* 486 (2000) 6
- [14] S.A. Voloshin, Y.Z. Zhang, *Z. Phys. C* 70 (1996) 665
- [15] P. Danielewicz and G. Odyniec, *Phys. Lett. B* 157 (1985) 146
- [16] J.Y. Ollitrault, *nucl-ex/9711003*, (1997)
- [17] J. Ritman *et al.* [FOPI collaboration], *Z. Phys. A* 352 (1995) 355
- [18] N. Herrmann, J.P. Wessels, T. Wienold, *Ann. Rev. Nucl. Part. Sci.*, 49, 581 (1999)
- [19] M. Merschmeyer *et al.* [FOPI collaboration], *Phys. Rev. C* 76 (2007) 02490
- [20] X. Lopez *et al.* [FOPI Collaboration], *Phys. Rev. C* 76 (2007) 052203
- [21] X. Lopez [FOPI Collaboration], *J. Phys. G* 35 (2008) 044020
- [22] N. Herrmann [FOPI Collaboration], *Prog. Part. Nucl. Phys.* 62 (2009) 445
- [23] O. Braun *et al.*, *Nucl. Phys. B* 124 (1977) 45
- [24] C. Pigeot *et al.*, *Nucl. Phys. B* 249 (1985) 172
- [25] R. Siebert *et al.*, *Nucl. Phys. A* 567 (1994) 819