



**HAL**  
open science

## Dose-Responsive Gene Expression Changes in Juvenile and Adult Mummichogs () After Arsenic Exposure

Horacio O. Gonzalez, Jianjun Hu, Kristen M. Gaworecki, Jonathan A. Roling, William S. Baldwin, Jorge L. Gardea-Torresdey, Lisa J. Bain

► **To cite this version:**

Horacio O. Gonzalez, Jianjun Hu, Kristen M. Gaworecki, Jonathan A. Roling, William S. Baldwin, et al.. Dose-Responsive Gene Expression Changes in Juvenile and Adult Mummichogs () After Arsenic Exposure. *Marine Environmental Research*, 2010, 70 (2), pp.133. 10.1016/j.marenvres.2010.04.003 . hal-00602615

**HAL Id: hal-00602615**

**<https://hal.science/hal-00602615>**

Submitted on 23 Jun 2011


**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Accepted Manuscript

Title: Dose-Responsive Gene Expression Changes in Juvenile and Adult Mummichogs (*Fundulus heteroclitus*) After Arsenic Exposure

Authors: Horacio O. Gonzalez, Jianjun Hu, Kristen M. Gaworecki, Jonathan A. Roling, William S. Baldwin, Jorge L. Gardea-Torresdey, Lisa J. Bain



PII: S0141-1136(10)00054-1

DOI: [10.1016/j.marenvres.2010.04.003](https://doi.org/10.1016/j.marenvres.2010.04.003)

Reference: MERE 3441

To appear in: *Marine Environmental Research*

Received Date: 24 June 2009

Revised Date: 9 September 2009

Accepted Date: 10 April 2010

Please cite this article as: Gonzalez, H.O., Hu, J., Gaworecki, K.M., Roling, J.A., Baldwin, W.S., Gardea-Torresdey, J.L., Bain, L.J. Dose-Responsive Gene Expression Changes in Juvenile and Adult Mummichogs (*Fundulus heteroclitus*) After Arsenic Exposure, *Marine Environmental Research* (2010), doi: 10.1016/j.marenvres.2010.04.003

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1  
2  
3 **Dose-Responsive Gene Expression Changes in Juvenile and Adult Mummichogs (*Fundulus heteroclitus*)**  
4  
5 **After Arsenic Exposure**  
6

7  
8  
9  
10  
11 Horacio O. Gonzalez<sup>a</sup>, Jianjun Hu<sup>b</sup>, Kristen M. Gaworecki<sup>c</sup>, Jonathan A. Roling<sup>c</sup>, William S. Baldwin<sup>c</sup>, Jorge L.  
12 Gardea-Torresdey<sup>d</sup>, and Lisa J. Bain<sup>c\*</sup>  
13  
14

15  
16  
17  
18  
19  
20 <sup>a</sup>Department of Biological Sciences, University of Texas at El Paso, El Paso, TX, USA  
21

22 <sup>b</sup>Department of Computer Science and Engineering, University of South Carolina, Columbia, SC, USA  
23

24 <sup>c</sup>Department of Biological Sciences, Clemson University, Clemson, SC, USA  
25

26 <sup>d</sup>Department of Chemistry, University of Texas at El Paso, El Paso, TX, USA  
27

28 <sup>e</sup>Department of Biological Sciences, Bridgewater State College, Bridgewater, MA, USA  
29  
30

31  
32  
33  
34  
35  
36  
37  
38 \*To whom correspondence should be addressed: Phone: +1 864 646 2187; FAX: +1 864 646 2277; Email:  
39

40 lbain@clemson.edu  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**ABSTRACT**

The present study investigated arsenic's effects on mummichogs (*Fundulus heteroclitus*), while also examining what role that gender or exposure age might play. Adult male and female mummichogs were exposed to 172ppb, 575ppb, or 1,720ppb arsenic as sodium arsenite for 10 days immediately prior to spawning. No differences were noted in the number or viability of eggs between the groups, but there was a significant increase in deformities in 1,720ppb arsenic exposure group. Total RNA from adult livers or 6-week old juveniles was used to probe custom macroarrays for changes in gene expression. In females, 3% of the genes were commonly differentially expressed in the 172 and 575ppb exposure groups compared to controls. In the males, between 1.1-3% of the differentially expressed genes were in common between the exposure groups. Several genes, including apolipoprotein and serum amyloid precursor were commonly expressed in either a dose-responsive manner or were dose-specific, but consistent across genders. These patterns of regulation were confirmed by QPCR. These findings will provide us with a better understanding of the effects of dose, gender, and exposure age on the response to arsenic.

**Keywords:** Arsenic, *Fundulus heteroclitus*, fish, apolipoprotein, serum amyloid precursor protein, arrays, gene expression

## 1. Introduction

Arsenic is an element present in water bodies throughout the world, due to both natural and anthropogenic processes (Mandal and Suzuki, 2002; Smedley and Kinniburgh, 2002). Because of its widespread presence, both the U.S. Agency for Toxic Substances and Disease Registry and the World Health Organization have classified arsenic as the number one substance of health concern in the world (ATSDR, 2007; NRC, 1999). Long-term exposure to arsenic is toxic to organisms. For example, epidemiological studies have shown correlations between disease and human exposure to arsenic in drinking water in Taiwan, Chile, Mexico, Japan, Germany and Bangladesh (reviewed in Bates et al., 1992; Engel et al., 1994; Col et al., 1999; NRC, 1999; Tseng et al., 2000; Yoshida et al., 2004; Chen et al., 2007; Díaz-Villaseñor et al., 2007; Hays et al., 2008; NRC, 1999; Raqib et al., 2009). Disease outcomes reported include increases in hyperpigmentation, keratosis, cancer, diabetes, cardiovascular disease, respiratory illnesses, and developmental and reproductive problems (Bates et al., 1992). In light of the epidemiological findings, the U.S. Environmental Protection Agency has placed arsenic as one of the top pollutants of environmental concern, and implemented a reduction in the drinking water standard from 50ppb to 10ppb starting in 2006 (EPA, 2001). However, the effects of arsenic at these low exposure levels are not yet clearly understood (NRC, 1999).

In aquatic organisms, sublethal exposure to arsenic results in several deleterious effects, such as immune system suppression in zebrafish (*Danio rerio*) and in the Indian catfish (*Clarias batrachus*) (Datta et al., 2009), hepatocyte proliferation in the Indian catfish (Datta et al., 2007), altered ability of mummichogs (*Fundulus heteroclitus*) to adapt to their environment (Bears et al., 2006; Shaw et al., 2007a; Shaw et al., 2007b), and changes in antioxidant enzymes in zebrafish (Ventura-Lima et al., 2009). Arsenic exposure has also caused mutations in the p53 gene in salamanders (*Hynobius leechii*) (Chang et al., 2009).

Additionally, recent studies in fish link changes in reproduction and development to arsenic exposure. For example, exposure to arsenic resulted in a reduction in 11-ketotestosterone-induced spermatogenesis in catfish (*Pangasianodon hypophthalmus*) (Yamaguchi et al., 2007), while zebrafish fed metal-contaminated

1  
2  
3 oligochaetes for 42-68 days had a significant reduction in the cumulative number of eggs, number of spawns, and  
4  
5 percent of hatching success (Boyle et al., 2008). Zebrafish exposed to high concentrations of arsenic during  
6  
7 development (64,950-259,800ppb) showed dorsal curvature, cardiac edema along with cardiac malformations,  
8  
9 inappropriate apoptosis and methylation patterns, and altered development of the neuro-muscular system (Li et  
10  
11 al., 2009). These findings corroborate studies in which offspring of mummichogs whose parents were exposed to  
12  
13 230ppb arsenic had a 2.8-fold increase in trunk curvatures. These abnormalities were correlated with differential  
14  
15 expression of genes important in cellular and organismal structure, such as myosin light chain 2, tropomyosin,  
16  
17 parvalbumin and type II keratin genes (Gonzalez et al., 2006). These effects are supported by the fact that arsenic  
18  
19 has been shown to be transferred from the mother to the offspring in a variety of species (Concha et al., 1998;  
20  
21 Kubota et al., 2005; Fängström et al., 2009). There also does appear to be gender-based differences in diseases  
22  
23 caused by arsenic, although whether males or females are more sensitive seems to depend on the specific type of  
24  
25 effect (reviewed in (Vahter et al., 2007)).  
26  
27  
28  
29  
30

31  
32 Thus, there is a need to investigate arsenic's effects on aquatic organisms at low concentrations, while  
33  
34 also examining what modifying role that gender or age of exposure may play. One way to do this is through the  
35  
36 use of microarrays. Arsenic exposure can alter the expression of genes including those involved in stress  
37  
38 response, proto-oncogenes, signaling molecules, transcription factors, chemokine receptors, and DNA repair  
39  
40 enzymes in a variety of cell lines at concentrations ranging from 0.125  $\mu\text{M}$  to 25 $\mu\text{M}$  (Chen et al., 2001; Zheng et  
41  
42 al., 2003; Snow et al., 2005; Posey et al., 2008; Yamamoto et al., 2008). A number of studies have demonstrated  
43  
44 the utility of microarrays in environmental toxicogenomics. For example, a sheepshead minnow estrogen  
45  
46 responsive array was developed to monitor the action of xenoestrogens in aquatic environments (Larkin et al.,  
47  
48 2002). Arrays have also been developed for *Fundulus* to examine differential gene expression after chromium  
49  
50 exposure to both adults and juveniles (Roling et al., 2006). Microarrays were used to identify biomarkers of  
51  
52 cadmium exposure in the European flounder (*Platichthys flesus*) (Shedder et al., 2006), and in rainbow trout  
53  
54 (Koskinen et al., 2004). Clusters of rainbow trout genes involved in energy metabolism, protein synthesis, and  
55  
56 metal ion transport were upregulated at high exposures (0.5mg/L cadmium), while downregulated at medium or  
57  
58 low exposures (0.25mg/L, 0.05mg/L respectively). In contrast, other genes involved in stress response, receptor  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 95 signaling, G-protein coupled receptor, lipid biosynthesis and sulfur metabolism were upregulated at low doses and  
4  
5 96 repressed at high doses (Koskinen et al., 2004).  
6  
7 97  
8 98

9 99  
10 Only a few studies have used microarrays to examine arsenic's effects on aquatic organisms. Using  
11 arrays, zebrafish exposed to 15ppm arsenic had decreased hepatic glycogen, increased incidence of cholestasis,  
12 and changes in overall hepatocyte morphology. Concurrent with these findings, the authors found genes involved  
13 in carbohydrate catabolism, DNA repair, and oxidant status-related proteins to be differentially expressed (Lam et  
14 al., 2006). Consequently, the present study used microarrays to investigate changes in the patterns of gene  
15 expression after adult and parentally-exposed juvenile mummichogs were exposed to three arsenic concentrations.  
16 These findings will provide us with a better understanding of dose-response relationships and the effects of  
17 gender and age on the response to arsenic.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

## 33 2. Methods

### 34 2.1. Fish exposures

35  
36 100 Mummichogs (2 males and 5 females) were randomly assigned to one of 10 different aquaria, 5 of which  
37 contained 20ppt saltwater (Coral Life, Burbank, CA) and 5 of which contained sodium arsenite in 20ppt saltwater.  
38 Three separate studies were conducted using the same set-up, but one exposure was with 172ppb (1.32 $\mu$ M), the  
39 second with 575ppb (4.43 $\mu$ M), and the third with 1,720ppb (13.2 $\mu$ M) arsenic, provided as sodium arsenite. Fish  
40 were exposed for a total of 10 days, maintained on a natural photoperiod, and fed daily with AquaTox flaked fish  
41 food (Zeigler Brothers, Gardners, PA) supplemented with freeze-dried brine shrimp. All studies were conducted  
42 using a static-renewal exposure, with 80% of the water being replaced every 48 hours. Although arsenic  
43 concentrations in the control aquaria water were below the detection limits, arsenic concentrations in the food  
44 (0.03ng/g) are similar to what others have reported (Shaw et al., 2007b).  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

56 101 Spawning substrates were placed in each tank on the three days surrounding the full moon to collect eggs.  
57 The total number of eggs produced was counted and their viability was determined. Eggs from each of the  
58 aquaria on each day were placed into petri dishes containing 20ppt saltwater to monitor their viability,  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 22 development, and time to hatch. After completion of spawning, the parents were euthanized in 1g/L buffered MS-  
4  
5 222 and two female fish from each group were stored at -80°C to determine arsenic body burdens. For the  
6  
7 224 remaining adults, livers were removed, placed in TriReagent, and stored at -80°C. The hatchlings from each  
8  
9 225 exposure group were transferred to an individual 1 gallon aquarium containing 20ppt seawater, maintained on a  
10  
11 226 16-h light/8-h dark photoperiod at 25°C, and fed recently hatched brine shrimp. The juveniles were observed  
12  
13 227 over the next six weeks to determine whether additional morphological changes, developmental delays, or death  
14  
15 228 would occur, but no additional phenotypes were noticed. Six weeks after hatching, the juveniles were euthanized  
16  
17 229 with an overdose of buffered MS-222 and were stored at -80°C.

## 20 21 230 22 23 231 2.2. *Chemical residue analysis.*

24  
25 232 One intact adult female fish carcass from each aquaria in 1,720ppb experiment was freeze-dried  
26  
27 233 (Labconco, Kansas City, MO), microwave digested (CEM, Mathews, NC) following the USEPA 3051 method  
28  
29 234 (Kingston and Jassie, 1988) and arsenic levels analyzed using an Inductively Coupled Plasma Optical Emission  
30  
31 235 Spectrometer (ICP/OES) Optima 4300 DV (Perkin-Elmer Instruments, Edgewood, NM). Arsenic was detected at  
32  
33 236 189nm, which has a detection limit of 3ppb. A standard curve for sodium arsenite was used to determine body  
34  
35 237 burdens in the fish, and statistical differences between the groups were determined using Student's t-test.

## 38 39 238 40 41 239 2.3. *Morphometric analyses of the vertebrae*

42  
43 240 Ten control and ten 575ppb parentally-exposed juveniles were examined for changes in vertebral and fin  
44  
45 241 bone size and number. Briefly, the fish were fixed in 10% formalin for 7 days and placed in Alcian Blue dye in  
46  
47 242 80% ethanol/20% glacial acetic acid solution for 12 hours. After neutralization with potassium hydroxide, the  
48  
49 243 specimens were bleached in 3% hydrogen peroxide. The fish were then digested in a sodium borate/trypsin  
50  
51 244 solution until bones and cartilage were visible, and stored in glycerol (Taylor and Van Dyke, 1985). A confocal  
52  
53 245 microscope (Carl Zeiss, Thornwood, NY) was used to quantify the area of each vertebra, which was normalized to  
54  
55 246 the weight of each fish.

## 58 59 247 60 61 248 2.4. *Gene Expression Changes using Macroarrays.*



1  
2  
3 149 Targeted mummichog cDNA arrays were constructed using genes from subtractive hybridization or  
4  
5 150 differential display experiments from fish exposed in the laboratory or collected from field sites containing  
6  
7 151 Cr(VI), Cr(III), anthracene, pyrene, and arsenic (Bain, 2002; Maples and Bain, 2004; Peterson and Bain, 2004;  
8  
9 152 Roling et al., 2004; Roling et al., 2006; Gonzalez et al., 2006). Each membrane contained 270 duplicate clones,  
10  
11 153 including 13 blanks or plasmid controls. Specific information on the array platform is available on the Gene  
12  
13 154 Expression Omnibus website (GEO-GPL2535 ) (Roling et al., 2006).  
14  
15  
16 155

17  
18 156 For each experiment, five arrays were probed using RNA samples from control groups and five were  
19  
20 157 probed with samples from the arsenic-exposed groups. Array experiments were conducted for the 172, 575, and  
21  
22 158 1,720ppb males and for the 172 and 575ppb females. RNA was extracted from a pool of 7 to 10 juveniles per  
23  
24 159 aquarium using TriReagent (Sigma, St. Louis, MO) and treated with DNase I. The pools of RNA were obtained  
25  
26 160 from the 172, 575, and 1,720ppb exposure groups, plus their respective controls. Probes were generated by  
27  
28 161 reverse transcription using 2µg of RNA, 0.1µM oligo dT primer, 1.2mM dNTPs lacking dATP, [<sup>33</sup>P]dATP  
29  
30 162 (3000Ci/mmol, Perkin Elmer, Boston, MA) and 200U MMLV-RT at 37°C for 2 hours. Meanwhile, the array  
31  
32 163 membranes were prehybridized for 4-6 hour in Express hybridization solution (Clontech, Palo Alto, CA)  
33  
34 164 containing 100mg/ml denatured salmon sperm. Each probe was purified using Amersham's Probe Quant-50  
35  
36 165 columns (Piscataway, NJ), denatured, and incubated overnight with the array at 68°C. Then, the arrays were  
37  
38 166 washed four times with 2X SSC/0.5%SDS at 64°C and four times with 0.5X SSC/0.5%SDS at 64°C. The  
39  
40 167 membranes were wrapped in plastic, placed on a phosphorimaging screen, and developed using a Bio-Rad  
41  
42 168 Molecular Imager FX (Hercules, CA). The intensity of each spot was quantified using the Invitrogen ResGen  
43  
44 169 Pathways 4-Universal Microarray Analysis Software (Carlsbad, CA). The data for this series of experiments has  
45  
46 170 been deposited in NCBI's Gene Expression Omnibus (Edgar et al., 2002) and can be accessed using GEO series  
47  
48 171 accession number GSE16772 (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE16772>).  
49  
50  
51  
52 172

### 53 54 173 2.5. *Microarray analysis.* 55 56 57

58  
59 174 The CLEAR-test method was used to analyze the microarray data for identifying differentially expressed  
60  
61 175 genes between normal state and different degrees of arsenic exposure for each fish group (Valls et al., 2008). This  
62  
63  
64  
65

1  
2  
36 method combines the z-test, which focuses on large changes, with a  $\chi^2$  test to evaluate variability, and has the  
4  
57 advantage of not reporting genes with small changes and low variances as differentially expressed. For all the  
6  
78 duplicate spots, the raw pixel intensity values were first averaged before running CLEAR-test. Average fold  
8  
99 expression changes for each gene were obtained for each of the three arsenic concentrations in the male, female,  
10  
110 and juvenile groups.  
12

## 13 14 15 16 182 2.6. Changes in RNA abundance by real-time PCR 18

19  
20 Total RNA from the liver of 5 adult fish of each gender per exposure group was isolated using TRI-  
21  
22 Reagent (Sigma) and then treated with RNase-free DNase. Total RNA from a pooled sample of 3 whole juveniles  
23  
24 was also isolated after homogenization (n=5 pools per group). To prepare cDNA, total RNA (2 $\mu$ g) was incubated  
25  
26 with 50ng random hexamers, RNasin, 10mM dNTP mix, and 200U Moloney murine leukemia virus (MMLV)  
27  
28 reverse transcriptase at 37°C for 1 hour. Real-time PCR was performed in Bio-Rad's I-Cycler (Hercules, CA)  
29  
30 using RT<sup>2</sup> SYBR Green/Fluorescein qPCR master mix (SABiosciences, Frederick, MD), along individual sets of  
31  
32 primers for the EST 9-7 (#CO897482; forward primer 5'-GCCAACACCATGTGTGCAAGTTCT-3'; reverse  
33  
34 primer 5'-TCTGGACACAGTCACACACAGACT-3'), apolipoprotein (8-56) (#DN474949; forward primer 5'-  
35  
36 ATCTTGCAGCTGCTGTTCTGTTGC-3'; reverse primer 5'-GTGTCAAAGTGGGTCTTGGCCTTT-3'), and  
37  
38 serum amyloid precursor (1-31) (#AY735160, forward primer 5'-AGGTTATCAGCGATGCCAGAGAGT-3';  
39  
40 reverse primer 5'-ATTTGCTTCCTGATCGGCCTCTGA-3'). 18S rRNA (#X00686; forward primer 5'-  
41  
42 TTTCTCGATTCTGTGGGTGGTGGT-3'; reverse primer 5'-TAGTTAGCATGCCGGAGTCTCGTT-3') was  
43  
44 used as a housekeeping gene to normalize the total number of molecules in each sample. All PCR products had a  
45  
46 denaturing step of 95°C for 15 seconds, an annealing/extension step at 57°C for 1 minute for a total of 40 cycles.  
47  
48 The cycle threshold values obtained from the real-time PCR were converted into starting number of molecules per  
49  
50 100ng cDNA using known concentrations of the specific gene product, which was normalized to the number of  
51  
52 18S rRNA molecules (Gonzalez et al., 2006). The standards were prepared by RT-PCR and sequenced at  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

### 3. Results

#### 3.1. Arsenic body burdens

At the conclusion of spawning, female fish from each control and 1,720ppb arsenic aquaria were used to determine the body burdens of arsenic. In the adults, arsenic levels were 2.6-fold higher in the exposed fish than in the controls after 10 days of exposure (Figure 1). This level of uptake is consistent with other studies in *Fundulus*. For example, adult mummichogs exposed to 8,000ppb arsenic for 4 days had approximately 6 times more arsenic than controls while those fish exposed to 100ppb arsenic for 14 days had approximately 3 times more arsenic than the controls (Miller et al., 2007). Similarly, mummichogs exposed to 787 ppb for 14 days had 2.8-fold higher hepatic arsenic levels than controls (Bears et al., 2006).

#### 3.2. Hepatic gene expression changes in adult *Fundulus* exposed to arsenic

Microarrays were used to determine both the dose-dependent patterns of gene expression changes in livers following arsenic exposure and whether the response to arsenic differed between genders. Replicated test array experiments were conducted for 2 different exposure concentrations in the females or 3 different exposure concentrations in the males. Of the 257 genes on the array, between 4.7-8.2% were differentially expressed after arsenic exposure amongst the 5 groups of experiments using the CLEAR test method. Analyzing the data for significantly differential changes in gene expression in a dose-responsive manner indicated that in the two female exposure groups, 172ppb and 575ppb arsenic, 16 and 19 hepatic genes were significantly changed compared to the controls, respectively. There were 7 common differentially expressed genes (Figure 2A). In comparing the livers of male fish exposed to arsenic with the controls, 21, 12, and 13 hepatic genes were differentially expressed in 172, 575, and 1,720ppb exposures. There was only 1 gene shared in common between all three male groups, which is an EST termed 10-15. Shared genes between two of the three male exposures ranged from 3 to 7 (Figure 2A). When examining the data from Table 1 for the number of common genes changed after arsenic exposure based upon gender, a total of 9 differentially expressed genes were common between males and females. The differential expression was more likely to occur in the 172ppb exposure group than the 575ppb group (Figure 2B).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Several genes were expressed in a dose-responsive manner. Only genes with statistically different expression are shown in Table 1. These included apolipoprotein, which was significantly downregulated in both female exposure groups and in two of the three male exposure groups. Apolipoprotein is represented by two partially overlapping clones on the array, 8-56 and Ab35, and both were changed in the same manner and the same magnitude (Table 1). Lysozyme precursor is also represented by two clones, 1-11 and 1-163, with both consistently demonstrating downregulation in the 172ppb males but upregulation in the 1,720ppb males exposed to arsenic (Table 1). Other clones show differences based upon gender, such as tributyltin (TBT)-binding protein, which is downregulated in females but upregulated in males (Table 1). A second overlapping clone for TBT-binding protein termed C3C61 (accession #[BM084929](#)) was also significantly downregulated (0.33-fold) in the 575ppb arsenic-exposed females (data not shown). Several other genes were differentially expressed in only one exposure group, but many of these genes are associated with cell stress, damage, and repair, such as heat shock proteins, glutathione S-transferases, transcription elongation factors, and ATP synthase (Table 2).

### 3.3. Reproductive effects, developmental effects, and gene expression changes in the juveniles

In addition to changes in hepatic gene expression, we were interested in investigating the effects of arsenic on embryonic development. After the exposure period, the fish were mated and eggs collected to determine changes in embryonic development and hatchling success. There were no differences in the number of eggs laid per tank between the control and the arsenic-exposed groups at any concentration, nor any differences in egg viability (Figure 3a). After hatching, any deformities in the embryos were recorded. In the offspring whose parents were exposed to 1,720ppb arsenic, there was a significant increase in deformities including embryos with abnormal trunk curvatures, and head and eye deformities (Figure 3b). Because of these abnormalities, it was hypothesized that arsenic could be altering the structure of bone or cartilage. However, no differences were observed in the number, orientation, or position of the vertebrae in parentally-exposed offspring compared to control offspring (Figure 3c).

Unlike the adult hepatic gene expression, we used RNA extracted from whole 6-week old juveniles. There were very few genes in common between the three different parental exposure groups, except for two

ESTs, 2-80 and 9-7 (Table 1). There was only one additional gene, complement component C8, which was shared between the juveniles and the adults. However, this gene was only significantly differentially expressed in the highest juvenile exposure group.

#### 3.4. Confirmation of gene expression patterns

To confirm the expression patterns of several of the genes, we used quantitative PCR (QPCR) to examine expression levels of several selected genes in the livers of the adult males and females. Expression levels of the EST shared in common with the adults, 9-7, was also examined in the juveniles (Figure 4). Apolipoprotein expression on the arrays was downregulated in all adult arsenic exposed groups, regardless of gender or concentration. This was validated in four of the five QPCR reactions. The exception was the 172ppb female group, which based upon QPCR expression, showed no change in expression (fold change=1.03). The EST termed 9-7 showed a bit more variability, but the arrays and QPCR validated one another in 5 of the 7 groups (Figure 4). The expression of 9-7 in the 172ppb female group was downregulated in the arrays but upregulated in the QPCR, while the 575ppb male group was upregulated in the arrays but downregulated based upon QPCR. The levels of 9-7 in two juvenile groups examined showed similar expression patterns, being upregulated in both the arrays and when using QPCR (Figure 4).

Serum amyloid precursor expression was downregulated in all of the male exposures and the 172ppb female exposure, but upregulated in the 575ppb female exposure group in the arrays. Again, the direction of change using QPCR matched in four of the five groups, but this time, the 172ppb male group showed no changes in expression based upon the QPCR data (fold change=1.07). For the higher exposure groups, both the directionality and magnitude of change were similar. The expression of serum amyloid precursor in the 575ppb males was downregulated 0.83-fold in the arrays and 0.67-fold in the QPCR, while the expression in the 1,720ppb males was downregulated 0.32-fold in the arrays and 0.28-fold in the QPCR. Interestingly, serum amyloid precursor expression was upregulated in the 575ppb female group, and the magnitude of change between the arrays and QPCR was also similar (2.5-fold in arrays; 3.7-fold in the QPCR). Overall, the QPCR data showed similar fold-changes and directionality, validating the array data.

#### 4. Discussion

##### 4.1. Arsenic-mediated changes in gene expression

The use of fish toxicogenomics is suited for studying contaminants such as arsenic, as fish can act as both models for human diseases as well as models for the health of the aquatic environment ( Williams et al., 2003; Gorman and Breden, 2007). In the present study, adult mummichogs were exposed to three different arsenic concentrations, and the differential expression of genes was investigated by examining the effects of gender and maternal transfer to offspring. Several genes were differentially expressed as a result of the arsenic exposure, including apolipoprotein, serum amyloid precursor, tributyltin-binding protein, and cytochrome c oxidase. Overall, arsenic appears to perturb genes involved in energy utilization.

Apolipoprotein was one gene that was downregulated in the adult livers at all exposure concentrations tested. Using QPCR, the underexpression of apolipoprotein and the magnitude of downregulation were in concordance with the array data. Apolipoproteins are lipid binding proteins important in lipid secretion, lipolytic enzyme activation, and transport and binding of lipoproteins in cells (Atkinson, 1992). Several isoforms have been identified, which bind to and act as ligands for low density or very low density lipoprotein receptors (Atkinson, 1992). In our experiment, both clones of the apolipoprotein C-1 were down-regulated similarly in females as well as males. The reduction in apolipoprotein expression is in concordance with other arsenic exposure studies. For example, arsenic downregulated apolipoprotein in the lung fluid of male mice exposed to 50ppb sodium arsenite for four weeks (Lantz et al., 2008), and was differentially expressed in zebrafish embryos exposed to arsenic (Mattingly et al., 2009).

Another protein, serum amyloid precursor, was also downregulated in the adult males and in the 172ppb female exposure group. Its altered expression was examined by QPCR, which confirmed a dose-responsive reduction in the male 172, 575, and 1720ppb groups. Serum amyloid precursor proteins (APPs) are also a family of apolipoproteins (Zheng and Koo, 2006). Although most often associated with the development of Alzheimer's disease (reviewed in: (Esler and Wolfe, 2001; Jaeger and Pietrzik, 2008), APPs are thought to function to control

1  
2  
3 cholesterol transport, ApoE metabolism, and help maintain cholesterol homeostasis (Yao and Papadopoulos,  
4  
5 2002; Liu et al., 2007). Additionally, cellular cholesterol can modify the both the processing and transcription of  
6  
7 APP (Bodovitz and Klein, 1996; Fassbender et al., 2001; Kiyosawa et al., 2004). In general, the downregulation  
8  
9 of APP in the adult mummichogs mimics the reduction in apolipoprotein expression.  
10

11  
12  
13 Both apolipoprotein and APPs are involved in lipid homeostasis, and in other studies, exposure to arsenic  
14  
15 does appear to alter lipid levels. Mice exposed to arsenic via drinking water from weaning to 1 year of age, or  
16  
17 from 8 weeks to 1 year of age had significant reductions in serum triglycerides, total cholesterol, and HDL  
18  
19 cholesterol compared to controls (Ahlborn et al., 2009). The authors hypothesized that the reduction in lipids  
20  
21 could be due to alterations in fatty acid production, as there was a reduction in the mRNA expression of stearoyl-  
22  
23 CoA desaturase 1, the rate limiting enzyme in monounsaturated fatty acid synthesis (Ahlborn et al., 2009).  
24  
25 Interestingly, the mice in their study exposed to arsenic only *in utero* did not have alterations in serum lipids  
26  
27 (Ahlborn et al., 2009). These findings are in concordance with our studies showing a marked reduction in both  
28  
29 apolipoprotein and APP proteins in the adults, but not in the offspring. Thus, arsenic exposure may impair  
30  
31 cholesterol homeostasis, which could have adverse physiological consequences for fish (Heath, 1995).  
32  
33 Examining lipid levels in the livers and serum of arsenic-exposed fish, and how that may impact adult  
34  
35 health and egg quality would be worthy of future investigation.  
36  
37  
38  
39  
40  
41  
42  
43

44  
45 Indeed the present study, along with many others, shows that arsenic often exhibits a complex gene  
46  
47 expression profile. For example, arsenic exposure in zebrafish embryos resulted in the differential expression of  
48  
49 several immune-responsive genes following a 10µg/L exposure, but no change in expression at 100µg/L  
50  
51 (Mattingly et al., 2009). In additional microarray studies using cell lines or rodent models, arsenic exposure  
52  
53 results in very few commonly differentially expressed genes between the exposure groups (Ahlborn et al., 2008),  
54  
55 or can alter the expression of genes in opposite directions at low versus high concentrations (Andrew et al., 2003;  
56  
57 Lau et al., 2004; Snow et al., 2005; Andrew et al., 2007). This may indicate that differing transcriptional  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
335 networks are activated at low arsenic levels versus high arsenic levels, resulting in differing biological processes  
4  
336 being altered ( Bodwell et al., 2004; Bodwell et al., 2006).

#### 337 8 338 4.2. *Gender differences in gene expression*

339 We also found differing responses depending on the gender of the fish. For example, serum amyloid  
12  
340 precursor was upregulated in the livers of the highest female exposure group, but downregulated in the livers of  
13  
341 the males. In contrast, several ESTs were upregulated in the males by downregulated in the females. Gender-  
15  
342 based differences in disease outcomes after arsenic exposure have been reported in the literature. For example,  
17  
343 men have a higher incidence of pathological liver changes after arsenic exposure, while women are at greater risk  
19  
344 of arsenic-related kidney, lung, and bladder cancers than men ( Vahter et al., 2007; Lindberg et al., 2008b). The  
21  
345 same increase in bladder lesions in females has also been reported in studies with rodents (Shen et al., 2006;  
22  
346 Waalkes et al., 2006), while increases in liver tumors after arsenic exposure are seen in male rodents ( Waalkes et  
24  
347 al., 2003; Waalkes et al., 2004a; Waalkes et al., 2004b; Ahlborn et al., 2009). It is known that hepatic  
26  
348 metabolism of contaminants in some organisms can vary by gender (reviewed in Burger et al., 2007) and it has  
28  
349 been speculated that sex hormones affect in arsenic metabolism, altering its methylation, excretion, and thus its  
30  
350 pathology ( Shen et al., 2006; Lindberg et al., 2008a; Lindberg et al., 2008b). As the fish in our study were  
32  
351 reproductively active, hormonal changes may have influenced the effects of arsenic on gene expression.  
34  
352

#### 353 4.3. *Effect of arsenic on development*

354 The second part of this study was to examine changes in development and gene expression in the  
355 offspring whose parents were exposed to arsenic. Viability was not different between the control and arsenic-  
356 exposed groups at any exposure concentration. However, parental exposure to arsenic increased the incidence of  
357 developmental deformities, such as spinal curvatures and smaller heads, by 17.4-fold at the highest concentration  
358 as compared to controls. In a previous experiment using a concentration of 230ppb arsenic, there was a 2.8-fold  
359 increase in the incidence of spinal deformities in the hatchlings of exposed parents (Gonzalez et al., 2006). A  
360 study with zebrafish revealed various similar morphological abnormalities including dorsal curvature, flattened  
361 head, and pericardial edema (Li et al., 2009), and mice exposed *in utero* via maternal oral treatment also



1  
2  
362 exhibited skeletal abnormalities (Hill et al., 2008). In humans, a decreased head and chest size was noted in  
4  
363 children exposed to arsenic *in utero* (Rahman et al., 2009).  
6  
364

9  
365 It has been well documented that arsenic can cross the maternal blood supply and be detected in the fetus  
10  
366 (Concha et al., 1998; Kubota et al., 2005; Fångström et al., 2009). A study looking at the maternal transfer of  
13  
367 arsenic from organisms such as adult black-tailed gulls noted that 11.3% of the arsenic burden was transferred to  
15  
368 eggs (Kubota et al., 2002), fetal porpoise arsenic burden was about 36% of that measured in the mothers (Kubota  
17  
369 et al., 2005), and newborn mouse livers had 10% of the average arsenic concentration in maternal livers (Xie et  
20  
370 al., 2007). Thus, even the low amounts of arsenic transferred to the offspring may impact its gene expression and  
22  
371 subsequent development.  
24  
372

27  
373 It is interesting to note that offspring exposed to arsenic *in utero* have very different pathologies based  
28  
374 upon their gender. In mice exposed to arsenic *in utero*, males have increase in liver and adrenal tumors, while  
31  
375 females have increases in urogenital tumors, but no changes in the incidence of liver or adrenal tumors (Waalkes  
33  
376 et al., 2003; Waalkes et al., 2004a; Waalkes et al., 2004b; Ahlborn et al., 2009). We did not determine the sex of  
35  
377 the juveniles in our study before using them in the microarray study. Given that in rodents, very different arsenic  
37  
378 metabolite profiles and pathologies are seen, not sexing our juvenile fish might have contributed to the lack of  
40  
379 gene expression changes and the high amount of variability associated with the offspring.  
42  
380

45  
381 In summary, we observed changes in lipid responsive genes at environmentally-relevant concentrations,  
47  
382 indicating that arsenic exposure may impair cholesterol homeostasis, which could have adverse effects on  
49  
383 organismal health. These findings will provide us with a better understanding of the effects of dose, gender, and  
51  
384 exposure age on the response to arsenic.  
53  
54  
55  
385  
57  
58  
386  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Acknowledgements:**

This work was supported by NCRR #2G12 RR008124, NSF/EPA #EF 0830117, and CSREES/USDA, under project number #SC-1700380. It is technical contribution No. 5647 of the Clemson University Experiment Station. The authors would also like to thank Marian Viveros and Judi Ellzey (UTEP) for technical assistance.

## References:

- Ahlborn, G.J., Nelson, G.M., Grindstaff, R.D., Waalkes, M.P., Diwan, B.A., Allen, J.W., Kitchin, K.T., Preston, R.J., Hernandez-Zavala, A., Adair, B., Thomas, D.J., Delker, D.A., 2009. Impact of life stage and duration of exposure on arsenic-induced proliferative lesions and neoplasia in C3H mice. *Toxicology* 262, 106-113.
- Ahlborn, G.J., Nelson, G.M., Ward, W.O., Knapp, G., Allen, J.W., Ouyang, M., Roop, B.C., Chen, Y., O'Brien, T., Kitchin, K.T., Delker, D.A., 2008. Dose response evaluation of gene expression profiles in the skin of K6/ODC mice exposed to sodium arsenite. *Toxicol. Appl. Pharmacol.* 227, 400-416.
- Andrew, A.S., Bernardo, V., Warnke, L.A., Davey, J.C., Hampton, T., Mason, R.A., Thorpe, J.E., Ihnat, M.A., Hamilton, J.W., 2007. Exposure to arsenic at levels found in U.S. drinking water modifies expression in the mouse lung. *Toxicol. Sci.* 100, 75-87.
- Andrew, A.S., Warren, A.J., Barchowsky, A., Temple, K.A., Klei, L., Soucy, N.V., O'Hara, K.A., Hamilton, J.W., 2003. Genomic and proteomic profiling of responses to toxic metals in human lung cells. *Environ. Health Perspect.* 111, 825-835.
- ATSDR, 2007. 2007 CERCLA priority list of hazardous substances that will be the subject of toxicological profiles and support document. In. Agency for Toxic Substances and Disease Registry, SUDHHS, PHS, City.
- Bain, L.J., 2002. cDNA cloning, sequencing, and differential expression of a heart-type fatty acid-binding protein in the mummichog (*Fundulus heteroclitus*). *Mar. Environ. Res.* 54, 379-383.
- Bears, H., Richards, J.G., Schulte, P.M., 2006. Arsenic exposure alters hepatic arsenic species composition and stress-mediated gene expression in the common killifish (*Fundulus heteroclitus*). *Aquat. Toxicol.* 77, 257-266.
- Bodovitz, S., Klein, W.L., 1996. Cholesterol modulates alphasecretase cleavage of amyloid precursor protein. *J. Biol. Chem.* 271, 4436-4440.
- Bodwell, J.E., Gosse, J.A., Nomikos, A.P., Hamilton, J.W., 2006. Arsenic disruption of steroid receptor gene activation: Complex dose-response effects are shared by several steroid receptors. *Chem. Res. Toxicol.* 19, 1619-1629.
- Bodwell, J.E., Kingsley, L.A., Hamilton, J.W., 2004. Arsenic at very low concentrations alters glucocorticoid receptor (GR)-mediated gene activation but not GR-mediated gene repression: complex dose-response effects are closely correlated with levels of activated GR and require a functional GR DNA binding domain. *Chem. Res. Toxicol.* 17, 1064-1076.
- Boyle, D., Brix, K.V., Amlund, H., Lundebye, A.K., Hogstrand, C., Bury, N.R., 2008. Natural arsenic contaminated diets perturb reproduction in fish. *Environ. Sci. Technol.* 42, 5354-5360.
- Burger, J., Fossi, C., McClellan-Green, P., Orlando, E.F., 2007. Methodologies, bioindicators, and biomarkers for assessing gender-related differences in wildlife exposed to environmental chemicals. *Environ. Res.* 104, 135-152.
- Chang, J.S., Gu, M.B., Kim, K.W., 2009. Effect of arsenic on p53 mutation and occurrence of teratogenic salamanders: their potential as ecological indicators for arsenic contamination. *Chemosphere* 75, 948-954.
- Chen, C.J., Wang, S.L., Chiou, J.M., Tseng, C.H., Chiou, H.Y., Hsueh, Y.M., Chen, S.Y., Wu, M.M., Lai, M.S., 2007. Arsenic and diabetes and hypertension in human populations: A review. *Toxicol. Appl. Pharmacol.* 222, 298-304.
- Chen, H., Liu, J., Merrick, B.A., Waalkes, M.P., 2001. Genetic events associated with arsenic-induced malignant transformation: applications of cDNA microarray technology. *Mol. Carcinogen.* 30, 79-87.
- Col, M., Col, C., Soran, A., Sayli, B.S., Ozturk, S., 1999. Arsenic-related Bowen's disease, palmar keratosis, and skin cancer. *Environ. Health Perspect.* 196, 687-689.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Concha, G., Vogler, G., Lezcano, D., Nermell, B., Vahter, M., 1998. Exposure to inorganic arsenic metabolites during early human development. *Toxicol. Sci.* 44, 185-190.
- Datta, S., Ghosh, D., Saha, D.R., Bhattacharaya, S., Mazumder, S., 2009. Chronic exposure to low concentration of arsenic is immunotoxic to fish: role of head kidney macrophages as biomarkers of arsenic toxicity to *Clarias batrachus*. *Aquat. Toxicol.* 92, 86-94.
- Datta, S., Saha, D.R., Ghosh, D., Majumdar, T., Bhattacharya, S., Mazumder, S., 2007. Sub-lethal concentration of arsenic interferes with the proliferation of hepatocytes and induces *in vivo* apoptosis in *Clarias batrachus L.* *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 145, 339-349.
- Díaz-Villaseñor, A., Burns, A.L., Hiriart, M., Cebrián, M.E., Ostrosky-Wegman, P., 2007. Arsenic-induced alteration in the expression of genes related to type 2 diabetes mellitus. *Toxicol. Appl. Pharmacol.* 225, 123-133.
- Engel, R.R., Hopenhayn-Rich, C., Receveur, O., Smith, A.H., 1994. Vascular effects of chronic arsenic exposure: a review. *Epidemiol. Rev.* 16, 184-209.
- EPA, 2001. National primary drinking water regulations; arsenic and clarifications to compliance and new source contaminants monitoring from the environmental protection agency; Final Rule. *Fed Reg* 66, 6976-7066.
- Esler, W.P., Wolfe, M.S., 2001. A portrait of Alzheimer secretases—new features and familiar faces. *Science* 293, 1449–1454.
- Fängström, B., Hamadani, J., Nermell, B., Grandér, M., Palm, B., Vahter, M., 2009. Impaired arsenic metabolism in children during weaning. *Toxicol. Appl. Pharmacol.* 239, 208-214.
- Fassbender, K., Simons, M., Bergmann, C., Stroick, M., Lutjohann, D., Keller, P., Runz, H., Kuhl, S., Bertsch, T., von Bergmann, K., Hennerici, M., Beyreuther, K., Hartmann, T., 2001. Simvastatin strongly reduces levels of Alzheimer's disease beta-amyloid peptides Abeta 42 and Abeta 40 *in vitro* and *in vivo*. *Proc. Natl. Acad. Sci. USA* 98, 5856–5861.
- Gonzalez, H.O., Roling, J.A., Baldwin, W.S., Bain, L.J., 2006. Physiological changes and differential gene expression in mummichogs (*Fundulus heteroclitus*) exposed to arsenic. *Aquat. Toxicol.* 77, 43-52.
- Gorman, K.F., Breden, F., 2007. Teleosts as models for human vertebral stability and deformity. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 145, 28-38.
- Hays, A.M., Lantz, R.C., Rodgers, L.S., Sollome, J.J., Vaillancourt, R.R., Andrew, A.S., Hamilton, J.W., Camenisch, T.D., 2008. Arsenic-induced decreases in the vascular matrix. *Toxicol. Pathol.* 36, 805-817.
- Heath, A.G., 1995. *Water pollution and fish physiology*. CRC Press, Boca Raton, Fl. .
- Hill, D.S., Wlodarczyk, B.J., Finnell, R.H., 2008. Reproductive consequences of oral arsenate exposure during pregnancy in a mouse model. *Birth Defects Res. (Part B)* 88, 40-47.
- Jaeger, S., Pietrzik, C.U., 2008. Functional role of lipoprotein receptors in Alzheimer's disease. *Curr Alzheimer Res.* 5, 15-25.
- Kingston, H.M., Jassie, L.B. (Eds.), 1988. *ACS Professional Reference Book Series*. Am. Chem. Soc., Washington, DC.
- Kiyosawa, N., Ito, K., Niino, N., Sakuma, K., Kanbori, M., Yamoto, T., Manabe, S., Matsunuma, N., 2004. Effect of serum cholesterol on the mRNA content of amyloid precursor protein in rat livers. *Toxicol. Lett.* 150, 157-166.
- Koskinen, H., Pehkonen, P., Vehniäinen, E., Krasnov, A., Rexroad, C., Afanasyev, S., Mölsa, H., Oikari, A., 2004. Response of rainbow trout transcriptome to model chemical contaminants. *Biochem. Biophys. Res. Comm.* 320, 745-753.
- Kubota, R., Kunito, T., Fujihara, J., Tanabe, S., Yang, J., Miyazaki, N., 2005. Placental transfer of arsenic to fetus of Dall's porpoises (*Phocoenoides dalli*). *Mar. Pollut. Bull.* 51, 845-849.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 488 Kubota, R., Kunito, T., Tanabe, S., Ogi, H., Shibata, Y., 2002. Maternal transfer of arsenic to eggs of  
489 blacktailed gull (*Larus crassirostris*) from Rishiri Island, Japan. *Appl. Organomet. Chem.* 16, 463-  
490 468.
- 491 Lam, S.H., Winata, C.L., Tong, Y., Korzh, S., Lim, W.S., Korzh, V., Spitsbergen, J., Mathavan, S.,  
492 Miller, L.D., Liu, E.T., Gong, Z., 2006. Transcriptome kinetics of arsenic-induced adaptive  
493 response in zebrafish liver. *Physiol. Genomics* 27, 351-361.
- 494 Lantz, R.C., Chau, B., Sarihan, P., Witten, M.L., Pivniouk, V.I., Chen, G.J., 2008. *In utero* and postnatal  
495 exposure to arsenic alters pulmonary structure and function. *Toxicol. Appl. Pharmacol.* 235, 105-  
496 113.
- 497 Larkin, P., Sabo-Attwood, T., Kelso, J., Denslow, N.D., 2002. Gene expression analysis of largemouth  
498 bass exposed to estradiol, nonylphenol, and p,p'-DDE. *Comp. Biochem. Physiol. B Biochem. Mol.*  
499 *Biol.* 133, 543-557.
- 500 Lau, A.T., Li, M., Xie, R., He, Q.Y., Chiu, J.F., 2004. Opposed arsenite-induced signaling pathways  
501 promote cell proliferation or apoptosis in cultured lung cells. *Carcinogenesis* 25, 21-28.
- 502 Li, D., Lu, C., Wang, J., Hu, W., Cao, Z., Sun, D., Xia, H., Ma, X., 2009. Developmental mechanisms of  
503 arsenite toxicity in zebrafish (*Danio rerio*) embryos. *Aquat. Toxicol.* 91, 229-237.
- 504 Lindberg, A.L., Ekström, E.C., Nermell, B., Rahman, M., Lönnerdal, B., Persson, L.A., Vahter, M.,  
505 2008a. Gender and age differences in the metabolism of inorganic arsenic in a highly exposed  
506 population in Bangladesh. *Environ. Res.* 106, 110-120.
- 507 Lindberg, A.L., Rahman, M., Persson, L.A., Vahter, M., 2008b. The risk of arsenic induced skin lesions  
508 in Bangladeshi men and women is affected by arsenic metabolism and the age at first exposure.  
509 *Toxicol. Appl. Pharmacol.* 230, 9-16.
- 510 Liu, Q., Zerbinatti, C.V., Zhang, J., Hoe, H.S., Wang, B., Cole, S.L., Herz, J., Muglia, L., Bu, G., 2007.  
511 Amyloid precursor protein regulates brain apolipoprotein E and cholesterol metabolism through  
512 lipoprotein receptor LRP1. *Neuron* 56, 66-78.
- 513 Mandal, B., Suzuki, T., 2002. Arsenic around the world: a review. *Talanta* 58, 201-235.
- 514 Maples, N.L., Bain, L.J., 2004. Trivalent chromium alters gene expression in the mummichog (*Fundulus*  
515 *heteroclitus*). *Environ. Tox. Chem.* 23, 626-631.
- 516 Mattingly, C.J., Hampton, T.H., Brothers, K.M., Griffin, N.E., Planchart, A., 2009. Perturbation of  
517 defense pathways by low-dose arsenic exposure in zebrafish embryos. *Environ. Health Perspect.*  
518 117, 981-987.
- 519 Miller, D.S., Shaw, J.R., Stanton, C.R., Barnaby, R., Karlson, K.H., Hamilton, J.W., Stanton, B.A.,  
520 2007. MRP2 and acquired tolerance to inorganic arsenic in the kidney of killifish (*Fundulus*  
521 *heteroclitus*). *Toxicol. Sci.* 97, 103-110.
- 522 NRC, 1999. Arsenic in drinking water. In. National Research Council, Washington, D.C..
- 523 Peterson, J.S.K., Bain, L.J., 2004. Differential gene expression in anthracene-exposed mummichogs  
524 (*Fundulus heteroclitus*). *Aquat. Toxicol.* 66, 345-355.
- 525 Posey, T., Weng, T., Chen, Z., Chintagari, N.R., Wang, P., Jin, N., Stricker, H., Liu, L., 2008. Arsenic-  
526 induced changes in the gene expression of lung epithelial L2 cells: implications in carcinogenesis.  
527 *BMC Genomics* 9, 115.
- 528 Rahman, A., Vahter, M., Smith, A.H., Nermell, B., Yunus, M., El Arifeen, S., Persson, L.A., Ekström,  
529 E.C., 2009. Arsenic exposure during pregnancy and size at birth: A prospective cohort study in  
530 Bangladesh. *Am. J. Epidemiol.* 169, 304-312.
- 531 Raqib, R., Ahmed, S., Sultana, R., Wagatsuma, Y., Mondal, D., Hoque, A.M., Nermell, B., Yunus, M.,  
532 Roy, S., Persson, L.A., Arifeen, S.E., Moore, S., Vahter, M., 2009. Effects of in utero arsenic  
533 exposure on child immunity and morbidity in rural Bangladesh. *Toxicol. Lett.* 185, 197-202.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 534 Roling, J.A., Bain, L.J., Baldwin, W.S., 2004. Differential gene expression in mummichog (*Fundulus*  
535 *heteroclitus*) following treatment with pyrene and fluoranthene: comparison to a creosote  
536 contaminated site. *Mar. Environ. Res.* 57, 377-395.
- 537 Roling, J.A., Bain, L.J., Gardea-Torresdey, J., Bader, J., Baldwin, W.S., 2006. Hexavalent chromium  
538 reduces larval growth and alters gene expression in mummichogs (*Fundulus heteroclitus*). *Environ.*  
539 *Tox. Chem.* 25, 2725-2733.
- 540 Shaw, J.R., Gabor, K., Hand, E., Lankowski, A., Durant, L., Thibodeau, R., Stanton, C.R., Barnaby, R.,  
541 Coutermarsh, B., Karlson, K.H., Sato, J.D., Hamilton, J.W., Stanton, B.A., 2007a. Role of  
542 glucocorticoid receptor in acclimation of killifish (*Fundulus heteroclitus*) to seawater and effects of  
543 arsenic. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 292, R1052-R1060.
- 544 Shaw, J.R., Jackson, B., Gabor, K., Stanton, S., Hamilton, J.W., Stanton, B.A., 2007b. The influence of  
545 exposure history on arsenic accumulation and toxicity in the killifish, *Fundulus heteroclitus*.  
546 *Environ. Toxicol. Chem.* 26, 2704-2709.
- 547 Sheader, D.L., Williams, T.D., Lyons, B.P., Chipman, J.K., 2006. Oxidative stress response of European  
548 flounder (*Platichthys flesus*) to cadmium determined by a custom cDNA microarray. *Mar. Environ.*  
549 *Res.* 62, 33-44.
- 550 Shen, J., Wanibuchi, H., Waalkes, M.P., Salim, E.I., Kinoshita, A., Yoshida, K., Endo, G., Fukushima,  
551 S., 2006. A comparative study of the sub-chronic toxic effects of three organic arsenical compounds  
552 on the urothelium in F344 rats; gender-based differences in response. *Toxicol. Appl. Pharmacol.*  
553 210, 171-180.
- 554 Smedley, P., Kinniburgh, D., 2002. A review of the source, behaviour and distribution of arsenic in  
555 natural waters. *Appl. Geochem.* 17, 517-568.
- 556 Snow, E.T., Sykora, P., Durham, T.R., Klein, C.B., 2005. Arsenic, mode of action at biologically  
557 plausible low doses: what are the implications for low dose cancer risk? *Toxicol. Appl. Pharmacol.*  
558 207, 557-564.
- 559 Taylor, W.R., Van Dyke, G.C., 1985. Revised procedures for staining and clearing small fishes and  
560 other vertebrates for bone and cartilage study. *Cybiurn* 9, 107-119.
- 561 Tseng, C.H., Tai, T.Y., Chong, C.K., Tseng, C.P., Lai, M.S., Lin, B.J., Chiou, H.Y., Hsueh, Y.M., Hsu,  
562 K.H., Chen, C.J., 2000. Long-term arsenic exposure and incidence of non-insulin-dependent  
563 diabetes mellitus: a cohort study in arseniasis-hyperendemic villages in Taiwan. *Environ. Health*  
564 *Perspect.* 108, 847-851.
- 565 Vahter, M., Akesson, A., Lidén, C., Ceccatelli, S., Berglund, M., 2007. Gender differences in the  
566 disposition and toxicity of metals. *Environ. Res.* 104, 85-95.
- 567 Ventura-Lima, J., de Castro, M.R., Acosta, D., Fattorini, D., Regoli, F., de Carvalho, L.M., Bohrer, D.,  
568 Geracitano, L.A., Barros, D.M., Marins, L.F., da Silva, R.S., Bonan, C.D., Bogo, M.R., Monserrat,  
569 J.M., 2009. Effects of arsenic (As) exposure on the antioxidant status of gills of the zebrafish *Danio*  
570 *rerio* (Cyprinidae). *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 149, 538-543.
- 571 Waalkes, M.P., Liu, J., Ward, J., Diwan, B.A., 2004a. Animal models for arsenic carcinogenesis:  
572 inorganic arsenic is a transplacental carcinogen in mice. *Toxicol. Appl. Pharmacol.* 198, 377-384.
- 573 Waalkes, M.P., Liu, J., Ward, J.M., Powell, D.A., Diwan, B.A., 2006. Urogenital carcinogenesis in  
574 female CD1 mice induced by in utero arsenic exposure is exacerbated by postnatal diethylstilbestrol  
575 treatment. *Cancer Res.* 66, 1337-1345.
- 576 Waalkes, M.P., Ward, J.M., Diwan, B.A., 2004b. Induction of tumors of the liver, lung, ovary and  
577 adrenal in adult mice after brief maternal gestational exposure to inorganic arsenic: promotional  
578 effects of postnatal phorbol ester exposure on hepatic and pulmonary, but not dermal cancers.  
579 *Carcinogenesis* 25, 133-141.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 580 Waalkes, M.P., Ward, J.M., Liu, J., Diwan, B.A., 2003. Transplacental carcinogenicity of inorganic  
581 arsenic in the drinking water: induction of hepatic, ovarian, pulmonary, and adrenal tumors in mice.  
582 *Toxicol. Appl. Pharm.* 186, 7-17.
- 583 Williams, T.D., Gensberg, K., Minchin, S.D., Chipman, J.K., 2003. A DNA expression array to detect  
584 toxic stress response in European flounder (*Platichthys flesus*). *Aquat. Toxicol.* 65, 141-157.
- 585 Xie, Y., Liu, J., Benbrahim-Tallaa, L., Ward, J.M., Logsdon, D., Diwan, B.A., Waalkes, M.P., 2007.  
586 Aberrant DNA methylation and gene expression in livers of newborn mice transplacentally exposed  
587 to a hepatocarcinogenesis dose of inorganic arsenic. *Toxicology* 236, 7-15.
- 588 Yamaguchi, S., Miura, C., Ito, A., Agusa, T., Iwata, H., Tanabe, S., Tuyen, B.C., Miura, T., 2007.  
589 Effects of lead, molybdenum, rubidium, arsenic and organochlorines on spermatogenesis in fish:  
590 monitoring at Mekong Delta area and *in vitro* experiment. *Aquat. Toxicol.* 83, 43-51.
- 591 Yamamoto, M., Hirano, S., Vogel, C.F., Cui, X., Matsumura, F., 2008. Selective activation of NF-  
592 kappaB and E2F by low concentration of arsenite in U937 human monocytic leukemia cells. *J.*  
593 *Biochem. Mol. Toxicol.* 136-146.
- 594 Yao, Z., Papadopoulos, V., 2002. Function of beta-amyloid in cholesterol transport: a lead to  
595 neurotoxicity. *FASEB J.* 16, 1677-1679.
- 596 Yoshida, T., Yamauchi, H., Fan Sun, G., 2004. Chronic health effects in people exposed to arsenic via  
597 the drinking water: dose-response relationships in review. *Toxicol. Appl. Pharmacol.* 198, 243-252.
- 598 Zheng, H., Koo, E., 2006. The amyloid precursor protein: beyond amyloid. *Mol. Neurodegener.* 1, 5.
- 599 Zheng, X.H., Watts, G.S., Vaught, S., Gandolfi, A.J., 2003. Low-level arsenite induced gene expression  
600 in HEK293 cells. *Toxicology* 187, 39-48.

1  
2  
3 **Figure Legend**

4  
5 **Fig. 1. Arsenic body burdens.** Arsenic concentrations were measured in the adult females after the 10 day  
6 exposure. Data is the average  $\pm$  standard deviation of 5 intact fish per group. Significant differences (\*) were  
7 determined using Student's t-test ( $p < 0.05$ ).  
8

9  
10  
11 **Fig. 2. Gene expression profiles for males and female *Fundulus* exposed to different concentrations of**  
12 **arsenic.** Hepatic gene expression was examined using the *Fundulus* cDNA array (GEO-GPL2535) and analyzed  
13 for differential expression using the CLEAR Test Method. The numbers of genes with significantly different  
14 expression ( $p \leq 0.05$ ) are indicated, and those changed by more than one exposure are found in the overlapping  
15 areas. **A.** Dose-responsive changes in hepatic gene expression in male and female fish **B.** Gender-responsive  
16 changes in hepatic gene expression in male and female fish.  
17  
18

19  
20 **Fig. 3. Percentage viability and morphological abnormalities following arsenic exposure.** Adults were  
21 exposed to 172, 575 or 1,720ppb arsenic for 10 days prior to spawning. Eggs were collected for three days  
22 following arsenic exposure, and their viability, along with any developmental abnormalities observed after 7 days,  
23 was recorded. **A.** Viability and deformity percentages are expressed as the average  $\pm$  standard deviation for  
24 offspring from 4 –5 tanks, with statistical differences (\*) determined by Mann-Whitney ( $p < 0.05$ ). **B.**  
25 Representative photographs from normal and deformed offspring. **C.** Morphological analysis of the vertebrae.  
26 The total size (area) for individual vertebrae were quantified using the LSM5 Pascal software. Values are the  
27 average area in square micrometers divided by the weight of the 6-week old fish. Significant differences (\*) were  
28 determined using Student's t-test ( $p < 0.05$ ).  
29  
30

31  
32 **Fig. 4. Comparisons of gene expression changes between the arrays and quantitative real-time PCR**  
33 **(QPCR).** Expression of three genes from the arrays that were differentially expressed in almost all genders,  
34 ages, and exposure groups were verified by QPCR using specific primers ( $n=5$  for each group). Array fold  
35 changes were plotted against QPCR fold changes to determine their directionality and magnitude. Genes  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

examined were apolipoprotein, serum amyloid precursor, and an EST termed 9-7. The letter indicates females (F), males (M), or juveniles (J), while the number (172, 575, or 1,720) indicates the exposure group.


ACCEPTED MANUSCRIPT


Table 1. Gene Expression Changes in Fundulus Exposed to Arsenic. Fold-changes in gene expression were calculated and significant differences were determined using the CLEAR Test method ( $p < 0.05$ ). Only genes that are differentially expressed and shared between more than 1 exposure group are listed.




Clone ID	Gene Name	GenBank #	Fold change (Arsenic/control)								
			Females		Males			Juveniles			
			172ppb	575ppb	172ppb	575ppb	1,720ppb	172ppb	575ppb	1,720ppb	
8--56	apolipoprotein	<a href="#">DN474949</a>	0.59	0.75	0.85		0.85				
Ab35	apolipoprotein	<a href="#">CN992696</a>	0.59	0.79	0.85		0.84				
9--7	EST	<a href="#">CO897482</a>	0.61	1.22	1.14	1.71		4.1		1.45	
9--12	TBT-binding protein	<a href="#">AY725225</a>	0.56	0.6		1.95					
8-6-T7	EST	<a href="#">DN596277</a>	5.71	0.45							
Bb56	EST	<a href="#">DN596273</a>	0.67	1.79							
1--31	serum amyloid precursor	<a href="#">AY735160</a>	0.63	2.52	0.77		0.32				
2--80	EST	<a href="#">DN474945</a>	0.54		0.81	2.13		1.54		1.29	
9--16	EST	<a href="#">CO897478</a>	0.7		0.74	1.39					
pa_1--15	complement component C8	<a href="#">AY521664</a>	0.7		0.82	1.93				1.58	
Bb53	cytochrome c oxidase subunit II	<a href="#">CV821001</a>		1.57	0.7						
8--40	cytochrome c oxidase subunit II	<a href="#">CN976180</a>		0.8	0.6						
10--15	EST	<a href="#">CO897479</a>			1.14	1.8	0.87				
10--57	EST	<a href="#">CO897481</a>			1.41		0.55				
1--11	lysozyme precursor	<a href="#">AY735156</a>			0.3		2.65				
1--163	lysozyme precursor	<a href="#">AY735156</a>			0.29		2.85				
8--46	EST	<a href="#">DN474948</a>				1.37	0.82				
8--8	EST	<a href="#">CO897478</a>				1.68	0.82				

**Table 2. Genes showing unique differential expression.** Fold-changes in hepatic gene expression in the males and females were calculated and significant differences were determined using the CLEAR Test method ( $p \leq 0.05$ ). Only genes that were differentially expressed in only 1 exposure group are listed. GenBank accession numbers are indicated in parentheses.

<b>Clone Identity and GenBank Accession Number</b>					
<b><u>172ppb females</u></b>	<b><u>575ppb females</u></b>		<b><u>172ppb males</u></b>	<b><u>575ppb males</u></b>	<b><u>1,720ppb males</u></b>
Small inducible cytokine A4 (AY735153)	EST (CX700381)		GLUT2 (AY521663)	Elongation factor 1 $\alpha$ (AY735180)	EST (CO897478)
Myosin light chain 2 (CX700395)	Coagulation factor XIII B precursor (CN988088)		Hemopexin precursor (AY735182)	70kDa HSP (AY735159)	Warp65 (AY735166)
EST (CX700386)	14-3-3 (AY725221)		EST (CO897478)	GST $\mu$ (AY725220)	EST (DN596272)
Map kinase kinase (BI993599)	Deiodinase (U70869)		transferrin (AY735165)	EST (CO897478)	EST (CX700393)
EST (DN474942)	Vitellogenin (U07055)		Cytochrome b (AY725226)		
EST (CO897490)	ATP synthase F <sub>o</sub> complex (AY735178)		fucolectin (AY735152)		
	EST (CO897485)		EST (CX700390)		





**B.**

	% viability	% deformities	Fold increase in deformities
Control	71.2 ± 21.4	0	
172ppb	82.5 ± 6.9	0	0
Control	91.0 ± 6.5	0	
575ppb	88.3 ± 8.3	0	0
Control	55.7 ± 32.6	0.7 ± 1.1	
1720ppb	77.7 ± 25.2	12.2 ± 6.0*	17.4*

