



Assessment of the acrylamide intake of the Belgian population and the effect of mitigation strategies

Wendie Liliane Claeys, Katleen Baert, Frédéric Mestdag, Jan Vercammen,
Paul Daenens, Bruno de Meulenaer, Guy Maghuin-Rogister, André
Huyghebaert

► To cite this version:

Wendie Liliane Claeys, Katleen Baert, Frédéric Mestdag, Jan Vercammen, Paul Daenens, et al..
Assessment of the acrylamide intake of the Belgian population and the effect of mitigation strategies.
Food Additives and Contaminants, 2010, 27 (09), pp.1199-1207. 10.1080/19440049.2010.489577 .
hal-00604049

HAL Id: hal-00604049

<https://hal.science/hal-00604049>

Submitted on 28 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Assessment of the acrylamide intake of the Belgian population and the effect of mitigation strategies

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2010-046.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	19-Apr-2010
Complete List of Authors:	Claeys, Wendie; FASFC, Policy Control Baert, Katleen; FASFC, Policy Control Mestdagh, Frédéric; Ghent University, Department of Food Safety and Food Quality Vercammen, Jan; FASFC, Laboratories Daenens, Paul; Catholic University Leuven De Meulenaer, Bruno; Ghent University, Department of Food Safety and Food Quality maghuin-rogister, guy; University of Liège, CART Huyghebaert, André; Ghent University, Department of Food Technology and Nutrition
Methods/Techniques:	Risk assessment, Exposure assessment, Probabilistic modelling, Chromatography - LC/MS
Additives/Contaminants:	Process contaminants – acrylamide
Food Types:	Processed foods

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

**ASSESSMENT OF THE ACRYLAMIDE INTAKE OF THE BELGIAN POPULATION
AND THE EFFECT OF MITIGATION STRATEGIES**

Wendie Claeys^{1,}, Katleen Baert¹, Frédéric Mestdagh², Jan Vercammen³, Paul Daenens⁴,
Bruno De Meulenaer², Guy Maghuin-Rogister⁵, and André Huyghebaert²*

¹ DG Control Policy, Federal Agency for the Safety of the Food Chain, Belgium
² nutriFOODchem unit, Department of Food Safety and Food Quality, Faculty of Bioscience
Engineering, Ghent University, Belgium
³ Laboratory of Gentbrugge, DG Laboratories, Federal Agency for the Safety of the Food Chain,
Belgium
⁴ Faculty of Pharmaceutical Sciences, Catholic University Leuven, Belgium
⁵ Centre d'analyse des résidus en Traces (CART), Faculty of Veterinary Medicine, University of
Liège, Belgium

*Author to whom correspondence should be addressed:
DG Control Policy, Federal Agency for the Safety of the Food Chain,
Boulevard du Jardin botanique 55, B-1000 Brussels, Belgium
Tel: +32-(0)2-211.87.02; Fax: +32-(0)2-211.87.22
E-mail : wendie.claeys@favv.be

Abstract

The acrylamide (AA) intake of the Belgian consumer was calculated based on AA monitoring data of the Belgian Federal Agency for the Safety of the Food Chain (FASFC) and consumption data of the Belgian food consumption survey coordinated by the Scientific Institute for Public Health (3214 participants of 15 years or older). The average AA exposure, calculated probabilistically, was 0.4 µg/kg bw/day (P97.5 = 1.6 µg/kg bw/day) with as main contributors to the average intake chips (23%), coffee (19%), biscuits (13%) and bread (12%). Additionally, the impact of a number of AA mitigation scenarios were evaluated (German minimisation concept, scenarios for mitigation from the literature, signal values), which is an important issue for public health as well as for policymakers. Specific actions in cooperation with the food industry to reduce the AA content of foods seems to be a more efficient strategy than mere implementation of signal values. Considering that an important share of the AA intake is due to prepared meals, the catering industry as well as the consumer need to be better informed on the various possibilities for keeping the AA content of meals as low as possible.

Keywords: acrylamide; risk assessment; exposure; food; mitigation; probabilistic

Introduction

Acrylamide (AA, $\text{CH}_2=\text{CHCONH}_2$, CAS No. 79-0601, EC No. 201-173-7) is neurotoxic, and probably genotoxic and carcinogenic to humans as well. Recently, the European Chemicals Agency (ECHA) announced to include AA on the European Union's candidate list of Substances of Very High Concern (ECHA 2009). Under the ECHA proposal, AA has been listed as a category 2 carcinogen and a category 2 mutagen. The chemical has a large number of industrial applications including the production of polyacrylamides, which are among others used in cosmetics, pesticides, paints, as a flocculant in water treatment and in the paper industry. AA is also present in cigarette smoke (JECFA 2005; IARC 1994).

In 2002 it was demonstrated that relatively high levels of AA are formed during heating of certain foods (baking, roasting, frying). Since then, intensive research has been conducted regarding the various formation mechanisms and the toxicology of AA (e.g. bio-availability), the development of accurate analytical methods (e.g. cheap and fast screening methods), factors that influence the AA content of foods (e.g. storage), etc. (Tardiff et al. 2010; EFSA 2008a; Friedman and Levin 2008; Claeys et al. 2005; Stadler and Scholz 2004; Taeymans et al. 2004).

The observed AA levels in food are a result of competitive, complex processes of formation and elimination or degradation (Mottram et al. 2002). The largest amount of AA is accumulated during the last stages of baking, roasting or frying when the moisture content of the food drops and the surface temperature rises, except for coffee where the AA level drops significantly during the later stages of the roasting process. AA appears to be stable in the large majority of foodstuffs. Grounded coffee, where the AA content decreases during storage, is once more an exception (Lanz et al. 2006; Hoenicke and Gattermann 2005).

Until present, there are no legal regulations regarding the AA content of food, at a national nor at a European level. Germany is the only country with a systematic mitigation strategy (BVL, 2005) and the only legal action undertaken so far, was the withdrawal of child biscuits containing > 1000 μg AA/kg from the Swiss market in 2005 (Grob 2007). In California, a court agreement was reached with a number of important chips and crisp producers and some fast food chains to reduce the AA content of their products significantly and to put warning labels regarding the presence of AA on the package (Hill 2008).

The Belgian Federal Agency for the Safety of the Food Chain (FASFC) monitors already for a number of years the AA content of different foodstuffs. When AA is detected at a level above 1000 µg/kg, the producer is urged to take possible mitigation actions. The vast amount of data acquired through the monitoring programme as well as the availability of a nationwide Belgian food consumption survey allows the assessment of the dietary exposure of the Belgian population to AA. The present study is the first large-scale study conducted with respect to the AA intake of the Belgian consumer.

Materials and Methods

Food consumption data

Food consumption data were obtained from the Belgian Food Consumption Survey (BFCS) in 2004 performed by the Scientific Institute of Public Health (Devriese et al. 2005). The survey involved 3214 participants of 15 years or older, which were interviewed twice about their consumption during the last 24 hours (repeated non-consecutive 24h recall) in combination with a self-administered food frequency questionnaire. The fieldwork was spread over one year to anticipate seasonal effects and was carried out by trained dietitians. The selection of interviewed people and the moment of the interview were chosen in order to obtain a representative consumption profile of the Belgian population over 1 year anticipating seasonal effects.

In this study, the total data sets, including zero intakes ('zero consumption days'), was used as part of an 'average' diet. It is remarked that the duration of the consumption survey affects the distribution of consumption data, particularly the upper percentiles or the group of high or frequent consumers. As such, a brief survey often underestimates the consumption of less frequently consumed foods, but at the same time overestimates the quantities of frequently consumed foods. High percentiles based on a one to two days consumption survey are often an overestimation compared to high percentiles obtained on the basis of a seven-day survey. The reliability of the high percentiles depends not only on the duration of the survey, but also on the number of people or data on which their calculation is based. Percentiles that are calculated

based on a limited amount of data, have a higher uncertainty and give only a rough indication of the higher consumption levels. According to Kroes et al. (2002), a high percentile P (> P75) can only be evaluated with sufficient accuracy when the sample size n satisfies with $n(1-P) \geq 8$. The minimum amount of data thus required for P95, P97.5 and P99 can be estimated as 160, 320 and 800 respectively. The same restrictions apply to the lower percentiles (EFSA 2008b).

Acrylamide concentration data

AA levels were measured in various foodstuffs on the Belgian market within the framework of the monitoring programme of the FASFC. The AA content of the samples was determined by a LC-MS accredited method in the FASFC laboratory of Gentbrugge, with a limit of quantification (LOQ) of 50 µg/kg and a limit of detection (LOD) of 25 µg/kg. After AA extraction, defatting and further clean-up of the sample over an Oasis HLB SPE column, 10 µl was injected on an Sequant ZIC-HILIC column (2.1mm x 150mm 5µm, equipped with a pre-column) with acetonitrile as mobile phase, coupled to a triple quadrupole mass spectrometer operating in APCI mode (atmospheric pressure chemical ionization) (TSQ Quantum Ultra, Thermo Fisher Scientific, USA). The content was measured in MS² on the ions m/z 72 and 52 (AA) and 75 and 58 (labelled AA, used as internal standard). Depending on the matrix, the recovery is situated between 81 and 100%. The extended measurement uncertainty is situated between 26% (clean matrix) and 52% (dirty matrix).

Statistical comparison of the AA levels measured between 2002 and 2007 showed no decrease or increase of the AA level as a function of time. Consequently, data from 2002 until 2007 could be pooled for calculating the intake, which resulted in a data set of 759 data. The statistical analysis of the data was performed with SPSS[®] 11.0 for Windows (SPSS Inc., USA). The Kolmogorov-Smirnov test was used to test normality. Means were compared using analysis of variance (one-way ANOVA) and Post Hoc Multiple Comparison tests (Tukey when variances were equal or Games-Howell when variances were unequal). Homogeneity of variances was tested using the Levene test.

For the exposure assessment, foods were grouped in twelve categories: chips, crisps, coffee (liquid), coffee surrogate (liquid), bread, bread rolls (including pastry), toast, biscuits,

gingerbread, almond or sweet spiced biscuits, chocolate, popcorn, breakfast cereals and cereal bars. The classification is based on the AA levels in different foodstuffs and is in line with the grouping applied in other studies (Mestdagh et al. 2007; Boon et al. 2005; Matthys et al. 2005).

The AA data were further expanded with German data from the AA European Monitoring Database, which is compiled by the Institute for Reference Materials and Measurements (IRMM 2006). Based on a statistical analysis showing no significant differences, the Belgian AA database could be supplemented with German data for chips, crisps, chocolate, coffee, biscuits and sweet spiced biscuits. In total, the AA dataset contained more than 3000 results.

Estimation of the acrylamide intake

The AA intake was determined per food group and overall by a probabilistic approach considering all data or the full distribution of the different variables (i.e. AA content and consumption). Hereto, a (1-dimensional) Monte Carlo simulation was performed with 100,000 iterations. To evaluate the uncertainty of the sampling, 'bootstrap sampling' (2-dimensional Monte Carlo model) was applied. In this method n observations (AA concentration and consumption of the relevant food) were theoretically resampled from the original dataset resulting in a 'bootstrap' dataset with n observations. By repeating this process 500 times, 500 'bootstrap' datasets are obtained, on which the same statistical calculations (e.g. 97.5th, 99.9th percentile, etc.) can be applied as on the original data set. As such a 'bootstrap' distribution of 500 97.5th, 99.9th percentiles, etc. is created that characterizes the uncertainty of the original data set (Vose 2006). The model input distributions were randomly sampled by the Latin Hypercube method. Calculations were performed by the software @Risk[®] (Palisade Corporation, Version 4.5.5, NY, USA).

AA concentrations below the quantification limit (LOQ) were replaced by LOQ/2 ('middle bound scenario'). For the conversion of the AA level of roasted or grounded coffee beans (coffee surrogate) to liquid coffee (coffee surrogate) a conversion factor of 0.046 was applied (van Dooren et al., 1995). Regarding the AA level of chocolate biscuits, chocolate and biscuit were considered separately. Based on chocolate levels mentioned on the package it was assumed that 40% of the biscuit consists of chocolate. Variability in preparation conditions (e.g.

deep-frying of chips, toasting of bread, etc.) was not taken into account.

Results and Discussion

Dietary exposure of the Belgian consumer to acrylamide

Table 1 shows the AA intake per food group and the overall AA intake of the Belgian population. The contribution of the most relevant food groups to the AA intake (contribution > 1%) is also shown in Figure 1.

The average AA intake was calculated as 0.35 µg/kg bw/day (P50 = 0.2 µg/kg bw/day). For high consumers the intake may be several times higher, amounting to 1.58 µg/kg bw/day for the 97.5th percentile or P97.5. Overall, about 6% of the population has an intake higher or equal to the average intake of 1 µg/kg bw estimated by the Joint FAO/WHO Expert Committee on Food Additives (JECFA 2005).

[Table 1]

On average, chips contributed the most to the AA intake (23%), followed by coffee (19%), biscuits (13%) and bread (12%) (Figure 1). For the lower intake percentiles, coffee and bread appeared to be most important sources of AA, whereas for the higher percentiles (≥ P85) chips, crisps and biscuits became more important sources. This reflects the difference in dietary pattern according to the AA intake (Dybing et al. 2005; Matthys et al. 2005).

The AA level, and as such the contribution to the AA intake, of different food groups and of foods within the same food group can vary greatly depending on the formulation of the food, manufacturing conditions, etc. The contribution of a food or food group to the intake is however, not only defined by its AA level, but also by the amount consumed. For example, gingerbread and coffee surrogate contributed relatively little to the exposure despite their relatively high mean AA level (for Belgium, the mean AA level is 692 ± 566 µg/kg for gingerbread and 2531 ± 825 µg/kg for coffee surrogate powder). With respect to bread the opposite is observed (mean AA level for bread = 30 ± 9 µg/kg) due to its high consumption compared to e.g. gingerbread

and coffee surrogate powder.

[Figure 1]

The AA intake given in Table 1, corresponds well to the range of values calculated previously for Belgium (restricted target population of adolescents and visitors of a university canteen) (Mestdagh et al. 2007; Matthys et al. 2005) and reported for other European countries (Boon et al. 2005; Dybing et al. 2005; Svensson et al. 2003). In general, values between 0.3 to 2.0 μg AA/kg bw/day are reported for the average AA intake of adults. The high intake percentiles (P90 to P97.5) range from 0.6 to 3.5 $\mu\text{g}/\text{kg}$ bw/day with 5.1 $\mu\text{g}/\text{kg}$ bw/day reported for P99. Food that contribute most to the AA intake are in general chips (16-30%), crisps (6-46%), coffee (13-39%), fine bakery and sweet biscuits (10-20%), and bread and rolls/toast (10-30%). Other relevant foodstuffs contribute less than 10% to the intake (WHO 2006; JECFA 2005). The absolute figure of the exposure and the relative contribution of each food group to the exposure may differ from study to study, depending on the number and nature of the food groups considered, the methodology applied for the calculation, the type of consumption survey, etc. (Dybing et al. 2005; JECFA 2005).

Any exposure assessment is confronted with a number of uncertainties (Kroes et al. 2002). In this study, the long-term intake was predicted based on a two-day consumption survey and consumption data were considered to be independent. The linkage between the analyzed and consumed food items, the categorization into food groups, the conversion factors applied (i.e. for chocolate in chocolate biscuits and for liquid coffee), and the fact that variability in preparation conditions was not accounted for and that values below the LOQ were unknown and replaced by LOQ/2, are additional sources of uncertainty in the estimated AA intake.

Risk characterization

Given that AA is probably genotoxic and carcinogenic to humans (IARC group 2A) (JECFA 2005; IARC 1994), it is recommended to keep the AA exposure 'as low as reasonably achievable' (ALARA). Nevertheless, to prioritise risks associated with unavoidable contaminants that are both genotoxic and carcinogenic, the 'margin of exposure' (MOE) concept has been put

forward (EFSA 2005). The MOE is the ratio between a particular point on the dose-response curve leading to tumours in experimental animals and the exposure. The size of the MOE gives an indication about the possible extent of the risk. The higher the MOE, the lower the risk of exposure to the component concerned. The Scientific Committee of the European Food Safety Authority (EFSA) considers the figure of 10,000 or higher as (in general) being of low concern from a public health point of view and might reasonably be considered as a low priority for risk management actions. For AA MOE values between 50 and 2000 have been reported, depending on the exposure data and toxicological values used (Table 2). The JECFA calculated a MOE of 300 based on an average exposure of 1 µg AA/kg bw/day and a BMDL10 (the lower confidence limit on the benchmark dose associated with 10% response) of 300 µg AA/kg bw/day for the induction of breast tumours in rats. For high consumers with an intake of 4 µg AA/kg bw/day, the MOE was 75 (JECFA 2005). The average (0.2 µg/kg bw/day) and the P97.5 (1.6 µg/kg bw/day) intake calculated in this study, would thus correspond to MOE values of 1500 and 188 respectively. Such low MOE values imply that additional efforts have to be taken for reducing the AA content in foods and that AA is a process contaminant of high priority. (For comparison: MOE values between 17,900 and 9500 are given for the polycyclic aromatic hydrocarbons (EFSA 2008c)).

[Table 2]

Possible strategies to reduce the acrylamide exposure

Various methods are reported in the literature to reduce the AA content of food, such as the selection of potato, wheat and other plant cultivars with low levels of the AA precursors asparagine and glucose, the removal of the AA precursors (e.g. soaking of potatoes, hydrolysis of asparagine to aspartic acid using asparaginase), adaptation of process and storage conditions (temperature, time, water activity (a_w) and pH), adding ingredients that inhibit the AA formation (acids, amino acids, antioxidants, non-reducing sugars, chitosan, garlic components, protein hydrolysates, proteins, multivalent ions), the removal or "capture" of AA through chromatography, evaporation or polymerization (Friedman and Levin 2008; Claeys et al. 2005;

Stadler and Scholz 2004; Taeymans et al. 2004).

Table 3 shows the effect of some of the specific AA mitigation possibilities reported in literature on the dietary exposure. It should be noted however, that the reduction percentages mentioned are indicative values and that the yield of the proposed mitigation possibilities might be different in practice (for one because there are different types of biscuits, breads, etc. with their own recipe, process conditions, etc.). Furthermore, the reductions mentioned are often based on laboratory (model) experiments, of which the industrial applicability has in most cases not been evaluated yet.

For 50% of the population, the AA intake is primarily due to the consumption of bread and coffee. The AA content of bread is for > 99% located in the crust (Surdyk et al. 2004). Some of the AA mitigation options for bread mentioned in literature are amongst others the addition of Ca^{2+} or Mg^{2+} to the dough (resulting in a potential reduction of 20% of the AA content), a prolonged yeast fermentation of the dough (50 to 77-78% reduction), and addition of a high dose of glycine (80% reduction) (Claus et al. 2004; Fredriksson et al. 2004). Assuming that a reduction of 60% of the AA content in bread is possible (scenario 1), the P50 and P75 of the AA intake could be significantly reduced with almost 30% (Table 3).

With respect to coffee, only a few options for reducing the AA level are put forward such as the selection of coffee bean species, darker roasting of the beans, a prolonged shelf life and the use of asparaginase (Lanz et al. 2006; Hoenicke and Gattermann 2005). If the AA level of coffee could be reduced with 30% (scenario 2), the P50 and P75 of the AA intake would be decreased with ~ 15-16%.

Other major sources of AA in the diet are chips, crisps and biscuits. The AA content in biscuits could be reduced with 70% by adjusting the baking process and by replacing glucose and fructose with sucrose (Gökmen et al. 2007; Graf et al. 2006) (scenario 3). Application of these measures would reduce the average intake with 9%. For potato products, a significant reduction of the AA level is reported when potatoes are soaked or blanched in an acetic acid solution (40-80%) or at 70°C for 10 to 15 min (65-96%) before frying (Mestdagh et al. 2008a,b; Kita et al. 2004). Reducing the baking temperature from 185°C to 175°C can reduce the AA level of potato products with 35% (scenario 4). Such an adjustment of the baking process

269 results in a ~10% reduction of the average AA intake, which is of the same order of magnitude
270 as the average reduction of 13% determined by Boon et al. (2005) for the Dutch consumer. For
271 a high consumer of fried potato products, the intake can even be reduced with 20%.

272 In addition, Boon et al. (2005) calculated that the AA intake decreases on average with about
273 4% when the AA level in gingerbread is reduced with 60%, which can be obtained when sodium
274 bicarbonate is used instead of ammonium carbonate as baking agent (scenario 5). In this study,
275 a similar reduction of the intake was obtained for the higher percentiles. Boon et al. (2005)
276 determined also the reduction that would be achieved when the two latter mitigation options are
277 combined (scenario 4 and 5) and obtained an average reduction of 17%. A comparable
278 reduction of 9 to 25% for the higher intake percentiles was calculated in this study.

279 Seal et al. (2008) addressed the effect of some mitigation options on the AA intake as well,
280 discussing not only the benefits (reduction of the AA intake) but also the risks (health, quality or
281 nutritional implications). The mitigation scenarios studied by Seal et al. (2008) decreased the
282 exposure by 1 to 14%, which is the same range as observed in this study. Application of all
283 mitigation measures simultaneously resulted in a reduction of 31-39%, depending on the
284 percentile of exposure considered.

[Table 3]

288 A possible policy measure for reducing the AA intake is by establishing signal values (action
289 values). In this respect, Germany introduced in 2002 the 'minimisation concept', which pursues
290 a gradual reduction of the AA levels in foodstuffs while retaining the product's properties (BVL,
291 2005). In this concept the lowest AA level of 10% of the food with the highest AA concentration
292 (P90) is taken as a signal value, with a ceiling of 1000 µg AA/kg food product. The German P90
293 signal value is revised each year and reduced when necessary. A signal value can be
294 maintained, but never increased. When a signal value is exceeded, it is discussed with the
295 companies involved which actions might be taken to lower the AA level.

296 Table 4 shows the effect of the German 2008 signal values on the total AA intake of the
297 Belgian consumer (see bottom line of the table). Hereto, AA levels above the German signal

values were replaced by the P90 signal value. It should be noted that there are no German signal values given for chocolate, bread, bread rolls, popcorn and cereal bars. For biscuits the signal value for diet biscuits was applied. For toast the signal value of knäckebröd was considered, but this value turned out to be higher than the levels measured for toast. The effect of applying the P90 signal value concept on the Belgian data (P90 of Belgian AA concentration data as signal value) is also presented in Table 4, (i) for each food group separately and (ii) for all food groups together.

[Table 4]

Application of the German signal values on the Belgian data results in a significant reduction of the intake for the higher ($\geq$ P95), but not for the lower intake percentiles. This can partly be explained by the fact that the signal values are P90 values limiting the higher AA concentration data. In addition, the intake at these percentiles is mainly determined by bread and coffee; no German signal value is presented for bread and in the case of coffee less than 2% of the samples exceeds the German signal value. Regarding the hypothetical Belgian signal values, a similar observation is made.

As expected, the introduction of P90 signal values affects the intake mainly through the food groups that contribute most to the exposure, namely chips, bread and biscuits.

Conclusion

AA is formed mainly during heating of carbohydrate-rich food. AA has always been present in food and is as such not a new contaminant. The derived MOE values however, indicate that additional efforts to reduce the AA exposure are a prerequisite.

In this study, the effect of some mitigation strategies on the AA exposure was examined. The most efficient approach for reducing the AA exposure seems to be the reduction of the AA content in foodstuffs that are potentially important contributors to the AA intake rather than the introduction of signal values (action limits). When starting from food with a high AA level, an already significant reduction of the dietary exposure can be achieved with some relatively minor

measures. Signal values or action limits can be a useful tool for policymakers to encourage the food industry to take action by means of a specific AA value. It is nevertheless the task of the food industry to ensure that the AA content of their products is as low as possible. In literature several possibilities are proposed for reducing the AA level in food. For example, the CIAA (Confederation of the food and drink industries of the EU) developed the “AA Toolbox”, which contains a brief description of possible intervention steps evaluated in cooperation with industry for reducing the AA content of food products and has published AA pamphlets for five different sectors (CIAA 2009 a & b). However, also the consumer has to take responsibility for reducing his AA intake. It is estimated that about half of the AA intake results from home-made meals or meals prepared in restaurants (Grob 2007). Even by means of some simple measures such as avoiding excessive browning when frying, baking or roasting potato and cereal products, the AA intake can be significantly reduced. In addition, the recommendations of having a diversified diet including sufficient vegetables and fruit, without exaggerating the consumption of fried food, are still valid and help to reduce the AA intake (Mestdagh et al. 2007).

Finally, it must be stressed that awareness of the food industry and the consumer as well as a clear information campaign, are indispensable for dealing with the AA issue.

Acknowledgements

The authors would like to acknowledge the Belgian Institute of Public Health for access to the consumption data, and the Scientific Committee of the Belgian Federal Agency for the Safety of the Food Chain (FASFC) for their guidance of this study. Frédéric Mestdagh is post-doctoral researcher funded by the Research Foundation Flanders (FWO Vlaanderen).

References

Boon P, de Mul A, van der Voet H, van Donkersgoed G, Brette M, van Klaveren J. 2005. Calculations of dietary exposure to acrylamide. *Mut. Res.* 580: 143–155.

BVL. 2005. Bundesamt für Verbraucherschutz und Lebensmittelsicherheit.

- 355 Minimierungskonzept zur Senkung der Acrylamidgehalte in Lebensmitteln [Internet].
356 Germany: BVL; [cited 2005 Nov 28]. Available from:
357 http://www.bvl.bund.de/cln_027/nn_592438/DE/01__Lebensmittel/03__UnerwStoffeUndOrga
358 [nismen/04__Acrylamid/00__Minimierungskonzept/minimierungskonzept__node.html__nnn=t](http://www.bvl.bund.de/cln_027/nn_592438/DE/01__Lebensmittel/03__UnerwStoffeUndOrga)
359 [rue](http://www.bvl.bund.de/cln_027/nn_592438/DE/01__Lebensmittel/03__UnerwStoffeUndOrga)
- 360 CIAA. 2009a. Confederation of the food and drink industries of the EU. The CIAA acrylamide
361 “toolbox”. February 2009, Rev. 12 [Internet]. EU: CIAA ; [cited 2009 Feb 16]. Available from:
362 http://www.ciaa.be/asp/documents/brochures_form.asp?doc_id=65
- 363 CIAA. 2009b. Confederation of the food and drink industries of the EU. Acrylamide
364 Pamphlets[Internet]. EU: CIAA ; [cited 2009 Feb 16]. Available from:
365 http://www.ciaa.be/asp/documents/l1.asp?doc_id=822
- 366 Claeys W, De Vleeschouwer K, Hendrickx M. 2005. Quantifying the formation of carcinogens
367 during food processing: acrylamide. Trends Food Sci. Technol. 16: 181-193.
- 368 Claus A, Carle R, Schieber A. 2008. Acrylamide in cereal products: A review. J. Cereal Sci. 47:
369 118–133.
- 370 Devriese S, De Backer G, De Henauw S, Huybrechts I, Kornitzer K, Leveque A, Moreau M, Van
371 Oyen H. 2005. The Belgian food consumption survey: aims, design and methods. Arch.
372 Public Health 63: 1-16.
- 373 Dybing E, Farmer P, Andersen M, Fennell T, Lalljie S, Muller DJ, Olin S, Peterson B, Schlatter J,
374 Scholz G, Scimeca J, Slimani N, Törnqvist M, Tuijelaars S, Verger P. 2005. Human exposure
375 and internal dose assessments of acrylamide in food. Food Chem. Toxicol. 43: 365-410.
- 376 ECHA. 2009. Press release: 15 new substances of very high concern to be added to the
377 Candidate List [Internet]. Helsinki (Finland): ECHA; [cited: 2009 Dec 7]. Available from:
378 http://echa.europa.eu/doc/press/pr_09_15_msc_svhc_20091207.pdf
- 379 EFSA. 2008a. European Food Safety Authority. EFSA's 11th Scientific Colloquium - Acrylamide
380 carcinogenicity - New evidence in relation to dietary exposure - 22 and 23 May 2008. Tabiano
381 (Italy): EFSA; [cited 2008 May 22]. Available from:
382 <http://www.efsa.europa.eu/en/events/event/colloque080522.htm>
- 383 EFSA. 2008b. Concise European Food Consumption Database. Tabiano (Italy): EFSA; [cited

- 2009 January 22]. Available from: <http://www.efsa.europa.eu/en/datex/datexfooddb.htm>.
- EFSA. 2008c. European Food Safety Authority. Polycyclic aromatic hydrocarbons in food. Scientific opinion of the Panel on Contaminants in the Food Chain (Question No EFSA-Q-2007-136). EFSA J. 724 : 1-114. Available from: http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1211902034842.htm
- EFSA. 2005. European Food Safety Authority. Opinion of the Scientific Committee on a request from EFSA related to a harmonised approach for risk assessment of substances which are both genotoxic and carcinogenic (Request No EFSA-Q-2004-020). EFSA J. 280 : 1-31. Available from: http://www.efsa.europa.eu/EFSA/efsa_locale-1178620753812_1178620763354.htm
- Fredriksson H, Tallving J, Rosén J, Åman P. 2004. Fermentation reduces free asparagine in dough and acrylamide content in bread. Cereal Chem. 81: 650-653.
- Friedman M, Levin C. 2008. Review of methods for the reduction of dietary content and toxicity of acrylamide. J. Agric. Food Chem. 56: 6113-6140.
- Gökmen V, Açar Ö, Köksel H, Acar J. 2007. Effects of dough formula and baking conditions on acrylamide and hydroxymethylfurfural formation in cookies. Food Chem. 104: 1136-1142.
- Graf M, Amrein T, Graf S, Szalay R, Escher F, Amadò R. 2006. Reducing the acrylamide content of semi-finished biscuit on industrial scale. Food Sci. Technol. 39: 724-728.
- Granvogl M, Koehler P, Latzer L, Schieberle P. 2008. Development of a stable isotope dilution assay for the quantitation of glycidamide and its application to foods and model systems. J. Agric. Food Chem. 56: 6087-6092.
- Grob K. 2007. Options for legal measures to reduce acrylamide contents in the most relevant foods. Food Addit. Contam. 24: 71-81.
- Hills S. 2008 Aug 04. Chips companies settle acrylamide lawsuit. Food USA [Internet]. Available from: <http://www.foodnavigator-usa.com/layout/set/print/content/view/print/215432>
- Hoenicke K, Gatermann R. 2005. Studies on the stability of acrylamide in food during storage. J. AOAC Int. 88: 268-273.
- IARC. 1994. International Agency for Research on Cancer. Summaries & Evaluations: Acrylamide (Group 2A). Vol. 60, p. 389. Available from: 15/17

- 1
2 413 <http://www.inchem.org/documents/iarc/vol60/m60-11.html>
3
4 414 IRMM. 2006. Institute for Reference Materials and Measurements. Acrylamide Monitoring
5
6 415 Database (update June 2006) [Internet]. Geel (Belgium): IRMM; [cited 2006 May 31]
7
8 416 Available from: <http://www.irmm.jrc.be/html/activities/acrylamide/database.htm>
9
10 417 JECFA. 2005. Joint FAO/WHO Expert Committee on Food Additives. Summary and conclusions
11
12 418 of the sixty-fourth meeting of the joint FAO/WHO Expert Committee on Food Additives.
13
14 419 Rome, 8-17 February 2005. Available from:
15
16 420 http://www.who.int/ipcs/food/jecfa/summaries/summary_report_64_final.pdf
17
18 421 Kita A, Erland B, Knutsen S, Wicklund T. 2004. Effective ways of decreasing acrylamide content
19
20 422 in potato crisps during processing. *J. Agric. Food Chem.* 52: 7011-7016.
21
22 423 Kroes R, Muller D, Lambe J, Lowik M, van Klaveren J, Kleiner J, Massey R, Mayer S, Urieta I,
23
24 424 Verger P, Visconti A. 2002. Assessment of intake from the diet. *Food Chem. Toxicol.* 40:
25
26 425 327-385.
27
28 426 Lanz I, Ternité R, Wilkens J, Hoenicke K, Guenther H, van der Stegen G. 2006. Studies on
29
30 427 acrylamide levels in roasting, storage and brewing of coffee. *Mol. Nutr. Food Res.* 50: 1039-
31
32 428 1046.
33
34 429 Matthys C, Bilau M, Govaert Y, Moons E, De Henauw S, Willems J. 2005. Risk assessment of
35
36 430 dietary acrylamide intake in Flemish adolescents. *Food Chem. Toxicol.* 43: 271-278.
37
38 431 Mestdagh F, De Wilde T, Delporte K, Van Peteghem C, De Meulenaer B. 2008a. Impact of
39
40 432 chemical pre-treatments on the acrylamide formation and sensorial quality of potato crisps.
41
42 433 *Food Chem.* 106: 914-922.
43
44 434 Mestdagh F, De Wilde T, Fraselle S, Govaert Y, Ooghe W, Degroodt JM, Verhe R, Van
45
46 435 Peteghem C, De Meulenaer B. 2008b. Optimization of the blanching process to reduce
47
48 436 acrylamide in fried potatoes. *Food Sci. Technol.* 41: 1648-1654.
49
50 437 Mestdagh F, Lachat C, Baert K, Moons E, Kolsteren P, Van Peteghem C, De Meulenaer B. 2007.
51
52 438 Importance of a canteen lunch on the dietary intake of acrylamide. *Mol. Nutr. Food Res.* 51:
53
54 439 509-516.
55
56 440 Mottram D, Wedzicha B, Dodson A. 2002. Acrylamide is formed in the Maillard reaction. *Nature*
57
58 441 419: 448-449.
59
60

- O'Brien J, Renwick A, Constable A, Dybing E, Müller D, Schlatter J, Slob W, Tueting W, van Benthem J, Williams G, Wolfreys A. 2006. Approaches to the risk assessment of genotoxic carcinogens in food: a critical appraisal. *Food Chem. Toxicol.* 44: 1613-1635.
- Seal C, de Mul A, Eisenbrand G, Haverkort A, Franke K, Lalljie S, Mykkänen H, Reimerdes E, Scholz G, Somoza V, Tuijtelaars S, van Boekel M, van Klaveren J, Wilcockson S, Wilms L. 2008. Risk-benefit considerations of mitigation measures on acrylamide content of foods - A case study on potatoes, cereals and coffee. *Br. J. Nutr.* 99 Suppl 2: S1-S46.
- Stadler R, Scholz G. 2004. Acrylamide: an update on current knowledge in analysis, levels in food, mechanisms of formation, and potential strategies of control. *Nutr. Rev.* 62: 449-467.
- Surdyk N, Rosén J, Andersson R, Åman P. 2004. Effects of asparagine, fructose, and baking conditions on acrylamide content in yeast-leavened wheat bread. *J. Agric. Food Chem.* 52: 2047-2051.
- Svensson K, Abramsson L, Becker W, Glynn A, Hellenäs KE, Lind Y, Rosén J. 2003. Dietary intake of acrylamide in Sweden. *Food Chem. Toxicol.* 41: 1581-1586.
- Taeymans D, Wood J, Ashby P, Blank I, Studer A, Stadler R, Gondé P, Van Eijck P, Lalljie S, Lingnert H, Lindblom M, Matissek R, Müller D, Tallmadge D, O'Brien J, Thompson S, Silvani D. 2004. A review of acrylamide: An industry perspective on research, analysis, formation, and control. *Crit. Rev. Food Sci. Nutr.* 44: 323-347.
- Tardiff R, Gargas M, Kirman C, Carson L, Sweeney L. 2010. Estimation of safe dietary intake levels of acrylamide for humans. *Food Chem Toxicol.* 48: 658-667.
- van Dooren M, Boeijen I, van Klaveren J, van Donkersgoed G. 1995. Conversion of foods to primary agriculturally products. Report 95.17 [internet]. Wageningen (The Netherlands): Rikilt; [cited 1995 March]. Available from: <http://library.wur.nl/way/bestanden/clc/912082.pdf>
- Vose D. 2006. Risk analysis – a quantitative guide. Chichester (UK): Wiley.
- WHO. 2006. World Health Organisation. Evaluation of certain food contaminants. Sixty-fourth report of the Joint FAO/WHO Expert Committee on Food Additives. Technical Report Series 930: 8-26 [internet]. Geneva (Switzerland): WHO; [cited 2006]. Available from: http://whqlibdoc.who.int/trs/WHO_TRS_930_eng.pdf

Figure 1. Contribution of the most important food groups to the estimated dietary exposure to acrylamide in terms of % (inner circle: P25 < P50 < P95 < outer circle: mean)




Table 1. Contribution of the different food groups to the dietary exposure of the Belgian population to acrylamide (µg/kg bw/day; best estimation [95% confidence interval]).

Food group	P25	P50	P75	P90	P95	P97.5	P99
Chips	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.15 [0.10 - 0.22]	0.44 [0.32 - 0.59]	0.77 [0.54 - 1.06]	1.33 [0.89 - 2.21]
Crisps	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.10 [0.06 – 0.17]	0.32 [0.21 – 0.47]	0.67 [0.43 – 1.03]
Bread	0.01 [0.01 – 0.02]	0.03 [0.03 – 0.04]	0.06 [0.05 – 0.06]	0.09 [0.08 – 0.10]	0.11 [0.10 – 0.13]	0.14 [0.12 – 0.16]	0.18 [0.15 – 0.22]
Bread rolls	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.02 [0.01 – 0.02]	0.05 [0.05 – 0.06]	0.08 [0.07 – 0.09]	0.10 [0.09 – 0.12]	0.13 [0.11 – 0.17]
Biscuits	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.07 [0.05 – 0.10]	0.18 [0.13 – 0.26]	0.34 [0.24 – 0.51]	0.64 [0.39 – 1.08]
Gingerbread	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.15 [0.08 – 0.24]	0.36 [0.21 – 0.68]
Sweet spiced biscuits	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.05 [0.03 – 0.08]	0.13 [0.09 – 0.20]	0.26 [0.17 – 0.44]
Chocolate	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.05 [0.03 – 0.07]	0.11 [0.07 – 0.17]	0.18 [0.11 – 0.32]	0.32 [0.17 – 0.55]
Coffee	0.00 [0.00 – 0.00]	0.03 [0.03 – 0.04]	0.09 [0.08 – 0.10]	0.16 [0.14 – 0.19]	0.23 [0.20 – 0.28]	0.31 [0.26 – 0.40]	0.45 [0.34 – 0.63]
Coffee surrogate	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.34 [0.20 – 0.75]
Breakfast cereals	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.01 [0.00 – 0.03]	0.09 [0.05 – 0.15]	0.21 [0.13 – 0.31]	0.37 [0.22 – 0.52]
Toast	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.01]	0.03 [0.01 – 0.05]	0.07 [0.03 – 0.13]
Popcorn	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]
Cereal bars	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]	0.00 [0.00 – 0.00]
Intake (total) (µg/kg bw per day)	0.10 [0.09 – 0.11]	0.19 [0.18 – 0.21]	0.40 [0.35 – 0.44]	0.76 [0.67 – 0.88]	1.12 [0.94 – 1.35]	1.54 [1.24 – 1.97]	2.26 [1.68 – 3.25]

Table 2. Comparison of 'Margins of exposure' (MOEs) determined in the present study with MOE values given in literature for acrylamide

T25 (µg/kg bw/day)	BMDL ₁₀ (µg/kg bw/day)	Exposure (µg/kg bw/day)	MOE		Remarks	Ref.
			T25	BMDL		
650	310	0.41	1600	760	Mean exposure ♂ (Norway)	(1)
		0.42	1600	740	Mean exposure ♀ (Norway)	(1)
		0.43	1500	720	Mean exposure (USA)	(1)
		0.92	710	340	P90 exposure (USA)	(1)
		2.31	280	130	P90 exposure 2-5 years old (USA)	(1)
	200 ^a			200 / 50	Mean / high exposure	(2)
	2000 ^a	1 / 4		2000 / 500		
	300	1 / 4		300 / 75	Mean / high exposure	(2)
	300	0.2 / 1.6		1500 / 188	Mean / P97.5 exposure	Present study

Notes: (1): O'Brien et al. 2006; (2): JECFA 2005

^a: NOAEL ('No observed adverse effect level') instead of BMDL₁₀

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Effect of some specific mitigation scenarios on the acrylamide intake of the Belgian population (expressed as % reduction of the intake assuming the % decrease in AA content as stated in the scenario).

Scenario	Average	P50	P75	P90	P95	P97.5	P99	P99.9
1 bread & bread rolls -60%	9.7	29.6	27.6	15.4	8.0	5.1	3.3	1.7
2 coffee -30%	5.6	15.2	16.2	8.7	4.7	3.3	2.4	1.9
3 biscuit -70%	9.1	0.0	0.0	7.4	9.9	10.6	11.2	12.4
4 chips -35%	7.7	0.0	0.0	9.4	11.0	9.4	8.2	7.9
crisps -35%	2.7	0.0	0.0	0.0	2.5	3.6	4.3	12.4
chips & crisps -35%	10.4	0.0	0.0	9.4	13.6	13.0	12.4	20.3
5 gingerbread -60%	2.2	0.0	0.0	0.0	0.0	3.1	4.2	4.1
4+5 chips & crisps -35% + gingerbread -60%	12.6	0.0	0.0	9.4	13.6	16.1	16.6	24.5

Table 4. Application of signal values ($\mu\text{g/kg}$) on the acrylamide intake of the Belgian population (expressed as percentage intake reduction); (1) effect of applying Belgian P90 signal values on each food group separately and on all food groups together, and (2) effect of applying German 2008 signal values on all food groups together (BVL 2005).

Food group	P90 Signal value ($\mu\text{g/kg}$)		Average	P75	P90	P95	P97.5	P99	P99.9
	German 2008	Belgian							
Breakfast cereals	80	430	0.2	0.0	0.0	0.0	0.0	0.6	0.4
Crisps	1000 (1063) ^a	1000 (1053)	0.8	0.0	0.0	0.2	0.2	0.9	2.3
Coffee	937 ^b	506 ^b	0.0	0.1	0.1	0.0	0.0	0.0	0.0
Chips	530 (589)	622	3.5	0.0	0.3	0.3	2.2	4.4	13.3
Biscuits	545 (738)	650	2.7	0.0	0.1	0.6	1.5	2.7	8.1
Chocolate	/	414	0.9	0.0	0.0	0.3	0.4	1.4	2.4
Gingerbread	1000 (1262)	1000 (1698)	0.8	0.0	0.0	0.0	0.1	1.4	2.7
Sweet spiced biscuits	416 (563)	719	0.3	0.0	0.0	0.1	0.2	0.4	0.5
Toast	496 (661)	230	0.1	0.0	0.0	0.0	0.0	0.1	0.3
Coffee surrogate	801 (1370) ^b	1000 (3440) ^b	1.9	0.0	0.0	0.0	0.0	3.9	5.5
Bread	/	50	11.9	35.3	15.7	7.9	4.9	3.1	1.6
Bread rolls	/	50	0.4	0.3	0.6	0.7	0.4	0.4	0.3
Popcorn	/	494	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cereal bars	/	135	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Total: Belgian signal values ^c			11.6	0.3	1.0	1.8	4.7	16.1	35.3
Total: German signal values ^c			16.3	0.0	0.0	37.4	59.3	72.9	84.2

Notes: ^a: values between brackets are the measured but not the applied values.

^b: powder

^c: the effect of P90 signal values applied on all food groups considered