



HAL
open science

A preliminary study of habitat and resource partitioning among co-occurring tropical dolphins around Mayotte, south-west Indian Ocean

Alexandra Gross, Jeremy Kiszka, Olivier van Canneyt, Pierre Richard,
Vincent Ridoux

► To cite this version:

Alexandra Gross, Jeremy Kiszka, Olivier van Canneyt, Pierre Richard, Vincent Ridoux. A preliminary study of habitat and resource partitioning among co-occurring tropical dolphins around Mayotte, south-west Indian Ocean. *Estuarine, Coastal and Shelf Science*, 2009, pp.367-374. hal-00606243

HAL Id: hal-00606243

<https://hal.science/hal-00606243>

Submitted on 5 Jul 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **A preliminary study of habitat and resource partitioning among co-occurring**
2 **tropical dolphins around Mayotte, south-west Indian Ocean**

3

4

5

6 Gross Alexandra¹, Kiszka Jeremy^{1,2}, Van Canneyt Olivier³, Richard Pierre¹, Ridoux Vincent^{1,3*}

7

8

9 Running title: Tropical dolphin niche segregation

10

11

12

13 1: Littoral ENvironnement et Sociétés (LIENSs), UMR6250, Université de la Rochelle, 2 rue
14 Olympe de Gouges, 17000 La Rochelle, France.

15 2: Direction de l'Environnement et du Développement Durable, Collectivité Départementale de
16 Mayotte, BP 101, 97600 Mamoudzou, Mayotte, France. Email : jeremy.kiszka@wanadoo.fr

17 3: Centre de Recherche sur les Mammifères Marins, Université de la Rochelle, 23 avenue Albert
18 Einstein, F-17071 La Rochelle cedex, France.

19

20 * Corresponding author: vincent.ridoux@univ-lr.fr

21 Tel.: +33(0)5.4644.9910

22 **Abstract**

23

24 Mayotte in the southwest Indian Ocean is characterized by high dolphin diversity. They may
25 coexist within a fairly small area around the island because they exploit neither the same
26 preferential habitats nor the same resources. This preliminary study aimed to investigate
27 ecological niche segregation among these delphinid communities: the Indo-Pacific bottlenose
28 dolphin, *Tursiops aduncus*, the pantropical spotted dolphin, *Stenella attenuata*, the spinner
29 dolphin, *Stenella longirostris*, and the melon-headed whale, *Peponocephala electra*. Two
30 approaches were used. Habitat preferences were investigated by analysing dolphin sighting data
31 and associated physiographical characteristics. Resource partitioning was explored by analysing
32 C and N stable isotopes in skin and blubber biopsies. Only *T. aduncus*, which showed clear
33 association with coastal habitats in the lagoon, differed from the others in terms of habitat
34 preferences, characterised by shallow depth and slope, and proximity to the coast. All other
35 species shared similar oceanic habitats immediately outside the lagoon, these being of higher
36 depth and slope, greater distance from the coast and were not discernable by discriminant
37 analysis. The two *Stenella* species and the melon-headed whale displayed very high overlap in
38 habitat physiographic variables. The analysis of stable isotopes confirmed the ecological isolation
39 of *T. aduncus* and revealed a clear segregation of *P. electra* compared to the two *Stenella* that
40 was not apparent in the habitat analysis. This may reflect ecological differences that were not
41 observable from diurnal surface observations.

42

43 **Key-words:** dolphins, ecological niche, habitats, stable isotopes, south-west Indian Ocean,
44 Mayotte.

45

46 **1. Introduction**

47 In biological communities, each species has its own unique niche, which provides the conditions
48 and resources needed for its survival. A shared resource in limited supply will bring about
49 competition between members of the same species (intra-specific competition) or between
50 individuals of different species (inter-specific competition). Competition can take two different
51 forms: interference, which is a direct, often aggressive, interaction between individuals, or
52 exploitation-competition, in which individuals interact with each other indirectly, by responding
53 to a resource level which has been depressed by the activity of competitors (Begon et al., 1986).

54 A niche occupied by a species in the absence of competitors is termed its fundamental niche,
55 whereas in the presence of competitors, species may be confined to a realized niche, which is
56 shaped by the presence of competing sympatric species (Begon et al., 1986). Hence, the
57 coexistence of potentially competing species is often made possible by the differentiation of their
58 realized ecological niches. The first mechanism that allows for niche differentiation is resource
59 partitioning. In this case, different species living in the same habitat exploit the resources
60 differently. For example, predators of different size may feed on prey of different size, hence
61 minimising the overlap between the various predators' prey size ranges. Prey specialization
62 presumably allows for niche partitioning in areas of sympatry (Ballance, 2002). The second
63 mechanism involves spatial (microhabitat differentiation) or temporal separation in the
64 availability of the different resources (these become available at different times of the day or
65 different seasons of the year; Huisman and Weissing, 2001).

66 Small delphinids belong to numerous species which have similar morphological characteristics.
67 This situation suggests that fine-scale mechanisms allow for the partitioning of habitats and
68 resources when and where the different species live in sympatry. A study of the cetacean
69 community of Great Abaco, Bahamas, has shown that the ecological niches of the four species

70 that permanently live there do not overlap (MacLeod et al., 2004). These species capture prey at
71 different depths of the water column. Other cetacean species are observed in the study area only
72 during the season when prey abundance is sufficiently high to support their presence, while they
73 are competitively excluded for the rest of the year (MacLeod et al., 2004). Indeed, the spatial
74 distribution of marine predators is mainly determined by the distribution and availability of their
75 prey, these in turn varying according to physical, chemical and biological characteristics of the
76 water masses (Forcada, 2002).

77 The dietary ecology of marine mammals and their trophic level can be determined using different
78 methods: traditional methods analyse faeces and regurgitated food of living animals, as well as
79 the stomach contents of dead animals (from strandings or by-catch; e.g. Pusineri et al., 2007). A
80 more recent method for studies of dietary ecology is stable isotope analysis of blubber, skin or
81 muscle samples (Bearhop et al., 1993). The carbon and nitrogen isotope ratios ($^{13}\text{C}/^{12}\text{C}$ and
82 $^{15}\text{N}/^{14}\text{N}$, expressed hereafter as $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) of a consumer reflect those of its diet, with a slight
83 retention of the heavier isotope and excretion of the lighter one (Das et al., 2003). As a
84 consequence, tissues will be enriched with heavy isotopes at every trophic level. The minor
85 stepwise trophic enrichment of the carbon-isotope ratio limits its use in assessing trophic levels,
86 but enhances its use in tracking carbon sources through a food chain. The carbon isotope ratio of
87 secondary and tertiary consumers should thus reflect the source of carbon at the base of their food
88 chain (Kelly, 2000). The higher enrichment of the nitrogen isotope ratio of consumers compared
89 to their prey makes it very useful for the determination of their trophic level (Kelly, 2000). Thus,
90 both ratios help elucidate trophic relations and habitat use.

91 Mayotte, a volcanic island in the northern Mozambique Channel (southwest Indian Ocean), is
92 characterized by the permanent presence of more than 20 species of cetaceans (Kiszka et al.,
93 2007). Of these, the most important in coastal waters are the Indo-Pacific bottlenose dolphin,

94 *Tursiops aduncus* (Ehremberg, 1833), the pan-tropical spotted dolphin, *Stenella attenuata* (Gray,
95 1846), the spinner dolphin, *S. longirostris* (Gray, 1828), and the melon-headed whale,
96 *Peponocephala electra* (Gray, 1846). The island has a great variety of marine ecosystems
97 offering a large diversity of habitats: coasts, mangroves, an extended lagoon (1100 km²),
98 different kinds of reefs (fringing, pinnacles, and barrier), a steep insular slope with many
99 submarine canyons and seamounts, and the open ocean (Quod et al., 2000). At least twelve
100 species of delphinids may coexist in a fairly small area around the island because they exploit
101 neither the same habitats nor the same resources (Kiszka et al., 2007). The present study aimed to
102 investigate ecological niche segregation among the resident dolphin community of Mayotte,
103 especially the Indo-Pacific bottlenose dolphin, the pantropical spotted dolphin, the spinner
104 dolphin and the melon-headed whale. We concentrated on these four species as they are of fairly
105 similar size and can be found within the same proximity around Mayotte, in closely-related
106 habitats within a small area and at all seasons (Kiszka et al., 2007). This is particularly so for the
107 two *Stenella* species and the melon-headed whale which are all encountered immediately outside
108 the barrier-reef and in the channels, whereas existing literature suggests that they would be more
109 differentiated habitat-wise, with the spinner dolphin feeding offshore but resting inshore, the
110 melon headed whale being an oceanic squid-eater and the Indo-Pacific bottlenose dolphin
111 dwelling in nearshore coastal habitats (Norris et al., 1994; Silva et al., 2005; Perryman, 2002;
112 Wells and Scott, 2002). Two approaches were used for the study: a comparison of habitat by the
113 analysis of dolphin sighting data and associated behavioural and physiographic characteristics
114 (group size, depth, slope, distance to the coast and proximity to the different kinds of reefs), and
115 the study of habitat and resource partitioning by the analysis of C and N stable isotopes from skin
116 and blubber biopsies.

117

118 **2. Material and Methods**

119 2.1 Study area

120 The volcanic island of Mayotte (45°10'E, 12°50'S), which is part of the Comoros archipelago, is
121 located in the northern Mozambique Channel (Indian Ocean) between Madagascar and Southeast
122 Africa. Its surface area is 376 km² and it is composed of two main islands and about 30 islets
123 scattered within and around a lagoon. Mayotte is surrounded by a large lagoon-reef complex,
124 whose width varies from 3 to 15 km. Fringing reefs surround the archipelago, an inner double-
125 reef is present off the south-west end of Mayotte, and the barrier reef, which is interrupted by
126 numerous channels, separates the lagoon itself (maximum depth 80 m) from the external slope
127 and more oceanic habitats.

128 The four species of interest, the Indo-Pacific bottlenose dolphin, the pantropical spotted dolphin,
129 the spinner dolphin, and the melon-headed whale, range in size from about 200 cm and 90 kg for
130 the spinner dolphin to 250 cm and 250 kg for the melon-headed whale (Perrin, 2002a and b,
131 Perryman, 2002, Wells and Scott, 2002).

132

133 2.2 Data and sample collection

134 Data were collected from 1997 to 2005, during small boat-based surveys dedicated to studying
135 marine mammals (Figure 1). Several types of boats were used: a 7 m catamaran equipped with
136 two, four-stroke, 60-hp outboard engines; a 7 m boat equipped with two, two-stroke, 40-hp
137 outboard engines; a 6.4 m cabin boat equipped with an inboard four-stroke and 150-hp outboard
138 engine. Surveys were conducted during daylight hours, i.e. between 0700 h and 1800 h, in sea
139 conditions not exceeding Beaufort 3. The survey vessels did not follow pre-defined transects but
140 sampling covered all habitats within the lagoon and over the external insular slope (Figure 1).
141 Effort varied according to month (Figure 2), with more effort being applied in the austral summer

142 (November to January). When dolphins were encountered, preliminary information records
143 included group size (maximum, minimum, best estimate), geographic position, activity
144 (travelling, resting, foraging/feeding, socialising, milling, play), group classification on the basis
145 of the relative size of individuals (adults, sub-adults, calves), research boat disturbance (bowride,
146 approach, avoidance, no response) as well as group formation (tight, loose, dispersed, variable,
147 convergent; Shane, 1990; Würsig et al., 1998). This study is still on-going and, therefore, only
148 the sighting locations and associated physiographic variables are analysed here.

149 When conditions were optimal (good weather and sea state, dolphins closely approaching the
150 boat), biopsies were collected using a cross-bow (*BARNETT Veloci-Speed® Class*) with Finn
151 Larsen bolts and tips (20-mm). The dolphins were hit below the dorsal fin, when close (3-10
152 meters) to the research boat. Sampling periods spanned all seasons but sample sizes did not allow
153 seasonal comparisons (January, August and December for *T. aduncus*, N=4; January, February,
154 March and October for *S. attenuata*, N=4; February, March, May and October for *S. longirostris*,
155 N=5; March, July and December for *P. electra*, N=4). It was not possible to determine sex, size
156 or age of the individuals biopsied. Blubber and skin biopsy samples were preserved separately in
157 90° ethanol before shipping and subsequent analysis. Biopsy sampling was conducted under
158 scientific permit #78/DAF/2004.

159

160 2.3 Database

161 We constituted a database in which every dolphin observation was associated with the
162 physiographic characteristics (distance to the coast, to the different reefs and to the closest
163 channel, as well as depth and slope of seafloor and the variance of these two parameters)
164 corresponding to the GPS (Global Positioning System) fixes of the observation. The distance data
165 were obtained using GIS (Geographic Information System) software ArcView (ArcGIS 8.2) by

166 ESRI (Environmental Systems Research Institute). Bathymetric data were obtained from *Service*
167 *Hydrographique et Océanographique de la Marine* (SHOM). Interpolation of bathymetry data,
168 needed to generate depth and slope data for each sighting, was undertaken with the extension
169 Spatial Analyst by kriging transformation of the raster file into an interpolated data file. This was
170 obtained by calculating the mean value of the twelve points closest to every bathymetric point in
171 a 1 km² grid. This interpolated data file thus allowed us to generate depth and slope data over the
172 whole study area and therefore to associate bathymetry data to any dolphin observation in the
173 area.

174

175 2.4 Data analysis

176 The environmental data were first compared between species using basic methods (non-
177 parametric analyses and ANOVA). Then we used multivariate statistical methods including
178 discriminant linear and quadratic analyses to examine how well sightings were assigned to the
179 correct species from the combination of associated physiographic variables. All methods were
180 implemented with the software R (R-2.2.1, R Development Core Team, 2005).

181

182 2.5 Stable isotope analyses

183 Blubber and skin were separated for each biopsy. The ethanol they contained was evaporated at
184 45°C over 48 h and the samples were ground and freeze-dried (Hobson et al., 1997). Lipids were
185 extracted using cyclohexane (C₆H₁₂) prior to analysis because they are depleted in ¹³C; if they
186 were not extracted, this would cause a bias in the isotopic signature of ¹³C (De Niro and Epstein,
187 1978; Tieszen et al., 1983). Small sub-samples (0.35 to 0.45 mg ± 0.001 mg) were prepared for
188 analysis. Stable isotope measurements were performed with a continuous-flow isotope-ratio mass
189 spectrometer (Isoprime, Micromass) coupled to an elemental analyser (Eurovector EA 3024).

190
191 Results are expressed in δ notation relative to PeeDee Belemnite and atmospheric N_2 for $\delta^{13}C$ and
192 $\delta^{15}N$, respectively, according to the equation $\delta X = [(R_{\text{sample}}/R_{\text{standard}}) - 1] \times 10^3$, where X is ^{13}C or
193 ^{15}N and R is the isotope ratio $^{13}C/^{12}C$ or $^{15}N/^{14}N$, respectively. Replicate measurements of internal
194 laboratory standards (acetanilide) indicated that measurement errors were $<0.15\%$ and $<0.20\%$
195 for $\delta^{13}C$ and $\delta^{15}N$, respectively. Percent C and N elemental composition of tissues were obtained
196 using the elemental analyzer and used to calculate the sample C:N ratio, indicating a good lipid
197 removal efficiency when <4 .

198 199 **3. Results**

200 3.1 General
201 Our cetacean sightings comprised 394 positional data for *S. longirostris* ($n = 208$), *S. attenuata* (n
202 $= 88$), *T. aduncus* ($n = 83$) and *P. electra* ($n = 15$) in all sectors around Mayotte (Figure 2) and all
203 seasons (Figure 3). Indo-Pacific bottlenose were mostly found within the lagoon, very often in
204 the immediate proximity of the fringing reef, and to a lesser extent over the North West bank
205 located outside the main barrier reef. The two *Stenella* species were observed all along the barrier
206 reef on its outer side and made only a few incursions into the lagoon. Finally, the melon-headed
207 whale was only seen outside the barrier reef.

208
209 3.2 Habitat use and niche partitioning

210 A preliminary Principal Component Analysis (PCA plot not shown) allowed us to select group
211 size, distance to coast, depth and slope as the set of physiographic variables with least
212 redundancy. Distances to the various reefs and to the nearest channel were heavily redundant

213 relative to distance from coast and therefore not retained. A simple comparison of the distribution
214 of these variables for the four focal species was performed using Kruskal-Wallis tests (Figure 4).
215 *T. aduncus* is observed at short distances from the coast with shallow bathymetry and moderate
216 slopes, whereas the other three species cannot be significantly differentiated in terms of habitat
217 characteristics. In particular, the two *Stenella* species manifested similar characteristics in group
218 size and environmental preferences. On the other hand, group size differed strongly between *T.*
219 *aduncus* that lives in small groups (median = 6) and *P. electra* that is mostly found in groups of
220 several hundred individuals (median = 300).

221 Linear discriminant analysis separated the species according to their environmental
222 characteristics (Figure 5). The predictive power of the linear discriminant analysis was good only
223 for *T. aduncus* which can be explained by its habitat preferences (Table 1, upper part). The other
224 species' habitat characteristics were too similar to permit acceptable predictions. The good
225 prediction for *S. longirostris* might be attributable only to the considerable number of sighting
226 data for this species. Finally, the quadratic discriminant analysis provided slightly better
227 predictive results, especially for *P. electra* (Table 1, lower part).

228 The different methods used to compare the preferred habitats of the four delphinid species under
229 study all clearly showed that *T. aduncus* differs from the other species in its preferred
230 environmental parameters. *S. attenuata* and *S. longirostris* share similar environmental
231 characteristics. *P. electra* is characterised by a larger group size and deeper depth preference, but
232 resembles *Stenella* in terms of the majority of its other environmental preferences (distance and
233 slope).

234

235 3.3 Resource partitioning

236 Stable isotope ratios were lower in blubber than in skin but the pattern of differences observed
237 between species was similar in both tissues (Figure 6). Skin and blubber of *T. aduncus* have the
238 highest values of $\delta^{13}\text{C}$. The two *Stenella* species are not discernable from each other as shown by
239 the extensive overlap in standard deviation for both $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$, and have the lowest values for
240 $\delta^{13}\text{C}$. *P. electra* has the highest $\delta^{15}\text{N}$ and a $\delta^{13}\text{C}$ values that are intermediate between those for the
241 two *Stenella* species and *T. aduncus*. The intra-specific variance is represented by the standard
242 deviation, which is more important for the $\delta^{13}\text{C}$ values than for the $\delta^{15}\text{N}$, except in the blubber
243 samples of *T. aduncus* where intra-specific variance in the $\delta^{15}\text{N}$ is prevalent.

244

245 **4. Discussion**

246

247 The present study comprises a preliminary analysis of habitat and resource use among an
248 assemblage of co-existing tropical delphinids living around Mayotte in the southwest Indian
249 Ocean. The principal finding is that, among the four species of interest, the Indo-Pacific
250 bottlenose dolphin is clearly differentiated from the other species in terms of both habitat
251 preference and stable isotope analyses. From the variables tested here, the other three species can
252 hardly be separated in terms of their preferred habitats but stable isotope analysis revealed a
253 dietary segregation between the melon-headed whale and the two species of the genus *Stenella*.
254 This was not initially evident in the sighting data analyses that described the dolphins' diurnal
255 use of habitats. However, some limitations render these findings preliminary. The sightings
256 constituted presence-only data as the observation effort could not be readily quantified and,
257 hence, the data cannot provide significant information concerning dolphin absence. In this work,
258 we tried to characterise the habitats where dolphins were found, not assess the overall distribution
259 of each species around the island. Nevertheless, the field surveys comprehensively covered the

260 study area such that all habitats were visited and the sighting data of the four focal species were
261 considered representative of their habitat preference.

262
263 The Indo-Pacific bottlenose dolphin, *Tursiops aduncus*, occupies an ecological niche that clearly
264 differs from those occupied by the other study species. *T. aduncus* was observed only inside the
265 lagoon except in the northern part of the island where the reef system is in the form of an open
266 bank outside the lagoon. This species is generally found close to the fringing reef which
267 constitutes the ecosystem where prey of this coastal dolphin may be concentrated. The high $\delta^{13}\text{C}$
268 value in its tissues indicates a benthic carbon source that dolphins more easily access in coastal
269 habitats (Hobson, 1999). This species has indeed been observed feeding near mangroves, along
270 the fringing reefs or over seagrass beds. Its isotopic signature of $\delta^{15}\text{N}$ is similar to that of the two
271 *Stenella*, but this cannot be interpreted as an indication of similar trophic levels as the $\delta^{15}\text{N}$ values
272 of the local primary producers *versus* those outside the lagoon have not been investigated at this
273 stage.

274 *T. aduncus* lives and forages individually or in small groups inside the lagoon, where the water is
275 shallow and large predators are absent, and probably feeds on prey that would not aggregate in
276 large schools (Mann et al., 2000). Indo-Pacific bottlenose dolphins forage over reefs or soft
277 bottom substrata and near the shore relatively close to the island of Mayotte and around Zanzibar
278 (Tanzania) (Amir et al., 2005).

279 The pantropical spotted dolphin and the spinner dolphin share a barrier reef-associated habitat
280 outside the lagoon. They are also found where the water depth rapidly attains a depth of 100 m.
281 In comparison to the Indo-Pacific bottlenose dolphin, their tissues had a lower $\delta^{13}\text{C}$, suggesting it
282 was derived from an oceanic carbon source; oceanic phytoplankton is reported to be ^{13}C -depleted

283 relative to marine phanerogams (Hobson, 1999). Their isotopic ratios suggested their diet
284 comprised oceanic prey.

285 *S. attenuata* and *S. longirostris* can occasionally be observed inside the lagoon where they might
286 take advantage of the safety it provides for resting. This kind of behaviour has been observed in
287 Hawaiian spinner dolphin populations (Norris et al., 1994). *S. longirostris* generally lives in
288 single-species groups from several dozens to several hundreds of individuals, e.g. around the
289 main Hawai'i island (Norris et al., 1994). Their aggregation in large groups might offer some
290 protection against predators but it might also facilitate feeding through communal hunting on
291 large pelagic fish schools. In the present work, *S. attenuata* was rarely observed in single-species
292 groups but was generally found in association with *S. longirostris*, forming important mixed-
293 species groups. Both species might take advantage of a larger group size for safety and foraging.
294 Nevertheless, a large group size increases the potential for intra-specific competition as well as
295 inter-specific competition if the two species feed together. A study in the southwest Atlantic has
296 shown that distributions of spotted dolphins and spinner dolphins may partially overlap (Moreno
297 et al., 2005). Associations of spotted dolphins and spinner dolphins are frequently found in
298 Hawai'i, where the two species travel together while showing many aggressive interactions, but
299 they do not seem to feed together (Psarakos et al., 2003). Conversely, in the Azores, common
300 dolphins, *Delphinus delphis*, and Atlantic spotted dolphins, *Stenella frontalis*, associate and
301 forage together (Clua and Grosvalet, 2001). As *S. attenuata* and *S. longirostris* have similar
302 habitat preferences and similar isotopic signatures, both for carbon and nitrogen, there is no
303 evidence of ecological niche differentiation in these two species. Two options could be
304 investigated in the future: either the two dolphins share the same resources in the same habitats,
305 presumably because food is not limiting there, or they segregate when feeding on different

306 resources but their prey have similar isotopic signatures because they rely on the same carbon
307 source at the same trophic level.

308 The melon-headed whale, *Peponocephala electra*, prefers steep slopes and slightly deeper water
309 than the two *Stenella* species. *Peponocephala electra* forms large groups of several hundred
310 individuals. As it was always observed in a habitat that overlaps with that of the spinner and
311 spotted dolphins, analysis of its habitat characteristics failed to distinguish between it and the two
312 *Stenella* species. This might be due to the fact that melon-headed whale sightings were only made
313 on groups that occasionally approached the island, as the surveys were conducted in a limited
314 area around Mayotte and groups living farther offshore in the open ocean would not have been
315 observed. In this case, our understanding of the preferred habitat of this species remains marginal,
316 being limited to its nearshore fringe. Stable isotope analysis, on the other hand, clearly showed
317 that there is an ecological differentiation between *P. electra* and the two *Stenella* species. Values
318 of $\delta^{13}\text{C}$ for *P. electra*'s were intermediate between those of the *Stenella* species and *T. aduncus*.
319 This would suggest that carbon isotopic composition of *P. electra* is influenced more by benthic
320 primary production than that of the two *Stenella* species. *Peponocephala electra* is known for its
321 oceanic habitat (Perryman, 2002), and possible interpretations for its $\delta^{13}\text{C}$ include the possibility
322 that it is a deep-diving species that feeds on prey derived from a detritus-based food web unlike
323 the *Stenella* that feed on a phytoplankton-based food web. In addition, *P. electra* clearly differs in
324 its $\delta^{15}\text{N}$, this being indicative of a higher trophic level, suggesting its diet includes more
325 carnivorous fish and squid than the spotted and spinner dolphins. Again, as in *T. aduncus*, all
326 sources of primary production should be investigated concerning their $\delta^{15}\text{N}$ to adequately
327 interpret the trophic levels.

328

329 If there is high inter-individual variance in the $\delta^{15}\text{N}$ (*versus* $\delta^{13}\text{C}$) values within a species, the
330 interpretation is that the species is composed of individuals that have varied feeding habits and
331 prey on organisms at different trophic levels ($\delta^{15}\text{N}$) or are found in different habitats ($\delta^{13}\text{C}$). A
332 low variance indicates that all the individuals are specialist feeders on similar prey or within the
333 same habitat. Thus, stable isotope variance is sometimes considered a measure of niche width
334 (Bearhop et al., 2004) or at least the inter-individual measure of niche width. Results of our stable
335 isotope analysis indicate that *P. electra* is a fairly specialized feeder, whereas *S. attenuata*, *S.*
336 *longirostris* and especially *T. aduncus* would be more eclectic feeders, which, in the latter case, is
337 in line with previous studies (Mann et al., 2000).

338
339 Detectable differences between the standard deviations in the stable isotope composition in the
340 skin and blubber samples might be attributable to their differential rates in tissue renewal: this
341 takes a few days for epidermis but several months for the collagen matrix of the blubber (Abend
342 and Smith, 1995). The residence time of elements in tissues depends on metabolic turnover rates
343 (Rubenstein and Hobson, 2004). This enables tracking of an animal's foraging history as
344 demonstrated by a study on pilot whales (*Globicephala melas*) in the North Atlantic (Abend and
345 Smith, 1995). *T. aduncus* and the two *Stenella* species clearly differ in the respective isotopic
346 carbon signatures in their skin and blubber, indicating that they feed on trophic webs with
347 different carbon sources over the long term.

348 Each species' habitat preference is presumably based on their prey distributions (Baumgartner et
349 al., 2000) which, in turn, are related to water depth (Hastie et al., 2005) and, indirectly,
350 bathymetric features that influence currents and productivity (Fiedler, 2002). Foraging behaviour
351 seems to be closely related to submarine habitat characteristics (Hastie et al., 2004). It has been
352 demonstrated that a variable bathymetry contributes to global delphinid abundance, promoting

353 the regional abundance of different species (Gannier, 2005). The rich marine biodiversity of
354 Mayotte is possibly related to its variety of habitats and these provide numerous ecological niches
355 for delphinid prey.

356

357 **5. Conclusions**

358 The present study provides preliminary ecological indications of niche differentiation and
359 resource partitioning within the Mayotte delphinid communities. The only species that differs
360 from the others in terms of habitat characteristics is the coastal-dwelling Indo-Pacific bottlenose
361 dolphin, whereas pantropical spotted and spinner dolphins and the melon-headed whale share
362 similar oceanic habitats immediately outside the lagoon. Stable isotope analysis confirmed the
363 ecological specialisation of *T. aduncus* and, in addition to this, revealed a clear segregation of *P.*
364 *electra*, from to the two *Stenella* species in terms of their feeding that was not apparent in the
365 habitat analysis. This may reflect behavioural differences that were not detectable from diurnal
366 surface observations. Further work should strengthen these conclusions through dedicated
367 sampling of habitat preference and an expansion of the stable isotope studies. These should focus
368 on seasonal changes in food partitioning and the isotopic composition of a series of putative prey
369 species and primary producers characteristic of the inshore-offshore gradient of habitats found
370 around Mayotte.

371

372 **Acknowledgements**

373 Sightings were collected from 1997 to 2004 during the austral winter (July to October) as part of
374 the humpback whale (*Megaptera novaeangliae*) surveys conducted by the *Observatoire des*
375 *Mammifères Marins de Mayotte (Direction de l'Agriculture et de la Forêt, DAF)* and the
376 Cetacean Conservation and Research Program (American Museum of Natural History-Wildlife

377 Conservation Society). The other sightings were collected from November 2004 to August 2005
378 during a dolphin research project conducted by the *Office National de la Chasse et de la Faune*
379 *Sauvage* (ONCFS) and the Agriculture and Forestry Office (*Direction de l'Agriculture et de la*
380 *Forêt*). We thank Peter Ersts (American Museum of Natural History, Center for Biodiversity and
381 Conservation) for providing the map showing tracks of the cetacean surveys conducted around
382 Mayotte. We are particularly grateful to Gaël Guillou (*Université de la Rochelle*) for undertaking
383 the mass spectrometer analyses. Funding for field work was provided by the *Ministère de*
384 *l'Energie, l'Ecologie, le Développement Durable et de l'Aménagement du Territoire*
385 (MEEDDAT) and the *Collectivité Départementale de Mayotte* (CDM). We thank Robin Rolland,
386 Alban Jamon, Wilfrid Fousse, Ismaël Ousseni (DAF), Sarah Caceres, Franck Charlier, Denis
387 Girou (ONCFS), Didier Fray (CDM) and the personnel of *Brigade Nature* (CDM) for assistance
388 in the field in Mayotte.

389

390 **References**

391

392 Abend, A.G., Smith, T.D., 1995. Differences in ratios of stable isotopes of nitrogen in long-
393 finned pilot whales (*Globicephala melas*) in the western and eastern North Atlantic. *ICES Journal*
394 *of Marine Science* 52: 837-841.

395

396 Amir, O.A., Berggren, P., Ndaró, S.G.M., Jiddawi, N.S., 2005. Feeding ecology of the Indo-
397 Pacific bottlenose dolphin (*Tursiops aduncus*) incidentally caught in the gillnet fisheries off
398 Zanzibar, Tanzania. *Estuarine Coastal and Shelf Science* 63: 429-437.

399

400 Ballance, L.T. 2002. Cetacean Ecology. *In*: Perrin, W.F., Würsig, B., Thewissen, J.G.M.
401 (Editors), Encyclopedia of Marine Mammals, Academic Press, San Diego, pp 208-214.
402

403 Baumgartner, M.F., Mullin, K.D., May, L.N., Leming, T.D., 2001. Cetacean habitats in the
404 northern Gulf of Mexico. *Fishery Bulletin* 99: 219-239.
405

406 Bearhop, S., Adams, C.E., Waldron, S., Fuller, R.A., Macleod, H., 2004. Determining trophic
407 niche width: A novel approach using stable isotope analysis. *Journal of Animal Ecology* 73:
408 1007-1012.
409

410 Begon, M., Harper, J.L., Townsend, C.R., 1986. *Ecology – Individuals, Populations and*
411 *Communities*. Blackwell Scientific Publications, Boston, 876 pp.
412

413 Chapman, J.L., Reiss, M.J., 1999. *Ecology – Principles and Applications*. Cambridge University
414 Press, Cambridge, 336 pp.
415

416 Clua, E., Grosvalet, F., 2001. Mixed-species feeding aggregation of dolphins, large tunas and
417 seabirds in the Azores. *Aquatic Living Resources* 14: 11-18.
418

419 Das, K., Lepoint, G., Leroy, Y., Bouquegneau, J.M., 2003. Marine mammals from the southern
420 North Sea: feeding ecology data from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements. *Marine Ecology Progress*
421 *Series* 263: 287-298.
422

423 De Niro, M.J., Epstein, S., 1978. Influence of diet on the distribution of carbon isotopes in
424 animals. *Geochimica Cosmochimica Acta* 42: 495-506.

425

426 Estep, M.F., Dabrowski, H., 1980. Tracing food webs with stable hydrogen isotopes. *Science*
427 209: 1537–1538.

428

429 Fiedler, P.C., 2002. Ocean environment. *In*: Perrin, W.F., Würsig, B., Thewissen, J.G.M.
430 (Editors), *Encyclopedia of Marine Mammals*, Academic Press, San Diego, pp 824-830.

431

432 Forcada, J., 2002. Distribution. *In*: Perrin, W.F., Würsig, B., Thewissen, J.G.M. (Editors),
433 *Encyclopedia of Marine Mammals*, Academic Press, San Diego, pp 325-333.

434

435 Gannier, A., 2005. Summer distribution and relative abundance of delphinids in the
436 Mediterranean Sea. *Revue d'Ecologie (Terre et Vie)* 60: 223-238.

437

438 Hastie, G.D., Swift, R.J., Slessor, G., Thompson, P.M., Turrell, W.R., 2005. Environmental
439 models for predicting oceanic dolphin habitat in the Northeast Atlantic. *ICES Journal of Marine*
440 *Science* 62: 760-770.

441

442 Hastie, G.D, Wilson, B., Wilson, L.J., Parsons, K.M., Thompson, P.M. 2004. Functional
443 mechanisms underlying cetacean distribution patterns: hotspots for bottlenose dolphins are linked
444 to foraging. *Marine Biology* 144: 397-403.

445

446 Hobson, K.A., 1999. Tracing origins and migration of wildlife using stable isotopes: a review.
447 *Oecologia* 120: 314-326.
448

449 Hobson, K.A., Gibbs, H.L., Gloutney, M.L. 1997. Preservation of blood and tissue samples for
450 stable-carbon and stable-nitrogen isotope analysis. *Canadian Journal of Zoology* 75:1720-1723.
451

452 Huisman, J., Weissing, F.J., 2001. Biological conditions for oscillations generated by
453 multispecies oscillations. *Ecology* 82: 2682-2695.
454

455 Kelly, J.F., 2000. Stable isotopes of carbon and nitrogen in the study of avian and mammalian
456 trophic ecology. *Canadian Journal of Zoology* 78: 1-27.
457

458 Kiszka, J., Ersts, P.J., Ridoux, V., 2007. Cetacean diversity around the Mozambique Channel
459 island of Mayotte (Comoros Archipelago). *Journal of Cetacean Research and Management* 9:
460 105-109.
461

462 Macleod, C.D., Hauser, N., Peckham, H., 2004. Diversity, relative density and structure of the
463 cetacean community in summer months east of Great Abaco, Bahamas. *Journal of the Marine*
464 *Biological Association of the U.K.* 84: 469-474.
465

466 Mann, J., Connor, R.C., Tyack, P.L., Whitehead, H., 2000. *Cetacean Societies – Field Studies of*
467 *Dolphins and Whales.* University of Chicago Press, Chicago, 433 pp.
468

469 Moreno, I.B., Zerbini, A.N., Danilewicz, D., De Oliveira Santos, M.C., Simoes-Lopes, P.C.,
470 Lailson-Brito, Jr. J., Azevedo, A.F., 2005. Distribution and habitat characteristics of dolphins of
471 the genus *Stenella* (Cetacea: Delphinidae) in the southwest Atlantic Ocean. *Marine Ecology*
472 *Progress Series* 300: 229–240.

473
474 Norris, K.S., Würsig, B., Wells, R. S., Würsig, M., 1994. *The Hawaiian Spinner Dolphin*,
475 University of California Press, Berkeley, 408 pp.

476
477 Perrin, W.F., 2002a. Pantropical spotted dolphin *Stenella attenuata*. *In*: Perrin, W.F., Würsig, B.,
478 Thewissen, J.G.M. (Editors), *Encyclopedia of Marine Mammals*, Academic Press, San Diego, pp
479 865-867.

480
481 Perrin, W.F., 2002b. Spinner dolphin *Stenella longirostris*. *In*: Perrin, W.F., Würsig, B.,
482 Thewissen, J.G.M. (Editors), *Encyclopedia of Marine Mammals*, Academic Press, San Diego, pp
483 1174-1178.

484
485 Perrin, W.F., Würsig, B., Thewissen, J.G.M., 2002. *Encyclopedia of Marine Mammals*,
486 Academic Press, San Diego, 1414 pp.

487
488 Perryman, W.L., 2002. Melon-headed whale *Peponocephala electra*. *In*: Perrin, W.F., Würsig,
489 B., Thewissen, J.G.M. (Editors), *Encyclopedia of Marine Mammals*, Academic Press, San Diego,
490 pp 733-735.

491

492 Psarakos, S., Herzing, D.L., Marten, K., 2003. Mixed-species associations between Pantropical
493 spotted dolphins (*Stenella attenuata*) and Hawaiian spinner dolphins (*Stenella longirostris*) off
494 Oahu, Hawaii. *Aquatic Mammals* 29: 390–395.

495

496 Quod, J.P., Naim, O., Abdourazi, F., 2000. The Comoros archipelago. *In*: Sheppard, C. (Editors),
497 Seas at the Millennium: An Environmental Evaluation. Pergamon Press, Oxford, pp 243-52.

498

499 R Development Core Team, 2005. R: A language and environment for statistical computing. R
500 Foundation for Statistical Computing. Vienna, Austria. (<http://www.Rproject.org>)

501

502 Pusineri, C., Magnin, V., Meynier, L., Spitz, J., Hassani, S., Ridoux, V., 2007. Food and feeding
503 ecology of the common dolphin (*Delphinus delphis*) in the northeast Atlantic and comparison
504 with its diet in neritic areas. *Marine Mammal Science* 23: 30-47.

505

506 Rubinstein, D.R., Hobson, K.A., 2004. From birds to butterflies: animal movement patterns and
507 stable isotopes. *Trends in Ecology and Evolution* 19: 256-263.

508

509 Silva, J.M., Silva, F.J.L., Sazima, I., 2005. Rest, nurture, sex, release, and play: diurnal
510 underwater behaviour of the spinner dolphin at Fernando de Noronha Archipelago, SW Atlantic.
511 *Journal of Ichthyology and Aquatic Biology* 9: 161–176.

512

513 Tieszen, L.L., Boutton, T.W., Tesdahl, K.G., Slade, N.A., 1983. Fractionation and turnover of
514 stable carbon isotopes in animal tissues: implications for $\delta^{13}\text{C}$ analysis of diet. *Oecologia* 57: 32-
515 37.

516

517 Shane, S.H., 1990. Behaviour and ecology of the bottlenose dolphin at Sanibel Island, Florida. In:
518 S. Leatherwood, R.R. Reeves (Editors.). *The Bottlenose Dolphin*. Academic Press, San Diego,
519 pp. 256-266.

520

521 Wells, R.S., Scott, M.D., 2002. Bottlenose dolphins *Tursiops truncatus* and *T. aduncus*. In:
522 Perrin, W.F., Würsig, B., Thewissen, J.G.M. (Editors), *Encyclopedia of Marine Mammals*,
523 Academic Press, San Diego, pp 122-128.

524

525 Würsig, B., Lynn, S.K., Jefferson, T.A., Mullin, K.D., 1998. Behaviour of cetaceans in the
526 Northern Gulf of Mexico relative to survey ships and aircraft. *Aquatic Mammals* 24: 41-50.

527

528

529 **Figure captions**

530

531 **Figure 1:** Location of the study area and observation routes from July 2004 to August 2005.

532 Isobath (left) and GPS tracks of sighting surveys (right) are shown around Mayotte Island with
533 barrier reef.

534 **Figure 2:** General locations of delphinid sightings around Mayotte and its reefs.

535 **Figure 3:** Temporal distribution of effort expressed as number of sightings per three-month
536 period.

537 **Figure 4:** Delphinid sighting median values, 50, 75 percentiles and outliers of (a) group size, (b)
538 distance to the coast, (c) depth and (d) seafloor slope. Kruskal Wallis tests showed that *T.*
539 *aduncus* differed significantly in all cases with p -values $<10^{-6}$.

540 **Figure 5:** Sighting density for melon-headed whale (dotted line), pan-tropical spotted (black
541 line), spinner (dashed line) and Indo-Pacific bottlenose dolphin (dashed and dotted line) along the
542 first discriminant axis.

543 **Figure 6:** Stable isotope ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in ‰) values in delphinid blubber (right-hand graph)
544 and skin (left-hand graph). Graphs show average values and standard deviations. Black squares
545 represent *S. longirostris* (N=5), white squares *S. attenuata* (N=4), black diamonds *T. aduncus*
546 (N=4) and white diamonds *P. electra* (N=4).

547 **Table 1:** Discriminant analysis: linear and quadratic predictions for the four species. Each
 548 sighting was assigned to one of the four study species on the basis of its associated physiographic
 549 characteristics. Only *T. aduncus* data were correctly assigned to the right species whereas all
 550 others were mostly assigned to *S. longirostris* as they all share similar habitat characteristics.

551 **Linear discriminant analysis**

	<i>Peponocephala electra</i>	<i>Stenella attenuata</i>	<i>Stenella longirostris</i>	<i>Tursiops aduncus</i>
555 <i>P. electra</i>	2	3	5	0
556 <i>S. attenuata</i>	1	3	1	2
557 <i>S. longirostris</i>	12	79	194	19
558 <i>T. aduncus</i>	0	2	8	62
560 Good prediction	14%	3%	93%	77%

562 **Quadratic discriminant analysis**

	<i>Peponocephala electra</i>	<i>Stenella attenuata</i>	<i>Stenella longirostris</i>	<i>Tursiops aduncus</i>
566 <i>P. electra</i>	11	4	4	0
567 <i>S. attenuata</i>	1	18	18	1
568 <i>S. longirostris</i>	3	46	131	4
569 <i>T. aduncus</i>	0	19	55	78
571 Good prediction	79%	20%	63%	96%

574

575 Fig.1

576

577

578

579

580

581


582

583

584

585

586



587

588

589

590

591

592

593

594


595

596

597

598 Fig.2

599




600

601 Fig.3

602

603




604

605

606


607 Fig.4

608



609 Fig. 5

610



611

612


613

614

615 Fig. 6

616

617



618

619

620