



HAL
open science

Spherical structures on torus knots and links

Alexander Kolpakov, Alexander Mednykh

► **To cite this version:**

Alexander Kolpakov, Alexander Mednykh. Spherical structures on torus knots and links. *Siberian Mathematical Journal*, 2009, 50 (5), pp.856-866. <10.1007/s11202-009-0096-2>. <hal-00606856v2>

HAL Id: hal-00606856

<https://hal.science/hal-00606856v2>

Submitted on 7 Jul 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ETALAB - Open licence

Spherical structures on torus knots and links *

Alexander Kolpakov, Alexander Mednykh

Abstract

The present paper considers two infinite families of cone-manifolds endowed with spherical metric. The singular strata is either the torus knot $t(2n+1, 2)$ or the torus link $t(2n, 2)$. Domains of existence for a spherical metric are found in terms of cone angles and volume formulæ are presented.

Key words: Spherical geometry, cone-manifold, knot, link.

1 Introduction

A three-dimensional cone-manifold is a metric space obtained from a collection of disjoint simplices in the space of constant sectional curvature k by isometric identification of their faces in such a combinatorial fashion that the resulting topological space is a manifold (also called the underlying space for a given cone-manifold).

Such the metric space inherits the metric of sectional curvature k on the union of its 2- and 3-dimensional cells. In case $k = +1$ the corresponding cone-manifold is called spherical (or admits a spherical structure). By analogy, one defines euclidean ($k = 0$) and hyperbolic ($k = -1$) cone-manifolds.

The metric structure around each 1-cell is determined by a cone angle that is the sum of dihedral angles of corresponding simplices sharing the 1-cell under identification. The singular locus of a cone-manifold is the closure of all its 1-cells with cone angle different from 2π . For the further account we suppose that every component of the singular locus is an embedded circle with constant cone angle along it.

A particular case of cone-manifold is an orbifold with cone angles $2\pi/m$, where m is an integer (cf. [1]).

The present paper considers two infinite families of cone-manifolds with underlying space the three-dimensional sphere \mathbb{S}^3 . The first family consists of cone-manifolds with singular locus the torus knot $t(2n+1, 2)$ with $n \geq 1$. In the rational census [2] these knots are denoted by $(2n+1)/1$. The second family of cone-manifolds consists of those with singular locus a two-component torus link $t(2n, 2)$ with $n \geq 2$. These links are two-bridge and correspond to the links

*The work is performed under auspices of the Swiss National Science Foundation no. 200020-113199/1, "Scientific Schools"-5682.2008.1 and RFBR no. 06-01-00153.

$2n/1$ in the rational census. The simplest examples of such the knots and links are the trefoil knot $3/1$ and the link $4/1$. In the Rolfsen table [2] one finds them as the knot 3_1 and the link 4_1^2 .

By the Theorem of W. Thurston [3], the manifold $\mathbb{S}^3 \setminus 3_1$ does not admit a hyperbolic structure. However, it admits two other geometric structures [4]: $\mathbb{H}^2 \times \mathbb{R}$ and $\widetilde{\text{PSL}}(2, \mathbb{R})$. It follows from the paper [5] that the spherical dodecahedron space (i.e. Poincaré homology sphere) is a cyclic 5-fold covering of \mathbb{S}^3 branched over 3_1 . Thus, the orbifold $3_1(\frac{2\pi}{5})$ with singular locus the trefoil knot and cone angle $\frac{2\pi}{5}$ is spherical. Due to the Dunbar's census [6], orbifold $3_1(\frac{2\pi}{n})$ is spherical if $n \leq 5$, Nil-orbifold if $n = 6$ and $\widetilde{\text{PSL}}(2, \mathbb{R})$ -orbifold if $n \geq 7$. Spherical structure on the cone-manifold $3_1(\alpha)$ with underlying space the three-dimensional sphere \mathbb{S}^3 is studied in [7].

The consideration of two-bridge torus links is carried out starting with the simplest one possessing non-abelian fundamental group, namely 4_1^2 .

The previous investigation on spherical structures for cone-manifolds is carried out mainly in the papers [8, 9, 10]. The present paper develops a method to analyse existence of a spherical metric for two-bridge torus knot and link cone-manifolds. Also, the lengths of singular geodesics are calculated and the volume formulæ are obtained (cf. Theorem 1 and Theorem 2).

2 Projective model \mathbb{S}_λ^3

The purpose of the present section is to construct the projective model \mathbb{S}_λ^3 that one can use to study geometric properties of two-bridge torus knots and links and to build up holonomy representation for the corresponding cone-manifolds. Other projective models for homogeneous geometries are described in [11].

Consider the set $\mathbb{C}^2 = \{(z_1, z_2) : z_1, z_2 \in \mathbb{C}\}$ as a four-dimensional vector space over \mathbb{R} . We denote it by $\mathbb{C}_\mathbb{R}^2$ and equip with Hermitian product

$$\langle (z_1, z_2), (w_1, w_2) \rangle_{\mathbb{H}} = (z_1, z_2) \mathcal{H} \overline{(w_1, w_2)}^T,$$

where

$$\mathcal{H} = \begin{pmatrix} 1 & \lambda \\ \lambda & 1 \end{pmatrix}$$

is a symmetric matrix with $-1 < \lambda < +1$.

The natural inner product is associated to the Hermitian form above:

$$\langle (z_1, z_2), (w_1, w_2) \rangle = \text{Re} \langle (z_1, z_2), (w_1, w_2) \rangle_{\mathbb{H}}$$

and the respective norm is

$$\|(z_1, z_2)\| = |z_1|^2 + |z_2|^2 + \lambda(z_1 \bar{z}_2 + \bar{z}_1 z_2).$$

Call two elements (z_1, z_2) and (w_1, w_2) in $\overset{\circ}{\mathbb{C}}_\mathbb{R}^2 = \mathbb{C}_\mathbb{R}^2 \setminus (0, 0)$ equivalent if there is $\mu > 0$ such that $(z_1, z_2) = (\mu w_1, \mu w_2)$. We denote this equivalence relation as $(z_1, z_2) \sim (w_1, w_2)$.

Identify the factor-space $\overset{\circ}{\mathbb{C}}_{\mathbb{R}}^2 / \sim$ with the three-dimensional sphere

$$\mathbb{S}_{\lambda}^3 = \{(z_1, z_2) \in \mathbb{C}_{\mathbb{R}}^2 : \|(z_1, z_2)\| = 1\},$$

endowed with the Riemannian metric

$$ds_{\lambda}^2 = |dz_1|^2 + |dz_2|^2 + \lambda(dz_1 d\bar{z}_2 + d\bar{z}_1 dz_2).$$

By means of equality

$$ds_{\lambda}^2 = \frac{1+\lambda}{2} |dz_1 + dz_2|^2 + \frac{1-\lambda}{2} |dz_1 - dz_2|^2,$$

the linear transformation

$$\xi_1 = \sqrt{\frac{1+\lambda}{2}} (z_1 + z_2), \quad \xi_2 = \sqrt{\frac{1-\lambda}{2}} (z_1 - z_2)$$

provides an isometry between $(\mathbb{S}_{\lambda}^3, ds_{\lambda}^2)$ and (\mathbb{S}^3, ds^2) , where $ds^2 = |d\xi_1|^2 + |d\xi_2|^2$ is the standard metric of sectional curvature +1 on the unit sphere $\mathbb{S}^3 = \{(\xi_1, \xi_2) \in \mathbb{C}^2 : |\xi_1|^2 + |\xi_2|^2 = 1\}$.

Let P, Q be two points in \mathbb{S}_{λ}^3 . The spherical distance between P and Q is a real number $d_{\lambda}(P, Q)$ that is uniquely determined by the conditions $0 \leq d_{\lambda}(P, Q) \leq \pi$ and $\cos d_{\lambda}(P, Q) = \langle P, Q \rangle$.

3 Torus knots \mathbb{T}_n

Let $\mathbb{T}_n, n \geq 1$ be the torus knot $t(2n+1, 2)$ embedded in \mathbb{S}^3 . The knot \mathbb{T}_n is the two-bridge knot $(2n+1)/1$ in the rational census (Fig. 1). Let $\mathbb{T}_n(\alpha)$ denote a cone-manifold with singular locus \mathbb{T}_n and the cone angle α along it.




Figure 1: Knot $(2n+1)/1$

The aim of the present section is to investigate cone-manifolds $\mathbb{T}_n(\alpha)$, $n \geq 1$ to find out the domain of sphericity in terms of the cone angle and to derive the volume formulæ.

Two lemmas precede the further exposition:

Lemma 1 For every $0 < \alpha < 2\pi$ and $-1 < \lambda < +1$ the linear transformations

$$A = \begin{pmatrix} 1 & 0 \\ -2i e^{i\frac{\alpha}{2}} \lambda \sin \frac{\alpha}{2} & e^{i\alpha} \end{pmatrix}$$

and

$$B = \begin{pmatrix} e^{i\alpha} & -2i e^{i\frac{\alpha}{2}} \lambda \sin \frac{\alpha}{2} \\ 0 & 1 \end{pmatrix}$$

are isometries of \mathbb{S}_λ^3 .

Proof. For the further account let us assume that the multiplication of vectors by matrices is to the right. A linear transformation L of the space $\mathbb{C}_\mathbb{R}^2$ preserves the corresponding Hermitian form if and only if for every pair of vectors $P, Q \in \mathbb{C}_\mathbb{R}^2$ it holds that

$$\langle P, Q \rangle_{\mathbb{H}} = P\mathcal{H}\overline{Q}^T = PL\mathcal{H}\overline{L}^T\overline{Q}^T = \langle PL, QL \rangle_{\mathbb{H}}.$$

The condition above is equivalent to

$$\mathcal{H} = L\mathcal{H}\overline{L}^T.$$

In particular,

$$\cos d_\lambda(P, Q) = \langle P, Q \rangle = \langle PL, QL \rangle = \cos d_\lambda(PL, QL),$$

that means L preserves the spherical distance between P and Q .

Let $L = A$ and $L = B$ in series, one verifies that A and B preserve the Hermitian norm on $\mathbb{C}_\mathbb{R}^2$ and, consequently, the spherical distance on \mathbb{S}_λ^3 . \square

Lemma 2 Let A and B be the same matrices as in the affirmation of Lemma 1. Then for all integer $n \geq 1$ one has

$$(AB)^n A - B(AB)^n = 2U_{2n}(\Lambda) e^{i\frac{(2n+1)(\pi+\alpha)}{2}} \sin \frac{\alpha}{2} M,$$

where M is a non-zero 2×2 -matrix and $U_{2n}(\Lambda)$ is the second kind Chebyshev polynomial of power $2n$ in variable $\Lambda = \lambda \sin \frac{\alpha}{2}$.

Proof. As far as $-1 < \lambda < +1$, one obtains

$$-1 < \Lambda = \lambda \sin \frac{\alpha}{2} < +1.$$

Substitute

$$\Lambda = \cos \theta,$$

with the unique $0 < \theta < \pi$.

Then matrices A and B are rewritten in the form

$$A = \begin{pmatrix} 1 & 0 \\ -2i e^{i\frac{\alpha}{2}} \cos \theta & e^{i\alpha} \end{pmatrix},$$

$$B = \begin{pmatrix} e^{i\alpha} & -2i e^{i\frac{\alpha}{2}} \cos \theta \\ 0 & 1 \end{pmatrix}.$$

On purpose to diagonalize the matrix AB , use

$$V = \begin{pmatrix} i e^{-i\frac{\alpha}{2}} e^{-i\theta} & i e^{-i\frac{\alpha}{2}} e^{i\theta} \\ 1 & 1 \end{pmatrix},$$

and obtain

$$D = V^{-1}(AB)V = \begin{pmatrix} -e^{i\alpha} e^{2i\theta} & 0 \\ 0 & -e^{i\alpha} e^{-2i\theta} \end{pmatrix}.$$

Note, that V might be not an isometry, but it is utile for computation.

Thus

$$\begin{aligned} (AB)^n A - B(AB)^n &= (V D^n V^{-1})A - B(V D^n V^{-1}) = \\ &= 2 \frac{\sin(2n+1)\theta}{\sin \theta} e^{i\frac{(2n+1)(\pi+\alpha)}{2}} \sin \frac{\alpha}{2} \begin{pmatrix} -1 & \lambda \\ -\lambda & 1 \end{pmatrix} = \\ &= 2 U_{2n}(\cos \theta) e^{i\frac{(2n+1)(\pi+\alpha)}{2}} \sin \frac{\alpha}{2} M = 2 U_{2n}(\Lambda) e^{i\frac{(2n+1)(\pi+\alpha)}{2}} \sin \frac{\alpha}{2} M, \end{aligned}$$

with the matrix

$$M = \begin{pmatrix} -1 & \lambda \\ -\lambda & 1 \end{pmatrix}$$

as the present Lemma claims. \square

The main theorem of the section follows:

Theorem 1 *The cone-manifold $\mathbb{T}_n(\alpha)$, $n \geq 1$ is spherical if*

$$\frac{2n-1}{2n+1} \pi < \alpha < 2\pi - \frac{2n-1}{2n+1} \pi.$$

The length of its singular geodesic (i.e. the length of the knot \mathbb{T}_n) equals

$$l_\alpha = (2n+1)\alpha - (2n-1)\pi.$$

The volume of $\mathbb{T}_n(\alpha)$ is

$$\text{Vol } \mathbb{T}_n(\alpha) = \frac{1}{2n+1} \left(\frac{2n+1}{2} \alpha - \frac{2n-1}{2} \pi \right)^2.$$

Proof. The fundamental group of the knot \mathbb{T}_n is presented as

$$\pi_1(\mathbb{S}^3 \setminus \mathbb{T}_n) = \langle a, b | (ab)^n a = b(ab)^n \rangle,$$

with generators a and b as at Fig. 1.

Since the cone-manifold $\mathbb{T}_n(\alpha)$ admits a spherical structure, then there exists a holonomy mapping [1], that is a homomorphism

$$h : \pi_1(\mathbb{S}^3 \setminus \mathbb{T}_n) \mapsto \text{Isom } \mathbb{S}_\lambda^3.$$

We will choose h in respect with geometric construction of the cone-manifold. All the further computations to find the length of the knot \mathbb{T}_n and the volume of the cone-manifold $\mathbb{T}_n(\alpha)$ are performed making use of the corresponding fundamental polyhedron \mathcal{P}_n (Fig. 2). The construction algorithm for the polyhedron is given in [12].

The combinatorial polyhedron \mathcal{P}_n has vertices P_i , $i \in \{1, \dots, 4n+2\}$ and edges $P_i P_{i+1}$, $i \in \{1, \dots, 4n+2\}$, with $P_{4n+3} = P_1$, also $P_1 P_{2n+2}$ and $P_2 P_{2n+3}$. Let N , S denote the middle points (the North and the South poles of \mathcal{P}_n) on the edges $P_1 P_{2n+2}$ and $P_2 P_{2n+3}$, respectively. Then, consider also edges NP_i , SP_i , $i \in \{1, \dots, 4n+2\}$.

Without loss in generality, choose the holonomy representation such that

$$h(a) = A, \quad h(b) = B,$$

where A and B are matrices from Lemma 1.

The generators of the fundamental group for \mathbb{T}_n under the holonomy mapping h correspond to isometries acting on \mathcal{P}_n . These isometries identify its faces by means of rotation about the edge $P_1 P_{2n+2}$ for the top ‘‘cupola’’ of \mathcal{P}_n and rotation about $P_2 P_{2n+3}$ for the bottom one (see, Fig. 2). Then the edges $P_1 P_{2n+2}$ and $P_2 P_{2n+3}$ knot itself to produce \mathbb{T}_n (cf. [12, 13]).

In order to construct the polyhedron \mathcal{P}_n assume that its edge $P_1 P_2$ is given by

$$P_1 = (1, 0), \quad P_2 = (0, 1).$$

Then one has

$$\cos d_\lambda(P_1, P_2) = \langle P_1, P_2 \rangle = \lambda,$$

i.e. the spherical distance between the points P_1 and P_2 can vary from 0 to π . Thus, prescribing certain coordinates to the end-points of the edge $P_1 P_2$ we do not loss in generality of the consideration.

Note, that the axis of the isometry A from Lemma 1 contains P_1 and the axis of B contains P_2 . The aim of the construction for the polyhedron \mathcal{P}_n is to bring its edges $P_1 P_{2n+2}$ and $P_2 P_{2n+3}$ to be axes of the respective isometries A and B . The other vertices P_i has to be images of P_1 and P_2 under action of A and B . The polyhedron \mathcal{P}_n is said to be proper if

- (a) inner dihedral angles along $P_1 P_{2n+2}$ and $P_2 P_{2n+3}$ are equal to α ;
- (b) the following curvilinear faces are identified by A and B :

$$A : NP_1 P_2 \dots P_{2n+2} \rightarrow NP_1 P_{4n+2} \dots P_{2n+3} P_{2n+2},$$

$$B : SP_2 P_1 P_{4n+2} \dots P_{2n+3} \rightarrow SP_2 P_3 \dots P_{2n+3};$$

- (c) sum of the inner dihedral angles ψ_i along $P_i P_{i+1}$, $i \in \{1, \dots, 4n+1\}$ equals 2π ;
- (d) sum of the dihedral angles ϕ_i for corresponding tetrahedra $NSP_i P_{i+1}$, $i \in \{1, \dots, 4n+1\}$ at their common edge NS is 2π ;




Figure 2: Fundamental polyhedron \mathcal{P}_n for $\mathbb{T}_n(\alpha)$

(e) all the tetrahedra NSP_iP_{i+1} with $i \in \{1, \dots, 4n+2\}$, $P_{4n+3} = P_1$ are non-degenerated and coherently oriented.

By the orientation of a tetrahedron NSP_iP_{i+1} one means the sign of the Gram determinant $\det(S, N, P_i, P_{i+1})$ for corresponding quadruple $S, N, P_i, P_{i+1} \in \mathbb{C}_{\mathbb{R}}^2$, where $i \in \{1, \dots, 4n+2\}$, $P_{4n+3} = P_1$. A tetrahedron is non-degenerated if $\det(S, N, P_i, P_{i+1}) \neq 0$. Thus, claim (e) is satisfied if all the Gram determinants are non-zero and of the same sign.

If $\alpha = \frac{2\pi}{m}$, $m \in \mathbb{N}$, then due to the Poincaré Theorem [14, Theorem 13.5.3] claims (a) – (e) imply that the group generated by the isometries A and B is discrete and its presentation is

$$\Gamma = \langle A, B | (AB)^n A = B(AB)^n, A^m = B^m = \text{id} \rangle.$$

The metric space $\mathbb{S}_\lambda^3/\Gamma \cong \mathbb{T}_n(\frac{2\pi}{m})$ is a spherical orbifold, and \mathcal{P}_n is its fundamental polyhedron. If $m \notin \mathbb{N}$ then the group generated by A and B might be non-discrete. However, the identification for the faces of \mathcal{P}_n is of the same fashion as if it were $m \in \mathbb{N}$ and as the result one obtains the cone-manifold $\mathbb{T}_n(\alpha)$. By means of Lemma 1 and construction of \mathcal{P}_n claims (a) and (b) are satisfied. For the holonomy mapping h to exist the following relation should be satisfied:

$$h((ab)^n a) - h(b(ab)^n) = (AB)^n A - B(AB)^n = 0.$$

By Lemma 2, the condition above is satisfied if and only if

$$U_{2n}(\Lambda) = 0,$$

where $\Lambda = \lambda \sin \frac{\alpha}{2}$.

Thus, the parameter λ of the metric ds_λ^2 is determined completely by a root of the polynomial $U_{2n}(\Lambda)$. From the above formula, λ is related to the cone angle α by means of the equality

$$\lambda = \frac{\Lambda}{\sin \frac{\alpha}{2}}.$$

The roots of $U_{2n}(\Lambda)$ are given by the following formula:

$$\Lambda_k = \cos \frac{k\pi}{2n+1},$$

with $k \in \{1, \dots, 2n\}$.

The parameter λ for the metric ds_λ^2 has to be chosen in order the polyhedron \mathcal{P}_n be proper and the metric itself be spherical.

Note, that the edges $P_i P_{i+1}$, $i \in \{1, \dots, 4n+2\}$, $P_{4n+3} = P_1$ are equivalent under action of the group $\Gamma = \langle A, B \rangle$. Thus, the relation $(AB)^n A = B(AB)^n$ implies the equality

$$\sum_{i=1}^{2(2n+1)} \psi_i = 2k\pi,$$

where k is an integer.

Show that one can choose λ for the equality $k = 1$ to hold for all α in the affirmation of the Theorem. Due to the paper [15], every two-bridge knot cone-manifold with cone angle π is a spherical orbifold. In this case all the vertices P_i of the fundamental polyhedron belong to the same circle and all the dihedral angles ψ_i and ϕ_i are equal to each other [12]:

$$\phi_i = \psi_i = \frac{\pi}{2n+1}.$$

As far as $\cos d_\lambda(N, S) = \cos d_\lambda(P_i, P_{i+1}) = \lambda$, then in case $\alpha = \pi$ one obtains

$$\lambda = \frac{\Lambda_k}{\sin \frac{\pi}{2}} = \cos \theta$$

for certain $k \in \{1, \dots, 2n\}$ and then

$$\sum_{i=1}^{2(2n+1)} \psi_i = 2(2n+1)\theta.$$

Using the formula for the roots of $U_{2n}(\Lambda)$ obtain that

$$\sum_{i=1}^{2(2n+1)} \psi_i = 2k\pi$$

if $\alpha = \pi$. Thus, claim (c) for the polyhedron \mathcal{P}_n with $\alpha = \pi$ is satisfied if $k = 1$. As far as the parameter α varies continuously and sum of the angles ψ_i

represents a multiple of 2π , one has that

$$\sum_{i=1}^{2(2n+1)} \psi_i = 2\pi$$

for all α .

By analogy, show that with

$$\lambda = \frac{\Lambda_1}{\sin \frac{\alpha}{2}}$$

the equality

$$\sum_{i=1}^{2(2n+1)} \phi_i = 2\pi$$

holds, that means claim (d) is also satisfied.

Verify that under conditions of the Theorem the metric ds_λ^2 is spherical. This claim is equivalent to the inequality

$$-1 < \lambda < +1.$$

Note, that for

$$\frac{2n-1}{2n+1} \pi < \alpha < 2\pi - \frac{2n-1}{2n+1} \pi$$

it follows

$$\sin \frac{\alpha}{2} > \sin \frac{(2n-1)\pi}{2(2n+1)}.$$

As far as $\sin \frac{\alpha}{2} > 0$ and $\Lambda_1 = \sin \frac{(2n-1)\pi}{2(2n+1)} > 0$, one has

$$0 < \lambda < 1.$$

By analogy with Lemma 1 verify that

$$C = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

is an isometry of ds_λ^2 .

Fixed point sets of A and B in \mathbb{S}_λ^3 are circles

$$\text{Fix } A = \{(z_1, 0) : z_1 \in \mathbb{C}, |z_1| = 1\}$$

and

$$\text{Fix } B = \{(0, z_2) : z_2 \in \mathbb{C}, |z_2| = 1\},$$

correspondingly. The geometric meaning of C is that it maps the first fixed circle to the other. Thus, the relation $B = CAC^{-1}$ holds.

The following equalities

$$P_{2k+1} = P_1(AB)^k, \quad k \in \{0, \dots, n\},$$

$$P_{2k} = P_2(AB)^{k-1}, k \in \{1, \dots, n+1\};$$

and

$$\begin{aligned} P_{2k+1} &= P_1(BA)^{2n-k+1}, k \in \{n+1, \dots, 2n\}, \\ P_{2k} &= P_2(BA)^{2n-k+2}, k \in \{n+2, \dots, 2n+1\}, \end{aligned}$$

follow from the identification scheme of the edges of \mathcal{P}_n .
Define the auxiliary function

$$\varepsilon(m) = \frac{m}{2} \alpha - \frac{4n-m}{2} \pi.$$

By analogy with the proof of Lemma 2 it follows that

$$\begin{aligned} (AB)^k &= C(BA)^k C^{-1} = \\ &= \begin{pmatrix} -\frac{\sin(2k-1)\theta}{\sin\theta} e^{i\varepsilon(2k)} & -\frac{\sin 2k\theta}{\sin\theta} e^{i\varepsilon(2k-1)} \\ \frac{\sin 2k\theta}{\sin\theta} e^{i\varepsilon(2k+1)} & \frac{\sin(2k+1)\theta}{\sin\theta} e^{i\varepsilon(2k)} \end{pmatrix}, \end{aligned}$$

where $\theta = \frac{\pi}{2n+1}$.

Suppose N and S to be middle-points of the edges P_1P_{2n+2} and P_2P_{2n+3} , respectively. Then

$$N = (e^{i\frac{\varepsilon(2n+1)}{2}}, 0), S = (0, e^{i\frac{\varepsilon(2n+1)}{2}}).$$

For the lengths l_α of the singular geodesic one has

$$\cos \frac{l_\alpha}{4} = \langle P_1, N \rangle = \langle P_1 C, N C \rangle = \langle P_2, S \rangle.$$

Thus

$$\cos \frac{l_\alpha}{4} = \cos \frac{(2n+1)\alpha - (2n-1)\pi}{4}.$$

By construction of the polyhedron \mathcal{P}_n , the inequality $0 < l_\alpha < 4\pi$ holds. Then it follows

$$l_\alpha = (2n+1)\alpha - (2n-1)\pi.$$

Given the coordinates of the vertices P_i and the poles N and S of the polyhedron \mathcal{P}_n , verify claim (e).

For every four points $A, B, C, D \in \mathbb{C}_{\mathbb{R}}^2$, where

$$A = (A_1, A_2), B = (B_1, B_2), C = (C_1, C_2), D = (D_1, D_2),$$

their Gram determinant is

$$\det(A, B, C, D) := \det \begin{pmatrix} \operatorname{Re} A_1 & \operatorname{Im} A_1 & \operatorname{Re} A_2 & \operatorname{Im} A_2 \\ \operatorname{Re} B_1 & \operatorname{Im} B_1 & \operatorname{Re} B_2 & \operatorname{Im} B_2 \\ \operatorname{Re} C_1 & \operatorname{Im} C_1 & \operatorname{Re} C_2 & \operatorname{Im} C_2 \\ \operatorname{Re} D_1 & \operatorname{Im} D_1 & \operatorname{Re} D_2 & \operatorname{Im} D_2 \end{pmatrix}.$$

Each tetrahedron NSP_iP_{i+1} with $i \in \{1, \dots, 2n+1\}$ is isometric to $NSP_{2n+i+1}P_{2n+i+2}$, $i \in \{1, \dots, 2n+1\}$, $P_{4n+3} = P_1$ by means of the isometry C defined above. Thus, we consider only the tetrahedra NSP_iP_{i+1} with $i \in \{1, \dots, 2n+1\}$. Split them into two groups: the tetrahedra $NSP_{2k+1}P_{2k+2}$ with $k \in \{0, \dots, n\}$ and the tetrahedra $NSP_{2k}P_{2k+1}$ with $k \in \{1, \dots, n\}$. Substitute $\alpha = \beta + \pi$ and proceed with straightforward calculations:

$$\begin{aligned}\Delta_k^{(1)}(\beta) &= \det(S, N, P_{2k+1}, P_{2k+2}) = \cos^2 \frac{L_1 \beta}{4} - U_{2k-1}^2(\cos \theta) \sin^2 \frac{\beta}{2} = \\ &= T_{L_1}^2(\cos \frac{\beta}{4}) - U_{2k-1}^2(\cos \theta) \sin^2 \frac{\beta}{2},\end{aligned}$$

where $k \in \{0, \dots, n\}$, $L_1 = |2n - 4k + 1|$, $\theta = \frac{\pi}{2n+1}$, $\beta \in [-2\theta, 2\theta]$;

$$\begin{aligned}\Delta_k^{(2)}(\beta) &= \det(S, N, P_{2k}, P_{2k+1}) = \cos^2 \frac{L_2 \beta}{4} - U_{2k-2}^2(\cos \theta) \sin^2 \frac{\beta}{2} = \\ &= T_{L_2}^2(\cos \frac{\beta}{4}) - U_{2k-2}^2(\cos \theta) \sin^2 \frac{\beta}{2},\end{aligned}$$

where $k \in \{1, \dots, n\}$, $L_2 = |2n - 4k + 3|$, θ and β the same as above. The first kind Chebyshev polynomial of degree $k \geq 0$ is denoted by T_k . Assume that

$$U_{-1}(\cos \theta) = 0, \quad U_0(\cos \theta) = 1$$

for the sake of brevity.

All the functions $\Delta_k^{(j)}(\beta)$, $j \in \{1, 2\}$ are even on the interval $[-2\theta, 2\theta]$. Then one considers them only on the interval $[0, 2\theta]$. Note, that the polynomial $T_{L_j}^2(\cos \beta)$ monotonously decreases and the function $\sin^2 \frac{\beta}{2}$ monotonously increases with $\beta \in [0, 2\theta]$. Moreover, $T_{L_j}^2(\cos 0) = T_{L_j}^2(1) = 1$. Then it follows that $\Delta_k^{(j)}(\beta) > 0$ with $\beta \in (-2\theta, 2\theta)$. Also, one has $\Delta_k^{(j)}(\pm 2\theta) = 0$. Then for all $\beta \in (-2\theta, 2\theta)$ (i.e. for all α in the affirmation of the Theorem)

$$\det(S, N, P_i, P_{i+1}) > 0$$

where $i \in \{1, \dots, 4n+2\}$, $P_{4n+3} = P_1$. Thus, claim (e) for the polyhedron \mathcal{P}_n is satisfied.

Use the Schläfli formula [16] to obtain the volume formula for $\mathbb{T}_n(\alpha)$. One has

$$d\text{Vol } \mathbb{T}_n(\alpha) = \frac{l_\alpha}{2} d\alpha = \frac{(2n+1)\alpha - (2n-1)\pi}{2} d\alpha.$$

Note, that $\text{Vol } \mathbb{T}_n(\alpha) \rightarrow 0$ with $\alpha \rightarrow \frac{2n-1}{2n+1}\pi$. In this case $d_\lambda(P_i, P_{i+1}) \rightarrow 0$, where $i \in \{1, \dots, 4n+2\}$, $P_{4n+3} = P_1$ and the fundamental polyhedron collapses to a point. Thus

$$\text{Vol } \mathbb{T}_n(\alpha) = \frac{1}{2n+1} \left(\frac{2n+1}{2} \alpha - \frac{2n-1}{2} \pi \right)^2.$$

□

Remark 1 *The domain of the spherical metric existence in Theorem 1 was indicated before in [10, Proposition 2.1].*

4 Torus links \mathbb{L}_n

Let $\mathbb{L}_n, n \geq 2$ be a torus link $t(2n, 2)$ with two components. The corresponding link in the rational census is $2n/1$ (Fig. 3). The fundamental group of \mathbb{L}_n is presented as

$$\pi_1(\mathbb{S}^3 \setminus \mathbb{L}_n) = \langle a, b | (ab)^n = (ba)^n \rangle.$$




Figure 3: Link $2n/1$

Let $\mathbb{L}_n(\alpha, \beta)$ denote a cone-manifold with singular locus the link \mathbb{L}_n and the cone angles α, β along its components.

For every $\alpha, \beta \in (0, 2\pi)$ and $\lambda \in (-1, +1)$, we denote

$$A = \begin{pmatrix} 1 & 0 \\ -2i e^{i\frac{\alpha}{2}} \lambda \sin \frac{\alpha}{2} & e^{i\alpha} \end{pmatrix}$$

and

$$B = \begin{pmatrix} e^{i\beta} & -2i e^{i\frac{\beta}{2}} \lambda \sin \frac{\beta}{2} \\ 0 & 1 \end{pmatrix}.$$

By Lemma 1, linear transformations A and B are isometries of \mathbb{S}_λ^3 .

Lemma 3 *For every integer $n \geq 2$ the following equality holds*

$$(AB)^n - (BA)^n = 4U_{n-1}(\Lambda) \lambda e^{i(\frac{\alpha+\beta}{2} + \pi)n} \sin \frac{\alpha}{2} \sin \frac{\beta}{2} M,$$

where M is a non-zero 2×2 matrix and $U_{n-1}(\Lambda)$ is the second kind Chebyshev polynomial of degree $n - 1$ in variable

$$\Lambda = (1 - \lambda^2) \cos \frac{\alpha - \beta}{2} + \lambda^2 \cos \frac{\alpha + \beta}{2}.$$

Proof. By analogy with Lemma 2. \square

With Lemma 3 the main theorem of the section follows:

Theorem 2 *The cone-manifold $\mathbb{L}_n(\alpha, \beta)$, $n \geq 2$ is spherical if*

$$\begin{aligned} -2\pi \left(1 - \frac{1}{n}\right) &< \alpha - \beta < 2\pi \left(1 - \frac{1}{n}\right), \\ 2\pi \left(1 - \frac{1}{n}\right) &< \alpha + \beta < 2\pi \left(1 + \frac{1}{n}\right). \end{aligned}$$

The lengths l_α, l_β of its singular geodesics (i.e. lengths of the components for \mathbb{L}_n) are equal to each other and

$$l_\alpha = l_\beta = \frac{\alpha + \beta}{2} n - \pi(n - 1).$$

The volume of $\mathbb{L}_n(\alpha, \beta)$ is

$$\text{Vol } \mathbb{L}_n(\alpha, \beta) = \frac{1}{2n} \left(\frac{\alpha + \beta}{2} n - (n - 1)\pi \right)^2.$$

Proof. One continues the proof by analogy with Theorem 1. Suppose that $\mathbb{L}_n(\alpha, \beta)$ is spherical. Then there exists a holonomy mapping [1]:

$$\begin{aligned} h : \pi_1(\mathbb{S}^3 \setminus \mathbb{L}_n) &\longmapsto \text{Isom } \mathbb{S}_\lambda^3, \\ h(a) &= A, \quad h(b) = B. \end{aligned}$$

Also,

$$h((ab)^n) - h((ba)^n) = (AB)^n - (BA)^n = 0.$$

By means of Lemma 3 the equality above holds either if $\lambda = 0$, or if

$$\Lambda = (1 - \lambda^2) \cos \frac{\alpha - \beta}{2} + \lambda^2 \cos \frac{\alpha + \beta}{2}$$

is a root of the equation $U_{n-1}(\Lambda) = 0$.

In case $\lambda = 0$ the image of h is abelian, because of the additional relation $AB = BA$. With $n \geq 2$ this leads to a degenerate geometric structure. Thus, one has to choose the parameter λ for the metric ds_λ^2 using roots of the Chebyshev polynomial $U_{n-1}(\Lambda)$.

The fundamental polyhedron \mathcal{F}_n for the cone-manifold $\mathbb{L}_n(\alpha, \beta)$ is depicted at Fig. 4. Suppose its vertices P_1 and P_2 to be

$$P_1 = (1, 0), \quad P_2 = (0, 1).$$

The axes of isometries A and B correspond to the edges P_1P_{2n+1} and P_2P_{2n+2} . Points N and S are respective middles of the edges P_1P_{2n+1} and P_2P_{2n+2} . Those are called North and South poles of the polyhedron. The polyhedron \mathcal{F}_n is said to be proper if

- (a) respective inner dihedral angles along the edges P_1P_{2n+1} and P_2P_{2n+2} are equal to α and β ;




Figure 4: The fundamental polyhedron \mathcal{F}_n for $\mathbb{L}_n(\alpha, \beta)$

(b) curvilinear faces of the polyhedron are identified by A and B :

$$A : NP_1P_2 \dots P_{2n+1} \rightarrow NP_1P_{4n} \dots P_{2n+2}P_{2n+1},$$

$$B : SP_2P_1P_{4n} \dots P_{2n+2} \rightarrow SP_2P_3 \dots P_{2n+2};$$

(c) sum of the inner dihedral angles ψ_i along the edges P_iP_{i+1} , $i \in \{1, \dots, 4n-1\}$ equals 2π ;

(d) sum of the dihedral angles ϕ_i for tetrahedra NSP_iP_{i+1} , $i \in \{1, \dots, 4n-1\}$ at their common edge NS equals 2π ;

(e) all the tetrahedra NSP_iP_{i+1} with $i \in \{1, \dots, 4n\}$, $P_{4n+1} = P_1$ are non-degenerated and coherently oriented.

In order to choose the parameter λ for the corresponding metric consider the fundamental polyhedron \mathcal{F}_n with $\alpha = \beta = \pi$. Then all its vertices belong to the same circle and all the dihedral angles ψ_i of the tetrahedra NSP_iP_{i+1} along the edges P_iP_{i+1} are equal to $\psi = \frac{\pi}{2n}$ [12]. Also the dihedral angles ϕ_i of the tetrahedra NSP_iP_{i+1} along their common edge NS are equal to each other:

$$\phi_i = \phi = \frac{\pi}{2n}.$$

In this case $\lambda = \langle P_1, P_2 \rangle = \cos \phi$ and

$$\Lambda = -\cos 2\phi = \cos \frac{(n-1)\pi}{n}.$$

All the roots of $U_{n-1}(\Lambda)$ are given by the formula

$$\Lambda_k = \cos \frac{k\pi}{n}, \quad k \in \{1, \dots, n-1\},$$

so one choose the root Λ_k with $k = n-1$. Then, by analogy with Theorem 1, equalities

$$\sum_{i=1}^{4n} \psi_i = 2\pi$$

and

$$\sum_{i=1}^{4n} \phi_i = 2\pi$$

are satisfied at the point $\alpha = \beta = \pi$ of the domain

$$\mathcal{D} = \left\{ (\alpha, \beta) : |\alpha - \beta| < 2\pi \left(1 - \frac{1}{n}\right), |\alpha + \beta - 2\pi| < \frac{2\pi}{n} \right\},$$

depicted at Fig. 5.

In terms of the parameter λ , that defines the metric ds_λ^2 , one has

$$\lambda^2 = \frac{\cos \frac{\alpha-\beta}{2} + \cos \frac{\pi}{n}}{\cos \frac{\alpha-\beta}{2} - \cos \frac{\alpha+\beta}{2}}.$$

As for all $(\alpha, \beta) \in \mathcal{D}$ the inequality $0 < \lambda^2 < 1$ is satisfied, the metric ds_λ^2 is spherical regarding the corresponding domain. By analogy with Theorem 1 one can show that claims (a) – (d) for the polyhedron \mathcal{F}_n are satisfied in the interior of \mathcal{D} .

The lengths l_α and l_β of singular geodesics for the cone-manifold $\mathbb{L}_n(\alpha, \beta)$ meet the relations

$$\begin{aligned} \cos \frac{l_\alpha}{2} &= \langle P_1, N \rangle, \\ \cos \frac{l_\beta}{2} &= \langle P_2, S \rangle. \end{aligned}$$

By analogy with the proof of Theorem 1 one obtains

$$l_\alpha = l_\beta = \frac{\alpha + \beta}{2} n - \pi(n-1).$$

Given the coordinates of the vertices for the fundamental polyhedron verify claim (e) for all (α, β) in the domain \mathcal{D} .

Make use of the Schläfli formula [16] to obtain the volume of $\mathbb{L}_n(\alpha, \beta)$:

$$d \text{Vol } \mathbb{L}_n(\alpha, \beta) = \frac{l_\alpha}{2} d\alpha + \frac{l_\beta}{2} d\beta = \left(\frac{\alpha + \beta}{2} n - \pi(n-1) \right) d \left(\frac{\alpha + \beta}{2} \right).$$

Note, that with

$$\alpha = \beta \rightarrow \pi \frac{n-1}{n}$$

the fundamental polyhedron \mathcal{F}_n collapses to a point (i.e. the volume tends to 0). The last affirmation of the Theorem follows. \square




Figure 5: The domain \mathcal{D} of sphericity for $\mathbb{L}_n(\alpha, \beta)$

Remark 2 Under condition $\alpha = \beta$ the inequality from the affirmation of Theorem 2 coincides with the inequality from [10, Proposition 2.2].

Remark 3 Note, that the lengths of the singular geodesics for $\mathbb{L}_n(\alpha, \beta)$ are equal even if $\alpha \neq \beta$.

References

- [1] Thurston W. P. The geometry and topology of 3-manifolds. Princeton: Lecture Notes, 1977–78. Available on-line.
- [2] Rolfsen D. Knots and links. Berkeley: Publish or Perish Inc., 1976. Google Books.
- [3] Thurston W. P. Hyperbolic geometry and 3-manifolds. Cambridge: Cambridge Univ. Press, 1982. (London Math. Soc. Lect. Note Ser.; 48, 9–25).
- [4] Neumann W. P. Notes on geometry and 3-manifolds, with appendix by Paul Norbury, in: Low Dimensional Topology, Böröczky, Neumann, Stipsicz, Eds. // Bolyai Society Mathematical Studies. 1999. V. 8. P. 191–267. Available on-line.
- [5] Seifert H., Weber C. Die beiden Dodecaederräume // Math. Z. 1933. V. 37. P. 237–253. Available on-line.
- [6] Dunbar W. D. Geometric orbifolds // Rev. Mat. Univ. Complut. Madrid. 1988. V. 1. P. 67–99. Available on-line.

- [7] *Derevnin D., Mednykh A., Mulazzani M.* Geometry of trefoil cone-manifold // Preprint. 2007.
- [8] *Hilden H. M., Lozano M. T., Montesinos-Amilibia J. M.* On a remarkable polyhedron geometrizing the figure eight cone manifolds // J. Math. Sci. Univ. Tokyo. 1995. V. 2. P. 501–561. Available on-line.
- [9] *Mednykh A., Rasskazov A.* Volumes and degeneration of cone-structures on the figure-eight knot // Tokyo J. of Math. 2006. V. 29. N. 2. P. 445–464.
- [10] *Porti J.* Spherical cone structures on 2-bridge knots and links // Kobe J. of Math. 2004. V. 21. N. 1. P. 61–70. Available on-line.
- [11] *Molnár E.* The projective interpretation of the eight 3-dimensional homogeneous geometries // Beiträge zur Algebra und Geometrie. 1997. V. 38. N. 2. P. 261–288. Available on-line.
- [12] *Mednykh A., Rasskazov A.* On the structure of the canonical fundamental set for the 2-bridge link orbifolds // Preprint. 1996. Available on-line.
- [13] *Minkus J.* The branched cyclic coverings of 2-bridge knots and links // Mem. Amer. Math. Soc. 1982. V. 35. N. 255. Google Books.
- [14] *Ratcliffe J.* Foundations of hyperbolic manifolds. New York: Springer-Verlag, 1994. (Graduate Texts in Math.; 149). Google Books.
- [15] *Hodgson C., Rubinstein J. H.* Involutions and isotopies of lens spaces, Knot theory and manifolds (Vancouver, B.C., 1983). Berlin: Springer-Verlag, 1985. (Lecture Notes in Math.; 1144, 60–96).
- [16] *Hodgson C.* Degeneration and regeneration of hyperbolic structures on three-manifolds // Princeton: Thesis, 1986.

Alexander Kolpakov
Novosibirsk State University
630090, Pirogova str., bld. 2
Novosibirsk, Russia
 kolpakov.alexander@gmail.com

Alexander Mednykh
Sobolev Institute of Mathematics, SB RAS
630090, Koptyug avenue, bld. 4,
Novosibirsk, Russia
 mednykh@math.nsc.ru