



HAL
open science

REVIEW ARTICLE: ENDPOINTS USED IN FUNCTIONAL DYSPEPSIA DRUG THERAPY TRIALS

Daphne Ang, Nicholas J Talley, Magnus Simrén, Pieter Janssen, Guy E
Boeckxstaens, Jan Tack

► **To cite this version:**

Daphne Ang, Nicholas J Talley, Magnus Simrén, Pieter Janssen, Guy E Boeckxstaens, et al.. REVIEW ARTICLE: ENDPOINTS USED IN FUNCTIONAL DYSPEPSIA DRUG THERAPY TRIALS. *Alimentary Pharmacology and Therapeutics*, 2011, 33 (6), pp.634. 10.1111/j.1365-2036.2010.04566.x . hal-00608029

HAL Id: hal-00608029

<https://hal.science/hal-00608029>

Submitted on 12 Jul 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**REVIEW ARTICLE: ENDPOINTS USED IN FUNCTIONAL
DYSPEPSIA DRUG THERAPY TRIALS**

Journal:	<i>Alimentary Pharmacology & Therapeutics</i>
Manuscript ID:	APT-0544-2010.R1
Wiley - Manuscript type:	Review Article
Date Submitted by the Author:	11-Dec-2010
Complete List of Authors:	Ang, Daphne; Changi General Hospital, Gastroenterology Talley, Nicholas; University of Newcastle, Faculty of Health Simrén, Magnus; Section of Gastroenterology and Hepatology, Dept of Internal Medicine Janssen, Pieter; KULeuven Boeckxstaens, Guy; University of Leuven, Pathophysiology Tack, Jan; University Hospital, Center for Gastroenterological Research
Keywords:	Functional dyspepsia < Disease-based, Functional GI diseases < Disease-based, Stomach and duodenum < Organ-based, Symptom score or index < Topics

1
2
3 **REVIEW ARTICLE: ENDPOINTS USED IN FUNCTIONAL DYSPEPSIA**
4
5 **DRUG THERAPY TRIALS**
6
7

8
9
10 **D. Ang¹, N.J. Talley², M. Simren³, P. Janssen¹, G. Boeckxstaens¹, J.**
11 **Tack¹**
12
13

14
15
16
17 ¹Center for Gastroenterological Research, K.U.Leuven, Belgium.

18 ²Department of Internal Medicine, Mayo Clinic Jacksonville

19 ³Department of Internal Medicine, Sahlgrenska University Hospital, Göteborg,
20 Sweden,
21
22

23
24
25 Correspondence to: Jan Tack, M.D., Ph.D.

26 Department of Pathophysiology,

27 Division of Gastroenterology

28 University Hospital Gasthuisberg

29 Herestraat 49, B-3000 Leuven, Belgium.

30 Phone: +32 16 345751

31 Fax: + 32 16 345939

32 E-mail: Jan.Tack@med.kuleuven.ac.be.
33
34

35 Key words: Functional dyspepsia, endpoints, clinical trials, patient reported
36 outcomes, quality of life, questionnaires
37
38

39
40 Conflict of interest: none declared
41
42

43
44 Concept: J. Tack, NJ Talley
45

46
47 Literature source selection: all authors
48

49
50 First draft: D. Ang
51

52
53 Corrections and text finalisation, all authors.
54
55
56
57
58
59
60

ABSTRACT

Background: How to measure patient reported outcomes (PRO) in treatment trials for functional gastrointestinal disorders is a matter of controversy. This review focuses on instruments and endpoints that have been used to evaluate the efficacy of therapeutic agents in functional dyspepsia (FD) trials, also considering the newly defined Rome III FD criteria. Methods: A Medline search was conducted to identify relevant studies pertaining to FD treatment, with particular emphasis on the studies to date which have used validated outcome measures. Results: Currently available outcome measures are heterogeneous across studies. They include global binary endpoints, analog or categorical scoring scales, uni- or multi-dimensional disease specific questionnaires, global outcome evaluations, and quality of life questionnaires. Across the available outcome measures, substantial heterogeneity is found, not only in the type of endpoint measure, but also in the number and types of symptoms that are considered to be part of the FD symptom complex. Especially based on content validity, none of the existing questionnaires or endpoints can be considered sufficiently validated to be recommended unequivocally as the primary outcome measure for FD trials according to the Rome III criteria. On the other hand, existing well-validated multi-dimensional questionnaires that include many non-FD symptoms can be narrowed down to evaluate only the cardinal symptoms according to Rome III. In conclusion, there is an urgent need to develop Rome III-based PROs for functional dyspepsia. Well-validated multi-dimensional questionnaires may serve as a

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

guidance for this purpose, and could also be considered for use in ongoing clinical trials.

For Peer Review

INTRODUCTION

Functional dyspepsia (FD), is one of the most prevalent functional gastrointestinal disorders (FGIDs). Over the last 20 years, the definition of FD has undergone major changes from the 1988 working party¹ to the consecutive Rome consensus documents²⁻⁴, in line with changing understanding of the pathophysiological basis of this disorder. Based on Rome III criteria⁴, FD is defined as the presence of symptoms thought to originate in the gastroduodenal region (early satiation, postprandial fullness, epigastric pain or burning), in the absence of any organic, systemic or metabolic disease that is likely to explain the symptoms. FD is further subdivided into 2 diagnostic categories of meal induced dyspeptic symptoms (postprandial distress syndrome [PDS], characterized by postprandial fullness and early satiation) and epigastric pain syndrome ([EPS], characterized by epigastric pain and burning). Whereas the previous Rome II definition³ for FD excluded patients with predominant heartburn and was unclear on non-predominant heartburn, the Rome III definition states that heartburn is not a gastroduodenal symptom, although it often occurs simultaneously with FD symptoms.⁴ Similarly, although there is frequent overlap with the irritable bowel syndrome, the symptom pattern of both entities is distinct.

FD is one of the most important categories of the FGID, in view of its prevalence and impact on the general population.⁵⁻⁸ At present, there is no treatment with established efficacy for FD.¹ The Food and Drug Administration (FDA) guidance released in February 2006 provides recommendations for the

1
2
3 use of validated instruments to assess treatment outcomes, and describes the
4 proper development and psychometric validation of patient reported outcomes
5 (PRO) before endorsing a clinical product.⁹ The Rome III committee¹⁰ has
6 also provided guidelines for clinical trial design in FGIDs, with a similar
7 emphasis on individual patient assessment and the use of validated outcome
8 measures. The aim of the present study is to evaluate the currently available
9 endpoints for FD drug development, in line with the Rome III definitions, the
10 published FDA guidelines and Rome III clinical trials recommendations.^{4,9,10}
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

MATERIALS AND METHODS

Identification of relevant studies

To identify relevant studies, both computerized (Medline) and manual searches were performed, using the cited references of the retrieved articles. MeSH and free-text terms for FD therapies were combined with the terms dyspepsia, clinical trials, symptom assessment, questionnaires, patient reported measures and randomized for searches conducted for the time period between January 1979 and December 2008. FD therapeutic trials were retrieved and studied, with emphasis on the method used for evaluating outcome measures. Article reference lists were examined for relevant papers. Analysis of both full papers and abstracts was conducted, with particular emphasis on outcome measures. The literature search was performed independently by 2 of the authors, and their retrievals were merged.

Inclusion criteria

The following criteria were used to select trials for analysis: (i) randomized double blind controlled trials (RCT), (ii) parallel or single cross-over trial designs, (iii) adult patients with FD, (iv) baseline gastroscopy to exclude structural pathology; (v) comparison of therapy vs. active or placebo control; (vi) clear description of the method of assessing outcome measures for FD symptoms, and (vi) articles in English.

Outcome measures

One of the main difficulties encountered with therapeutic trials in FGIDs, and FD in particular where efficacy has not been established for any treatment, is the lack of objectively measurable outcome measures.¹¹⁻¹⁴ In FD, symptom patterns do not correlate well with putative pathophysiological mechanisms such as gastric accommodation, sensitivity to distension and gastric emptying rates.¹⁵ In the absence of quantifiable surrogate markers for symptom improvement, evaluation of treatment response has to rely on patient's reporting of symptom intensities.

Several types of outcome measures have been used in FD clinical trials. These can be broadly classified into global outcomes, generic instruments and disease specific instruments.¹⁶ Global outcomes are measured either by a dichotomous binary type response (e.g. yes/no to symptom improvement) or a scoring method which can either be a Likert scale (categorical) or a visual analogue scale (VAS). Likert scales allow easier interpretation for the physician compared to the VAS, with 5-point or 7-point scales providing greater sensitivity than 4-point scales.¹⁷ The VAS and 7-point Likert scales exhibit comparable responsiveness, although the ease of administration and interpretation of the 7-point scale recommend its use in clinical trials.¹⁸ The Likert scale is often incorporated into global scales, such as the overall treatment effect (OTE),¹⁹ generic instruments and disease specific instruments. The OTE uses a graded scale to assess overall symptom improvement or deterioration. The subjective nature of this approach allows the individual to integrate all aspects of his condition into a single treatment

1
2
3 outcome, and is particularly suitable to show deterioration. Unlike generic
4 instruments, disease specific instruments focus on a particular medical
5 condition and are likely to be more useful in quantifying changes in quality of
6 life in FD therapeutic trials.²⁰ Disease specific instruments in FD can be
7 unidimensional (evaluating gastrointestinal symptoms) or multidimensional
8 (evaluating both gastrointestinal symptoms and other domains such as
9 emotional or social functioning and impact of symptoms on daily activities).
10
11
12
13
14
15
16
17
18
19
20
21

22 **Validity of outcome measures**

23
24
25
26
27 In 1999, the Rome II Working Group emphasised individual patient
28 assessment as a primary outcome, allowing for the integration of multiple
29 symptoms into a single global endpoint for FGID studies.²¹ An alternative
30 method was the use of a disease specific questionnaire to evaluate relevant
31 aspects of the patient's symptoms and disease related quality of life. The
32 Rome III guidelines and the FDA February 2006 guidance provide further
33 recommendations on the proper development and validation of PRO
34 measures.^{9,10} Psychometric validation of a symptom based measure
35 incorporates several components, including evidence that the instrument
36 addresses all the patient's symptoms that are indicative of the disorder
37 (content validity); is related to other measures of the same or similar concepts
38 such as symptom improvement and symptom free days (construct validity);
39 produces similar results when re-administered to patients whose health status
40 has not changed (reliability); detects clinically meaningful change in health
41 status when a change has occurred (longitudinal construct validity or
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 responsiveness); and is associated with changes in score that can be related
4
5 to clinical indicators that are meaningful to clinicians (predictive or criterion
6
7 validity). Responsiveness is an important component of validity²² and forms a
8
9 major criterion for selecting an optimal measure of outcome for a randomized
10
11 controlled trial.²³ Patient involvement in the development of PRO
12
13 measurement is emphasised by the FDA guidelines, and this can be aided by
14
15 structured interview sessions, focus groups and quality research methods.
16
17
18 The outcome measure should have an effective measurement range²⁴,
19
20
21 allowing the instrument to detect changes in outcomes during the course of
22
23
24 the clinical trial without the limitations of the ceiling or floor effects.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

Description of studies

Amongst the therapeutic trials for FD, 117 studies were initially obtained. After excluding 31 studies which did not satisfy the inclusion criteria, 86 studies were selected for analysis of the outcome measures. The outcome measures that were employed in these studies can be classified broadly under the following categories: (a) binary outcome measures (adequate, satisfactory or sufficient relief), (b) individual symptom assessment with Likert or VAS scales, (c) disease specific questionnaires, (d) global outcome evaluations, and (e) quality of life questionnaires. Table 1 summarizes the types of endpoints used in the FD studies that were selected for the analysis.

A. Adequate relief or satisfactory relief (Binary Outcome) as a primary endpoint

Evaluating the relief of FD symptoms using a binary outcome approach (yes/no response) was first employed in a study which was designed to compare reflux episodes in FD patients who responded to omeprazole with non-responders.²⁵ Responders were defined as patients who considered themselves to have achieved sufficient relief of symptoms at the end of study period. A significantly higher number of reflux episodes on 24 hour oesophageal pH measurement were detected in omeprazole responders. Based on extensive experience from irritable bowel syndrome (IBS) studies,²⁶⁻

1
2
3 ³⁷ adequate relief and satisfactory relief have been the most commonly used
4
5
6 outcome measures in treatment trials for FGIDs, although the Rome III
7
8 committee on design of treatment trials indicated that further validation may
9
10 be desirable, and that alternative outcome measures such as integrative
11
12 symptom questionnaires are also attractive.¹⁰
13
14
15
16

17
18 The endpoint of adequate relief is responsive, reproducible and demonstrates
19
20 good construct validity,^{38,39} and has been used in both IBS²⁶⁻³³ and FD⁴⁰⁻⁴²
21
22 clinical trials. It allows the patient to integrate all relevant symptoms and
23
24 normalises the assessment to the patient's own internal reference system.^{38,39}
25
26

27
28 On the other hand, the endpoint does not reflect the magnitude of
29
30 improvement needed to reach the endpoint (which depends in part on the
31
32 baseline severity level) and does not detect worsening of symptoms³⁹.
33

34
35 Adequate relief was used as a primary endpoint in a phase 2b RCT of
36
37 alosetron for FD⁴⁰. During the 12 weeks of treatment with alosetron or
38
39 placebo, patients responded yes or no to the weekly question: "In the past 7
40
41 days, have you had adequate relief of your upper abdominal pain or
42
43 discomfort?" A 1 mg dose of alosetron showed benefit over placebo and 0.5
44
45 or 2 mg doses for the 12-week average rate of adequate relief. When
46
47 stratified by gender, alosetron 1.0 mg showed significant benefit in females
48
49 (p=0.03) in achieving the primary endpoint of adequate relief. Secondary
50
51 efficacy endpoints were patients' daily self rating of symptom frequency
52
53 (absence or presence of early satiety, bloating, nausea, belching, pain and
54
55 postprandial fullness) and severity (on a 5-point Likert scale). When the
56
57 primary global outcome of adequate relief was compared with the secondary
58
59
60

1
2
3 outcome of individual symptoms, patients with adequate relief of upper
4 abdominal pain or discomfort also had significantly greater reductions in
5 severity of pain, nausea and bloating, and % of days with pain, early satiety,
6 bloating and nausea compared to patients who did not have adequate relief
7 (all $p < 0.001$). Thus, the adequate relief endpoint showed good correlation
8 when individual symptoms were analysed, but no further validation in FD has
9 occurred. The adequate relief endpoint was also used in a phase 2 U.S. study
10 with Acotiamide, a muscarinic receptor antagonist with fundus-relaxing
11 properties.⁴¹ During the first 4 weeks of treatment, the number of subjects
12 reporting adequate relief for more than 50% of time was significantly higher
13 with acotiamide 300 mg t.i.d. compared to placebo.⁴² This was associated with
14 significant improvement of symptoms of bloating, postprandial nausea, and
15 stomach pain before meals, and with significant improvement of several
16 domains of the Short Form-36 quality of life scale and of the Short form
17 Nepean Dyspepsia Index.⁴²

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41 The satisfactory relief endpoint has a construct that is similar to adequate
42 relief, although less extensive validation data are available in the literature for
43 the former.³⁹ Using satisfactory relief as an endpoint, the beneficial effects of
44 tegaserod in IBS have been demonstrated.³⁴⁻³⁷ The binary satisfactory relief
45 endpoint was also used in two phase 3 studies with tegaserod in dysmotility-
46 type FD, and demonstrated significant efficacy in one of these.⁴³ In the trial
47 that achieved a significantly higher proportion of days with satisfactory relief
48 from tegaserod, patients also had significant improvement of overall and
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 individual symptom severity measured on a Likert scale, and a significantly
4
5 better global assessment of change in dyspepsia symptoms.
6
7
8
9

10 **B. Categorical (Likert) Scales**

11
12
13
14
15 Likert scales have been widely used both in assessing individual and global
16
17 symptom severity as well as symptom frequency in numerous FD trials.⁴⁴⁻⁹⁷
18
19 Veldhuyzen and colleagues⁹³ used a 5-point Likert scale (0=no problem,
20
21 4=very severe problem) to score the severity of 8 gastrointestinal symptoms
22
23 (epigastric pain, burping, heartburn, bloating, flatulence, sour taste, nausea
24
25 and halitosis). A validation study was performed in non ulcer dyspepsia (NUD)
26
27 and *Helicobacter pylori* associated gastritis (HPAG). Its reproducibility was
28
29 demonstrated by comparing symptom scores in the 2 groups of patients
30
31 before treatment, and responsiveness was demonstrated in both NUD and
32
33 HPAG patients, where cumulative scores were significantly reduced after
34
35 therapy. However, the validity of this scale as a measure of improvement was
36
37 only assessed by comparing changes in symptom scores with changes in the
38
39 patients' overall health status (deteriorated, stayed the same or improved
40
41 during the trial) rather than against an established questionnaire. In addition, it
42
43 combined FD symptoms with symptoms suggestive of GERD, nausea and
44
45 intestinal dysfunction. Finally, the cumulative symptom score attributed equal
46
47 weight to all symptoms, although some of them may be related. Nevertheless,
48
49 the results from this study support the applicability of a Likert-scale in
50
51 measuring the severity of FD symptoms.
52
53
54
55
56
57
58
59
60

1
2
3 Likert scores have been employed in numerous studies to date (Table 1),
4 including the BOND and OPERA studies⁸⁸ which utilised a 4-point Likert scale
5 to evaluate the efficacy of omeprazole in FD. A 7-point Likert scale was used
6 in 2 very similar multicentre, multinational, RCTs: the OCAY⁸⁹ and ORCHID
7 studies,⁹⁰ which compared the efficacy of *Helicobacter pylori* eradication
8 therapy with that of either omeprazole (OCAY study) or placebo (ORCHID
9 study) in relieving dyspeptic symptoms 12 months after eradication therapy.
10 Using a stringent definition of treatment success (absence of symptoms or the
11 presence of only minimal symptoms (Likert score 1 or 2) during the seven
12 days preceding the final visit) no significant symptomatic benefit of
13 *Helicobacter pylori* eradication therapy in FD was found.^{89,90}
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 In addition, Likert scales have been used for global evaluation. Using a Likert-
37 type scale with 5 grades (symptom-free, markedly improved, moderately
38 improved, not changed, and deteriorated) to measure patients' global
39 assessment of efficacy, a statistically significant benefit was found in a Phase
40 IIb trial for itopride over placebo in the number of patients who reported being
41 symptom-free or with marked improvement.⁹¹ However, this effect was not
42 seen when the same global patient assessment of efficacy was evaluated in 2
43 phase III trials.⁹²
44
45
46
47
48
49
50
51
52
53
54
55
56

57 Another method of assessing global outcome is the "overall treatment effect"
58 (OTE) approach.¹⁸ At intervals during or at the completion of treatment, the
59
60

1
2
3 patient is asked to decide whether symptoms have remained the same,
4
5 improved or deteriorated compared to pre-treatment phase, by means of a
6
7 Likert scale. The advantage of this endpoint is that it closely resembles the
8
9 way physicians evaluate treatment benefit in clinical practice, but the main
10
11 disadvantage is the inherent recall of pre-treatment symptom severity which
12
13 may lead to bias. OTE was used as a secondary outcome measure in the
14
15 OCAY study, and in a number of studies of acid suppression in FD.^{89,94-97}
16
17 Overall treatment evaluation was used most recently in the tegaserod FD
18
19 studies, where a weekly global assessment of change was rated on a 7-point
20
21 Likert scale, and this endpoint generated the most consistent improvements
22
23 with tegaserod over placebo.⁴³ OTE was also used in the U.S. and Japanese
24
25 Acotiamide trials as a secondary or primary endpoint respectively.^{42,98} In the
26
27 U.S. study, benefit using the OTE paralleled the adequate relief outcome
28
29 result, with significance during the first 4 to 6 weeks.⁴² In the Japanese study,
30
31 significant benefit was found using the OTE for the 100 mg dose in meal-
32
33 related FD symptoms, and this was associated with a higher rate of
34
35 disappearance of postprandial fullness and upper abdominal bloating.⁹⁸ VAS
36
37 scores, which may be more difficult to interpret in terms of magnitude of
38
39 response, have been used in only a few, mainly less recent studies.⁹⁹⁻¹⁰⁴
40
41
42
43
44
45
46
47
48
49

50 **C. Disease specific questionnaires (Unidimensional)**

51
52
53
54

55 Integrative symptom questionnaires incorporate the frequency and/or severity
56
57 of one or a group of symptoms pertaining to the FGID in question both at
58
59 baseline and at the end of therapy. A review of the questionnaires that have
60

1
2
3 been used as outcome measures in FD drug therapy trials is pertinent, as
4 they serve an important tool in addressing the effectiveness of pharmaceutical
5 agents. We highlight several instruments that have been developed for
6 assessing outcome measures in FD. Where appropriate, FD trials that have
7 utilised the respective outcome measures are described.
8
9
10
11
12
13
14
15
16

17 C. 1. Global Overall Symptom scale (GOS)

18
19
20
21
22 The Global Overall Symptoms (GOS) scale is a validated outcome measure
23 for dyspepsia treatment trials,¹⁰⁵ adapted from the previously validated 5-point
24 scale.⁹³ Using a 7-point Likert scale, patients are asked to grade the overall
25 severity of 10 upper gastrointestinal symptoms (epigastric pain; epigastric
26 discomfort; heartburn; acid regurgitation; upper abdominal bloating; excessive
27 belching; nausea; early satiety; postprandial fullness and persistent fullness
28 after a meal) [specific symptom subtypes (SSS)] over a certain retrospective
29 period of time, either 28 days (GOS-28) or 2 days (GOS-2). Validation of the
30 GOS was tested within the CADET-HN study⁹⁴ and the Confirmatory Acid
31 Suppression Test (CAST) study.¹⁰⁶ The GOS showed construct validity,
32 reliability and responsiveness.¹⁰⁵ Construct validity was determined using
33 Spearman correlation coefficients, by correlating changes in the GOS to
34 changes in severity of individual and mean symptom scores; Quality of Life in
35 Reflux and Dyspepsia (QoLRAD)¹⁰⁷ overall score and dimensions the
36 Gastrointestinal Symptom Rating Scale (GSRS, see below)¹⁰⁸ overall score
37 and dimensions, and the Reflux Disease Questionnaire (RDQ)¹⁰⁹ overall
38 score and dimensions. Moderate to high Spearman correlation coefficient
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 values (0.41-0.80) were achieved for each of the above mentioned outcome
4
5 measures, demonstrating good construct validity. Test-retest reliability,
6
7 assessed by Intraclass Correlation Coefficient (ICC) was 0.62 for GOS-2 and
8
9 0.42 for GOS-28. In addition, there was a positive correlation between change
10
11 in GOS and change in symptom severity. To date, the GOS has been used as
12
13 a primary outcome measure in the CADET (Canadian Adult Dyspepsia
14
15 Empiric Treatment) study programmes as well as in the ENTER trial.⁹⁵⁻⁹⁷ In
16
17 this multicentre placebo-controlled trial carried out across Canada, patients
18
19 with FD of moderate severity (defined as a GOS score ≥ 4) were randomized
20
21 to receive esomeprazole or placebo once daily for 8 weeks. For the primary
22
23 outcome measure of symptom relief (GOS score ≤ 2 at 8 weeks), no
24
25 statistically significant difference was found between both groups. However,
26
27 the GOS combines FD symptoms with symptoms suggestive of GERD or IBS,
28
29 and the mean score attributes equal weight to all contributing symptoms,
30
31 which might be considered to be a drawback.
32
33
34
35
36
37
38
39
40

41 C.2. Gastrointestinal Symptom Rating Score (GSRS)

42
43
44
45
46 The GSRS¹⁰⁸ was developed in the early 1980s as an outcome measure for
47
48 peptic ulcer disease and IBS. It comprises 15 items incorporating 5 symptom
49
50 clusters (gastroesophageal reflux, abdominal pain, indigestion, diarrhoea and
51
52 constipation) over the prior 2 weeks. Although it was originally designed to be
53
54 interview based with a 4-point adjectival scale, it was subsequently modified
55
56 to be self administered with a 7-point Likert scale. The GSRS has been well
57
58 validated and shown to be responsive.¹⁰⁹⁻¹¹¹ A shortcoming of the GSRS is
59
60

1
2
3 that it is not specific for dyspepsia. In addition, it measures only the severity of
4 GI related symptoms and not their impact on quality of life. The GSRS has
5 been used as a primary outcome measure in a study evaluating the efficacy of
6 rebamipide¹¹² and as a secondary outcome measure in various studies.<sup>88,94-
7
8
9
10
11
12 97,113,114</sup>

13 14 15 16 17 C.3. Leeds Dyspepsia Questionnaire (LDQ)

18
19
20
21
22 The LDQ¹¹⁵ is administered by an investigator during an interview, and
23 evaluates the frequency and severity of 8 symptoms (epigastric pain,
24 retrosternal pain, regurgitation, nausea, vomiting, belching, early satiety and
25 dysphagia) The scoring system uses the frequency of the first 5 symptoms to
26 determine the presence of dyspepsia, whilst all 8 symptoms are used to
27 access severity of dyspepsia on a scale of 0 (absent) to 5 (most severe). The
28 severity of dyspepsia is assessed as a summary score (range 0-40). The LDQ
29 was shown to be reliable, valid and responsive to change in both primary and
30 secondary care populations in the U.K. Several potential drawbacks of the
31 LDQ include the administration of the questionnaire by a researcher, inclusion
32 of GERD and nausea symptoms, and a complex scoring system.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50
51 The psychometric properties of the LDQ were evaluated in 99 primary care
52 patients and 215 hospital referral patients. Physician diagnosis was the gold
53 standard used to validate the ability of the LDQ to detect dyspepsia. The
54 sensitivity and specificity of the LDQ at the primary care setting were 80% and
55 79%, respectively. In the hospital setting, the LDQ showed a sensitivity of
56
57
58
59
60

1
2
3 99% whilst the specificity was 53%. Based on κ statistics, there was moderate
4
5 to substantial agreement between the LDQ and physician assessment. Test-
6
7 retest reliability was evaluated in 107 patients by a research nurse during two
8
9 clinic visits within 4-7 days, yielding a κ statistic of 0.83 and internal
10
11 consistency by Cronbach's α of 0.68. A similar high κ statistic of 0.90 was
12
13 achieved when two different researchers administered the LDQ to 42 patients
14
15 within 30 minutes. Responsiveness to change was also detected, with the
16
17 median LDQ score decreasing from 22.5 to 4.5 one month after receiving
18
19 appropriate therapy. However, this was only assessed in 12 patients with
20
21 relatively severe symptoms. The LDQ has been used in phase 2 and 3
22
23 studies of itopride.^{91,92} The short-Form Leeds Dyspepsia Questionnaire (SF-
24
25 LDQ)¹¹⁶ was developed to be self completed and a shorter version of the
26
27 original LDQ. The SF-LDQ assesses the frequency and severity over the
28
29 preceding 2 months of 4 symptoms (indigestion, heartburn, regurgitation and
30
31 nausea). In the validation study involving both primary and secondary care
32
33 patients, Cronbach's α coefficient was 0.90, demonstrating a high level of
34
35 internal consistency. Pearson's correlation coefficient for test-retest reliability
36
37 2 days apart was 0.93. Validity was demonstrated by comparison with general
38
39 practitioners' diagnosis. Comparison of the summed total scores for patients
40
41 in the primary care and secondary care setting yielded statistically significant
42
43 differences, further confirmed the validity of the SF-LDQ. The SF-LDQ's
44
45 responsiveness to change was significant in 37 patients who received
46
47 treatment of known effectiveness.
48
49
50
51
52
53
54
55
56
57
58
59
60

C.4. Dyspepsia Symptom Severity Index (DSSI)

1
2
3
4
5
6 The DSSI¹¹⁷ is a 20-item self administered questionnaire divided into 3
7
8 subscales (reflux-like, ulcer-like and dysmotility-like symptoms), quantifying
9
10 the severity of dyspepsia symptoms over the past 2 weeks. The items were
11
12 derived from patient focus group interviews. One global item is included at the
13
14 conclusion of the questionnaire to assess the patient's overall impression of
15
16 dyspepsia severity. The severity questions are graded on a 5-point Likert
17
18 scale ranging from 0 (absent) to 4 (very severe). Subscale internal
19
20 consistency levels were high (Cronbach α 0.84-0.89) and reproducible (ICC
21
22 0.90-0.92). The DSSI correlated well with the patients' symptom diary and
23
24 discriminated between patients and age-matched controls. However, despite
25
26 its validity and reliability, its responsiveness was not assessed. Inclusion of
27
28 GERD and nausea symptoms is another drawback. The DSSI was used in the
29
30 Acotiamide U.S. Phase 2 program as a secondary endpoint.⁴² Significant
31
32 improvements in symptoms of bloating, postprandial nausea, and stomach
33
34 pain before meals with the 300 mg dose t.i.d., were associated with
35
36 significantly higher response rates on the adequate relief and OTE endpoints,
37
38 and with significant improvement of several domains of the Short Form-36
39
40 quality of life scale and of the Short form Nepean Dyspepsia Index during the
41
42 first weeks of the study.⁴²
43
44
45
46
47
48
49
50
51
52

53 C.5. Other uni-dimensional dyspepsia symptom questionnaires.

54
55
56

57
58 The only dyspepsia questionnaire to be validated in a Chinese population, the
59
60 Hong Kong Dyspepsia Index¹¹⁸ originally contained 24 gastrointestinal

1
2
3 symptoms that were self administered, but only the 12 most discriminating
4
5 symptoms (epigastric pain, upper abdominal bloating, upper abdominal dull
6
7 ache, epigastric pain before meals, epigastric pain when anxious, vomiting,
8
9 nausea, belching, acid regurgitation, heartburn, feeling of acidity in the
10
11 stomach and loss of appetite) were selected. Symptoms are graded on a 5
12
13 point Likert scale, ranging from 1(no symptoms) to 5 (incapacitating
14
15 symptoms resulting in an inability to perform daily activities and/or requiring
16
17 days off work). A cut-off score of ≥ 16 was shown to have a sensitivity of 0.82
18
19 and specificity of 0.83 for identification of dyspeptic symptoms. There was
20
21 good test-retest reliability and internal consistency, with an intraclass
22
23 correlation coefficient of 0.89 and Cronbach's alpha coefficient of 0.90.
24
25 Dyspepsia scores correlated negatively with all aspects of the SF-36 quality of
26
27 life scale with the exception of physical functioning, lending support for its
28
29 construct validity. It was also able to discriminate between patients who
30
31 reported a subjective improvement in symptoms and those who reported no
32
33 change or worsening. Although it assesses disease severity, this
34
35 questionnaire does not measure frequency of dyspeptic symptoms, excludes
36
37 some key meal-related symptoms (e.g. early satiety) and includes a number
38
39 of GERD symptoms. This scoring system has been used in a study evaluating
40
41 the efficacy of 4 weeks of lansoprazole treatment in patients who fulfilled
42
43 Rome II criteria for FD with a baseline score ≥ 16 .¹¹⁹
44
45
46
47
48
49
50
51
52
53
54

55 The Porto Alegre Dyspeptic Symptoms Questionnaire (PADYQ)¹²⁰ is an
56
57 investigator administered unidimensional 11-item instrument which evaluates
58
59 5 symptoms of NUD as defined by the Rome I consensus. The symptoms of
60

1
2
3 upper abdominal pain, nausea and upper abdominal bloating are accessed for
4 intensity, duration and frequency; whilst vomiting and early satiety are
5 accessed for frequency. Internal consistency of this questionnaire, as
6 measured by Cronbach's alpha was 0.82. It was shown to be reproducible
7 when submitted to the test-retest procedure, both by the same interviewer
8 (ICC 0.86) and by different interviewers (ICC 0.87). Its responsiveness was
9 demonstrated by a statistically significant reduction in the mean score
10 obtained after treatment compared with baseline. In addition, the PADYQ
11 demonstrated content, discriminant and criterion validity. However, the
12 questionnaire focuses on the pain spectrum of FD, and does not address
13 symptoms such as postprandial fullness and epigastric burning.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

D. Disease specific instruments (multidimensional)

D.1. Nepean Dyspepsia Index (NDI)

The NDI^{121,122} was developed primarily as a disease specific quality of life measure. From an initial 42-items to measure the impact of dyspepsia on a patient's overall quality of life, it was shortened to 25-items centred around 4 subscales: (i) interference with activities of daily living (13 items); (ii) lack of control over the illness (7 items); (iii) disturbance of eating or drinking (3 items) and (iv) sleep disturbance (2 items). A separate symptom checklist measures the frequency, severity and bothersomeness of 15 common upper gastrointestinal symptoms over a 2-week period. An expert panel documented high internal consistency for the quality of life part of the NDI (Cronbach α 0.81-0.96) as well as good discriminative and convergent validity. The NDI was used as a secondary outcome measure in phase 2 and 3 itopride studies, in tegaserod phase 2 and 3 studies and in the Acotiamide U.S. Phase 2.^{42,43,91,92,123} Test retest reliability was evaluated in a multi-center study in the United States.¹²⁴ The NDI has been translated and validated in several languages, including Arabic.¹²⁵ The NDI was further shortened to 10 quality-of-life items in 5 domains¹²⁶. This short form NDI was validated in a large multicentre trial in Europe and the United States. The short form NDI demonstrated adequate internal consistency (Cronbach α 0.71-0.76), correlated with the original long version and was highly responsive. To date,

1
2
3 one study¹²⁷ employed both the short and long versions of the NDI, but only
4
5 the results of the long version were presented.
6
7
8
9

10 D.2. Severity of Dyspepsia Assessment (SODA)

11
12
13

14
15 The SODA¹²⁸ is a multidimensional outcome measure for dyspepsia related
16
17 health that consists of 17 questions incorporating 3 dimensions of pain
18
19 intensity (6 items); 7 non-pain symptoms (belching, heartburn, bloating,
20
21 passing gas, sour taste, nausea and bad breath) and satisfaction scales (4
22
23 items). In this self administered questionnaire, patients are asked to assess
24
25 the preceding 7 days. Reliability was good (Cronbach's α were 0.97, 0.90 and
26
27 0.92 for pain intensity, non-pain symptoms and satisfaction scales
28
29 respectively) and mean change scores discriminated between patients who
30
31 reported improvement compared with those who were unchanged. The
32
33 psychometric properties of SODA were further evaluated in a large study of
34
35 arthritis patients receiving NSAIDS or celecoxib, confirming good internal
36
37 consistency.¹²⁹
38
39
40
41
42

43 Responsiveness as evaluated by receiver operating characteristic (ROC)
44
45 curves was high for pain intensity scale (area under the curve (AUC)=0.78);
46
47 non-pain symptoms (AUC =0.74) and satisfaction scales (AUC=0.75).
48
49 However, reproducibility was low, with ICC values of 0.49, 0.61 and 0.45 for
50
51 pain intensity, non-pain symptoms and satisfaction scales respectively.
52
53 Notably, SODA assesses the severity but not the frequency of symptoms, and
54
55 it includes a broad range of symptoms that are also suggestive for GERD or
56
57
58
59
60

1
2
3 IBS. The SODA was used as a secondary outcome measure in the ENTER
4
5 trial.⁹⁷
6
7
8
9

10 D.3. Patient assessment of upper gastrointestinal symptom severity index and
11 quality of life (PAGI-SYM and PAGI-QOL) Questionnaires.
12
13
14
15
16

17 The PAGI-SYM[®] questionnaire has been developed and validated for the
18 evaluation of symptom severity and treatment responsiveness in upper
19 gastrointestinal disorders such as FD, gastroparesis and GERD.¹³⁰ It is
20 composed of 20 items in 6 subscales: heartburn/regurgitation (7 items),
21 nausea/vomiting (3 items), postprandial fullness/early satiety (4 items),
22 bloating (2 items), upper abdominal pain (2 items), and lower abdominal pain
23 (2 items). Each item is scored from 0 (none) to 5 (very severe). The items
24 were derived from patient focus groups and expert opinion. The PAGI-SYM
25 questionnaire has good internal consistency, reliability and reproducibility over
26 2 weeks, and adequate content and construct validity in samples of subjects
27 with upper gastrointestinal disorders.¹³⁰ Responsiveness has been evaluated
28 in an 8-week study in GERD and FD, using OTE as a comparator.¹³¹ The
29 questionnaire has also been translated and validated in many different cultural
30 and linguistic settings. Disadvantages of the PAGI-SYM questionnaire are the
31 equal weighting of different symptoms when making up total or subscales,
32 and the inclusion of a broad range of symptoms that are not specific for FD.
33 However, subscales were proposed for use in specific disorders.¹³¹ The
34 PAGI-SYM has been used in a study evaluating the 5-HT_{1A} agonist R137696
35 in FD, where no efficacy was demonstrated.¹³²
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6 The PAGA-QOL questionnaire consists of 30 items in five subscales: daily
7
8 activities (10 items), clothing (2 items), diet and food (7 items), relationships (3
9
10 items) and psychological distress (8 items). Patients accord a score to each
11
12 item from 0 (absent) to 5 (severely impaired). It has been validated for
13
14 evaluating quality of life in FD, gastro-oesophageal reflux disease and
15
16 gastroparesis by comparing it with the OTE.¹³³
17
18

19 20 21 22 D.4. Glasgow Dyspepsia Severity Score (GDSS) 23 24

25
26
27 The GDSS¹³⁴ provides a tool for the global evaluation of dyspepsia during the
28
29 preceding six months, and was shown to be reliable, valid and responsive. It
30
31 records frequency of symptoms (maximum score,5); effect on daily activities
32
33 (2); the number of days off work because of dyspepsia (2), frequency of
34
35 medical consultations (2); home visits by a general practitioner (2); clinical
36
37 investigations performed (2); use of over the counter medications (2) and
38
39 prescription medications(3). Scores range from 0-20, and are significantly
40
41 lower in healthy volunteers (mean score 1.16) compared to patients with
42
43 duodenal ulcer (11.1) or FD (10.5). The coefficient of variation was 2% and
44
45 8% for intra- and inter-observer assessment respectively. Following
46
47 eradication of *Helicobacter pylori* in duodenal ulcer, the score changed from
48
49 11.4 to 1.3, compared with an average change of 10.5 to 8.5 in patients with
50
51 persistent *Helicobacter pylori* infection, confirming responsiveness of the
52
53 GDSS. A score of 0 or 1 on the GDSS corresponds to complete resolution of
54
55
56
57
58
59
60

1
2
3 dyspepsia. Limitations of the GDSS are administration by an investigator, and
4
5 lack of standardized definition of dyspepsia for the investigator to adhere to.
6
7

8
9
10 The GDSS was utilised by McColl et al¹³⁵ in a trial evaluating the efficacy of
11
12 *Helicobacter pylori* eradication therapy in non ulcer dyspepsia. A Spanish
13
14 translation of the GDSS and a Likert-scale symptomatic test were evaluated
15
16 for responsiveness to treatment, validity and reproducibility by means of
17
18 phone interview.¹³⁶ Both the GDSS and Likert-scales were valid (higher
19
20 scores in patients undergoing endoscopy than in healthy controls); had
21
22 adequate response after subjects completed *Helicobacter pylori* eradication
23
24 therapy, and showed low intraobserver variation whether the interviews were
25
26 conducted by phone or by a combination of phone and clinical interview. The
27
28 GDSS was superior to the Likert-scale in remaining reproducible even when
29
30 conducted by different observers.
31
32
33
34
35
36
37
38

39 D.5. Other disease specific multidimensional instruments.

40
41
42

43 The Clinical Dyspepsia Questionnaire is a condition specific self-administered
44
45 measure of dyspepsia and peptic-ulcer related symptoms. It assesses the
46
47 frequency and severity of symptoms and their impact on QOL.¹³⁷ Patient
48
49 scores on this questionnaire reviewed only small to moderate correlations
50
51 (0.20 to 0.66) with the SF-36 quality of life survey, but further validity tests
52
53 showed significant correlations with general practitioner perceptions of
54
55 symptom severity, family history of peptic ulcer disease and whether patients
56
57
58
59
60

1
2
3 were referred. Test-retest reliability yielded an intraclass coefficient value of
4
5 0.69. However, responsiveness was not assessed.
6
7

8
9
10 The Gastrointestinal Symptom Score (GIS) is a 10-item questionnaire
11 developed for the study of FD.¹³⁸ It was developed based on patient focus
12 groups, and validated by comparison with the Nepean Dyspepsia Index.
13
14 However, the questionnaire includes typical GERD symptoms like heartburn,
15 regurgitation and retrosternal discomfort. The GIS has been used in studies of
16
17 herbal medicines in FD.¹³⁹⁻¹⁴¹
18
19
20
21
22
23
24
25
26
27
28

29 **Quality of life measures**

30
31
32
33

34 Quality of life (QOL) measures can be assessed either by generic instruments
35 or by disease specific measures. The Short Form of General Health
36 Questionnaire (SF-36)¹⁴² is a generic instrument that has been widely used in
37 various disorders. To evaluate more specifically QOL in specific disorders,
38 such as in functional gastrointestinal disorders, questionnaires that are
39 available include the Nepean Dyspepsia Index (NDI)¹²¹⁻¹²⁶ for FD, the PAGI-
40 QOL for upper GI disorders¹³⁰ and the IBS-QOL¹⁴³ for IBS. Although these
41 quality of life measures have been strongly recommended as secondary
42 outcome measures, they are not utilised as primary outcome measures as
43 they are regarded by regulatory agencies as being insufficiently responsive to
44
45 treatment, although this may not necessarily be correct.¹⁰
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Various trials have measured QOL as primary or secondary outcome
4 measures (see Table 1). In a recent trial of tegaserod for dysmotility-type FD,
5
6 significant improvement in treatment satisfaction and reduced work and daily
7
8 activity impairment were reported in the treatment group compared with
9
10 placebo.¹⁴⁴
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

DISCUSSION

Evaluating the therapeutic efficacy of drugs in FD remains challenging due to the variable intensity of symptoms in one individual and the high placebo response rates.^{13,145-149} The choice of the endpoint for a clinical trial is a key factor in assessing therapeutic efficacy. However, there is a lack of consensus on the best method to measure clinical outcome in FD drug trials.^{13,14} In this setting, the most recent mandate emphasises the use of PRO in symptom reporting.⁹ The dichotomous endpoint of adequate or satisfactory relief has seen increasing use as primary endpoint in recent trials, while alternative outcome measures such as integrative questionnaires are also considered potentially acceptable, but these views seem largely driven by experience with IBS trials.²⁶⁻³⁷

Achieving the adequate relief endpoint in clinical drug trials was instrumental in obtaining FDA approval initially for ranitidine for the treatment of heartburn,¹⁵¹ for alosetron in IBS²⁶⁻²⁸, and subsequently for tegaserod in IBS (using satisfactory relief).³⁴⁻³⁷ Evidence supporting responsiveness of the adequate or satisfactory relief endpoints was documented mainly in IBS trials, where the binary endpoint correlated with other standard measures of treatment efficacy.^{26-37,152,153}

On the other hand, limitations of the binary endpoints include the lack of sensitivity and validation, inability to detect worsening of symptoms (unlike the OTE), lack of information on the magnitude of improvement needed to reach

1
2
3 adequate relief, the potential for failure to report adequate relief in subsequent
4
5 weeks once achieved, the influence of baseline severity of symptoms and
6
7 response outcomes and variable interpretation of the adequate relief
8
9 construct.³⁹ Based on an observational IBS study in a health maintenance
10
11 organization setting, it has been suggested that patients with mild symptoms
12
13 at baseline are more likely to achieve satisfactory relief compared to those
14
15 with severe symptoms at baseline.¹⁵⁴ Contrary to this suggestion, the greatest
16
17 improvement in percentage of days that FD patients reported satisfactory
18
19 relief of symptoms occurred in those who had severe baseline symptoms
20
21 compared with those who had mild symptoms ($p < 0.05$).⁴³ Hence, the use of
22
23 binary relief endpoints in FD deserves additional validation studies. Compared
24
25 to questionnaires that use specific FD symptoms, the binary or overall
26
27 treatment evaluation question may be more suitable to capture an overall
28
29 treatment impact in functional disorders, especially as this approach has the
30
31 potential to also contain changes in associated more global symptoms (e.g.
32
33 fatigue).

34
35
36
37
38
39
40
41
42
43
44
45
46 In addressing the use of an acceptable alternative outcome measure, a
47
48 questionnaire should undergo appropriate psychometric validation in
49
50 accordance with the FDA guidance regulations and Rome III guidelines on
51
52 clinical trials.^{9,10} In our opinion, existing questionnaires present some
53
54 methodological flaws, mainly due to the failure to satisfy the psychometric
55
56 property of content validity. In the light of the Rome III criteria⁴, none of the
57
58 existing questionnaires are sufficiently validated to be recommended
59
60

1
2
3 unequivocally as the primary outcome measure for FD trials. Central to the
4
5
6 problem of previously validated questionnaires was the inclusion of many
7
8 symptoms which are not considered cardinal FD symptoms according to
9
10 current criteria⁴. An adequate and valid questionnaire, i.e. one that
11
12 incorporates all the psychometric validation standards, in particular
13
14 incorporating the Rome III criteria for FD, would facilitate future therapeutic
15
16 trials in FD.
17
18

19
20
21
22 The absence of a validated instrument for the evaluation of treatment efficacy
23
24 in FD according to Rome III criteria and the lack of robust psychometric
25
26 validation data for the binary outcome measure in FD trials should not impair
27
28 ongoing or planned drug development in this area. One plausible solution is to
29
30 build upon the heritage of current instruments to serve as a useful framework
31
32 in designing a new questionnaire. Existing questionnaires that include many
33
34 non-FD symptoms can be narrowed down to evaluate only the cardinal
35
36 symptoms according to Rome III. Such an approach was used in the itopride
37
38 phase 2 and 3 studies, which used 2 FD-specific questions (severity of
39
40 epigastric pain and of postprandial fullness) from the LDQ as co-primary
41
42 endpoint, whereas the full LDQ score was a secondary endpoint.^{91,92} The
43
44 most valuable questionnaires are probably the ones that used patient focus
45
46 groups to identify relevant symptoms, such as the DSSI, PAGI-SYM or GIS
47
48 questionnaires.^{117,121,130,138} Another possibility is to incorporate the OTE with
49
50 the binary endpoints of adequate or satisfactory relief, for which there is
51
52 extensive experience from IBS trials²⁶⁻³⁷ and emerging experience from FD
53
54 trials.^{25,40-43} The OTE was extensively used in the OCAY, CADET and ENTER
55
56
57
58
59
60

1
2
3 trials^{89,94-97} and most recently in the tegaserod and acotiamide studies.^{42,43}

4
5 The subjective nature of the global outcome approach or OTE allows the
6 individual to integrate all aspects of his condition into a single treatment
7 outcome. Although the OTE is particularly suitable to show deterioration, it is
8 prone to recall bias as it compares current symptoms to pre-treatment
9 severity. Nevertheless, using both the OTE and binary outcome to evaluate
10 treatment efficacy would provide valuable information both from a global
11 perspective as well as in assessing symptom improvement and/or
12 deterioration.
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 Similar to other FGIDs, the magnitude of the placebo response is a major
28 concern in the evaluation of efficacy of drugs in FD. The available recent
29 studies do not show a major difference in responder rate when comparing
30 binary endpoints to OTE responses or to responder rates on composite
31 questionnaires like the LDQ or DSSI, provided the cut-off level for response
32 was adequately chosen.^{42,43,92} Indeed, these studies illustrate how the
33 magnitude of the placebo effect depends on the level of improvement used to
34 define a responder. For instance, in the itopride phase 3 trials, an
35 improvement of 1 point on the LDQ had a placebo response of 70 to 75%;
36 when a 2 point improvement was chosen as response definition, the placebo
37 response was 45 to 55%, quite similar to the placebo response on a global
38 patient assessment of efficacy.⁹² Similarly, requiring a 50% weekly response
39 rate over a 75% weekly response rate is associated with a higher placebo
40 rate.^{42,43} In addition, the tegaserod phase 3 studies showed a high placebo
41 response rate, and a low therapeutic gain of active drug over placebo, when
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 patients with mild symptoms were considered, while a bigger therapeutic gain
4
5 over placebo was obtained in those with moderate or severe symptoms.⁴³
6
7 Hence, in order to minimize the placebo effect, inclusion of patients with at
8
9 least moderate symptom severity, and the choice of a high enough threshold
10
11 for response definition is recommended.
12
13
14
15

16
17 Some studies in FD have used elimination of symptoms, usually measured on
18
19 Likert scales, as an endpoint.^{98,114,135} Although it is clear that becoming
20
21 asymptomatic is clinically relevant, the proportion of patients with functional
22
23 gastrointestinal disorders in whom this can be achieved by a pharmacological
24
25 intervention (and by the comparator placebo arm) is likely to be low. Placebo
26
27 responses are indeed low when this endpoint is considered.^{98,114,135}
28
29 Elimination of symptoms, therefore, probably sets a very high threshold for
30
31 response and focusing on this endpoint induces a risk of disregarding
32
33 clinically relevant degrees of symptom improvement in FD and other FGIDs.
34
35 International consensus guidelines, as well as FDA guidance, therefore focus
36
37 on identifying and recognizing significant symptom improvement using
38
39 validated scales,⁹⁻¹³ rather than symptom elimination which can still be
40
41 considered a useful secondary outcome variable.
42
43
44
45
46
47
48
49

50
51 No regulatory guideline exists on which magnitude of an active drug response
52
53 rate over placebo is needed to be considered clinically relevant. This was also
54
55 not specifically addressed by the Rome III working committee on design of
56
57 treatment trials in functional disorders.¹⁰ However, a group of experts
58
59 proposed a minimal range of efficacy of 10 to 15% over placebo to be
60

1
2
3 clinically relevant for functional disorders and for the irritable bowel syndrome
4
5 in particular.¹⁵⁵ It seems reasonable to assume that the same magnitude of
6
7 margin over placebo could also be considered clinically relevant in FD.
8
9

10
11
12 In conclusion, there are emerging data on the use of binary outcome
13
14 measures in FD clinical trials, although further validation is required. Most
15
16 existing questionnaires do not sufficiently fulfil psychometric validation criteria,
17
18 in view of the latest consensus definitions of FD by Rome III criteria. Hence,
19
20 evaluating and comparing the various options will need to go hand in hand
21
22 with the analysis of recent, ongoing or future pharmaceutical trials for the
23
24 treatment of FD, and optimization of potential endpoints that could be derived
25
26 from existing instruments.
27
28
29
30
31
32
33

34 **Grant support:**

35
36 Pieter Janssen is a research fellow of the FWO Flanders. This work was
37
38 supported by a Methusalem grant to Jan Tack, M.D., Ph.D.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Management of dyspepsia: Report of a working party. *Lancet* 1988;1:576-9.
2. Talley NJ, Koch KL, Koch M, et al. Functional dyspepsia: A classification with guidelines for diagnosis and management. *Gastrointest Int* 1991;4:145-60.
3. Talley NJ, Stanghellini V, Heading RC, Koch KL, Malagelada JR, Tytgat GNJ. Functional gastroduodenal disorders. *Gut* 1999; 45(Suppl II):37-42.
4. Tack J, Talley NJ, Camilleri M, Holtmann G, Hu PJ, Malagelada JR, Stanghellini V. Functional gastroduodenal disorders. *Gastroenterology* 2006;130:1466-79.
5. Drossman DA, Thompson G, Talley NJ, et al. Identification of subgroups of functional gastrointestinal disorders. *Gastroenterol Int* 1991;4:145-60.
6. Talley NJ, Phillips SF. Non-ulcer dyspepsia: Potential causes and pathophysiology. *Ann Intern Med* 1988;108:965-79.
7. Talley NJ, Weaver AL, Zinsmeister AR, et al. Onset and disappearance of gastrointestinal symptoms and functional gastrointestinal disorders. *Am J Epidemiol* 1992;136:165-77.
8. Shaib Y, El-Serag HB. The prevalence and risk factors of functional dyspepsia in a multiethnic population in the United States. *Am J Gastroenterol* 2004;99:2210-6.

- 1
2
3
4 9. US Food and Drug Administration (FDA). Guidance for Industry.
5 Patient reported outcome measures: use in medical product
6 development to support labeling claims. 2006.
7
- 8
9
10 10. Design of Treatment Trials Committee; Irvine EJ, Whitehead WE, Chey
11 WD, Matsueda K, Shaw M, Talley NJ, Veldhuyzen Van Zanten SJ.
12 Design of treatment trials for functional gastrointestinal disorders.
13 Gastroenterology 2006;130:1538-51.
14
- 15
16
17 11. Veldhuyzen van Zanten SJ. A systematic overview (meta-analysis) of
18 outcome measures in Helicobacter pylori gastritis trials and functional
19 dyspepsia. Scan J Gastroenterol 1993;28(Suppl 199):40-43.
20
- 21
22
23 12. Talley NJ. A critique of therapeutic trials in Helicobacter pylori-positive
24 functional dyspepsia. Gastroenterology 1994;106:1174-83
25
- 26
27
28 13. Veldhuyzen van Zanten SJ, Cleary C, Talley NJ, et al. Drug treatment
29 of functional dyspepsia: a systematic analysis of trial methodology
30 with recommendations for design of future trials. Am J Gastroenterol
31 1996;91:660-73.
32
- 33
34
35 14. Veldhuyzen van Zanten SJO. Assessment of outcome in dyspepsia:
36 has progress been made? Gut 2002;50(Suppl IV):23-25
37
- 38
39
40 15. Tack J, Bisschops R, Sarnelli G. (2004). Pathophysiology and
41 treatment of functional dyspepsia. Gastroenterology 127: 1239-55.
42
- 43
44
45 16. Guyatt GH, Veldhuyzen van Zanten SJO, Feeney DH, et al. Measuring
46 quality of life in clinical trials: a taxonomy and review. Can Med J
47 Assoc 1989;140:1441-8.
48
- 49
50
51 17. Talley NJ, Nyren O, Drossman DA, Heaton KW, Veldhuyzen van
52 Zanten SJO, Koch MM, Ransohoff DF. The irritable bowel syndrome:
53
54
55
56
57
58
59
60

- 1
2
3 toward optimal design of controlled treatment trials. *Gastroenterol Int*
4
5 1993;6:189-211.
6
7
- 8 18. Guyatt GH, Townsend M, Berman LB, Keller JL. A comparison of Likert
9 and Visual Analogue Scales for measuring change in function. *Chron*
10
11 *Dis* 1987;40:1129-33.
12
13
- 14 19. Jaeschke R, Singer J, Guyatt GH. Measurement of health
15 status:ascertaining the minimal clinically important difference. *Control*
16
17 *Clin Trials* 1989;10:407-15.
18
19
- 20 20. Guyatt G, Deyo RA, Charlson M, Levine MN, Mitchell A.
21 Responsiveness and validity in health status measurement: a
22
23 clarification. *J Clin Epidemiol* 1989;42:403-8.
24
25
- 26 21. Veldhuyzen van Zanten SJ, Talley NJ, Bytzer P, Klein KB, Whorwell
27
28 PJ, Zinsmeister AR. Design of treatment trials for
29
30 functionalgastrointestinal disorders. *Gut* 1999; 45(Suppl 2):1169-77.
31
32
- 33 22. Hays RD, Hadorn D. Responsiveness to change: an aspect of validity,
34
35 not a separate dimension. *Qual Life Res* 1992;1:73-5.
36
37
- 38 23. Kirshner B, Guyatt GH. A methodological framework for assessing
39
40 health indices. *J Chron Dis* 1985;38:27-36.
41
42
- 43 24. Cook KF, Rabeneck L, Campbell CJM, Wray NP. Evaluation of a
44
45 multidimensional measure of dyspepsia related health for use in a
46
47 randomized clinical trial. *J Clin Epidemiol* 1999;52:381-92.
48
49
- 50 25. Farup PG, Hovde O, Torp R, Wetterhus S. Patients with functional
51
52 dyspepsia responding to omeprazole have a characteristic gastro-
53
54 oesophageal reflux pattern. *Scan J Gastroenterol* 1999;34:575-9.
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
26. Camilleri M, Mayer EA, Drossman DA, Heath A, Dukes GE, McSorley D, Kong S, Mangel AW, Northcutt AR. Improvement in pain and bowel function in female irritable bowel patients with alosetron, a 5-HT₃ receptor antagonist. *Aliment Pharmacol Ther* 1999;13:1149-59.
27. Camilleri M, Northcutt SR, Kong S, Dukes GE, McSorley D, Mangel AW. Efficacy and safety of alosetron in women with irritable bowel syndrome: a randomized, placebo-controlled trial. *Lancet* 2000;355:1035-40.
28. Camilleri M, Chey WY, Mayer EA, Northcutt AR, Heath A, Dukes GE, McSorley D, Mangel AM. A randomized controlled clinical trial of the serotonin type 3 receptor antagonist alosetron in women with diarrhoea-predominant irritable bowel syndrome. *Arch Intern Med* 2001;161:1733-40.
29. Miner P, Stanton DB, Carter F, et al. Cilansetron in irritable bowel syndrome with diarrhoea predominance (IBS-D); efficacy and safety in a 3 month US study. *Am J Gastroenterol* 2004;99:S277.
30. Leventer SM, Kucharik RF, Keogh JC, et al. The potential dextofisopam for treatment of irritable bowel syndrome and the irritable bowel syndrome. *Am J Gastroenterol* 2004;99:S279.
31. Bradette M, Moennikes H, Carter F, Krause G, Caras S, Steinborn C. Cilansetron in irritable bowel syndrome with diarrhoea predominance (IBS-D): efficacy and safety in a 6 month global study. *Gastroenterology* 2004;126(Suppl. 2):A42.
32. Coremans G, Clouse RE, Carter F, et al. Cilansetron, a novel 5-HT₃ antagonist demonstrated efficacy in males with irritable bowel

- 1
2
3 syndrome with diarrhoea-predominance (IBS-D). *Gastroenterology*
4 2004;126(4 Suppl.2):A643.
5
6
7
8 33. Leventer S, Raudibaugh K, Frissora C, et al. The safety and efficacy of
9 dextofisopam in patients with diarrhoea-predominant or alternating
10 irritable bowel syndrome. *Gastroenterology* 2005;128(4 Suppl 2):A94.
11
12
13 34. Kellow J, Lee OY, Chang FY, et al. An Asia-Pacific, double-blind,
14 placebo controlled, randomized study to evaluate the efficacy, safety,
15 and tolerability of tegaserod on patients with irritable bowel syndrome.
16 *Gut* 2003;52:671-6.
17
18
19 35. Nyhlin H, Bang C, Elsborg L, Silvennoinen J, Holme I, Ruegg P, Jones
20 J, Wagner A. A double-blind, placebo-controlled, randomized study to
21 evaluate the efficacy, safety and tolerability of tegaserod in patients
22 with irritable bowel syndrome. *Scand J Gastroenterol* 2004;39:119-
23 126.
24
25
26 36. Muller-Lissner S, Holtmann G, Rueegg P, Weidinger G, Loffler H.
27 Tegaserod is effective in the initial and re-treatment of irritable bowel
28 syndrome with constipation. *Aliment Pharmacol Ther* 2005;21:11-20.
29
30
31 37. Tack J, Muller-Lissner S, Bytzer P et al. A randomized controlled trial
32 assessing the efficacy and safety of repeated tegaserod therapy in
33 women with irritable bowel syndrome with constipation. *Gut* 2005;
34 54:1707-13.
35
36
37 38. Mangel AW. Personal view: adequate relief as a primary endpoint in
38 irritable bowel syndrome. *Aliment Pharm Ther* 2006; 23:879-881.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
39. Camilleri M, Drossman DA, Fehnel S, Mangel A, Mayer EA, Talley NJ. Primary endpoints for FGID trials: A review of binary and integrated symptom assessments. *Clin Gastro Hepatol* 2007;5:534-40.
40. Talley NJ, Van Zanten SV, Saez LR, Dukes G, Perschy T, Heath M, Kleoudis C, Mangel AW. A dose ranging placebo controlled randomized trial of alosetron in patients with functional dyspepsia. *Aliment Pharmacol Ther* 2001;15:525-37.
41. Matsueda K, Hongo M, Sasaki D, Kusano M, Harasawa S, Arakawa T, Haruma K, Nakashima M, Miwa T and Saitou Y. Therapeutic Efficacy of Novel Agent(Z-338) in Functional Dyspepsia(FD). *Gastroenterology* 2006, A467 (abstract).
42. Talley NJ, Tack J., Kowalski D., Borton M.A., and Barve A. A Novel Acetylcholine Esterase Inhibitor Acotiamide Hydrochloride (YM443) in Functional Dyspepsia: Efficacy in a Randomized, Double-Blind, Placebo-Controlled Dose Ranging Trial. *Gastroenterology* 2008; 4, suppl. 1: A-157 (abstract).
43. Vakil N, Talley NJ, Zakko S, Laine L, Chey W, Tack J. Tegaserod treatment for functional dyspepsia dysmotility symptoms: results of two randomized, controlled trials. *Am J Gastroenterol.* 2008 Aug;103(8):1906-19.
44. Jian R, Ducrot F, Ruskone A, et al. Symptomatic, radionuclide and therapeutic assessment of chronic idiopathic dyspepsia. A double-blind placebo-controlled evaluation of cisapride. *Dig Dis Sci* 1989;34:657-64.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
45. Tatsuta M, Iishi H, Nakaizumi A, Okuda S. Effect of treatment with cisapride alone or in combination with domperidone on gastric emptying and gastrointestinal symptoms in dyspeptic patients. *Aliment Pharmacol Ther* 1992; 6:221-8.
46. Sarin SK, Sharma P, Chawla YK, et al. Clinical trial on the effect of domperidone on non-ulcer dyspepsia. *Indian J Med Res* 1986; 83:623-8.
47. Champion MC, MacCannell KI, Thomson AB, et al. A double-blind randomized study of cisapride in the treatment of non-ulcer dyspepsia. The Canadian Cisapride Nud Study Group. *Can J Gastroenterol* 1997;11:127-34.
48. De Groot GH, De Both PSM. Cisapride in functional dyspepsia in general practice. A placebo-controlled, randomized, double-blind study. *Aliment Pharmacol Ther* 1997;11:193-9.
49. Tack J, Masclee A, Heading R, et al. A dose-ranging, placebo-controlled, pilot trial of acotiamide in patients with functional dyspepsia. *Neurogastroenterol Motil.* 2009 Mar;21(3):272-80.
50. Gotthard R, Bodemar G, Brodin U, Jonsson KA. Treatment with cimetidine, antacid or placebo in patients with dyspepsia of unknown origin. *Scand J Gastroenterol* 1988;23:7-18.
51. Bekhti A, Rutgeerts L. Domperidone in the treatment of functional dyspepsia in patients with delayed gastric emptying. *Postgrad Med J* 1979;55(Suppl.1):30-32.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
52. Van de Mierop L, Rutgeerts L, Van den Langenberg B, Staessen A. Oral domperidone in chronic postprandial dyspepsia. *Digestion* 1979;19:244-50.
53. Haarmann K, Lebkuchner F, Widmann A, Kief W, Esslinger M. A double-blind study of domperidone in the symptomatic treatment of chronic post-prandial upper gastrointestinal distress. *Postgrad Med J* 1979;55(Suppl. 1):24-7.
54. Lance P, Wastell C, Schiller KFR. A controlled trial of cimetidine for the treatment of non-ulcer dyspepsia. *J Clin Gastroenterol* 1986;8:414-8.
55. Smith PM, Troughton AH, Gleeson F, Walters J, McCarthy CF. Pirenzepine in non-ulcer dyspepsia: a double-blind multicentre trial. *J Int Med Res* 1990;18:16-20.
56. Deruyttere M, Lepoutre L, Heylen H, Samain H, Pennoit H. Cisapride in the management of chronic functional dyspepsia: A multicentre, double-blind, placebo-controlled study. *Clin Ther* 1987;10:44-51.
57. Rosch W. Cisapride in non ulcer dyspepsia. Results of a placebo-controlled trial. *Scand J Gastroenterol* 1987;22:161-4.
58. Hannon R. Efficacy of cisapride in patients with nonulcer dyspepsia. A placebo-controlled study. *Curr Ther Res Clin Exp* 1987;42:814-22.
59. Goethals C, Van De Mierop L. Cisapride in the treatment of chronic functional dyspepsia. Results of a double-blind, placebo-controlled, crossover study. *Curr Ther Res* 1987;42:261-7.
60. Francois I, De Nutte N. Nonulcer dyspepsia: effect of the gastrointestinal prokinetic drug cisapride. *Curr Ther Res* 1987;41:891-8.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61. De Nutte N, Van Ganse W, Witterhulghé M, Defrance P. Relief of epigastric pain in nonulcer dyspepsia: controlled trial of the promotility drug cisapride. *Clin Ther* 1989;11:62-8.
62. Wood SF, Penney SC, Cochran KM. Cisapride in functional dyspepsia: A double-blind placebo-controlled randomized trial in general practice patients. *Scand J Gastroenterol* 1993;28(Suppl 195):5-10.
63. Chung JM. Cisapride in chronic dyspepsia: results of a double-blind, placebo-controlled trial. *Scand J Gastroenterol* 1993;195(Suppl)11-4.
64. Van Outryve M, De Nutte N, Van Eeghem P, Gooris JP. Efficacy of cisapride in functional dyspepsia resistant to domperidone or metoclopramide: A double-blind placebo-controlled study. *Scand J Gastroenterol* 1993;28(Suppl. 195):47-53.
65. Fumagalli I, Hammer B. Cisapride versus metoclopramide in the treatment of functional dyspepsia. A double-blind comparative trial. *Scand J Gastroenterol* 1994;29:33-7.
66. Al-Quorain A, Larbi EB, Al-Shedoki F. A double-blind, randomized, placebo-controlled trial of cisapride in Saudi Arabs with functional dyspepsia. *Scand J Gastroenterol* 1995;30:531-4.
67. Kellow JE, Cowan H, Shuter B, et al. Efficacy of cisapride therapy in functional dyspepsia. *Aliment Pharmacol Ther* 1995;9:153-60.
68. Carvalhinhos A, Fidalgo P, Freire A, Matos L. Cisapride compared with ranitidine in the treatment of functional dyspepsia. *Euro J Gastroenterol and Hepatol* 1995;7:411-7.

- 1
- 2
- 3
- 4 69. Yeoh KG, Kang JY, Tay HH, et al. Effect of cisapride on functional
- 5 dyspepsia in patients with and without histological gastritis: a double-
- 6 blind placebo controlled trial. *J Gastroenterol Hepatol* 1997;12:13-8.
- 7
- 8
- 9
- 10 70. Hansen JM, Bytzer P, Schaffalitzky de Muckadell O. Placebo-
- 11 controlled trial of cisapride and nizatidine in unselected patients with
- 12 functional dyspepsia. *Am J Gastroenterol* 1998;93:368-74.
- 13
- 14
- 15
- 16
- 17 71. Creytens G. Effect of the non-antidopaminergic drug cisapride on
- 18 postprandial nausea. *Curr Ther Res* 1999;36(5):1063-70.
- 19
- 20
- 21
- 22 72. Chen JD, Ke MY, Lin XM, Wang Z, Zhang M. Cisapride provides
- 23 symptomatic relief in functional dyspepsia associated with gastric
- 24 myoelectrical abnormality. *Aliment Pharmacol Ther* 2000;14:1041-7.
- 25
- 26
- 27
- 28
- 29 73. Fraitag B, Homerin M, Hecketsweiler P. Double-blind dose-response
- 30 multicenter comparison of fedotozine and placebo in treatment of
- 31 nonulcer dyspepsia. *Dig Dis Sci* 1994;39:1072-77.
- 32
- 33
- 34
- 35
- 36 74. Read NW, Abitol JI, Bardhan KD, Whorwell PJ, Fraitag B. Efficacy and
- 37 safety of the peripheral kappa agonist fedotozine versus placebo in
- 38 the treatment of functional dyspepsia. *Gut* 1997;41:664-8.
- 39
- 40
- 41
- 42
- 43 75. Delattre M, Malesky M, Prinzie A. Symptomatic treatment of non-ulcer
- 44 dyspepsia with cimetidine. *Curr Ther Res* 1985;37(5):980-1.
- 45
- 46
- 47
- 48 76. Kelbaek H, Linde J, Eriksen J, et al. Controlled clinical trial of treatment
- 49 with cimetidine for non-ulcer dyspepsia. *Acta Med Scand*
- 50 1985;217:281-7.
- 51
- 52
- 53
- 54
- 55 77. Singal AK, Kumar A, Broor SL. Cimetidine in the treatment of non-ulcer
- 56 dyspepsia: results of a randomized double-blind, placebo-controlled
- 57 study. *Curr Med Res Opin* 1989;11:390-7.
- 58
- 59
- 60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
78. Kairaluoma MI, Hentilae R, Alavaikko M, et al. Sucralfate versus placebo in treatment of non-ulcer dyspepsia. *Am J Med* 1987; 83:51-5.
79. Loffeld RJ, Potters HV, Stobberingh E, Flendrig JA, van Spreeuwel JP, Arends JW. *Campylobacter* associated gastritis in patients with non-ulcer dyspepsia: a double blind placebo controlled trial with colloidal bismuth subcitrate. *Gut* 1989;30:1206-12.
80. Vaira D, Holton J, Ainley C, Falzon M, Osborn J, D'Anna L, et al. Double blind trial of colloidal bismuth subcitrate versus placebo in *Helicobacter pylori* positive patients with non-ulcer dyspepsia. *Italian J Gastroenterol* 1992;24(7):400-4.
81. Hausken T, Stene-Larsen G, Lange O, et al. Misoprostol treatment exacerbates abdominal discomfort in patients with non-ulcer dyspepsia and erosive pyloric changes. A double-blind, placebo-controlled multicentre study. *Scand J Gastroenterol* 1990;25:1028-33.
82. Bolling-Sternevald E, Lauritsen K, Aalykke C, et al. Effect of profound acid suppression in functional dyspepsia: a double-blind, randomized, placebo-controlled trial. *Scand J Gastroenterol* 2002;37:1395-1402.
83. Peura DA, Kovacs TO, Metz D, Siepmann N, Pilmer BL, Talley NJ. Lansoprazole in the treatment of functional dyspepsia: Two double-blind, randomized, placebo-controlled trials. *Am J Med* 2004;116:740-8.
84. Blum AL, Arnold R, Stolte M, Fischer M, Koelz HR and the Frosch Study Group. Short course acid suppressive treatment for patients

- 1
2
3 with functional dyspepsia: results depend on *Helicobacter pylori*
4 status. *Gut* 2000;47:473-80.
5
6
7
- 8 85. Talley NJ, Verlinden M, Snape W, Beker JA, et al. Failure of a motilin
9 receptor agonist (ABT-229) to relieve the symptoms of functional
10 dyspepsia in patients with and without delayed gastric emptying: a
11 randomized double-blind placebo-controlled trial. *Aliment Pharmacol*
12 *Ther* 2000;14:1653-61.
13
14
15
16
17
18
19
- 20 86. Hallerback BI, Bommelaer G, Bredberg E, et al. Dose finding study of
21 mosapride in functional dyspepsia: a placebo-controlled, randomized
22 study. *Aliment Pharmacol Ther* 2002;16:959-67.
23
24
25
26
- 27 87. Holtmann G, Adam B, Haag S, Collet W, Grunewald E, Windeck T.
28 Efficacy of artichoke leaf extract in the treatment of patients with
29 functional dyspepsia: a six-week placebo-controlled, double-blind
30 multicentre trial. *Aliment Pharmacol Ther* 2003; 18:1099-1105.
31
32
33
34
35
- 36 88. Talley NJ, Meineche-Schmidt V, Pare P, et al. Efficacy of omeprazole
37 in functional dyspepsia: double-blind, randomized, placebo-controlled
38 trials (the Bond and Opera studies). *Aliment Pharmacol*
39 *Ther* 1998;12:1055-65.
40
41
42
43
44
45
- 46 89. Blum, AL, Talley NJ, O'Morain C, et al. Lack of effect of treating
47 *Helicobacter pylori* infection in patients with nonulcer dyspepsia.
48 Omeprazole plus Clarithromycin and Amoxicillin Effect One Year after
49 Treatment (OCAY) Study Group. *N Engl J Med* 1998;339:1875-81.
50
51
52
53
54
- 55 90. Talley NJ, Janssens J, Lauritsen K, et al. Eradication of *Helicobacter*
56 *pylori* in functional dyspepsia: Randomised double blind placebo
57 controlled trial with 12 months' follow up. *The Optimal Regimen Cures*
58
59
60

- 1
2
3 Helicobacter Induced Dyspepsia (ORCHID) Study group. BMJ
4 1999;318:833-7.
5
6
7
8 91. Holtmann G, Talley NJ, Liebrechts T, Adam B, Parow C. A placebo-
9 controlled trial of itopride in functional dyspepsia. N Eng J Med
10 2006;354:832-40.
11
12
13 92. Talley NJ, Tack J, Ptak T, Gupta R, Giguere M. Itopride in Functional
14 Dyspepsia: Results of two phase III multicenter, randomized, double-
15 blind, placebo-controlled trials. Gut 2008.
16
17
18
19
20
21
22 93. Veldhuyzen van Zanten SJO, Tytgat KMAJ, Pollak PT, Goldie J,
23 Goodacre RL, Riddell RH, Hunt RH. Can severity of symptoms be
24 used as outcome measures in trials of non-ulcer dyspepsia and
25 Helicobacter pylori associated gastritis? J Clin Epidemiol
26 1993;46:273-9.
27
28
29
30
31
32
33
34 94. Veldhuyzen van Zanten SJO, Chiba N, Armstrong D, et al. A
35 randomized trial comparing omeprazole, ranitidine, cisapride, or
36 placebo in Helicobacter pylori negative, primary care patients with
37 dyspepsia: The CADET-HN study. Am J Gastroenterol 2005;100:1477-
38 88.
39
40
41
42
43
44
45
46 95. Chiba N, Veldhuyzen van Zanten SJO, Sinclair P, et al. Treating
47 Helicobacter pylori infection in primary care patients with
48 uninvestigated dyspepsia: The Canadian adult dyspepsia empiric
49 treatment-Helicobacter pylori positive (CADET-HP) randomised
50 controlled trial. BMJ 2002;324:1012-6.
51
52
53
54
55
56
57
58 96. Armstrong D, Veldhuyzen van Zanten SJO, Barkun A, et al, the
59 CADET-HR Study Group. Heartburn-dominant, uninvestigated
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- dyspepsia: A comparison of 'PPI-Start' and 'H2-RA-Start' management strategies in primary care. The CADET-HR Study. *Aliment Pharmacol Ther* 2005;21:1189-202.
97. Veldhuyzen van Zanten S, Armstrong D, Chiba N, et al. Esomeprazole 40 mg once a day in patients with functional dyspepsia: The Randomized, Placebo-Controlled "ENTER" Trial. *Am J Gastroenterol*
98. Matsueda K., Hongo M., Sasaki D., Kusano M., Harasawa S, Arakawa T., Haruma K., Miwa T., and Nkashima M. Clinical trial: dose-dependent therapeutic efficacy of acotiamide hydrochloride (Z-338) in patients with functional dyspepsia - 100 mg t.i.d. is an optimal . *Neurogastroenterol Motil.* 2010 Jun;22(6):618-e173.
99. Weberg R, Berstad A. Low dose antacids and pirenzepine in the treatment of patients with nonulcer dyspepsia and erosive prepyloric changes. A randomized, double-blind, placebo-controlled trial. *Scand J Gastroenterol* 1988;23:237-43.
100. Johannessen T, Fjosne U, Kleveland PM, et al. Cimetidine responders in non-ulcer dyspepsia. *Scan J Gastroenterol* 1988;23:327-336.
101. Seno H, Nakase H, Chiba T. Usefulness of famotidine in functional dyspepsia patient treatment: comparison among prokinetic, acid suppression and antianxiety therapies. *Aliment Pharmacol Ther* 2005;21(Suppl2):32-36.
102. Hausken T, Berstad A. Cisapride treatment of patients with non-ulcer dyspepsia and erosive prepyloric changes. A double-blind, placebo-controlled trial. *Scan J Gastroenterol* 1992;27:213-7.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
103. May B, Kohler S, Schneider B. Efficacy and tolerability of a fixed combination of peppermint oil and caraway oil in patients suffering from functional dyspepsia. *Aliment Pharmacol Ther* 2000;14:1671-77.
104. Holtmann G, Gschossmann J, Mayr P, Talley NJ. A randomized placebo-controlled trial of simethicone and cisapride for the treatment of patients with functional dyspepsia. *Aliment Pharmacol Ther* 2002;16:1641-8.
105. Veldhuyzen van Zanten SJO, Chiba N, Armstrong D, et al. Validation of a 7-point global overall symptom (GOS) scale to measure the severity of dyspepsia symptoms in clinical trials. *Aliment Pharmacol Ther* 2006;23:521-9.
106. Armstrong D, Veldhuyzen van Zanten SJO, Barkun AN, et al. Symptom response at 1-week predicts responders at 4-weeks with esomeprazole (40 mg bid and od) in patients with uninvestigated dyspepsia. *Am J Gastroenterol* 2003;98(9 Suppl1):S48-9.
107. Wiklund IK, Junghard O, Grace E, et al. Quality of life in reflux and dyspepsia patients. Psychometric documentation of a new disease-specific questionnaire (QOLRAD). *Eur J Surg* 1998;583:41-9.
108. Svedlund J, Sjodin I, Dotevall G. GSRS-a clinical rating scale for gastrointestinal symptoms in patients with irritable bowel syndrome and peptic ulcer disease. *Dig Dis Sci* 1988;33:129-34.
109. Shaw MJ, Talley NJ, Beebe TJ, et al. Initial validation of a diagnostic questionnaire for gastroesophageal reflux disease. *Am J Gastroenterol* 2001;96:52-7.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
110. Dimenas E, Glise H, Hallerback B, et al. Quality of life in patients with upper gastrointestinal symptoms. An improved evaluation of treatment regimens? *Scand J Gastroenterol* 1993;28:681-7.
111. Revicki DA, Wood M, Wiklund I, et al. Reliability and validity of the Gastrointestinal Symptom Rating Scale in patients with gastroesophageal reflux disease. *Qual Life Res* 1998;7:75-83.
112. Talley NJ, Riff DS, Schwartz H, Marcuard SP. Double-blind placebo-controlled multicentre studies of rebamipide, a gastroprotective drug, in the treatment of functional dyspepsia with or without *Helicobacter pylori* infection. *Aliment Pharmacol Ther* 2001;15:1603-11.
113. Bolling-Sternevald E, Lauritsen K, Aalykke C, et al. Effect of profound acid suppression in functional dyspepsia: a double-blind, randomized, placebo-controlled trial. *Scand J Gastroenterol* 2002;37:1395-1402.
114. Miwa H, Nagahara A, Tominaga K, Yokoyama T, Sawada Y, Inoue K, Ashida K, Fukuchi T, Hojo M, Yamashita H, Tomita T, Hori K, Oshima T. Efficacy of the 5-HT_{1A} agonist tansospirone citrate in improving symptoms of patients with functional dyspepsia: a randomized controlled trial. *Am J Gastroenterol*. 2009 Nov;104(11):2779-87.
115. Moayyedi P, Duffett S, Braunholtz D, Mason S, Richards ID, Dowell AC, Axon AT. The Leeds Dyspepsia Questionnaire: a valid tool for measuring the presence and severity of dyspepsia. *Aliment Pharmacol Ther* 1998;12:1257-62.
116. Fraser A, Delaney BC, Ford AC, Qume M, Moayyedi P. The Short-Form Leeds Dyspepsia Questionnaire validation study. *Aliment Pharmacol Ther* 2007;25:477-86.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
117. Leidy NK, Farup C, Rentz AM, et al. Patient-based assessment in dyspepsia: Development and validation of Dyspepsia Symptom Severity Index (DSSI). *Dig Dis Sci* 2000; 45:1172-9.
118. Hu WM, Lam KF, Wong YH, et al. The Hong Kong index of dyspepsia: A validated symptom severity questionnaire for patients with dyspepsia. *J Gastroenterol Hepatol* 2002;17:545-51.
119. Wong WM, Wong BCY, Hung WK, et al. Double blind, randomised placebo controlled study of four weeks of lansoprazole for the treatment of functional dyspepsia in Chinese patients. *Gut* 2002;51:502-6
120. Sander GB, Mazzoleni LE, Francesconi CF, et al. Development and validation of a cross-cultural questionnaire to evaluate nonulcer dyspepsia: The Porto Alegre Dyspeptic Symptoms Questionnaire (PADYQ). *Dig Dis Sci* 2004;49:1822-9.
121. Talley NJ, Haque M, Wyeth JW, Stace NH, Tytgat GNJ, Stanghellini V, Holtmann G, Verlinden M, Jones M. Development of a new dyspepsia impact scale: the Nepean Dyspepsia Index. *Aliment Pharmacol Ther* 1999;13:225-35.
122. Talley NJ, Verlinden M, Jones MP. Validity of a new quality of life scale for functional dyspepsia: a United States multicentre trial of the Nepean Dyspepsia Index (NDI). *Am J Gastroenterol* 1999;94: 2390-7.
123. Functional dyspepsia, delayed gastric emptying, and impaired quality of life. *Gut*. 2006 Jul;55(7):933-9

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
124. Validity of a new quality of life scale for functional dyspepsia: a United States multicenter trial of the Nepean Dyspepsia Index. *Am J Gastroenterol.* 1999 Sep;94(9):2390-7
125. Initial linguistic and psychometric validation of the Arabic version of Nepean Dyspepsia Index. *Saudi Med J.* 2006 Oct;27(10):1554-60.
126. Talley NJ, Verlinden M, Jones M. Quality of life in functional dyspepsia: responsiveness of the Nepean Dyspepsia Index and development of a new 10-item short form. *Aliment Pharmacol Ther* 2001;15:207-16.
127. Holtmann G, Adam B, Haag S, Collet W, Grunewald E, Windeck T. Efficacy of artichoke leaf extract in the treatment of patients with functional dyspepsia: a six-week placebo-controlled, double-blind multicentre trial. *Aliment Pharmacol Ther* 2003; 18:1099-1105.
128. Rabeneck L, Cook KR, Wristers K, Soucek J, Menke T, Wray NP. SODA (Severity of Dyspepsia Assessment): A new effective outcome measure for dyspepsia-related health. *J Clin Epi* 2001;54:755-65.
129. Rabeneck L, Wristers K, goldstein JL, et al. Reliability, validity, and responsiveness of severity of dyspepsia assessment (SODA) in a randomized clinical trial of a COX-2-specific inhibitor and traditional NSAID therapy. *Am J Gastroenterol* 2002;97:32-9.
130. Rentz AM, Kahrilas P, Stanghellini V, Tack J, Talley NJ, de la Loge C, Trudeau E, Dubois D, Revicki DA. Development and psychometric evaluation of the patient assessment of upper gastrointestinal symptom severity index (PAGI-SYM) in patients with upper gastrointestinal disorders. *Qual Life Res* 2004; 13:1737-49.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
131. Revicki DA, Rentz AM, Tack J, Stanghellini V, Talley NJ, Kahrilas P, De La Loge C, Trudeau E, Dubois D. Responsiveness and interpretation of a symptom severity index specific to upper gastrointestinal disorders. *Clin Gastroenterol Hepatol* 2004;2:769-77.
132. Tack J, Van Elzen B, Tytgat G, Wajs E, Van Nueten L, De Ridder F, Boeckxstaens G. A placebo-controlled trial of the 5-HT_{1A} agonist R-137696 on symptoms, visceral hypersensitivity and on impaired accommodation in functional dyspepsia. *Neurogastroenterol Motil.* 2009 Jun;21(6):619-26, e23-4.
133. De La Loge C, Trudeau E, Marquis P, Revicki DA, Rentz AM, Stanghellini V, Talley NJ, Kahrilas P, Tack J, Dubois D. Responsiveness and interpretation of a quality of life questionnaire specific to upper gastrointestinal disorders. *Clin Gastroenterol Hepatol* 2004;2:778-86.
134. El-Omar EM, Banerjee S, Wirz A, McColl KEL. The Glasgow dyspepsia severity score-a tool for the global measurement of dyspepsia. *Eur J Gastroenterol Hepatol* 1996; 8:967-71.
135. McColl K, Murray L, El-Omar E, et al. Symptomatic benefit from eradicating *Helicobacter pylori* infection in patients with nonulcer dyspepsia. *N Eng J Med* 1998;339:1869-74
136. Calvet X, Bustamante E, Montserrat A, Roque M, Campo R, Gene B, Brullet E. Validation of phone interview for follow-up in clinical trials on dyspepsia: evaluation of the Glasgow Dyspepsia Severity Score and a Likert-scale symptoms test. *Eur J Gasatroenterol Hepatol* 2000;12:949-53.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
137. Garratt AM, Ruta DA, Russel I, et al. Developing a condition-specific measure of health for patients with dyspepsia and ulcer-related symptoms. *J Clin Epidemiol* 1996;49:565-71.
138. Adam B, Liebrechts T, Saadat-Gilani K, Vinson B, Holtmann G. Validation of the gastrointestinal symptom score for the assessment of symptoms in patients with functional dyspepsia. *Aliment Pharmacol Ther.* 2005 Aug 15;22(4):357-63.
139. Madisch A, Holtmann G, Mayr G, Vinson B, Hotz J. Treatment of functional dyspepsia with a herbal preparation. A doubleblind, randomized, placebo-controlled, multicenter trial. *Digestion* 2004; 69: 45–52.
140. Rosch W, Vinson B, Sassin I. A randomised clinical trial comparing the efficacy of a herbal preparation STW 5 with the prokinetic drug cisapride in patients with dysmotility type of functional dyspepsia. *Z Gastroenterol* 2002; 40: 401–8.
141. Madisch A, Melderis H, Mayr G, Sassin I, Hotz J. A plant extract and its modified preparation in functional dyspepsia. Results of a double-blind placebo controlled comparative study. *Z Gastroenterol* 2001; 39: 511–7.
142. Stewart AL, Hays RD, Ware JE Jr. The MOS short-form general health survey. Reliability and validity in a patient population. *Med Care* 1988;26:724-35.
143. Patrick DI, Drossman DA, Frederick HO, et al. Quality of life in persons with irritable bowel syndrome. Development and validation of a new measure. *Dig Dis Sci* 1998;43:400-411.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
144. Laine L, Tack J, Howden C, Omar M, Ligozio G, Kralstein J, Earnest DL. Work productivity and daily activity impairment in patients with functional dyspepsia: Double-blind comparison of tegaserod vs placebo. *Gastroenterology* 2007;132(Suppl. 2):A94.
145. Locke GR, III. Prevalence, incidence and natural history of dyspepsia and functional dyspepsia. In: Talley NJ, ed. *Balliere's Clinical Gastroenterology*, volume 12, No. 3. London:Balliere Tindall, 1998:435-42.
146. Lindell GH, Celebioglu F, Graffner HO. Non ulcer dyspepsia in the long term perspective. *Eur J Gastroenterol Hepatol* 1995;7:829-33.
147. Finney JS, Kinnersley N, Hughes M, et al. Meta analysis of antisecretory and gastrokinetic compounds in functional dyspepsia. *J Clin Gastroenterol* 1998; 26:312-20.
148. Klein KB. Controlled treatment trials in the irritable bowel syndrome: A critique. *Gastroenterology* 1988;95:232-41.
149. Moayyedi P, Soo S, Deeks J, Delaney B, Innes M, Forman D. Pharmacological interventions for non-ulcer dyspepsia. *Cochrane Database of Systematic Reviews* 2006; Issue 4:1-22
150. Pappa KA, Williams BO, Payne JE, Buaron KS, Mussari KL, Ciociola AA. A double-blind, placebo-controlled study of the efficacy and safety of non-prescription ranitidine 75 mg in the prevention of meal-induced heartburn. *Aliment Pharmacol Ther* 1999; 13(4):467-73.
151. FDA Gastrointestinal Drugs Advisory Committee. Transcript from FDA Gastrointestinal Drugs Advisory Committee. Lotronex 1999;November:119.

- 1
2
3
4 152. Mangel AW, Hahn BA, Heath AT, et al. Adequate relief as an endpoint
5
6 in clinical trials in irritable bowel syndrome. *J Int Med Res* 1998;26:76-
7
8 81.
9
- 10 153. Mangel AW, Northcutt AR. Review article: safety and efficacy of
11
12 alosetron, a 5-HT₃ receptor antagonist, in female irritable bowel
13
14 syndrome patients. *Aliment Pharmacol Ther* 1999;13(Suppl. 2):77-82.
15
16
- 17 154. Whitehead WE, Palasson OS, Levy RL, Feld AD, VonKorff M, Turner
18
19 M. Reports of "satisfactory relief" by IBS patients receiving usual
20
21 medical care are confounded by baseline symptom severity and do
22
23 not accurately reflect symptom improvement. *Am J Gastroenterol*
24
25 2006;101:1057-65.
26
27
- 28 155. Corazziari E, Bytzer P, Delvaux M, Holtmann G, Malagelada JR, Morris
29
30 J, Muller-Lissner S, Spiller RC, Tack J, Whorwell PJ. (2003). Clinical
31
32 trial guidelines for pharmacological treatment of irritable bowel
33
34 syndrome. *Aliment Pharmacol Ther.* 18: 569-80.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLES

Table 1. Summary of included trials and outcome measures employed.

Outcome measure	Scoring system	Limitations	References
Binary outcome measure	Adequate/satisfactory relief	Not Rome III-based Does not reflect magnitude of improvement Does not show deterioration May depend on baseline severity	25,40,42,43,53
Individual symptom assessment	(i) Likert scores	Not Rome III-based Inclusion of non-FD symptoms Equal weight to individual symptoms	40,43,44-98,
	(ii) VAS score	Not Rome III-based	99-103
Global evaluation	Likert scale / OTE	Not Rome III-based Recall bias for OTE	42,43, 48, 89, 91, 94-98 70-97,104-107,110,
Disease specific questionnaires	GOS	Not Rome III-based Inclusion of non-FD symptoms Equal weight to individual symptoms	94-96
	GSRS	Not Rome III-based Inclusion of non-FD symptoms Equal weight to individual symptoms	88,89,90,94- 97,112,113,122
	Leeds	Not Rome III-based Inclusion of non-FD symptoms Investigator-administered	91,92

	Complex scoring system	
DSSI	Not Rome III-based Inclusion of non-FD symptoms Equal weight to individual symptoms	42, 117
Hong Kong Index of Dyspepsia	Not Rome III-based Inclusion of non-FD symptoms Equal weight to individual symptoms	119
PADYQ	Not Rome III-based Inclusion of non-FD symptoms Equal weight to individual symptoms	120
Nepean Dyspepsia Index (NDI)	Not Rome III-based Inclusion of non-FD symptoms	42,43,91,92,123,131-135
SODA	Not Rome III-based Inclusion of non-FD symptoms Does not assess frequency of symptoms	97, 137
PAGI	Not Rome III-based Inclusion of non-FD symptoms Equal weight to individual symptoms	132
GDSS	Not Rome III-based Inclusion of non-FD symptoms Investigator-administered Heterogeneous scoring system	135
Quality of life	Not Rome III-based	102,108 (primary)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

assessments		Not validated as primary outcome measure	outcome measure), 66,70,72,73,78,98- 100,103-107 (secondary outcome measures)
-------------	--	---	---

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 I would like to see some further mention of the high placebo response rates in FD (the
4 authors mention this briefly in the Discussion) and the challenge in ascertaining a
5 significant difference between placebo and active management in disease specific
6 measures and overall improvement. How much of a difference is worthwhile? Many
7 researchers are of the view that in functional disorders improvements in specific
8 symptoms are not as important as an overall improvement in well being. Specific
9 symptoms measures may be the wrong trading currency in functional problems, which,
10 by their nature are likely to be part of a global disorder (e.g. tiredness included).
11
12
13
14
15
16
17
18
19
20

21 **Response:** We added a paragraph on the placebo response; the available data show
22 an influence of threshold level choice, but not of the type of endpoint chosen (binary vs
23 overall vs composite). We also, briefly, addressed the (lack of data and consensus on)
24 minimally relevant margin over placebo. We also added a statement on the limitations of
25 addressing only specific FD symptoms in assessment of therapeutic benefit.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60