

Design and preparation of market baskets of EU commercial baby foods for the assessment of infant exposure to food chemicals and to their effects

Raffaela Piccinelli, Marchela Pandelova, Cinzia Le Donne, Marika Ferrari,

Karl-Werner Schramm, Catherine Leclercq

▶ To cite this version:

Raffaela Piccinelli, Marchela Pandelova, Cinzia Le Donne, Marika Ferrari, Karl-Werner Schramm, et al.. Design and preparation of market baskets of EU commercial baby foods for the assessment of infant exposure to food chemicals and to their effects. Food Additives and Contaminants, 2010, 27 (10), pp.1337. 10.1080/19440049.2010.489913 . hal-00608722

HAL Id: hal-00608722 https://hal.science/hal-00608722

Submitted on 14 Jul 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Food Additives and Contaminants



Design and preparation of market baskets of EU commercial baby foods for the assessment of infant exposure to food chemicals and to their effects

Journal:	Food Additives and Contaminants
Manuscript ID:	TFAC-2010-013.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	23-Apr-2010
Complete List of Authors:	Piccinelli, Raffaela; National Research Institute on Food and Nutrition, Applied Sciences Pandelova, Marchela; Helmholtz Zentrum München Research, Center for Environmental Health, Institute of Ecological Chemistry Le Donne, Cinzia; National Research Institute on Food and Nutrition, Applied Sciences Ferrari, Marika; National Research Institute on Food and Nutrition, Applied Sciences Schramm, Karl-Werner; Helmholtz Zentrum München Research, Center for Environmental Health, Institute of Ecological Chemistry Leclercq, Catherine; National Research Institute on Food and Nutrition, Applied Sciences
Methods/Techniques:	Total diet, Exposure
Additives/Contaminants:	Hormones, Food contact materials, Metals, Pesticide residues
Food Types:	Baby food, Infant formulae



1	Design and preparation of market baskets of EU commercial baby
2	foods for the assessment of infant exposure to food chemicals and to
3	their effects
4	
5	R. Piccinelli ^{a1} , M. Pandelova ^b , C. Le Donne ^a , M. Ferrari ^a , KW. Schramm ^{b,c} and C.
6	Leclercq ^a
7	
8	^a National Research Institute on Food and Nutrition, Rome, Italy
9	^b Helmholtz Zentrum München, Research Center for Environmental Health, Institute of
10	Ecological Chemistry, Neuherberg, Germany
11	^c TUM, Wissenschaftszentrum Weihenstephan für Ernährung und Landnutzung,
12	Department für Biowissenschaften, Freising, Germany
13	
14	
15	
16	
l / 10	
10	
20	
20	
22	
$\frac{-}{23}$	
24	
25	
26	
27	
28	
	kamaeia Piccinelli. National Research Institute on Food and Nutrition, 00178 Rome. Italy.

¹Raffaela Piccinelli. National Research Institute on Food and Nutrition, 00178 Rome, Italy. E-mail:piccinelli@inran.it

1 Abstract:

The assessment of acute and chronic dietary exposure to contaminants in baby foods is needed to ensure healthy infant growth. Monthly EU market baskets of commercial baby foods were designed for the first 9 months of life by the "babyfood" study group of the CASCADE Network of Excellence for the specific purpose of assessing exposure to potentially toxic substances in infants fed with such foods. The present paper reports the different steps that led to the preparation of monthly pooled samples of commercial baby foods ("Infant formulae and follow-on formulae", and "Other baby foods") that may constitute the extreme case of the diet for an infant that would not be breast fed at all. Several market baskets were generated for an "average EU infant" and for infants of 4 selected countries (Italy, Sweden, Spain and Slovakia), fed with either milk infant formulae, soy infant formulae or hypoallergenic infant formulae and weaned (at the 5th month) with commercial baby foods and beverages available on the EU market. Market share data for 2007 for baby foods were used to design the baskets. Holding companies and the name of all their products were identified. Monthly diets for EU infants were elaborated in terms of food categories (e.g. infant cereals) of typologies of products (e.g. infant cereals without gluten) and of a specific product. The number of baskets generated was 30 for "Infant formulae and follow-on formulae" (including 62 products) and 13 for "Other baby foods" (including 35 products). These market baskets were designed to be used for the determination of certain contaminants and nutrients in the diet of EU infants and for the assessment of their effects on infant health.

23 Keywords: Market basket, baby foods, infant formulae, contaminants, EU

1 Introduction

Commercial food products specifically designed for infants (0-12 months) are available on the market in all EU countries. Their composition is regulated by two specific directives related to "infant formulae and follow-on formulae" (European Commission 2006a) and to "cereal-based foods and baby foods for infants and young children" (European Commission 2006b). In the present paper the terms "Infant formulae and follow-on formulae" and "Other baby foods" (including all others solid and liquid commercial products designed for infants) are used to refer to these two groups of products. According to the legislation, "Infant formulae" (sometimes called "starting formulae") are used during the first months of life and satisfying by themselves the nutritional requirements of such infants until the introduction of appropriate complementary feeding", whereas "follow-on formulae" are used by infants when appropriate complementary feeding is introduced" (European Commission 2006a).

The lifestyles changes encourage an increasing number of women to work while their children are young and parents have generally little time to prepare home-cooked foods for infants. This factor, together with manufacturer's innovation, probably contributed to the baby foods market growth which was observed in recent years (Research and Market 2006 [internet]). No more than 35% of infants worldwide are exclusively breastfed during the first four months of life (WHO 2003).

19 Therefore, in order to assess dietary exposure to potentially toxic substances in infants, it 20 appears crucial to monitor their exposure through "Infant formulae and follow-on formulae" and 21 "Other baby foods".

Infants are especially vulnerable to the acute, sub-acute and chronic effects of ingestion
 of chemical hazards. The nervous, reproductive, digestive, respiratory and immune systems of
 infants are developing.

Moreover, dietary exposure of infants to chemical hazards tends to be higher than that of other population groups. In fact, the average energy requirement of an infant aged 0 to 9 months ranges from 0.40 to 0.48 MJ /kg body weight i.e. three times that of a sedentary adult aged 30-59 years which is 0.14 MJ/kg bw in males (10.7 MJ for 75 kg) and 0.13 MJ/kg bw in females (8.3 MJ for 62 kg) (Commission of the European Communities 1993). Therefore, when the concentration of a chemical is similar in the diet of infants and adults, the dietary exposure of infants is three times that of adults.

In the EU legislation on "Infant formulae and follow-on formulae" and "Other baby foods" there are specific provisions about hygiene, use of food additives, presence of some contaminants and on the use of materials intended to come into contact with foods. These provisions are aimed to reduce the potential dietary exposure of infants to a number of
 microbiological and chemical hazards.

However there are no provisions in relation to a number of important endocrine disruptors(with the exception of provisions for pesticides).

5 Within the frame of the EU funded project CASCADE ("Chemicals as contaminants in 6 the food chain"; project website: <u>www.cascadenet.org</u>) a specific integrated study called 7 "Babyfood" was performed. The main objective of this study was to provide information on 8 dietary exposure through baby foods to a limited number of endocrine disruptors in the EU. The 9 target substances were bisphenol A (BPA), vinclozolin, polychlorinated dibenzo-p-dioxins and 10 dibenzofurans (PCDD/F), polychlorinated biphenyls (PCB), genistein, procimidone, iprodione 11 and cadmium.

The technique recommended by WHO (GEMS/Food 2005 [internet]) to assess dietary exposure to contaminants is the analytical determination of market baskets; this technique is very often used. The extreme case of an infant diet with the exclusive consumption of commercial products is not desirable since breast milk is the preferred food for infants. However, it provides a worst case for the assessment of potential exposure of children to food chemicals contained in these products.

18 The aim of the present paper is to illustrate the methods used within the CASCADE 19 project for the preparation of market baskets of commercial baby foods representing the average 20 diet of infants (aged 0- 9 months) who are not breast fed at all and are fed with commercial baby 21 foods only.

23 Materials and Methods

24 Market basket design and sample collection

In order to sample the food products, EU market baskets were created to represent the diet for each of the first 9 months of life of an infant who is not breast fed at all and is fed with commercial baby foods only. Such baskets, that aim to assess the average dietary exposure of infants in the EU, may not capture information on the variability of such exposure. Therefore, additional National market baskets were designed in order to assess the geographical variability. Four countries were selected: Sweden, Italy, Spain and Slovak Republic.

To design the baskets, "Infant formulae and follow-on formulae" and "Other baby foods" products were considered separately and monthly-based diets were elaborated. In the first group of products "Infant formulae" and "follow-on formulae" were considered. Both typologies were represented with three kinds of "Infant formulae and follow-on formulae": milk based formula, soy based formula and hypoallergenic formula, for a total of 6 kinds of products. The baskets

Food Additives and Contaminants

simulate the progressive introduction in the diet of "Other baby foods" from 5 months of age of an average infant weaned with commercial baby foods only. In the EU, some infants already receive little quantities of complementary foods at 4 months but the majority of infants receive such foods after 6 months of age, as suggested by World Health Organization (Michaelsen K et al. 2003). Moreover a recent opinion by EFSA (2009) states that the introduction of complementary food into the diet of healthy term infants in the EU between the age of 4 and 6 months is safe and does not pose a risk for adverse health effects. Month 5 was therefore chosen to represent the average between fourth and sixth month. From month 5 to month 9 the baskets of "Other baby foods" were varied every month with the addition of new "Other baby foods" products.

Market share data of year 2007 of 22 single countries, for "Infant formulae and follow-on formulae" and "Other baby foods", were purchased from the Company "Food for Thought" (FFT) ([Internet]). For each block of products taken separately, the overall market share of each Holding Company in the 22 countries was taken to represent its market share for these products in the whole EU. For the 4 specific countries, the national market share was used. These data were used to identify the brands of the products to be sampled. The name of the Holding Companies and of their main brands in 22 EU countries was obtained together with their overall market share.

The company FFT was able to provide data for three blocks of products:

Baby milks: including the 6 categories of baby milks to be sampled ("infant formulae" and
 "follow-on" of milk formula, soy formula and hypoallergenic formula respectively).

22 - Baby Jars including: baby food canned, in bottles or in jars, excluding freeze dried products.

Other Baby Foods including: juices and all dried products such as cereal powders with and
 without milk, biscuits, dry pasta and any freeze-dried product such as freeze dried meat,
 freeze dried fish, herbal tea.

26 The overall market shares for each of these three blocks of products were used as proxy 27 for the specific market shares of the specific products belonging to them.

Products belonging to the last two blocks, "Baby Jars" and "Other Baby Foods" are called
in this paper "Other baby foods". They include all liquid and solid commercial baby foods with
the exclusion of "Infant formulae and follow-on formulae".

To identify the products and to design the market baskets, different methods have been followed depending on the category of products ("Infant formulae and follow-on formulae" or "Other baby foods") and the typology of the sample (EU or National).

Holding Companies were ranked in decreasing order of market shares and the first
 companies that cumulatively reached over 80% were selected. These companies are called

hereafter "leading companies". A check of the websites of these companies showed that they produce and distribute the same (or similar) products in most EU countries. For logistic reasons, the collection of products was therefore tentatively organised in a unique country. Germany was selected. Consequently, the brands of the leading companies that were on the German market were identified. In all cases where a leading company did not have a brand in Germany, an alternative country was selected in which a brand of the same leading company was present. Therefore only a part of products were bought in Germany. The criterion used for the selection of the alternative country was "being the country with the highest estimated number of potential consumers within EU". Demographic data were downloaded from EUROSTAT (2009) ([Internet]) to assess the dimension of the population aged less than 1 year in each of the 22 EU countries. The number of potential consumers was estimated by weighing these data with the market shares of the leading company of interest in each of the 22 countries.

Regarding the National baskets, in order to capture the highest variability in chemical content of the diet, only one brand was identified in each of the four selected countries. It was the main brand from the leading company with the highest market share in that country. If certain categories of product were not produced by this leading company in that particular country, the most popular product (from another Company) sold in the same category was identified. This identification was performed through expert judgement.

21 "Infant formulae and follow-on formulae" *baskets*

In order to find the name of all "Infant formulae and follow-on formulae" products of the brands
identified, the websites of the brands were searched or the leading companies were contacted.

In term to EU baskets of "Infant formulae and follow-on formulae" the percentage of each product was simply the proportion of market share of each brand among the leading companies. In the event that more than one product was available for one brand, the market share of the specific brand was equally divided by the number of products.

For the National baskets, each typology of product was represented by a unique brand in each of the 4 selected countries and 100 was the percentage of each brand.

"Other baby foods" baskets

The name of the "Other baby foods" products were identified with the help of some CASCADE partners who searched products on their own country market. In order to find the names of the products to be sampled they were suggested to contact a Chemist's who would provide them a list of all products for the brand of interest or to search the websites of the companies or to

1 contact the companies. An appropriate database, organised by typology of products was filled 2 by CASCADE partners. The original name of each product and its English translation were 3 recorded together with a brief description of the product and some other information (typology 4 of packaging, month of introduction as suggested by the producer, main ingredients, ready to eat 5 or powder).

The following steps were followed to generate the baskets:

8 A) Identification and assessment of the elements necessary for the elaboration of the diet for EU
9 and National infants from month 5 to month 9:

• Calculation of energy requirements: the standard values of weight for age and sex established by WHO (WHO 2006) and the estimated energy requirements (kcal/kg of body weight/day) set by FAO (FAO 2004) for formula-fed infants were used to calculate the average energy requirements (kcal/day) of infants in the first 9 months of life (Table 1). For the reasons above explained, month 5 was chosen to insert little quantities of complementary foods. It was observed that the average energy intakes assessed by SCF (2003) from the 5th to the 7th months of age for infants fed with both complementary foods and follow-on formula were slightly higher than the energy requirements calculated from FAO (2004) and WHO (2006) (Table 1). Also average body weights were slightly higher. Therefore, in table 1, for infants of 5th months of age, the energy requirement for 5-6 month (622 kcal/day) was considered instead of that for 4-5 month (598 kcal/day). The same substitution was also applied for months 6th to 9th

23 (Table 1here)

• Identification of food categories included in the "Other baby foods" basket for each month of age: in order to establish the month of inclusion of each category of product, the guidelines for infant feeding developed by WHO Europe (Michaelsen K et al. 2003) were considered. However, these guidelines do not provide recommendations at a high level of detail in terms of product categories. The only paediatricians' guidelines providing recommendations at a high level of detail in the European region are those from Switzerland. Therefore, the guidelines of the Swiss Paediatric Society (Société Suisse de Pédiatrie 2002 [Internet]) were used to complement the WHO guidelines.

On these bases, a first theoretical schedule of the monthly inclusion of "Other baby
foods" was created. The availability of these products in the EU countries was then
investigated in order to produce a refined schedule (Figure 1). Thus, in the final

schedule, used in the design of the market baskets, the food category "infant cereals" was divided into "infant cereals without gluten", "infant cereals with gluten" and "infant cereals with milk" and the food category "fruit" was split into "pureed fruit" and "fruit juice". Eggs were not present in commercial baby foods for infants aged up to 9 months and therefore not considered in the final schedule. This is due to the caution required in the introduction of whole eggs due to the allergy risk (Michaelsen et al. 2003). Moreover several brands of baby foods were found to distribute "dairy based dessert with chocolate" that can be consumed by infants aged 6 to 9 months. The results of the food consumption survey performed in UK (Mills et al. 1992) suggest that these typologies of baby foods are effectively consumed in this period of the life. These products were therefore also included in the 9th month basket and the food category "dairy based dessert" was divided into "dairy based dessert with chocolate".

(Figure 1 here)

16 B) *Elaboration of the monthly diets:*

Amounts, energy and nutrient content of "follow-on formulae" and "Other baby foods": the estimated average amount of "follow-on formulae" consumed by infants during the 5th, 6t^h and 7th month (respectively 600 ml/day, 400 ml/day and 240 ml /day) were taken from a report of Scientific Committee on Food (SCF 2003). These amounts of liquid "follow-on formulae" (ml/day) were translated in gram/day of powder, according to the label of the sampled products, that suggests to use on average 30 ml of water to dilute 4.6 g of powder of "follow-on formulae" and based on an estimated volumetric density of milk of 1.03 g/ml (Di Thomas, 2003). The daily energy intake from "follow-on formulae" was calculated based on its average energy content according to the labels of the products of 8 brands (70.8 kcal/100 g). The daily energy intake from "Other baby foods" was then calculated by difference (Table 1).

To calculate the nutrient content of the elaborated diets, the average of values (by typologies of product) reported on labels of 6 brands among the most popular in EU was assessed.

- 33 The monthly diets were elaborated in three steps:
 - 1) in terms of main food categories (as infant cereals, pasta, fruits, meat etc...): the diets were designed in grams for each category of "Other baby foods", based on

Food Additives and Contaminants

the portion sizes suggested by producer labelling and on the meal daily frequency suggested by WHO Europe (Michaelsen et al. 2003), with the exception of "infant cereals" products. The amount of this latter category was calculated by difference from the total calculated energy intake from "Other baby foods" (Table 1). 2) in terms of typologies of foods products on the basis of specific features of the products (as Infant cereals without milk and with gluten, Infant cereals with milk, Pureed fruit, Fruit juice, Pulses with vegetables, Fish with vegetables etc...): the typologies of "Other baby foods" products to be included in the diets were chosen based on prepared by the CASCADE partner database. Table 2 shows the typologies of products and the specific features selected for inclusion in the EU baskets. The selection was based on the availability of similar products in different EU countries. Added fats (oil, butter, etc....) were not sampled, since they were present as ingredients in the mixed baby dishes and no such products specifically designed for babies were available in any EU country. Table 3 shows the elaborated monthly diets infants from 5 to 9 months of age (in terms of main category and typology of food products). For the National baskets of the 4 selected countries, the same typologies of products were chosen. If some typologies of products were not available in certain countries, the basket was modified for this country only. For example the typologies of products "Infant cereals without milk and with gluten", "Dairy based dessert with chocolate", "Biscuits", "Pulses with vegetables and fish" and "Pulses with vegetables and potatoes" were not available in Sweden. These typologies were therefore replaced with "Infant cereals with milk and with gluten", "Dairy based dessert with forest fruit", "Infant cereals with milk and with gluten" and "Fish, pulses with vegetables and potatoes", respectively. For similar reasons the National baskets of Slovakia were slightly modified with respect to the standard baskets. (Tables 2 and Table 3 here) 3) in terms of specific products identified: the tables filled in by the responsible person for shopping were used to choose the specific products to be inserted in the EU and National baskets, based on their ingredients.

For each food typology, the products chosen were the same for all ages of the infants; thus pureed fruit was the same at 5th, 6th, 7th, 8th and 9th. The National baskets were designed with a unique product from a unique brand.

The percentages of energy from proteins and lipids were calculated for the monthly diet elaborated and compared with the WHO recommendations (Michaelsen et al. 2003) for infants aged 5-9 months: 14-14.5 g/day of protein and 30-40% of energy from lipids.

- C) Creation of the EU and National baskets:
- Calculation of the percentage of the different products within the sample: the percentage of each brand within the EU and National baskets of "Other baby foods" was simply the proportion of market share of each of the selected leading company that were found to produce a specific product.
- With regard to the EU basket, when a typology of product was not available for a certain brand, "0" was assigned as percentage of market share for that leading company and the percentage of the other brands of the same typology were increased proportionally, leading to a "corrected market share". As far as the National baskets were concerned, also in this case, when the selected "leading company" did not distribute a certain category of product in the country, the most popular product of that category sold in the country was identified for sampling.

The preparation of the "Other baby foods" baskets, as they were designed initially and above described, would have required collection and sample preparation of a huge number of products and consequently lead to very high costs in terms of time and budget resources. Therefore, a random sampling of brands, weighed on market share, was performed in order to identify only one brand within each typology of product. For the National baskets of "Other baby foods", only the 5th month diet was prepared due to lower number products included compared with month 6^{th} , 7^{th} , 8^{th} and 9^{th} .

Table 4 shows an example of EU basket of "Infant formulae and follow-on formulae" and "Other baby foods" at month 5 with the products purchased only (the name of the brands were substituted with a code). The other baskets (with all the products selected and with the products purchased only) are not reported as tables in the present paper due to space restrictions but can be requested to the corresponding author.

(Table 4 here)

1	
2	D) Purchase of the products to be sampled
3	More than half of the products needed to constitute the EU baskets had to be bought in EU
4	countries other than Germany by the CASCADE partners (57% of "Infant formulae and
5	follow-on formulae" and 55% of "Other baby foods" products) and delivered to Helmholtz
6	Zentrum München. The other products were bought in Germany, where sample preparations
7	were performed prior analyses.
8	
9	Pooling, homogenisation, and lyophilisation of the baby foods samples
10	The pooling of the EU and National basket baby foods samples and related weighting
11	procedures were performed in a purified glovebox under nitrogen atmosphere. Prior to pooling,
12	the solid food and beverages were lyophilisated using the freeze drying equipment Type BETA
13	1-16 (Christ GmbH). Overall, the duration of the procedure for each food item was about 36
14	hours. After lyophilisation, the weight of baby food products was reduced with yield varying
15	from 12 to 22%. The proportion of each "Other baby foods" product in the pooled samples was
16	that reported in Table 4. The samples were homogenised in a rotoshaker, Type Reax 20/8
17	(Heidolph GmbH) for 8 hours at 10 rpm.
18	The samples were kept at room temperature and stored in brown glass jars protected from
19	direct light. In order to prevent any contamination, each further open and close of the sample
20	collector was performed under nitrogen atmosphere.
21	
22	Results
23	The number of baskets and of product sampled is reported in Table 5.
24	
25	"Infant formulae and follow-on formulae" baskets
26	The first 8 leading companies of "Infant formulae and follow-on formulae" in the EU were
27	selected, covering altogether 82% of the entire EU market. A total of 42 products were
28	purchased from 6 different countries in order to prepare 6 EU market baskets of "Infant
29	formulae and follow-on formulae".
30	On the other hand, 20 products were purchased to prepare 24 National market baskets of
31	"Infant formulae and follow-on formulae".
32	
33	Other baby foods baskets
34	With regard to the baskets of "Other baby foods", the 6 leading companies of the "baby jars"
35	market and the first 7 companies of the "other baby foods" market were selected, covering

altogether respectively 82% and 81% of the entire EU market. Nine EU market baskets of
 "Other baby foods" were designed and 112 products were identified. However because of the
 above mentioned reasons, only 22 randomly extracted products were sampled from 5 different
 countries.

5 As far as National baskets of "Other baby foods" are considered, 4 market baskets were 6 elaborated and 47 products were identified. As already mentioned only the basket for the 5th 7 months was prepared, based on a total of 13 products.

8 (Table 5 here)

11 Discussion and conclusion

Robust estimates of consumer exposure to food chemicals are needed for regulatory chemical risk assessment. Among the main approaches to food chemical intake assessment, WHO recommends the development of a diet model or market basket representing the average diet of the considered population. This model may then be analysed directly for food chemicals of interest (WHO 1999).

Survey market baskets are generally developed on the basis of national food consumption data. It is the case in the U.S (FDA[Internet]), in Australia (FSANZ [Internet]), and in the UK (MAFF [Internet]). A food consumption survey that could be considered representative at EU level is not available. Moreover, in many EU countries, information on exposure to chemical hazards in the diet is collected for the whole population, not specifically for children or infants. The available information is often incomplete or not comparable between EU countries. In order to assess exposure to hazardous chemicals through food, the data collection needs to be harmonized and data need to be collected regularly and to reflect the specific risks to children and infants in the region (ENHIS 2007). For these reasons the methodologies generally used to elaborate national markets baskets have been adapted to the needs of this study. Baskets were designed on the basis of theoretical schemes of insertion of baby foods and of EU market share data for "Infant formulae and follow-on formulae" and "Other baby foods".

The market share data purchased were expressed in value (money) because baby food data expressed in volume (amount of products) were not available at the same level of disaggregation. Although the price of products of the same typology can be different depending on the brand, one leading company can also produce more than one brand destined to different kinds of market (e.g. only for supermarkets with a low or medium price, only for chemists' with a high price). It was therefore assumed that on average the prices of similar products are

comparable among the companies and data purchased were used as a proxy for market shares in
 volumes.

Another limitation of the market basket illustrated in the present paper is that the market shares of the specific categories of "baby milks", "baby jars" and "other baby foods" were not available. Thus, since no specific market shares were available for "fruit baby jars" the assumption was made that the market shares of "baby jars" in general could apply to "fruit baby jars". Moreover, market share data used to identify the brands of the products to be sampled, did not contain private brands (supermarket) and baby foods of this kind of market were not sampled. However, the market share of each of these private brands is known to be low and they would not have been selected as leading brands. In fact, the overall market share of non identified brands of all the baby foods categories considered was 4.5% including private labels. Since the overall share of such private brands is growing, in future studies one might consider the random extraction of some private brands that would be included in the market basket in proportion to such overall share.

The amount of proteins in the monthly diets here elaborated and expressed in terms of typologies of products was higher than recommended by WHO (up to 11% versus 9%) (Table 4). However, this percentage of protein is lower than that observed in food consumption surveys performed in infants in some European countries: 13.2% of energy in the UK for infants aged 6-9 months (Mills et al 1992), 11.2-13.2% of energy in Germany for infants aged 3-9 months (Alexy et al. 1999), 13-14% of energy in Denmark for infants aged 9^h months (Hoppe et al. 2004), 13.3-15.7% of energy in Spain for infants aged 4-9 months (Capdevila et al. 1998). The amount of proteins in the diets considered in the present paper are therefore intermediate between that recommended by WHO and that observed in consumption surveys.

On the other hand, the percentage of energy from fat was slightly lower than recommended (24-35% versus 30-40%), due to the fact that 'added fats' were not considered in the elaboration of the monthly diets.

It is likely that some infants can consume honey since birth. This fact was taken in
consideration, but this food was not sampled because honey is not distributed by companies of
baby foods.

The lyophilisation of fruity baby foods in jar result a sticky material that caused difficulties in further homogenisation of the samples. As a consequence, these specific products were delivered separately from the other products belonging to the same basket. The further pooling was therefore performed in each testing laboratory itself according above mentioned weighting procedures. During the survey, the labels of infant formulae were checked and it appeared that the total energy intake of infants aged 0-4 months could be higher than recommended when the consumer followed instructions reported on the label. This observation has been reported by Pandelova et al., (2009a).

5 Several samples of "Infant formulae and follow-on formulae" and "Other baby foods", 6 both extracted and non extracted, were prepared and delivered to CASCADE partners for 7 further chemical target analysis, *in-vitro* and *in-vivo* testing. The results of some analyses are 8 published (Riu 2008, Pandelova 2009b).

9 The above described samples of baby foods have been used for the assessment of dietary 10 exposure to a number of substances within the CASCADE project, with the aim of obtaining a 11 realistic panorama of exposure to food chemicals in the infant population, useful for quantitative 12 risk assessment.

Market basket studies involve the analysis of groups of foods that reflect the average food consumption patterns of a given population. They may be used to establish specific maximum limits when there is evidence that the health of specific consumers may be at stake (CGSCTFF, 1999). In the present study, a market basket was designed to assess exposure in a very sensible population group: infants.

Given the time and resources available it was decided to distinguish only two groups of foods: "infant formulae and follow-on formulae" and "other products" and to focus on possible variability at the different ages of the babies and between countries rather than on variability between products categories. Thus, if a food chemical was to be found at a high level of concentration in one or more of the designed baskets it may not be possible to identify the incriminated product category. However, products were purchased in excess so that further analyses on single categories can be performed afterwards.

Despite some limitations, the method should allow to assess average dietary exposure of infants in the EU. The food items considered for the elaboration of the monthly diets are those normally consumed by this population group, the quantities were set based on nutritional requirements of infants and the products were purchased in seven different EU countries on the basis of EU market share data collected in 22 countries. The technique could be used again in the future and the exposure data obtained through this methodology could support regulatory actions over time and contribute to the promotion of health of infants fed with commercial foods during early development.

35 Acknowledgements

2 3								
4	1	The authors would like to thank all CASCADE partners for their assistance in the identification,						
5 6	2	collection and shipping of baby foods namely Ingemar Pongratz, Lars-Arne Haldosen, Stefan						
7 8	3	Rehnmark, Nicolas Olea, Jean-Pierre Cravedi and Július Brtko.						
9	4	The study was financially supported by the European Union network CASCADE (FOOD-CT-						
10	5	2003-506319).						
12 13	6							
14	7							
15 16	8	References						
17 18	9							
19	10	Alexy U, Kersting M, Sichert-Hellert W, Manz F, Schöch G. 1999. Macronutrient intake of 3-						
20 21	11	to 36-month-old German infants and children: results of the DONALD study. Ann Nutr Metab.						
22 23	12	43:14-22.						
24	13							
25 26	14	Capdevila F, Vizmanos B, and Marti ² -Henneberg C. 1998. Implications of the weaning pattern						
27	15	on macronutrient intake, food volume and energy density in non-breastfed infants during the						
20 29	16	first year of life. J Am Coll of Nutr. 17(3): 256–262.						
30 31	17							
32	18	CGSCTFF - Codex General Standard for Contaminants and Toxins in Food. Methodology and						
33 34	19	principles for exposure assessment in the Codex General Standard for Contaminants. 1999.						
35 36	20	(CX/FAC 99/13). Progress Report by WHO on the Revision of GEMS/Food Regional Diets						
37	21	(CX/PR 99/3) http://www.fao.org/docrep/meeting/005/X7137E/x7137e24.htm						
38 39	22							
40 41	23	Commission of the European Communities. 1993. Nutrient and energy intakes for the European						
42	24	Community, Reports of the scientific Committee for Food, thirty first series, Office for Official						
43 44	25	Publications of the European Communities, Luxembourg. (Available at:						
45 46	26	http://ec.europa.eu/food/fs/sc/scf/out89.pdf).						
47	27							
48 49	28	Di Thomas Kardos. 2003. Easy Science Demos & Labs: Physics. Edition 2. Walch Publishing.						
50 51	29	Available at:						
52	30	http://books.google.it/books?id=GBGS34Aew1gC&printsec=frontcover&source=gbs_summary						
53 54	31	<u>r&cad=0</u>						
55 56	32							
57	33	EFSA - European Environment and Health Information System. (2009). Panel on Dietetic						
58 59	34	Products, Nutrition and Allergies (NDA), Scientific Opinion on the appropriate age for						
60	35	introduction of complementary feeding of infants. EFSA Journal; 7(12): 1423 [38 pp.].						

Food Additives and Contaminants

4 5	1	doi:10.2903/j.efsa.2009.1423. (Available online:						
6	2	http://www.efsa.europa.eu/it/efsajournal/doc/1423.pdf)						
7 8	3							
9 10	4	ENHIS 2007. Exposure of children to chemical hazards in food 2007. European Environment						
11	5	and Health Information System .WHO. 8 May 2007. Copenhagen (Available at:						
12	6	http://www.enhis.org/object_document/o4736n27387.html).						
14 15	7							
16	8	European Commission. 2006a. Commission Directive 2006/141/EC of 22 December 2006 on						
17	9	infant formulae and follow-on formulae and amending Directive 1999/21/EC (OJ L 401,						
19 20	10	30.12.2006(Available at:						
21	11	http://ec.europa.eu/food/food/labellingnutrition/children/formulae_en.htm).						
22 23	12							
24 25	13	European Commission. 2006b Commission Directive 2006/125/EC of 5 December 2006 on						
26	14	processed cereal-based foods and baby foods for infants and young children (OJ L 339/16,						
27 28	15	6.12.2006). (Available at: <u>http://eur-</u>						
29	16	lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2006:339:0016:0035:EN:PDF).						
30 31	17							
32 33	18	EUROSTAT. Population and social conditions. [Internet]. Available at:						
34	19	http://epp.eurostat.ec.europa.eu/portal/page/portal/population/data/database . Acceded 20 July						
35 36	20	2009.						
37 38	21							
39	22	FAO. 2004. Food and nutrition technical report series 1: Human energy requirements Report of						
40 41	23	a Joint FAO/WHO/UNU Expert Consultation, Rome, 17-24 October. Available at:						
42	24	http://www.fao.org/docrep/007/y5686e/y5686e00.HTM .						
43 44	25							
45 46	26	FDA/Center for Food Safety & Applied Nutrition [Internet]. Available at:						
47	27	http://www.cfsan.fda.gov/~comm/tds-toc.htmlAcceded 15April 2009.						
48 49	28							
50 51	29	FFT - Food for Thought. Real-time Food & Drink Market Data Provider [Internet]. Geneva						
52	30	(Switzerland). Available at: http://www.fft.com/fftt/servlet/fftt/template/Index.vm. Acceded						
53 54	31	15 April 2009.						
55 56	32							
57	33	FSANZ - Food Standards Australia New Zealand. 22nd Australian Total Diet Study. [Internet].						
58 59 60	34	Available at:						

1 2									
3 4	1	http://www.foodstandards.gov.au/newsroom/publications/22ndaustraliantotaldietstudy/index.cf							
5 6	2	m. Acceded 5 September 2009							
7	3								
8 9	4	GEMS Food. Total diet studies: A recipe for safer food. [Internet] 2005. Geneva: World							
10 11	5	Health Organization. Available at:							
12	6	http://www.who.int/foodsafety/chem/TDS_recipe_2005_en.pdf Acceded 5 September 2009.							
13	7								
15 16	8	Hoppe C, Mølgaard C, Thomsen BL, Juul A and Michaelsen KF. 2004. Protein intake at 9							
17	9	month of age is associated with body size but not with body fat in 10-v-old Danish children Am							
18	10	J Clin Nutr. 79(3 March): 494-501.							
20 21	11								
22	12	MAFF. UK Food Surveillance Information Sheets, 1994 Index to maff, UK food surveillance							
23 24	13	information sheets [Internet]. 1994. Available at:							
25 26	14	http://archive.food.gov.uk/maff/archive/food/infsheet/1994/no34/34diet.htm. Acceded 16 April							
27	15	<u>2009</u> .							
28 29	16								
30 31	17	Michaelsen K, Weaver L, Branca F and Robertson A. 2003. Feeding and nutrition of infants and							
32	18	young children. Guidelines for the WHO European Region, with emphasis on the former Soviet							
33 34	19	countries. WHO Regional Publications, European Series, No. 87. Copenhagen: WHO -World							
35 26	20	Health Organization.							
30 37	21								
38 39	22	Mills A, Tyler H. 1992. Food and nutrient intakes of British infant aged 6-12 months. HMSO:							
40	23	MAFF. London (UK).							
41	24								
43 44	25	Pandelova ME, Levy Lopez W, Kasham S, Leclercq C, Piccinelli R & Schramm K–W. 2009a.							
45	26	Assessment of energy intake of infants exclusively fed with infant formulae available on the							
46 47	27	European market. Int J Food Sci Nutr. 60: 212-219.							
48 49	28								
50	29	Pandelova M, Kasham S, Henkelmann B, Leclercq C, Piccinelli R, Schramm K-W. 2009b.							
51 52	30	PCDD/F and dioxin-like PCB level in infant formulae available on the EU market. Proceeding							
53	31	of the 29th International Symposium on Halogenated Persistent Organic Pollutants (POPs),							
55	32	August 23-28, 2009, Beijing, China.							
56 57	33								
58 50									
59 60									

Research and Market. Baby Foods Market Assessment 2006. [Internet]. Oct. 2006. Dublin
 (Ireland). Available at: <u>http://www.researchandmarkets.com/reportinfo.asp?report_id=461819</u>.

3 Acceded 16 April 2009.

Riu A, Balaguer P, Perdu E, Pandelova M, Piccinelli R, Gustafsson J-A, Leclercq C, Schramm
K-W, Dagnino S, Debrauwer L, Cravedi J-P, Zalko D. 2008. Characterisation of bioactive
compounds in infant formulas using immobilised recombinant estrogen receptor-α affinity
columns. Food Chem Toxicol. 46: 3268-3278.

SCF- Scientific Committee on Food. 2003. SCF/CS/NUT/IF/65 Final 18 May 2003. Report of
the Scientific Committee on Food on the revision of essential requirements of infant formula
and follow-on formula.

Société Suisse de Pédiatrie. 2002. Plan of feeding for newborn baby, infant and toddler up to 36
months in good health and without allergy risk. rev.11.2002. [Internet]. Pediatrica. Vol.12 n. 5.
Available at: <u>http://www.swiss-paediatrics.org/paediatrica/vol12/n5/iplan-gfi.htm</u>. Acceded15
September 2009.

WHO -World Health Organization. 1999. Consultations and Workshops: GEMS/Food Total
Diet Studies, Report of a Joint USFDA/WHO, WHO/SDE/FOS/99.9. Papers presented at:
International Workshop on Total Diet Studies in cooperation with the Pan American Health
Organization - Food safety programme department of protection of the human environment
World Health Organization. 26 July- 6 August 1999; Kansas City, Missouri, USA.

WHO -World Health Organization. 2003. Global strategy for infant and young child
feeding. Geneva; (Available at:

27 <u>http://www.who.int/nutrition/publications/gs_infant_feeding_text_eng.pdf</u>)

WHO Multicentre Growth Reference Study Group. 2006.WHO Child Growth Standards:
Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass
index-for-age: Methods and development. Geneva: World Health Organization.

		MONTHS								
	FOOD CATEGORIES	1	2	3	4	5	6	7	8	9
	INFANT FORMULAE FOLLOW-ON									
<u>_</u>	INFANT CEREALS:									
1	INFANT CEREALS without gluten									
2 3	INFANT CEREALS with gluten								-	
5	INFANT CEREALS with milk									
	PASTA									T
	BISCUITS									T
	FRUIT:									
	PUREED FRUIT							-	-	
	FRUITJUICE									
	VEGETABLES/POTATOES									
	DIII SES						 		·	
	MEAT							-	-	╞
	FISH									
	DAIRY BASED DESSERT:									
	DAIRY BASED DESSERT with fruit									
	DAIRY BASED DESSERT with chocolate									
	ADDED FATS (OIL/BUTTER etc)	_								-
Figure 1. Time of inbaskets of "Other ba	clusion of baby food categories: final schec by foods".	dule	use	d in	the c	desig	gn of	f the	CA	SC.

Figure 1. Time of inclusion of baby food categories: final schedule used in the design of the CASCADE market baskets of "Other baby foods".

Table 1. Calculated average energy requirement (kcal/day) by age class in infants aged 0 to 10 months, SCF data from 5th to 9th month and calculated daily energy intake from "Infant formulae and follow-on formulae" (follow-on) and "Other baby foods"

Months	Calculated average body weight ⁽¹⁾ at 50^{th} percentile	Energy requirement in formula- fed infants (kcal/kg bw/day) ⁽²⁾	Calculated average energy requirement (kcal/day) ⁽³⁾	SCF ⁽⁴⁾ energy intake for infants fed with both complementary foods and follow- on formula	Amount (ml/day) of infant formulae (follow-on) (4)	Energy intake (kcal/day) from "infant formulae" (follow-	Energy intake (kcal/day) from "Other baby foods"
	(Kg)	Uw/uay)		(kcal/day)		(5) on) ⁽⁵⁾	
0-1	3.80	120	456	•			
1–2	4.85	109	529				
2–3	5.73	100	573				
3–4	6.40	87	557				
4–5	6.95	86	598				
5–6	7.40	84	622	650-690	600	425	197
6–7	7.78	81	630	650-730	400	283	347
7–8	8.10	81	656	768-816	240	170	486
8–9	8.40	81	680		240	170	510
9–10	8.70	81	705		240	170	535

(1) Average body weight in infants of age between month x and month x+1 was estimated by calculating
(1) the average of 50th percentile at month x and month x+1 in both females and males, according to WHO 2006 and 2) by calculating the average between the values obtained in females and males.
(2) EAO 2004

(2) FAO 2004.

(3) For infants of 5 months of age, calculated average energy requirement for 5-6 month (622 kcal/day) was used instead of that for 4-5 month (598 kcal/day). The same was performed for 6th to 9th months of age.
(4) SCF 2003.

(5) Energy intake from infant formulae was calculated multiplying the average of the energy content reported on the labels of sampled products (on average, 100 g of rehydrated infant formulae follow-on contain 71 kcal) by the amount of infant formula (ml/day) consumed by infants during the 5h, 6th and 7h month (SCF 2003); the amount of infant formulae follow-on consumed at 8th and 9 th month is assumed to be the same as that consumed at 7 th month (one baby bottle).

(6)Energy intake from "Other baby foods" were calculated by subtracting the energy intake deriving from "Infant formulae" from the calculated average energy requirements

Table 2. Categories and typologies of baby foods chosen to design baskets from 5th and 9th month.

FOOD CATEGORY	TYPOLOGY OF FOOD PRODUCTS	SPECIFIC FEATURES OF THE TYPOLOGIES OF PRODUCTS
INFANT FORMULAE AND FOLLOW-ON FORMULAE	Infant formula follow-on	Infant formula follow on - in powder
INFANT CEREALS:		
	Infant cereals without milk and without gluten	Rice - in powder
	Infant cereals without milk and with gluten	Multicereals or cereals with gluten- in powder
	Infant cereals with milk	Infant cereals with milk and fruit or honey - in powder
PASTA	Rice or pasta with meat and vegetables	Rice or pasta (around 20%) with meat (veal and/or chicken and/or turkey) (around 20%) and vegetables (around 60%) - where available with potatoes - in jars (ready to eat)
BISCUITS	Infant biscuits	Biscuits with gluten - No granular
FRUIT:		
	Pureed fruit	Fruit mix with two fruit among apple and/or pear and/or banana and/or apricot- in jars (ready to eat)
	Fruit juice	Apricot or other if not available - liquid (ready to eat)
PULSES/VEGETABLES	Pulses with vegetables	Pulses (around 40%) and vegetables (around 60%)- where available with potatoes - in jars (ready to eat)
FISH	Fish with vegetables	Fish (sole and/or salmon and/or codfish) (around 40%) with vegetables (around 60%) - where available with potatoes- in jars (ready to eat)
MEAT	Meat with vegetables	Meat (veal and/or chicken and/or turkey) (around 40%) with vegetables (around 60%) - where available with potatoes - in jars (ready to eat)
DAIRY BASED DESSERT	Dairy based dessert	Milk or cheese based dessert with forest fruit or chocolate - in jars (ready to eat)

Table 3. Theoretical diets elaborated in terms of main food categories and of typologies of commercial baby foods (months 5-9).

DIET IN TERM MAIN FOOD CATI	IS OF EGORIES		DIET IN TERMS OF TYPOLOGIES OF FOOD PRODUCTS				
CATEGORIES	Daily consumption (g/day) ⁽¹⁾	Energy (kcal)	TYPOLOGIES	Daily consumption (g/day)	Energy (kcal)		
5 th MONTH							
Infant formulae and follow-on formulae	89 (600) ⁽²⁾	425	Infant formulae follow-on (powder)	89 (600) ⁽²⁾	425		
Infant cereals ⁽³⁾	12	44	Infant cereals without milk and without gluten (powder)	20	74		
Fruit	80	59	Pureed fruit (jar)	80	59		
Vegetables Meat	40 40	19 33	Meat ⁽⁵⁾ +vegetables +potatoes (jar)	100	63		
Added fats (oil/butter etc) ⁽⁴⁾	5	41					
Total		622	Total		622		
6 th MONTH							
Infant formulae and follow-on formulae	60 (400) ⁽²⁾	283	Infant formulae follow-on (powder)	60(400) ⁽²⁾	283		
Infant cereals ⁽³⁾	33	126	Infant cereals without milk and without gluten (powder) Infant cereals without milk and with gluten (powder)	15 15	57 56		
Fruit	100	74	Pureed fruit (jar) Fruit juice (liquid)	75 25	56 14		
Vegetables/ Pulses	80	39	Meat ⁽⁵⁾ +vegetables +potatoes (jar)	200	126		
Meat	80	65	Pulses+vegetables+potatoes (jar)	75	38		
Added fats (oil/butter etc) ⁽⁴⁾	5	41					
Total		630	Total		630		

(1) The theoretical diets were designed in order to ensure that the energy intake from "Infant formulae and follow-on formulae" and from "Other baby foods" were in line with those reported in table 2.

(2) In parenthesis consumption of "Infant formulae" expressed in ml/day of product ready to be consumed.

(3) The amount of this food category was calculated by difference from the calculated energy intake from "Other baby foods" (Table 3).

(4) "Added fats (oil/butter etc..)" were not sampled, since they are present in commercial baby foods mixed dishes and no such products specifically designed for babies were found to be available in any EU country.

(5) The total quantity of jars containing meat was calculated to ensure the daily consumption of the portion size of meat considered in the theoretical diets elaborated in terms of main food categories.

1	
2	
3	
4	
5	
6	
7	
8	
0	
3	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
20	
29	
21	
31	
32	
33	
34	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
<u>4</u> 0	
73 50	
50	
51	
5Z	
53	
54	
55	
56	
57	
58	
59	
60	

Table 3 (continued). Theoretical diets elaborated in terms of main food categories and of typologies of commercial baby foods (months 5-9).

DIET IN TEI MAIN FOOD CA	RMS OF ATEGORIES	DIET IN TERMS OF TYPOLOGIES OF FOOD PRODUCTS			
CATEGORIES	Daily consumption (g/day) ⁽¹⁾	Energy (kcal)	TYPOLOGIES	Daily consumption (g/day)	Energy (kcal)
7 th MONTH					
Infant formulae and follow-on formulae	36 (240) ⁽²⁾	170	Infant formulae follow- on (powder)	36 (240) ⁽²⁾	170
			Infant cereals without milk and without gluten (powder)	10	38
Infant cereals ⁽³⁾	27	101	Infant cereals without milk and with gluten (powder)	20	75
			Infant cereals with milk and fruit (powder)	10	36
Fruit	160	118	Pureed fruit (jar) Fruit juice (liquid)	120 40	89 22
Pasta	20	71	Pasta + meat +vegetables +potatoes (jar)	100	63
Vegetables/ Pulses	100	49	Pulses +vegetables	75	38
Meat	80	65	+polatoes (Jar) Meat ⁽⁵⁾ +vegetables +potatoes (jar)	150	95
Dairy based dessert	40	40	Dairy based dessert (with fruit) (jar)	40	39
Added fats (oil/butter etc) ⁽⁴⁾	5	41			
Total		656	Total		665
8 th MONTH					
Infant formulae and follow-on formulae	36 (240) ⁽²⁾	170	Infant formulae follow- on (powder)	36 (240) ⁽²⁾	170
			Infant cereals without milk and without gluten (powder)	9	33
Infant cereal ⁽³⁾	26	99	Infant cereals without milk and with gluten (powder)	18	66
			Infant cereals with milk and fruit (powder)	9	32
Biscuits	8	33	Baby biscuits	8	33
Fruit	160	118	Pureed fruit (jar)	120	89
1 1011	100	110	Fruit juice (liquid)	40	22
Pasta	20	71	Pasta + meat +vegetables +potatoes (jar)	100	63
Vegetables/ Pulses	100	49	Pulses + vegetables + Potatoes (jar)	75	38

Total		680	Total		
Added fats (oil/butter etc) ⁽⁴⁾	5	41			
Dairy based dessert	40	40	Dairy based dessert (with fruit) (jar)	4	
Fish	30	18	Fish +vegetables +potatoes (jar)	7	
Meat	50	41	Meat ⁽⁵⁾ +vegetables +potatoes (jar)	7	

(1) The theoretical diets were designed in order to ensure that the energy intake from "Infant formulae and follow-on formulae" and from "Other baby foods" were in line with those reported in table 2.

(2) In parenthesis consumption of "Infant formulae" expressed in ml/day of product ready to be consumed.

(3) The amount of this food category was calculated by difference from the calculated energy intake from "Other baby foods" (Table 3).

(4) "Added fats (oil/butter etc..)" were not sampled, since they are present in commercial baby foods mixed dishes and no such products specifically designed for babies were found to be available in any EU country.

(5) The total quantity of jars containing meat was calculated to ensure the daily consumption of the portion size of meat considered in the theoretical diets elaborated in terms of main food categories.

Table 3 (continued). Theoretical diets elaborated in terms of main food categories and of typologies of commercial baby foods (months 5-9).

DIET IN TE MAIN FOOD C	RMS OF ATEGORIES	DIET IN TERMS OF TYPOLOGIES OF FOOD PRODUCTS				
CATEGORIES	Daily consumption (g/day) ⁽¹⁾	Energy (kcal)	TYPOLOGIES	Daily consumption (g/day)	Energy (kcal)	
9 ^h MONTH						
Infant formulae and follow-on formulae	36 (240) ⁽²⁾	170	infant formulae follow-on (powder)	36 (240) ⁽²⁾	170	
			Infant cereals without milk and without gluten (powder)	8	28	
Infant cereals ⁽³⁾	22	83	Infant cereals without milk and with gluten (powder)	15	56	
			Infant cereals with milk and fruit (powder)	8	27	
Biscuits	8	33	Baby biscuits	8	33	
F '4	1(0	110	Pureed fruit (jar)	120	89	
Fruit	160	118	Fruit juice (liquid)	40	22	
Pasta	20	71	Pasta + meat +vegetables +potatoes (jar)	100	63	
Vegetables/ Pulses	100	49	Pulses + vegetables + potatoes (jar)	75	38	
Meat	50	41	Meat ⁽⁵⁾ +vegetables +potatoes (jar)	75	47	
Fish	30	18	Fish +vegetables +potatoes (jar)	75	49	
	80	80	Dairy based dessert (with fruit) (jar)	60	59	
Dairy based dessert	80	80	Dairy based dessert (with chocolate) (jar)	20	20	
Oil/butter ⁽⁴⁾	5	41				
Total		705	Total		701	

(1) The theoretical diets were designed in order to ensure that the energy intake from "Infant formulae and follow-on formulae" and from "Other baby foods" were in line with those reported in table 2.

(2) In parenthesis consumption of "Infant formulae" expressed in ml/day of product ready to be consumed.

(3) The amount of this food category was calculated by difference from the calculated energy intake from "Other baby foods" (Table 3).

- (4) "Added fats (oil/butter etc..)" were not sampled, since they are present in commercial baby foods mixed dishes and no such products specifically designed for babies were found to be available in any EU country.
- (5) The total quantity of jars containing meat was calculated to ensure the daily consumption of the portion size of meat considered in the theoretical diets elaborated in terms of main food categories.

http://mc.manuscriptcentral.com/tfac Email: fac@tandf.co.uk

		DIET				Ν	MARKET BASKE	ET				
Typology	Grams	% Total diet	% "Other baby foods"	Sampling country	Sampled products	Description	Brand	Market share (%)	Corrected market share $(\%)^{(2)}$	Product by typology (%)	"Other baby foods" basket (%)	"Infant formulae"
Milk based infant formula	89	68.3										
tollow on				Germany Germany	Product 1 Product 2		BRAND A BRAND A	27.9	13.95 13.95	17.0 17.0		11 11
				Germany	Product 3		BRAND B	26.5	26.5	32.2		22
				France Sweden	Product 4 Product 5		BRAND D BRAND F	9.3 5.9	9.3 5.9	7.2		7 4 2
				Germany	Product 7		BRAND N	3.4	3.4	4.1		2
				France Italy	Product 8 Product 9		BRAND G BRAND I_1	3 2.7	3 0.9	3.6 1.1		2 0
				Italy Italy	Product 10 Product 11 Total		BRAND I_3 BRAND I_2	823	0.9 0.9 82 3	1.1 1.1 100.0		0 0 6!
Infant cereals without gluten and without milk	20	11.9	37.7	France	Product 12	rice vanilla flavour	BRAND F			100.0	37.8	12
Pureed Fruit	80	9.6	30.2			peaches and						
				Germany	Product 13	apple and apricots	BRAND B			50.0	15.1	4
				Germany	Product 14 Total	apricots, apple	BRAND B			50.0 100.0	15.1 30.2	4. 9
Meat +vegetabl es +potatoes	100	10.2	32.1					2				_
				Italy Italy Italy	Product 15 Product 16 Product 17 Product 18	beef/chicken beef/chicken beef/chicken mixed vagetables	BRAND I_1 BRAND I_3 BRAND I_2 BRAND I_1			16.7 16.7 16.7 50.0	5.3 5.3 5.3	1 1 1 5
				nary	Troduct 18	mixed vegetables	DRAND I_I			100.0	22.0	1

Table 4. EU basket of "Infant formulae and follow-on formulae" and "Other baby foods" at month 5, as sampled within the CASCADE project – with milk based infant formula follow-on.

(1) Data referred to 2007, purchased from the Company "Food for Thought" (FFT)

5 (<u>http://www.fft.com/fftt/servlet/fftt/template/Index.vm</u>).

(2) In the event that more than one product was available for one brand, the percentage of each brand was
equally divided by the number of products.
(2) Mustice of the product of the produc

(3) Meat jars of the sampled company do not contein vegetables. As a consequence the typologies "meat +

 $_{60}^{59}$ vegetables +potatoes" was obtained by combining meat jars and vegetables jars.

Table 5. Number of baskets created and of products purchased within the CASCADE project.

)	Types of baskets	Typologies of infant formulae	Number of BASKETS created	Number of PRODUCTS purchased	Country where the products were purchased
<u> </u>	"INFANT FORMULAE AND				
	FOLLOW_ON FORMULAE"				
	Infant formulae (0-4 months)	Milk based	1		
		Soya based	1		
		Hypo Allergenic	1		
	Follow-on (after 4 th month)	Milk based	1		
		Soya based	1		
		Hypo Allergenic	1		
		TOTAL	6	_	
				18	Germany
				4	UK
EU				9	France
				1	Portugal
				3	Sweden
				7	Italy
				42	TOTAL
	"OTHER BABY FOODS"		9	_	
				10	Germany
				9	Italy
				1	France
				1	Spain
				1	UK
				22	TOTAL
	"INFANT FORMULAE AND		9		
	FOLLOW_ON FORMULAE"				
	Infant formulae (0-4 months)	Milk based	4		
		Soya based	4		
		Hypo Allergenic	4		
	Follow-on (after 4 month)	Milk based	4		
		Hypo Allergenic	4		
		TOTAL	24		
NATION	AL			6	Italy
				5	Slovak Republic
				5	Spain Swadan
				4 20	TOTAL
	"OTHER BABY FOODS"		4	20	101111
				4	Italy
				3	Slovak Republic
				3	Spain
				3 13	Sweden TOTAI
				13	IUIAL