



HAL
open science

Immunomodulatory effects of Diclofenac in leukocytes through the targeting of Kv1.3 voltage-dependent potassium channels

Núria Villalonga, Miren David, Joanna Bielańska, Teresa González, David Parra, Concepció Soler, Núria Comes, Carmen Valenzuela, Antonio Felipe

► To cite this version:

Núria Villalonga, Miren David, Joanna Bielańska, Teresa González, David Parra, et al.. Immunomodulatory effects of Diclofenac in leukocytes through the targeting of Kv1.3 voltage-dependent potassium channels. *Biochemical Pharmacology*, 2010, 80 (6), pp.858. 10.1016/j.bcp.2010.05.012 . hal-00608940

HAL Id: hal-00608940

<https://hal.science/hal-00608940>

Submitted on 16 Jul 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Immunomodulatory effects of Diclofenac in leukocytes through the targeting of Kv1.3 voltage-dependent potassium channels

Authors: Núria Villalonga, Miren David, Joanna Bielańska, Teresa González, David Parra, Concepció Soler, Núria Comes, Carmen Valenzuela, Antonio Felipe



PII: S0006-2952(10)00351-5
DOI: doi:10.1016/j.bcp.2010.05.012
Reference: BCP 10567

To appear in: *BCP*

Received date: 7-4-2010
Revised date: 10-5-2010
Accepted date: 11-5-2010

Please cite this article as: Villalonga N, David M, Bielańska J, González T, Parra D, Soler C, Comes N, Valenzuela C, Felipe A, Immunomodulatory effects of Diclofenac in leukocytes through the targeting of Kv1.3 voltage-dependent potassium channels, *Biochemical Pharmacology* (2008), doi:10.1016/j.bcp.2010.05.012

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Immunomodulatory effects of Diclofenac in leukocytes through the targeting of
Kv1.3 voltage-dependent potassium channels**

Núria Villalonga^a, Miren David^b, Joanna Bielańska^a, Teresa González^b, David Parra^c,
Concepció Soler^c, Núria Comes^a, Carmen Valenzuela^b and Antonio Felipe^{a,*}

^aMolecular Physiology laboratory, Departament de Bioquímica i Biologia Molecular, Institut de Biomedicina, Universitat de Barcelona. Avda Diagonal 645, E-08028 Barcelona, Spain. ^bInstituto de Investigaciones Biomédicas “Alberto Sols”, CSIC-UAM. C/Arturo Duperier 4, E-28029 Madrid, Spain. ^cDepartament de Patologia i Terapèutica Experimental, Facultat de Medicina, Campus de Bellvitge, Universitat de Barcelona, Feixa Llarga s/n, E-08907, L'Hospitalet de Llobregat, Barcelona, Spain.

e-mail addresses: nvillalonga@gmail.com (N. Villalonga), mdavid@iib.uam.es (M. David), astygmata@gmail.com (J. Bielanska), tgonzalez@iib.uam.es (T. González), dparrago@gmail.com (D. Parra), concepciosoler@ub.edu (C. Soler), nuriacomés@ub.edu (N. Comes), cvalenzuela@iib.uam.es (C. Valenzuela), afelipe@ub.edu (A. Felipe).

***Correspondence should be addressed to:** Antonio Felipe, Departament de Bioquímica i Biologia Molecular, Universitat de Barcelona. Avda Diagonal 645, E-08028 Barcelona, Spain. Phone: 34-934034616; fax: 34-934021559; e-mail: afelipe@ub.edu

ABSTRACT

1
2
3
4
5 Kv1.3 plays a crucial role in the activation and proliferation of T-lymphocytes and
6
7 macrophages. While Kv1.3 is responsible for the voltage-dependent potassium current
8
9 in T-cells, in macrophages this K^+ current is generated by the association of Kv1.3 and
10
11 Kv1.5. Patients with autoimmune diseases show a high number of effector memory T
12
13 cells that are characterized by a high expression of Kv1.3 and Kv1.3 antagonists
14
15 ameliorate autoimmune disorders in vivo. Diclofenac is a non-steroidal anti-
16
17 inflammatory drug (NSAID) used in patients who suffer from painful autoimmune
18
19 diseases such as rheumatoid arthritis. In this study, we show that diclofenac impairs
20
21 immune response via a mechanism that involves Kv1.3. While diclofenac inhibited
22
23 Kv1.3 expression in activated macrophages and T-lymphocytes, Kv1.5 remained
24
25 unaffected. Diclofenac also decreased iNOS levels in Raw 264.7 cells, impairing their
26
27 activation in response to lipopolysaccharide (LPS). LPS-induced macrophage migration
28
29 and IL-2 production in stimulated Jurkat T-cells were also blocked by pharmacological
30
31 doses of diclofenac. These effects were mimicked by Margatoxin, a specific Kv1.3
32
33 inhibitor, and Charybdotoxin, which blocks both Kv1.3 and Ca^{2+} -activated K^+ channels
34
35 ($K_{Ca3.1}$). Because Kv1.3 is a very good target for autoimmune therapies, the effects of
36
37 diclofenac on Kv1.3 are of high pharmacological relevance.
38
39
40
41
42
43
44
45
46
47

48 **Key words:** Voltage-dependent K^+ channels, inflammation, leukocytes,
49
50 immunomodulation, non-steroidal anti-inflammatory drugs
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 **Abbreviations:** BMDM, bone marrow-derived macrophages; COX, cyclooxygenase;
2 CTX, Charybdotoxin; Kv, voltage-dependent K⁺ channels; LPS, lipopolysaccharide;
3
4 MgTx, Margatoxin; NSAID, non-steroidal anti-inflammatory drug; PHA,
5
6 phytohematogglutinin; PMA, phorbol ester; T_{EM}, Effector memory T-cells.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

1
2 Voltage-dependent potassium channels (Kv) play a crucial role in the
3 repolarization of membrane potential in nerve and muscle, controlling their excitability.
4
5 In addition, they are involved in the proliferation and activation of leukocytes [1, 2].
6
7 Particularly, Kv1.3 performs a key function in the immune system, controlling and
8 leading to proliferation and IL-2 synthesis [3]. Effector memory T-cells (T_{EM}), which
9 express high levels of Kv1.3, are present in autoimmune disorders [4]. In this context,
10 antagonizing the activity of Kv1.3 with highly specific peptides reverses and prevents
11 experimental autoimmune encephalomyelitis in rats [5]. Rheumatoid arthritis is a T-cell
12 mediated autoimmune disease that causes severe inflammation and pain in the
13 articulations and surrounding tissues. Kv1.3-high T_{EM} are abundant in these patients,
14 and selective blockage of Kv1.3 ameliorates arthritis in rats [6]. In addition, Kv1.3,
15 together with Kv1.5, is also present in macrophages, where it controls proliferation and
16 activation processes [2, 7-9]. Macrophages turn the immune response toward
17 inflammation or tolerance. These cells, which also act as antigen-presenting cells,
18 modify the cytokine milieu and the intensity of T-cell signaling. Therefore, the
19 inhibitory effects of Kv1.3 blockers on macrophages and microglia may account, in
20 part, for the benefits of Kv1.3-based therapies.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 Diclofenac (2-(2,6-dichloranilino) phenylacetic acid) is a non-steroidal anti-
44 inflammatory drug (NSAID) that inhibits COX-2 (cyclooxygenase) activity and
45 prevents the synthesis of inflammatory mediators. This compound is therapeutically
46 used for pain-suffering patients, including those with rheumatoid arthritis or
47 osteoarthritis. Diclofenac shows anti-nociceptive effects that are reversed by potassium
48 channel blockers [10]. This effect seems to be dependent on the mechanism of each
49 NSAID, as other compounds of this family do not involve potassium channel activity
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 [11]. ATP-dependent potassium channels (K_{ATP}) play a key role in the anti-nociceptive
2 effect of diclofenac on rat paw [12]. Moreover, diclofenac, and not other NSAIDs,
3 inhibits sodium currents in sensory neurons, which contributes to its analgesic activity
4 [13]. Similarly, diclofenac's anticonvulsant properties involve the activation of
5 heterotetrameric KCNQ2/KCNQ3 channels, which are responsible for the M current
6 that modulates neuronal excitability and firing [14]. Finally, diclofenac also has many
7 effects in the immune system, including impairing migration and accumulation of
8 leukocytes [15-18] and diminishing NO production by macrophages [19, 20]. We here
9 demonstrate that Kv1.3 is involved in the anti-inflammatory mechanism of diclofenac.
10 Our results have clinical relevance because Kv1.3 is considered a very important target
11 in therapies for anti-immune disorders. Our data provide new information about
12 additional benefits of the use of diclofenac in the pharmacological treatment of the
13 progression of rheumatoid arthritis.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 **2. Materials and methods**

32 *2.1. Animals and cell culture*

33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Murine bone marrow-derived macrophages (BMDM) were isolated as described elsewhere [2, 21]. Cells were cultured in DMEM containing 20% FBS and 30% supernatant of L-929 fibroblast (L-cell) conditioned media as a source of Macrophage-Colony Stimulating Factor (MCSF). Raw 264.7 macrophages and Jurkat T-lymphocytes were cultured in RPMI culture media, containing 10% FBS and supplemented with 10 U/mL penicillin and streptomycin and 2 mM L-glutamine. All animal handling was approved by the Ethics Committee of the University of Barcelona and was in accordance with EU regulations.

2.2. Chemicals

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Treatments were done in the culture media for 18-24 h. BMDM, Raw macrophages, and Jurkat T-lymphocytes were treated with 100 ng/ml, 1 µg/ml and 10 µg/ml of lipopolysaccharide (LPS, Sigma-Aldrich, Madrid, Spain), respectively. Jurkat T-cells were also stimulated with 5 µg/ml phytohematogglutinin (PHA, Sigma-Aldrich), with 80 nM phorbol ester (PMA, Sigma-Aldrich) or with combinations of both. Controls with DMSO, the vehicle for PMA, were performed to discard possible non-specific side effects. Diclofenac sodium salt (Sigma-Aldrich) was used at concentrations of 1.5 and 15 µM. Margatoxin (MgTx) (10 nM) and Charybdotoxin (CTX) (100 nM) were used to inhibit Kv1.3 alone or Kv1.3 and K_{Ca}3.1, respectively (Alomone, Israel).

2.3. RNA isolation and RT-PCR analysis

Total RNA was isolated by using Nucleospin RNA II (Invitrogen, Carlsbad, CA), which contains DNaseI. Transcriptor reverse transcriptase (Roche, Barcelona, Spain) was used for cDNA synthesis, according to manufacturer's instructions. Primers for PCR analysis were: Kv1.5, accession number NM012972 (F: 5'-GGATCACTCCATCACCAG-3'; R: 5'-GGCTTCCTCCTCCTTCCTTG-3'); and Kv1.3, accession number NM019270 (F: 5'-CTCATCTCCATTGTCATCTTCTGC-3'; R: 5'-TTGAAGTTGGAAACAATCAC-3'). In all cases, negative controls were performed in the absence of the reverse transcriptase reaction.

Real-time PCR primers were: Kv1.5 (F: 5'-TCCGACGGCTGGACTCAATAA-3'; R: 5'-CAGATGGCCTTCTAGGCTGTG-3'); and Kv1.3 (F: 5'-AGTATATGGTGA TCGAAGAGG-3'; R: 5'-AGTGAATATCTTCTTGATGTT-3'). LightCycler FastStart DNA MasterPLUS SYBR Green I (Roche) was used, and the reactions were performed under the following conditions: 95°C for 5 s, 55°C for 8 s, and 72°C for 9 s, preceded by

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

10 min at 95°C and followed by 10 min at 95°C. Melting curves were performed to verify the specificity of the product, and primers for 18S were included as an internal reference [8, 9].

2.4. Protein extracts and western blot

Cells were washed twice in cold phosphate-buffered saline (PBS) and lysed with lysis solution (50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1% Triton X-100, and 1 mM EDTA) supplemented with 0.1% pepstatin, 0.1% leupeptin, 0.1% PMSF, and 0.1% aprotinin as protease inhibitors. Cells were disrupted by passing the lysate 10 times through a 25 G syringe. Lysates were centrifuged at 14000 rpm for 10 min to discard cellular debris.

Total proteins (50 µg) were boiled in Laemmli SDS loading buffer and separated on a 10% SDS-PAGE gel. They were transferred to PVDF membranes (Immobilon-P, Millipore, Billerica, MA) and blocked in 5% dry milk supplemented with 0.05% Tween-20 PBS before immunoreaction. Polyclonal antibodies against Kv1.3 (1/200, Alomone), Kv1.5 (1/500, Alomone), and inducible Nitric Oxide Synthase (1/500, Santa Cruz Biotechnology, Santa Cruz, CA) were used. A monoclonal anti-β-actin antibody (Sigma) was used as a control. The specificity of the antibodies was tested with control antigen peptides provided by the manufacturer. Secondary antibodies were purchased from BioRad (Hercules, CA) (anti-rabbit, 1/3000; anti-mouse, 1/10000).

2.5. IL-2 production measurements

IL-2 production in Jurkat T-lymphocytes was measured with an ELISA kit (eBioscience, San Diego, CA), following the manufacturer's instructions. Cells were centrifuged at 1200 rpm for 4 min, and the supernatant was used to quantify the IL-2

1 concentration. Reactions were performed in 96-well plates coated with the capture
2 antibody. After blocking, standards and samples were incubated with Detection
3 Antibody followed by HRP-Streptavidin. H_3PO_4 (1M) was used as stop solution. Plates
4 were read at 450 nm.
5
6
7
8
9

10 11 12 *2.6. Migration assays*

13
14 Confluent BMDM plated in 6-well dishes were scratched with a single-edged
15 razor blade cut. The scratch began at the middle of the dish and extended over an area 7-
16 10 mm wide. The migrating cells were counted at 20x magnification at three distinct
17 sites along the wound (1 mm² each), beginning at the scratch line and extending as far
18 out as the cells had migrated. The mean distance between the wound edge and the nuclei
19 of migrating cells was also measured. The results are given as the percentage of
20 migrating cells and migrated distance (μM).
21
22
23
24
25
26
27
28
29
30
31
32
33

34 *2.7. Electrophysiological Recordings*

35
36 Whole-cell currents were recorded at room temperature (21-23 °C) using the
37 whole cell patch-clamp technique with an Axopatch 1C amplifier (Axon Instruments,
38 Foster City, CA). Micropipettes of 2-4 M Ω were pulled from borosilicate glass capillary
39 tubes (GD-1; Narishige, Tokyo, Japan) using a P-87 puller (Sutter Instruments, Novato,
40 CA). Electrodes were filled with the following solution (in mM): 80 K-aspartate, 42
41 KCl, 3 phosphocreatine, 10 KH_2PO_4 , 3 MgATP, 5 HEPES-K, and 5 EGTA-K (adjusted
42 to pH 7.25 with KOH). The extracellular solution contained (in mM) 136 NaCl, 4 KCl,
43 1.8 CaCl_2 , 1 MgCl_2 , 10 HEPES-Na, and 10 D-glucose, and it was adjusted to pH 7.40
44 with NaOH. Macrophages and Jurkat T-cells were clamped to a holding potential of -80
45 mV. The current-voltage (IV) protocol used consisted of 15 pulses ranging from -80 to
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 +60 mV, in 10 mV steps. Each pulse was 250 ms long. Peak amplitude (pA) was
2 normalized by capacitance values (pF). Data analysis was performed using the
3
4 CLAMPFIT utility of PCLAMP 9.2 and Origin 7.0.3 (Microcal Software, Beijing, China).
5
6
7
8

9 2.8. Statistical analysis

10
11 Real-time PCR, protein expression, migration and IL-2 concentration
12 experiments were repeated at least three times in duplicate. Electrophysiological
13 recordings were performed in 6-10 cells from at least 3 independent experiments.
14
15 Values are expressed as mean \pm S.E.M. The significance of any observed differences
16
17 was established by Student's *t* test (Graph Pad, PRISM 3.0, La Jolla, CA). A value of
18
19 $p < 0.05$ was considered significant.
20
21
22
23
24
25
26
27
28

29 3. Results

30
31 First we characterized Kv channels expressed in Raw 264.7 macrophages and Jurkat
32 T- lymphocytes. Delayed rectifier K^+ currents were evoked in Raw (Fig. 1A) and Jurkat
33 cells (Fig. 1B) by depolarizing pulses (250 ms) from a holding potential of -80 mV to +
34
35 60 mV in 10 mV steps. More depolarizing potentials were needed to open channels in
36
37 Raw cells (Fig. 1C). While the threshold potential for activation was -30 mV in T-cells,
38
39 it was approximately -10 mV in macrophages. Moreover, current density was 10-fold
40
41 higher in T-lymphocytes than in macrophages. RT-PCR experiments showed that while
42
43 Jurkat T-cells only expressed Kv1.3, Raw macrophages also presented Kv1.5 (Fig. 1D).
44
45
46 These results were further confirmed by western blot analysis (Fig. 1E). Furthermore, as
47
48 expected, Kv currents in macrophages and T-lymphocytes were fully inhibited by 10
49
50 nM MgTX and 100 nM CTX (not shown).
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Kv1.3 plays an important role in macrophage proliferation and activation, and LPS-induced activation increases Kv1.3 expression. Macrophages, with or without 1 $\mu\text{g/ml}$ LPS, were treated with diclofenac for 24 h and voltage-dependent K^+ currents were analyzed. The presence of 15 μM diclofenac led to a 50% decrease in the current amplitude in macrophages without LPS treatment (Fig 2A). Similar results were obtained in LPS-treated macrophages. However, lower doses of diclofenac (1.5 μM) did not produce apparent changes in activated cells. Surprisingly, not only LPS activation, but also the presence of diclofenac, resulted in a reduction in cell size (Fig 2B). Therefore, we calculated the peak amplitude normalized to cell size (pA/pF) (Fig. 2C). We observed that LPS increased peak current density in the absence of diclofenac ($p < 0.05$ vs. control). In addition, while diclofenac (15 μM) had no effects on control Raw cells, it markedly reduced the current density in LPS-stimulated macrophages ($p < 0.05$ vs. LPS). Therefore, 15 μM diclofenac led to a significant reduction of K^+ currents in LPS-activated macrophages.

Because the association between Kv1.3 and Kv1.5 generates the major voltage-dependent K^+ current in macrophages [7, 9], we wanted to check whether this diclofenac-dependent current decrease was due to changes in the expression of either of these Kv channels. Real-time PCR analysis and, as shown in Fig. 3A, Kv1.3 mRNA expression was increased 2.5-fold with LPS treatment. However, the presence of 15 μM diclofenac counteracted this increase. A lower concentration of diclofenac (1.5 μM) had no effect on either control or LPS-treated macrophages. Unlike Kv1.3, Kv1.5 expression was not affected by any of these treatments. We also undertook western blot experiments to check Kv1.3 and Kv1.5 protein expression (Fig. 3B-D). While 15 μM diclofenac reversed the LPS-induced increase in Kv1.3, no changes were observed for Kv1.5 (Fig. 3 B-D). iNOS expression was checked as a marker of cell activation (Fig. 3

1 B and E). As expected, LPS-induced activation led to an important increase in iNOS
2 expression. As with Kv1.3, 15 μ M diclofenac - but not 1.5 μ M - decreased iNOS levels
3 in LPS-stimulated Raw cells.
4
5

6
7 Macrophage activation was impaired by diclofenac. Therefore, we next wanted to
8 examine its effects on cell migration. To that end, we used primary cultures of bone
9 marrow-derived macrophages (BMDM) because their morphological changes during
10 activation allow for better measurements in this kind of studies. Representative pictures
11 of cell-migration assays are shown in Fig. 4 (A-F). The percentage of migrating
12 macrophages was calculated (Fig. 4G and H). Almost no migration was observed in
13 resting macrophages (Fig 4A and G), regardless of the presence of diclofenac (data not
14 shown). In contrast, LPS stimulated cell migration (Fig 4D and G), and this effect was
15 reversed by the presence of 15 μ M diclofenac (Fig 4F and G). No significant changes
16 were observed with 1.5 μ M diclofenac on LPS-stimulated BMDM (Fig 4E and G). We
17 also measured the migrated distance (Fig. 4H). In the presence of LPS, both 1.5 μ M and
18 15 μ M diclofenac reduced the travelled distance. Margatoxin (MgTx) potently blocks
19 Kv1.3, and Charybdotoxin (CTX) inhibits both Kv1.3 and intermediate-conductance
20 Ca^{2+} -activated K^+ channels ($\text{K}_{\text{Ca}3.1}$), which are present in leukocytes [22, 23]. Both
21 toxins markedly reduced the number of LPS-activated migrating cells (Fig 4B and C).
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 According to these results, diclofenac impairs macrophage activation and migration,
45 and these effects are mediated by Kv1.3. Macrophages are involved in the inflammatory
46 response by modifying the extracellular milieu and acting as antigen-presenting cells.
47 Therefore, the level of activation of macrophages strongly influences the immune
48 response carried out by T-lymphocytes. In this context, we also evaluated the effects of
49 diclofenac on T-lymphocytes. We measured IL-2 production, as a marker of T-cell
50 activation, in response to different stimuli (Fig. 5). Fig. 5A shows IL-2 production
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 induced by the treatment of Jurkat T cells with LPS or PMA. Neither of these stimuli
2 significantly enhanced IL-2 synthesis; on the contrary, PHA triggered a 10-fold increase
3 of IL-2 production (Fig. 5B). Diclofenac, as well as MgTx and CTX, inhibited T-cell
4 IL-2 production of PHA-stimulated lymphocytes. The effect of double stimulation with
5 PHA and PMA is shown in Fig. 5C. Although PMA by itself did not increase IL-2
6 production (Fig. 5A), it enhanced PHA-induced stimulation, as the double stimulation
7 produced a 450-fold induction of IL-2 synthesis. In this case, only 15 μ M diclofenac
8 significantly inhibited T-cell activation. Similar results were obtained with the Kv1.3-
9 blockers MgTx and CTX.

10
11
12
13
14
15
16
17
18
19
20
21
22 Our results with the Kv1.3-blockers suggested that this protein could be involved in
23 the T-cell immunosuppressive response generated by pharmacological relevant doses of
24 diclofenac. Therefore, we evaluated the effects of diclofenac on Kv1.3 expression in
25 Jurkat cells. Real-time PCR experiments indicated that a 15 μ M dose of diclofenac, but
26 not a 1.5 μ M dose, decreased gene expression of Kv1.3 in both PHA- and (PHA +
27 PMA)-stimulated cells (Fig. 6A). Similar to the effect observed in macrophages, 15 μ M
28 diclofenac showed no significant effect on non-stimulated T-cells. Protein levels were
29 also analyzed (Fig 6B-E). Representative western blots are shown in Fig. 6B and D,
30 together with densitometric quantifications of different experiments (Fig. 6C and E). As
31 can be seen, no changes were observed in PHA (Fig. 6B and C) or (PHA + PMA)-
32 treated cells (Fig. 6D and E). However, in all cases, 15 μ M diclofenac significantly
33 decreased Kv1.3 levels in stimulated Jurkat T-lymphocytes.

34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 **4. Discussion**

55
56
57 This study demonstrates that Kv1.3 contributes to the anti-inflammatory effects of
58 diclofenac, which could provide additional benefits in therapies against autoimmune
59
60
61
62
63
64
65

1 diseases associated with inflammation and pain. Moreover, our data also show that the
2 modulation of Kv1.3 in macrophages and T-lymphocytes could lead to important
3 alterations in the immune response in these disorders.
4
5

6
7 A relationship between potassium channels and diclofenac has been previously
8 described [10, 14]. However, this is the first time that this connection has been
9 established in the immune system. Diclofenac functions as a channel opener, activating
10 ATP-dependent K^+ channels and triggering anti-nociceptive effects by reducing pain
11 sensation in response to different stimuli [12]. This is in agreement with the diclofenac-
12 dependent abrogation of leukocyte activation that we found. In fact, unlike inward-
13 rectifier ATP-dependent K^+ channels, diclofenac inhibits the outward-rectifier K^+
14 channel Kv1.3. Similar opposite mechanisms occur between Kir2.1 and Kv1.3 during
15 pro-inflammatory insults [2, 24]. Diclofenac also activates KCNQ2/KCNQ3
16 heteromeric channels in rat cortical neurons, but again, the final effect is an attenuation
17 of their excitability, which is the basis for its anticonvulsant properties [14].
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33
34 Besides K^+ channels, Na^+ currents are also affected by diclofenac. Diclofenac
35 inhibits Na^+ currents in rat myoblasts [25], which are necessary for muscle development
36 and myoblast differentiation [26, 27]. Furthermore, diclofenac also blocks Na^+ currents
37 in sensory neurons producing analgesia [13]. All of these observations further support
38 the immunosuppressive role of diclofenac via Kv1.3.
39
40
41
42
43
44
45

46 We suggest that diclofenac modulates Kv1.3 via a still unidentified mechanism in
47 leukocytes. Unlike diclofenac, most NSAIDs involve no K^+ channel activity in their
48 anti-nociceptive effects, indicating distinct mechanisms of action [11]. Several NSAIDs
49 have been tested, and only flufenamic acid partially mimics diclofenac's effects on Na^+
50 currents in neurons [13]. Authors argue that the structure of NSAIDs, and not the
51 inhibition of COX, must be involved [13]. In this context, diclofenac specifically
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 decreases Kv1.3, but not Kv1.5, expression. Therefore, by changing the
2 heterotetrameric composition of the channelosome, leukocytes may be able to fine-tune
3 the immune response [9].
4
5

6
7 There is growing experimental support for the existence of a diclofenac-dependent
8 attenuation of the immune response. Diclofenac inhibits the accumulation of leukocytes
9 and the formation of lesions [16, 17]. In addition, it also blocks the migration of
10 leukocytes [15, 18]. Our results also indicated a reduction in macrophage migration.
11
12 With 15 μM diclofenac, the number of migrating LPS-activated BMDM decreased, as
13 did the distance they travelled. Paradoxically, some studies have shown that diclofenac
14 enhances B-16 murine melanoma cell migration [28]. However, these discrepancies
15 could be explained by the NSAID concentrations used. Liu et al. (2004) used 100 μM
16 diclofenac, whereas clinically available doses of diclofenac reach a maximum
17 concentration of 10 $\mu\text{g/ml}$ in blood, which is equivalent to 30 μM [20]. Similarly high
18 concentrations of diclofenac (100 μM) also affect rat cerebellar granule cells in ways
19 contrary to what our study shows, activating transient outward K^+ currents [29]. In our
20 study, we tested two concentrations of diclofenac, both of clinical relevance, which in
21 all cases trigger dose-dependent effects. Higher doses (150 μM) cause cell death, thus
22 making it impossible to perform functional studies (data not shown).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 MgTx and CTX, which block Kv1.3, mimicked the effects of diclofenac on LPS-
45 stimulated migrating cells. Kv1.3, which interacts with $\beta 1$ -integrin, is important in
46 leukocyte migration because its opening activates T-cell integrin function [30]. Our
47 results further support a Kv1.3-mediated cell migration mechanism and indicate that
48 diclofenac partially impairs macrophage migration via Kv1.3.
49
50
51
52
53
54
55
56

57 Similar to our results, diclofenac partially blocks NO production in Raw cells [19,
58 20]. Although both Kv1.3 and Kv1.5 are present in macrophages, Kv1.3 is the main
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

isoform responsible for cell activation [8, 9]. The incubation of BMDM with Margatoxin partially blocks iNOS induction in response to LPS or TNF α , which involve Kv1.3 in the activating response [2, 8]. We demonstrate that the decrease in iNOS is concomitant with a reduction in Kv1.3. Kv currents are generated mainly by Kv1.3 homomeric complexes and by heterotetrameric combinations of Kv1.3 and Kv1.5 in macrophages [7, 9, 31]. Low Kv1.3/Kv1.5 ratios, as found in Raw cells, impair pharmacological treatments based on Kv1.3 and dull the control of proliferation and activation [8, 9]. Therefore, our results show a potent effect of diclofenac on Kv1.3-driven macrophage activation. This is the first time that a relationship has been shown between a NSAID mechanism and Kv1.3 in the immune system.

Kv1.3 plays a key role in the initiation of the immune response in T-lymphocytes. Kv1.3 is the responsible for triggering the calcium influx necessary for IL-2 synthesis. The Ca²⁺-activated K⁺ channel K_{Ca}3.1, the main K_{Ca} channel in T-cells, sustains this calcium influx [1, 3]. Charybdotoxin, which inhibits Kv1.3 and K_{Ca}3.1, blocks IL-2 production and mitogenesis in T-cells [32]. However, K_{Ca}2.2, which is charybdotoxin-insensitive [33-35], is the main K_{Ca} channel involved in the Ca²⁺ signaling in Jurkat cells [36]. Therefore, CTX solely targets Kv1.3 in our model. Diclofenac inhibited IL-2 production in stimulated T-cells, leading to the attenuation of the immune response. Furthermore, both MgTx and CTX mimicked these effects, suggesting an important role of Kv1.3 in diclofenac-induced immunosuppression. We also demonstrated the inhibitory effect of diclofenac on Kv1.3 expression. However, PHA and PHA + PMA stimulation did not increase Kv1.3, possibly due to the cell's origin. Only effector memory T cells (T_{EM}) markedly increase their number of Kv1.3 channels when they receive activation stimuli, acquiring the Kv1.3^{high}K_{Ca}3.1^{low} phenotype. Naïve and central memory T-lymphocytes only show a very slight increase in Kv1.3 [35, 37].

1 Jurkat T cells come from a human leukemia, and they do not differentiate into T_{EM}. In
2 addition, unlike macrophages, Kv1.3 function in T-lymphocytes is not always
3 coincident with an increase in expression [38, 39]. However, our data show that Kv1.3
4 expression is reduced by diclofenac, impairing T-cell function. This result may be
5 important for rheumatoid arthritis patients who are treated with diclofenac because of its
6 anti-inflammatory and pain killing effects. Parallel actions of this NSAID on Kv1.3
7 could be attenuating the lymphocyte-enhanced immune response because Kv1.3 is a
8 very good target for T-cell-mediated autoimmune disorders [6]. Selective blockade of
9 Kv1.3 reverses these types of disorders *in vivo* [6].

10
11 In summary, we provide strong evidence of the inhibitory effects of diclofenac on
12 Kv1.3 expression in macrophages and T-lymphocytes. These changes are selective, as
13 Kv1.5 is not altered by this treatment. Kv1.3 inhibition is accompanied by impaired
14 inflammatory response in these cells. In macrophages, we observed a decrease in iNOS
15 expression and a blockage in their migration in response to LPS. In T-cells, IL-2
16 production is reduced. These changes were mimicked by treatment with Margatoxin and
17 Charybdotoxin, which block Kv1.3. Our results suggest that diclofenac ameliorates
18 rheumatoid arthritis, and probably other inflammatory autoimmune diseases, by
19 targeting Kv1.3 in leukocytes. Therefore, in addition to the well-known anti-
20 inflammatory and anti-nociceptive effects of diclofenac, this drug could also aid in the
21 prognosis of the disease by ameliorating its progression.

22 **Acknowledgements**

23
24 This work was supported by the Ministerio Ciencia e Innovación (MICINN),
25 Spain (BFU2008-00431 and CSD2008-00005 to AF, SAF2007-65868 and FIS
26 D06/0014/0006 to CV, and BFU2009-07501 to CS). NV and JB held a fellowship from
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 the Ministerio Ciencia e Innovación (MICINN). MD held a FIS RD06/0014/0006
2 contract. N.C. and T.G. supported by the Juan de la Cierva program (MICINN). TG
3 actually, holds a JAE-Doc contract from CSIC. We thank the editorial assistance of
4 American Journal Experts.
5
6
7
8
9

10 11 **REFERENCES**

- 12
13
14 [1] Cahalan MD, Wulff H, Chandy KG. Molecular properties and physiological
15 roles of ion channels in the immune system. *J Clin Immunol* 2001;21:235-52.
16
17 [2] Vicente R, Escalada A, Coma M, Fuster G, Sanchez-Tillo E, Lopez-Iglesias C,
18 et al. Differential voltage-dependent K⁺ channel responses during proliferation
19 and activation in macrophages. *J Biol Chem* 2003;278:46307-20.
20
21 [3] Cahalan MD, Chandy KG. Ion channels in the immune system as targets for
22 immunosuppression. *Curr Opin Biotechnol* 1997;8:749-56.
23
24 [4] Wulff H, Calabresi PA, Allie R, Yun S, Pennington M, Beeton C, et al. The
25 voltage-gated Kv1.3 K(+) channel in effector memory T cells as new target for
26 MS. *J Clin Invest* 2003;111:1703-13.
27
28 [5] Beeton C, Wulff H, Barbaria J, Clot-Faybesse O, Pennington M, Bernard D, et
29 al. Selective blockade of T lymphocyte K(+) channels ameliorates experimental
30 autoimmune encephalomyelitis, a model for multiple sclerosis. *Proc Natl Acad
31 Sci U S A* 2001;98:13942-7.
32
33 [6] Beeton C, Wulff H, Standifer NE, Azam P, Mullen KM, Pennington MW, et al.
34 Kv1.3 channels are a therapeutic target for T cell-mediated autoimmune
35 diseases. *Proc Natl Acad Sci U S A* 2006;103:17414-9.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [7] Vicente R, Escalada A, Villalonga N, Texido L, Roura-Ferrer M, Martin-Satue M, et al. Association of Kv1.5 and Kv1.3 contributes to the major voltage-dependent K⁺ channel in macrophages. *J Biol Chem* 2006;281:37675-85.
- [8] Villalonga N, Escalada A, Vicente R, Sanchez-Tillo E, Celada A, Solsona C, et al. Kv1.3/Kv1.5 heteromeric channels compromise pharmacological responses in macrophages. *Biochem Biophys Res Commun* 2007;352:913-8.
- [9] Villalonga N, David M, Bielanska J, Vicente R, Comes N, Valenzuela C, et al. Immunomodulation of voltage-dependent K⁺ channels in macrophages: molecular and biophysical consequences. *J Gen Physiol* 2010;135:135-47.
- [10] Ortiz MI, Torres-Lopez JE, Castaneda-Hernandez G, Rosas R, Vidal-Cantu GC, Granados-Soto V. Pharmacological evidence for the activation of K(+) channels by diclofenac. *Eur J Pharmacol* 2002;438:85-91.
- [11] Ortiz MI, Granados-Soto V, Castaneda-Hernandez G. The NO-cGMP-K⁺ channel pathway participates in the antinociceptive effect of diclofenac, but not of indomethacin. *Pharmacol Biochem Behav* 2003;76:187-95.
- [12] Alves DP, Tatsuo MA, Leite R, Duarte ID. Diclofenac-induced peripheral antinociception is associated with ATP-sensitive K⁺ channels activation. *Life Sci* 2004;74:2577-91.
- [13] Lee HM, Kim HI, Shin YK, Lee CS, Park M, Song JH. Diclofenac inhibition of sodium currents in rat dorsal root ganglion neurons. *Brain Res* 2003;992:120-7.
- [14] Peretz A, Degani N, Nachman R, Uziyel Y, Gibor G, Shabat D, et al. Meclofenamic acid and diclofenac, novel templates of KCNQ2/Q3 potassium channel openers, depress cortical neuron activity and exhibit anticonvulsant properties. *Mol Pharmacol* 2005;67:1053-66.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [15] Perianin A, Gougerot-Pocidallo MA, Giroud JP, Hakim J. Diclofenac sodium, a negative chemokinetic factor for neutrophil locomotion. *Biochem Pharmacol* 1985;34:3433-8.
- [16] Prescott MF, McBride CK, Venturini CM, Gerhardt SC. Leukocyte stimulation of intimal lesion formation is inhibited by treatment with diclofenac sodium and dexamethasone. *J Cardiovasc Pharmacol* 1989;14 Suppl 6:S76-81.
- [17] Marsolais D, Cote CH, Frenette J. Nonsteroidal anti-inflammatory drug reduces neutrophil and macrophage accumulation but does not improve tendon regeneration. *Lab Invest* 2003;83:991-9.
- [18] Martinez LL, Oliveira MA, Miguel AS, Rastelli VM, Cruz JW, Tostes RC, et al. Losartan attenuates the antimigratory effect of diclofenac in spontaneously hypertensive rats. *J Cardiovasc Pharmacol* 2005;46:190-9.
- [19] Berg J, Fellier H, Christoph T, Grarup J, Stimmeder D. The analgesic NSAID lornoxicam inhibits cyclooxygenase (COX)-1/-2, inducible nitric oxide synthase (iNOS), and the formation of interleukin (IL)-6 in vitro. *Inflamm Res* 1999;48:369-79.
- [20] Cho JY. Immunomodulatory effect of nonsteroidal anti-inflammatory drugs (NSAIDs) at the clinically available doses. *Arch Pharm Res* 2007;30:64-74.
- [21] Vicente R, Escalada A, Soler C, Grande M, Celada A, Tamkun MM, et al. Pattern of Kv beta subunit expression in macrophages depends upon proliferation and the mode of activation. *J Immunol* 2005;174:4736-44.
- [22] Garcia-Calvo M, Leonard RJ, Novick J, Stevens SP, Schmalhofer W, Kaczorowski GJ, et al. Purification, characterization, and biosynthesis of margatoxin, a component of *Centruroides margaritatus* venom that selectively

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- inhibits voltage-dependent potassium channels. *J Biol Chem* 1993;268:18866-74.
- [23] MacKinnon R, Reinhart PH, White MM. Charybdotoxin block of Shaker K⁺ channels suggests that different types of K⁺ channels share common structural features. *Neuron* 1988;1:997-1001.
- [24] Vicente R, Coma M, Busquets S, Moore-Carrasco R, Lopez-Soriano FJ, Argiles JM, et al. The systemic inflammatory response is involved in the regulation of K⁽⁺⁾ channel expression in brain via TNF-alpha-dependent and -independent pathways. *FEBS Lett* 2004;572:189-94.
- [25] Fei XW, Liu LY, Xu JG, Zhang ZH, Mei YA. The non-steroidal anti-inflammatory drug, diclofenac, inhibits Na⁽⁺⁾ current in rat myoblasts. *Biochem Biophys Res Commun* 2006;346:1275-83.
- [26] David M, Martinez-Marmol R, Gonzalez T, Felipe A, Valenzuela C. Differential regulation of Na^(v)beta subunits during myogenesis. *Biochem Biophys Res Commun* 2008;368:761-6.
- [27] Martinez-Marmol R, David M, Sanches R, Roura-Ferrer M, Villalonga N, Sorianello E, et al. Voltage-dependent Na⁺ channel phenotype changes in myoblasts. Consequences for cardiac repair. *Cardiovasc Res* 2007;76:430-41.
- [28] Liu LY, Hu CL, Ma LJ, Zhang ZH, Mei YA. ET-1 inhibits B-16 murine melanoma cell migration by decreasing K⁽⁺⁾ currents. *Cell Motil Cytoskeleton* 2004;58:127-36.
- [29] Liu LY, Fei XW, Li ZM, Zhang ZH, Mei YA. Diclofenac, a nonsteroidal anti-inflammatory drug, activates the transient outward K⁺ current in rat cerebellar granule cells. *Neuropharmacology* 2005;48:918-26.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [30] Levite M, Cahalon L, Peretz A, Hershkovich R, Sobko A, Ariel A, et al. Extracellular K(+) and opening of voltage-gated potassium channels activate T cell integrin function: physical and functional association between Kv1.3 channels and beta1 integrins. *J Exp Med* 2000;191:1167-76.
- [31] Vicente R, Villalonga N, Calvo M, Escalada A, Solsona C, Soler C, et al. Kv1.5 association modifies Kv1.3 traffic and membrane localization. *J Biol Chem* 2008;283:8756-64.
- [32] Price M, Lee SC, Deutsch C. Charybdotoxin inhibits proliferation and interleukin 2 production in human peripheral blood lymphocytes. *Proc Natl Acad Sci U S A* 1989;86:10171-5.
- [33] Grissmer S, Lewis RS, Cahalan MD. Ca(2+)-activated K⁺ channels in human leukemic T cells. *J Gen Physiol* 1992;99:63-84.
- [34] Hanselmann C, Grissmer S. Characterization of apamin-sensitive Ca(2+)-activated potassium channels in human leukaemic T lymphocytes. *J Physiol* 1996;496 (Pt 3):627-37.
- [35] Cahalan MD, Chandy KG. The functional network of ion channels in T lymphocytes. *Immunol Rev* 2009;231:59-87.
- [36] Fanger CM, Rauer H, Neben AL, Miller MJ, Rauer H, Wulff H, et al. Calcium-activated potassium channels sustain calcium signaling in T lymphocytes. Selective blockers and manipulated channel expression levels. *J Biol Chem* 2001;276:12249-56.
- [37] Chandy KG, Wulff H, Beeton C, Pennington M, Gutman GA, Cahalan MD. K⁺ channels as targets for specific immunomodulation. *Trends Pharmacol Sci* 2004;25:280-9.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
- [38] Ghanshani S, Wulff H, Miller MJ, Rohm H, Neben A, Gutman GA, et al. Up-regulation of the IKCa1 potassium channel during T-cell activation. Molecular mechanism and functional consequences. *J Biol Chem* 2000;275:37137-49.
- [39] Alizadeh AA, Staudt LM. Genomic-scale gene expression profiling of normal and malignant immune cells. *Curr Opin Immunol* 2000;12:219-25.

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

FOOTNOTES

NV and MD contributed equally. CV and AF contributed equally.

FIGURE LEGENDS

1
2
3
4
5 **Figure 1.** Macrophages and T lymphocytes express voltage-dependent K⁺ channels.
6
7 Representative traces of outward K⁺ currents evoked in Raw 264.7 macrophages (A)
8
9 and Jurkat T cells (B). Cells were held at -80 mV, and currents were elicited by
10
11 depolarizing pulses that increased in 10-mV steps (250 ms in duration) from -80 to +60
12
13 mV. (C) Current density *versus* voltage relationship of K⁺ currents in Jurkat T cells (○)
14
15 and Raw 264.7 macrophages (●). Current densities are much higher in Jurkat
16
17 lymphocytes. The inset highlights K⁺ currents in Raw. (D) mRNA expression of Kv1.3
18
19 and Kv1.5 in Raw 264.7 macrophages and Jurkat T lymphocytes. E, Kv1.3 and Kv1.5
20
21 protein expression in both cell types.
22
23
24
25
26
27
28

29 **Figure 2.** Diclofenac decreases current density in activated Raw 264.7 macrophages.
30
31 Cells were incubated for 18-24 h with LPS and/or diclofenac (DIC). (A) Representative
32
33 traces of outward K⁺ currents evoked in control and 15 μM diclofenac-treated Raw
34
35 264.7 cells (left). Right, representative traces of 1 μg/μl LPS-activated Raw cells with
36
37 and without diclofenac. Two concentrations of diclofenac were tested: 1.5 μM, grey
38
39 current record; 15 μM, black current record. Cells were held at -80 mV, and currents
40
41 were elicited by a depolarizing pulse of +60 mV lasting 250 ms. (B) Cell size calculated
42
43 as cell capacitance in pF for the analyzed populations of cells. *, p < 0.05 vs. control
44
45 without diclofenac. (C) Current density analysis of the different groups. Current
46
47 intensity (pA) at -60 mV was normalized by cell size (pF). Significant changes were
48
49 found for LPS (**, p<0.01 vs. control) and for 15 μM diclofenac treatment (*, p<0.05
50
51 vs. LPS) by Student's *t* test.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 3. Diclofenac selectively inhibits Kv1.3 expression in LPS-activated Raw 264.7 macrophages. Cells were incubated for 18-24 h in the presence or absence of 1 $\mu\text{g}/\mu\text{l}$ LPS with/without 1.5 and 15 μM diclofenac. (A) Kv1.3 (white) and Kv1.5 (black) mRNA expression by real-time PCR for the treatments indicated. Significant differences were found for Kv1.3 in LPS-activated macrophages (*, $p < 0.05$ vs. control) and for LPS-activated macrophages treated with 15 μM diclofenac (***, $p < 0.001$ vs. LPS). (B) Representative western Blot analysis of Kv1.3 and Kv1.5 protein expression for the treatments indicated. (C-E) Quantification of data in panel B: Kv1.3 (C), Kv1.5 (D) and iNOS (E). Significant differences were found for Kv1.3 in LPS-activated macrophages (*, $p < 0.05$ vs. control) and in LPS-activated macrophages treated with 15 μM diclofenac (***, $p < 0.001$ vs. LPS). Also iNOS showed significant changes after treatment with LPS (***, $p < 0.05$ vs. control) and 15 μM diclofenac (***, $p < 0.05$ vs. LPS). All values were normalized by β -actin expression and compared to the control in the absence of diclofenac (0). AU: arbitrary units. White bars, no addition; Grey bars, 1.5 μM diclofenac; black bars, 15 μM diclofenac.

40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 4. Diclofenac impairs migration of bone marrow derived macrophages (BMDM). Experimental procedures are described in the Materials and Methods section. Cells were incubated for 18-24 h with 100 $\text{ng}/\mu\text{l}$ of LPS in the presence or absence of 10 nM MgTx, 100 nM CTX or diclofenac (DIC) at the concentrations indicated. (A-F) Representative images of the different groups. Migrating cells are found at the right of the scratch. Bars represent 500 μm . (A) control cells without addition. (B) LPS-activated cells in the presence of MgTx. (C) LPS-activated cells in the presence of CTX. (D-F) LPS-activated cells in the absence (D) or the presence of 1.5 μM (E) and 15 μM (F) diclofenac. (G) Percentage of migrating cells. Significant differences were found in

1 LPS-activated BMDM treated with 15 μM diclofenac (**, $p < 0.01$ vs. LPS). (H)
 2 Distance travelled by the migrating macrophages (μm). Distance was measured between
 3 the scratch and the nucleus of the cell. Significant differences were found in LPS-
 4 activated BMDM treated with diclofenac at both concentrations (**, $p < 0.01$ and ***,
 5 $p < 0.001$ for LPS vs. 1.5 and 15 μM diclofenac, respectively). White bars, no diclofenac;
 6 Grey bars, 1.5 μM diclofenac; Black bars, 15 μM diclofenac.
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17

18 **Figure 5.** Inhibitory effects of diclofenac and K^+ channel blockers on IL-2 production
 19 by stimulated Jurkat T lymphocytes. Cells were stimulated with 10 $\mu\text{g}/\mu\text{l}$ LPS, 5 $\mu\text{g}/\text{ml}$
 20 PHA, 80 nM PMA and PHA (5 $\mu\text{g}/\text{ml}$) + PMA (80 nM) for 18-24 h. Simultaneously, 10
 21 nM Margatoxin (MgTx), 100 nM Charybdotoxin (CTX), or diclofenac (DIC) were
 22 added at the concentrations indicated. Supernatant was collected and processed for an
 23 ELISA analysis. (A) IL-2 production (pg/ml) of control (Ctrl), LPS and PMA-
 24 stimulated Jurkat cells. (B) IL-2 production (pg/ml) of control (Ctrl) and PHA-
 25 stimulated Jurkat T cells. Significant differences were found in PHA-stimulated cells
 26 (***, $p < 0.01$ vs. Ctrl), MgTx, CTX, 1.5 μM diclofenac and 15 μM diclofenac (***,
 27 $p < 0.001$; ***, $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$, respectively vs. PHA without
 28 diclofenac). (C) IL-2 production (ng/ml) of control (Ctrl) and (PHA + PMA)-stimulated
 29 Jurkat T lymphocytes. Significant differences were found in PHA + PMA-stimulated
 30 cells (***, $p < 0.001$ vs. Ctrl) and cells treated with MgTx, CTX and 15 μM diclofenac
 31 (*, $p < 0.05$; ***, $p < 0.001$; *, $p < 0.05$, respectively vs. PHA + PMA without diclofenac).
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54

55 **Figure 6.** Diclofenac decreases Kv1.3 expression in activated T-lymphocytes. Cells
 56 were stimulated with either 5 $\mu\text{g}/\text{ml}$ PHA or combination of PHA (5 $\mu\text{g}/\text{ml}$) + 80 nM
 57 PMA for 18-24 h. (A) Kv1.3 mRNA expression in control, PHA-stimulated and
 58
 59
 60
 61
 62
 63
 64
 65

1 PHA+PMA-stimulated Jurkat T cells in the presence or absence of diclofenac (DIC) at
2 the concentrations indicated. Significant differences were found with 15 μ M DIC in
3
4 PHA and PHA+PMA- stimulated lymphocytes (*, $p < 0.05$; ** $p < 0.01$ vs. PHA and
5
6 PHA+PMA in the absence of DIC (0), respectively). (B-E) Kv1.3 protein expression in
7
8 control, PHA-stimulated and PHA+PMA-stimulated Jurkat T cells in the presence or
9
10 absence of DIC. B, Representative western blot analysis in the presence or absence of
11
12 PHA. (C) Quantification of the relative Kv1.3 protein expression from experiments
13
14 presented in panel B. (D) Representative western blot analysis in the presence or
15
16 absence of PHA+PMA. (E) Quantification of the relative Kv1.3 protein expression from
17
18 experiments presented in panel D. All values were normalized by β -actin expression
19
20 and compared to the control in the absence of DIC (0). *, $p < 0.05$; **, $p < 0.01$ vs.
21
22 PHA+PMA and PHA in the absence (0) of DIC, respectively. White bars, no
23
24 diclofenac; Grey bars, 1.5 μ M diclofenac; Black bars, 15 μ M diclofenac. AU, arbitrary
25
26 units.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 1




Figure 2




Figure 3




Figure 4






Figure 5




Figure 6

