



HAL
open science

Hyperbolic Eisenstein series for geometrically finite hyperbolic surfaces of infinite volume.

Thérèse Falliero

► **To cite this version:**

Thérèse Falliero. Hyperbolic Eisenstein series for geometrically finite hyperbolic surfaces of infinite volume.. Publications of the Research Institute for Mathematical Sciences, In press. hal-00610249v4

HAL Id: hal-00610249

<https://hal.science/hal-00610249v4>

Submitted on 1 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hyperbolic Eisenstein series for geometrically finite hyperbolic surfaces of infinite volume.

Thérèse FALLIERO¹

classification: Primary 30F30, 32N10, 47A10; Secondary 53C20, 11M36, 11F12

keywords: Harmonic differential, Eisenstein series, Degenerating surfaces.

INTRODUCTION.

The spectrum of the Laplace–Beltrami operator for a compact Riemann surface is discrete, but this is no longer the case when M is non-compact. For example, when we remove one point from M , there arises a continuous part in the spectrum whose spectral measure is described by an Eisenstein series. The study of the limiting behaviour of the spectrum of the Laplace–Beltrami operator for a degenerating family of Riemann surfaces with finite area hyperbolic metric has been used to explain this phenomenon (see for example [27], [17], [14]). The present paper has one of its motivations in the general study of the approximation of Eisenstein series (see for example the question of Ji in [17], p. 308, line 15, concerning the approximation of Eisenstein series by suitable eigenfunctions of a degenerating family of hyperbolic Riemann surfaces). We hope to surround it via hyperbolic Eisenstein series (for results on degenerating Eisenstein series, see, for example [22], [23], [8], [9]). This article is organized as follows. In section 1 we provide the necessary background on geometrically finite hyperbolic surfaces of infinite volume. In section 2 we recall the definition and verify the convergence of hyperbolic Eisenstein series in the infinite volume case, like suggested in [20]. In section 3 we review the spectral decomposition for a geometrically finite hyperbolic surface of infinite volume. We obtain the analytic continuation of hyperbolic Eisenstein series and then the fact that this permits realizing a harmonic dual form to a simple closed geodesic on a geometrically finite hyperbolic surface of infinite volume (Theorem 3.1). In a similar way, in section 4, we realize a harmonic dual form to an infinite geodesic joining a pair of punctures (Theorem 4.2). Moreover in Section 5 we generalize the definition of hyperbolic Eisenstein series to the case of q -forms (see Section 5 formula (18)). After we study the limiting behavior of these q -forms on a degenerating family of geometrically finite hyperbolic surfaces of infinite volume (see Theorem 5.2 and Theorem 5.3). In particular we obtain a degeneration of hyperbolic Eisenstein series to horocyclic ones (Theorem 5.3(2)). Since this new result is at the heart of our motivation, we will be a little more precise.

Main Theorem

Let $(S_l)_l$ be a degenerating family of Riemann surfaces with infinite area hyperbolic metrics, S_l having a funnel F_l whose boundary geodesic is denoted c_l . We denote by $\Omega_{c_l} = \Omega_l$ the hyperbolic Eisenstein series of the Kudla–Millson theory associated to the pinching geodesic c_l (see Section 2), $S_0 = \Gamma \backslash H$ the component of the limiting surface containing the cusp ∞ of stabilizer Γ_∞ (see Section 5.1) and \mathcal{E}_∞ the horocyclic Eisenstein series associated to the limiting cusp defined, for $\operatorname{Re} s > 0$ by $\mathcal{E}_\infty(s+1, z) = \sum_{\Gamma_\infty \backslash \Gamma} \operatorname{Im}(\sigma z)^s d(\sigma z)$ (see (12)). For $s \in \{\operatorname{Re} s > -1/2\}$, except

¹Thérèse Falliero: Université d’Avignon et des Pays de Vaucluse, Laboratoire de mathématiques d’Avignon (EA 2151), F-84916 Avignon, France;

possibly a finite number of points in $(-1/2, 0)$, the family of C^∞ 1-forms on S_l depending meromorphically on s , $\frac{1}{l^{s+1}}\Omega_l(s, \cdot)$ converges uniformly on compact subsets of S_0 to $\frac{\Gamma(1+\frac{s}{2})}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{2}+\frac{s}{2})}\text{Im}\mathcal{E}_\infty(s+1, \cdot)$; where $\text{Im}\mathcal{E}_\infty(s+1, \cdot)$ is to be understood as $\text{Im}\mathcal{E}_\infty(s+1, z) = \frac{1}{2i}(\sum_{\Gamma_\infty \setminus \Gamma} \text{Im}(\sigma z)^s d(\sigma z) - \sum_{\Gamma_\infty \setminus \Gamma} \text{Im}(\sigma z)^s d(\overline{\sigma z}))$.

1. PRELIMINARY DEFINITIONS

Let us recall the standard analytic and geometric notations which will be used. In this paper, a surface is a connected orientable two-dimensional manifold, without boundary unless otherwise specified. We denote by H the hyperbolic upper half-plane endowed with its standard metric of constant Gaussian curvature -1 . A topologically finite (i.e. finite Euler characteristic) surface is a surface homeomorphic to a compact surface with finitely many points excised and a geometrically finite hyperbolic surface M is a topologically finite, complete Riemann surface of constant curvature -1 . We will require that M be of infinite volume, i.e. there exists a finitely generated, torsion free, discrete subgroup Γ of $PSL(2, \mathbb{R})$, unique up to conjugation, such that M is the quotient of H by Γ acting as Möbius transformations, Γ is a Fuchsian group of the second kind, and Γ has no elliptic elements. The group Γ admits a finite sided polygonal fundamental domain in H . We recall now the description of the fundamental domain of $M = \Gamma \backslash H$ (see [1]). Let $L(\Gamma)$ be the limit set of Γ , that is the set of limit points (in the Riemann sphere topology) of all orbits Γz for $z \in H$ and $O(\Gamma) = \mathbb{R} \cup \{\infty\} - L(\Gamma)$. As $L(\Gamma)$ is closed in $\mathbb{R} \cup \{\infty\}$, $O(\Gamma)$ is open and so can be written as a countable union $O(\Gamma) = \cup_{\alpha \in A} O_\alpha$ where the O_α are disjoint open intervals in $\mathbb{R} \cup \{\infty\}$. Now let $\Gamma_\alpha = \{\gamma \in \Gamma, \gamma(O_\alpha) = O_\alpha\}$. This is an elementary hyperbolic subgroup of Γ . The fixed points of Γ_α are exactly the end-points of O_α . There is a finite subset $\{\alpha(1), \alpha(2), \dots, \alpha(n_f)\} \subset A$ such that for $\alpha \in A$, O_α is conjugate to precisely one $O_{\alpha(j)}$ ($1 \leq j \leq n_f$). Let λ_α be the half-circle, lying in H , joining the end-points of O_α . Let Δ_α be the region in H bounded by O_α and λ_α . The Δ_α ($\alpha \in A$) are mutually disjoint. Let P be the set of parabolic vertices of Γ , and for $p \in P$, let Γ_p be the parabolic subgroup of Γ fixing p . There is a finite subset $\{p(1), p(2), \dots, p(n_c)\} \subset P$ such that Γ_p is conjugate to precisely one $\Gamma_{p(j)}$ ($1 \leq j \leq n_c$). A circle lying in H and tangent to ∂H at p is called a horocycle at p . We can construct an open disc C_p determined by a horocycle at $p \in P$ such that

- (i) if $p, q \in P, p \neq q$, then $C_p \cap C_q = \emptyset$,
- (ii) $\gamma(C_p) = C_{\gamma(p)}$ ($\gamma \in \Gamma$),
- (iii) $C_p \cap \Delta_\alpha = \emptyset$ ($p \in P, \alpha \in A$).

If we consider the set $H - (\cup_{p \in P} C_p \cup \cup_{\alpha \in A} \Delta_\alpha)$, we see that it is invariant under Γ . We can find a finite-sided fundamental domain D for the action of Γ on this set; D is relatively compact in H .

Proposition 1.1. *There is a fundamental domain D for Γ of the form*

$$D = K^* \cup \cup_{j=1}^{n_f} D_{\alpha(j)} \cup_{k=1}^{n_c} D_{p(k)}^*$$

where

- 1) K^* is relatively compact in H ,
- 2) $D_{\alpha(j)}$ is a standard fundamental domain of $\Gamma_{\alpha(j)}$ on $\Delta_{\alpha(j)}$,
- 3) $D_{p(k)}^*$ is a standard fundamental domain for $\Gamma_{p(k)}$ on $C_{p(k)}$.

We should note that $n_f \neq 0$ if and only if Γ is of the second kind.

The Nielsen region of the group Γ is the set $\tilde{N} = H - (\cup_{\alpha \in A} \Delta_\alpha)$, the truncated Nielsen region of Γ is $\tilde{K} = \tilde{N} - (\cup_{p \in P} C_p)$, $K = \Gamma \backslash \tilde{K}$ is called the compact core of M . So the surface $M = \Gamma \backslash H$ can be decomposed into a finite area surface with geodesic boundary N , called the Nielsen region, on which infinite area ends F_i are glued: the funnels. The Nielsen region N is itself decomposed into a compact surface K with geodesic and horocyclic boundary on which non compact, finite area ends C_i are glued: the cusps. We have $M = K \cup C \cup F$, where $C = C_1 \cup \dots \cup C_{n_c}$ and $F = F_1 \cup \dots \cup F_{n_f}$.

A hyperbolic transformation $T \in PSL(2, \mathbb{R})$ generates a cyclic hyperbolic group $\langle T \rangle$. The quotient $C_l = \langle T \rangle \backslash H$ is a hyperbolic cylinder of diameter $l = l(T)$. By conjugation, we can identify the generator T with the map $z \mapsto e^l z$, and we define Γ_l to be the corresponding cyclic group. A natural fundamental domain for Γ_l would be the region $\mathcal{F}_l = \{z \in H, 1 \leq |z| \leq e^l\}$. The y -axis is the lift of the only simple closed geodesic on C_l , whose length is l . The standard funnel of diameter $l > 0$, F_l , is the half hyperbolic cylinder $\Gamma_l \backslash H$, $F_l = (\mathbb{R}^+)_r \times (\mathbb{R}/\mathbb{Z})_x$ with the metric $ds^2 = dr^2 + l^2 \cosh^2(r) dx^2$.

We can always conjugate a parabolic cyclic group $\langle T \rangle$ to the group Γ_∞ generated by $z \mapsto z + 1$, so the parabolic cylinder is unique up to isometry. A natural fundamental domain for Γ_∞ is $\mathcal{F}_\infty = \{0 \leq \operatorname{Re} z \leq 1\} \subset H$. The standard cusp C_∞ is the half parabolic cylinder $\Gamma_\infty \backslash H$, $C_\infty = ([0, \infty]_r \times (\mathbb{R}/\mathbb{Z})_x$ with the metric $ds^2 = dr^2 + e^{-2r} dx^2$. The funnels F_i and the cusps C_i are isometric to the preceding standard models. We define the function r as the distance to the compact core K and the function ρ by

$$(1) \quad \rho(r) = \begin{cases} 2e^{-r} & \text{in } F \\ e^{-r} & \text{in } C \end{cases},$$

with ρ extended to a smooth non vanishing function inside K in some arbitrary way. We will adopt $(\rho, t) \in (0, 2] \times \mathbb{R}/l_j\mathbb{Z}$ as the standard coordinates for the funnel F_j , where t is the arc length around the central geodesic at $\rho = 2$.

For the cusp, our standard coordinates $(\rho, t) \in (0, 1] \times \mathbb{R}/\mathbb{Z}$ are based on the model defined by the cyclic group Γ_∞ . The cusp boundary is $y = 1$, so that $y = e^r$ and $\rho = 1/y$. We set $t \equiv x \pmod{\mathbb{Z}}$.

2. HYPERBOLIC EISENSTEIN SERIES ON A GEOMETRICALLY FINITE HYPERBOLIC SURFACE OF INFINITE VOLUME.

2.1. Return to the definition of Kudla and Millson of a hyperbolic Eisenstein series. In the following, M will denote an arbitrary Riemann surface and $L^2(M)$, the Hilbert space of square integrable 1-forms with inner product

$$(w_1, w_2) = \frac{1}{2} \int_M w_1 \wedge * \bar{w}_2,$$

and corresponding norm $\|\cdot\|_{L^2}$. The pointwise norm of a 1-form w is defined by $w \wedge * \bar{w} = \|w\|^2 * 1$ where $*1$ is the volume form.

Let c be a simple closed curve on M . We may associate with c a real smooth closed differential n_c with compact support such that

$$(2) \quad \int_c \omega = \int_M \omega \wedge n_c,$$

for all closed differentials ω . Since every cycle c on M is a finite sum of cycles corresponding to simple closed curves, we conclude that to each such c , we can associate a real closed differential n_c with compact support such that (2) holds.

Let a and b be two cycles on the Riemann surface M . We define the intersection number of a and b by

$$a.b = \int_M n_a \wedge n_b.$$

In [20], Kudla and Millson construct the harmonic 1-form dual to a simple closed geodesic on a hyperbolic surface of finite volume M in terms of Eisenstein series. Let us recall the definition:

Definition 2.1. *Let η be a simple closed geodesic or an infinite geodesic joining p and q . A 1-form α is dual to η if for any closed 1-form ω with compact support,*

$$\int_M \omega \wedge \alpha = \int_\eta \omega.$$

Or, equivalently, for any closed oriented cycle c' ,

$$(3) \quad \int_{c'} \alpha = \eta \cdot c'.$$

Kudla and Millson construct a meromorphic family of forms on M , called the hyperbolic Eisenstein series associated to an oriented simple closed geodesic c . Let \tilde{c} be a component of the inverse image of c in the covering $H \rightarrow M$ and Γ_1 the stabilizer of \tilde{c} in Γ . The hyperbolic Eisenstein series are expressed in Fermi coordinates in the following way for $\text{Re } s > 0$:

$$(4) \quad \Omega_c(s, z) = \Omega(s, z) = \frac{1}{k(s)} \sum_{\Gamma_1 \backslash \Gamma} \gamma^* \frac{dx_2}{(\cosh x_2)^{s+1}}, \quad k(s) := \frac{\Gamma(\frac{1}{2}) \Gamma(\frac{1}{2} + \frac{s}{2})}{\Gamma(1 + \frac{s}{2})}.$$

By applying an element of $SL(2, \mathbb{R})$, we may assume that \tilde{c} is the y -axis in H and that Γ_1 is generated by $\gamma_1 : z \mapsto e^l z$. The Fermi coordinates (x_1, x_2) associated to \tilde{c} are related to Euclidean polar coordinates by

$$\begin{aligned} r &= e^{x_1}, \\ \sin \theta &= \frac{1}{\cosh x_2}. \end{aligned}$$

We obtain

$$(5) \quad \Omega(s, z) = \frac{1}{k(s)} \sum_{\Gamma_1 \backslash \Gamma} \gamma^* (\sin \theta)^s d\theta.$$

Using the following equalities

$$\begin{aligned} \text{Im } \gamma z &= |\gamma z| \sin \theta(\gamma z) = |\gamma' z| \text{Im } z, \\ d \log \frac{\gamma z}{\gamma' z} &= 2id\theta(\gamma z), \end{aligned}$$

we find

$$(6) \quad \Omega(s, z) = \frac{1}{2ik(s)} (\text{Im } z)^s \left(\sum_{\Gamma_1 \setminus \Gamma} \left(\frac{\gamma'z}{|\gamma z|} \right)^s \frac{\gamma'z}{\gamma z} dz - \left(\frac{\gamma'z}{|\gamma z|} \right)^s \frac{\overline{\gamma'z}}{\gamma \bar{z}} d\bar{z} \right).$$

We denote, with a little misuse of notation, to simplify $\Omega(s, z) = \text{Im}(\Theta(s, z))$ with

$$(7) \quad \Theta(s, z) = \frac{1}{k(s)} \sum_{\Gamma_1 \setminus \Gamma} \gamma^* \left(\frac{y^s}{z|z|^s} dz \right) = i\Omega(s, z) - *\Omega(s, z),$$

keeping in mind that the complex conjugation is apply on z and “not on s ”.

At the end of their paper, they make the remark that “it is also interesting to consider the infinite volume case.”

2.2. The infinite volume case. We are going to verify that this definition is still meaningful in the case of a geometrically finite hyperbolic surface of infinite volume, $M = \Gamma \setminus H$, Γ being a Fuchsian group of the second kind without elliptic elements. We identify M locally with its universal cover H . By $d(z, w)$ we denote the hyperbolic distance from $z \in H$ to $w \in H$. For $z_0 \in H$ and $\epsilon > 0$, by $B(z_0, \epsilon) \subset H$ we denote the hyperbolic metric ball centered at z_0 with radius ϵ .

Proposition 2.1. *The hyperbolic Eisenstein series $\Omega(s, z)$ converges for $\text{Re } s > 0$, uniformly on compact subsets of H , is bounded on M , and represents a C^∞ closed form which is dual to c . Moreover, it is an analytic function of s in $\text{Re } s > 0$.*

The proof in the infinite volume case is as straightforward as in the finite volume case ([20], [10]), but for the convenience of the reader we give some details.

Proof. Recall first that if K is a compact subset of the fundamental domain D of Γ , then there exists $\eta > 0$ such that for all $z_0 \in K$, the balls $(B(\gamma z_0, \eta))_{\gamma \in \Gamma_1 \setminus \Gamma}$ are disjoint.

For a fundamental domain of Γ_1 , let us choose $\mathcal{D}_1 = \{z \in H : 1 \leq |z| \leq \beta\}$. After passing to ordinary Euclidean polar coordinates (r, θ) , with $\sigma = \text{Re } s$, where $\|\Omega\|$ denotes the pointwise norm of Ω , we obtain

$$\begin{aligned} \|\Omega(s, z)\| &\leq \frac{1}{|k(s)|} \sum_{\Gamma_1 \setminus \Gamma} \frac{1}{(\text{ch } x_2(\gamma z))^{\sigma+1}} \\ &\leq \frac{1}{|k(s)|} \sum_{\Gamma_1 \setminus \Gamma} \left(\frac{y}{r} \right)^{\sigma+1} (\gamma z) \leq \frac{1}{|k(s)|} \sum_{[\gamma] \in \Gamma_1 \setminus \Gamma, \gamma z \in \mathcal{D}_1} y^{\sigma+1}(\gamma z). \end{aligned}$$

Now y^s is an eigenfunction of all the invariant integral operators on H . Let $k(z, z')$ be the point-pair invariant defined by $k(z, z') = 1$ or 0 according as the distance between z and z' is less or no less than η . Then there exists a Λ_η independent of z_0 such that

$$\int_{B(z_0, \eta)} y^\sigma \frac{dx dy}{y^2} = \int_H k(z_0, z) y^\sigma \frac{dx dy}{y^2}$$

and

$$\int_{B(z_0, \eta)} y^\sigma \frac{dx dy}{y^2} = \Lambda_\eta y(z_0)^\sigma.$$

This is a particular case of a more general result we will need further on (see Proposition 5.2).

We write, as in [20], $R(T_1, T_2) = \{P \in \mathcal{D}_1 : T_1 < x_2(P) < T_2\}$. So for $T > 2\eta$,

$$\frac{1}{|k(s)|} \sum_{\gamma \in \Gamma_1 \setminus \Gamma, \gamma z \notin R(-T, T)} \frac{1}{(\text{ch}(x_2(\gamma z)))^{\sigma+1}} \leq \frac{1}{|k(s)|} \sum_{[\gamma] \in \Gamma_1 \setminus \Gamma, \gamma z \in \mathcal{D}_1 \setminus R(-T, T)} y^{\sigma+1}(\gamma z).$$

We need the following.

Lemma 2.1. *Let $\gamma \in \Gamma_1 \setminus \Gamma$, $z, \zeta \in H$ such that for $\gamma z \notin R(-T, T)$, $\gamma \zeta \in B(\gamma z, \eta)$. Then $\gamma \zeta \notin R(-T + 2\eta, T - 2\eta)$.*

Proof. Let $\pi : H \rightarrow \tilde{c}$ be the orthogonal projection on \tilde{c} . As π is 1-Lipschitzian, we have for the hyperbolic distance $d(\pi\gamma z, \pi\gamma \zeta) \leq d(\gamma z, \gamma \zeta) \leq \eta$. If $x_2(\gamma z) \geq T$,

$$T \leq d(\gamma z, \pi\gamma z) \leq d(\gamma z, \gamma \zeta) + d(\gamma \zeta, \pi\gamma \zeta) + d(\pi\gamma z, \pi\gamma \zeta).$$

Then

$$T - 2\eta \leq d(\gamma \zeta, \pi\gamma \zeta).$$

□

Then

$$\begin{aligned} \sum_{\gamma \in \Gamma_1 \setminus \Gamma, \gamma z \notin R(-T, T)} y^{\sigma+1}(\gamma z) &= \frac{1}{\Lambda_\eta} \sum_{\gamma \in \Gamma_1 \setminus \Gamma, \gamma z \notin R(-T, T)} \int_{B(\gamma z, \eta)} y^{\sigma+1} \frac{dx dy}{y^2} \\ &\leq \frac{1}{\Lambda_\eta} \int_{R^c(-T+2\eta, T-2\eta)} y^{\sigma+1} \frac{dx dy}{y^2} \end{aligned}$$

where $R^c(-T + 2\eta, T - 2\eta)$ is the complement in \mathcal{D}_1 of $R(-T + 2\eta, T - 2\eta)$. Note that if $\gamma z \notin R(-T, T)$, then $y(\gamma z) \leq \frac{\beta}{\text{ch} T}$, so

$$\begin{aligned} \sum_{\gamma \in \Gamma_1 \setminus \Gamma, \gamma z \notin R(-T, T)} y^{\sigma+1}(\gamma z) &\leq \frac{\beta}{\Lambda_\eta} \int_0^{\frac{\beta}{\text{ch}(T-2\eta)}} y^{\sigma-1} dy \\ &\leq \frac{\beta}{\Lambda_\eta \sigma} \left(\frac{\beta}{\text{ch}(T-2\eta)} \right)^\sigma. \end{aligned}$$

From this, there follows the uniform convergence of $\Omega(s, z)$ on compact subsets of H , uniformly on compact subsets of the half-plane $\text{Re } s > 0$.

We next show that $\Omega(s, z)$ is bounded on D . For this we use a very useful fundamental lemma (see [15], p. 178, [12], pp. 27, 214 note 30):

Proposition 2.2. *Suppose that $q > 1$. For any Fuchsian group Γ , there exists a $\mathcal{C}(q, \Gamma)$ such that for all $z \in H$,*

$$\sum_{\gamma \in \Gamma} \frac{y(\gamma z)^q}{[1 + |\gamma z|]^{2q}} \leq \mathcal{C}(q, \Gamma).$$

The constant $\mathcal{C}(q, \Gamma)$ depends only on q and Γ .

Proof. Let $z \in H$. Then there exists a system of representatives S of $\Gamma_1 \backslash \Gamma$ (depending on z) such that for all $\gamma \in S$, $|\gamma z| \leq \beta$. In effect, let $(\delta_1, \dots, \delta_n, \dots)$ be a system of representatives of $\Gamma_1 \backslash \Gamma$, then for all n there exists $p_n \in \mathbb{Z}$ such that $\gamma_1^{p_n} \delta_n(z) \in \mathcal{D}_1$.

Then

$$\begin{aligned} \sum_{\Gamma_1 \backslash \Gamma} \frac{y(\gamma z)^{\sigma+1}}{(1+\beta)^{2(\sigma+1)}} &\leq \sum_{\Gamma_1 \backslash \Gamma} \frac{y(\gamma z)^{\sigma+1}}{(1+|\gamma z|)^{2(\sigma+1)}} \\ &\leq \sum_{\Gamma} \frac{y(\gamma z)^{\sigma+1}}{(1+|\gamma z|)^{2(\sigma+1)}} \\ &\leq \mathcal{C}(\sigma+1, \Gamma) \end{aligned}$$

and the result follows. \square

Remark 2.1. *The constant $\mathcal{C}(q, \Gamma)$ depends solely on q and \mathcal{D} where \mathcal{D} is a tiny circular disk such that:*

$$\mathcal{D} \cup \partial \mathcal{D} \subset H \text{ and } T(\mathcal{D}) \cap \mathcal{D} = \emptyset \text{ for all } T \in \Gamma - \{I\}.$$

The fact that $\Omega(s, z)$ is dual to c follows straightforwardly from the construction of Kudla and Millson. \square

3. SPECTRAL DECOMPOSITION AND ANALYTIC CONTINUATION.

The aim is to realize the injection $H_c^1 \rightarrow \mathcal{H}^1$, where H_c^1 is the first de Rham cohomology group with compact support of M and \mathcal{H}^1 is the space of L^2 harmonic 1-forms of M . Recall that in our context, $\dim \mathcal{H}^1 = \infty$ (see [2], p. 27).

We are going to prove, as in [20], that the hyperbolic Eisenstein series have an analytic continuation. The essential difference from the finite volume case is the spectral decomposition of $L^2(M)$.

3.1. Spectral theory. For any non-compact geometrically finite hyperbolic surface M , the essential spectrum of the (positive) Laplacian Δ_M defined by the hyperbolic metric on M (the Laplacian on functions) is $[1/4, \infty)$ and this is absolutely continuous. The discrete spectrum consists of finitely many eigenvalues in the range $(0, 1/4)$. In the finite-volume case, one may also have embedded eigenvalues in the continuous spectrum, but these do not occur for infinite volume surfaces. Then if M has infinite volume, the discrete spectrum of Δ_M is finite (possibly empty). The exponent of convergence δ of a Fuchsian group Γ is defined to be the abscissa of convergence of the Dirichlet series

$$\delta = \inf \left\{ s > 0, \sum_{T \in \Gamma} e^{-sd(z, Tw)} < \infty \right\}$$

for some $z, w \in H$, where $d(z, w)$ again denotes the hyperbolic distance from $z \in H$ to $w \in H$.

Let Γ be a Fuchsian group of the second kind and $L(\Gamma)$ be its limit set. Then $0 < \delta < 1$ with $\delta > 1/2$ if Γ has parabolic elements. Patterson and Sullivan showed that δ is the Hausdorff dimension of the limit set when Γ is geometrically finite. Furthermore, if $\delta > 1/2$, then $\delta(1 - \delta)$ is the lowest eigenvalue of the Laplacian Δ_M . The connection with spectral theory was later extended to the case $\delta \leq 1/2$

by Patterson. In this case, the discrete spectrum of Δ_M is empty and δ is the location of the first resonance. For a detailed account of the spectral theory of infinite area surfaces, we refer the reader to [1].

3.2. Tensors and automorphic forms. This section introduces the notations used in the following subsection 3.3 and Section 5. Let M be a geometrically finite hyperbolic surface. Let z be a local conformal coordinate and $ds^2 = \rho|dz|^2$ the Poincaré metric. Let $\omega_M = T^*M$ be the holomorphic cotangent bundle of M and for any integers n and m , let $\mathcal{E}^{r,s}(M, \omega_M^n \otimes \bar{\omega}_M^m)$ be the vector space of smooth differential forms of type (r, s) on M with values in the line bundle $\omega_M^n \otimes \bar{\omega}_M^m$. For an integer q , a q -form (or q -differential) is an element of $T^q = \mathcal{E}^{0,0}(M, \omega_M^q)$, the space of tensors of type q on M , written locally as $f(z)(dz)^q$. M may be realized as $\Gamma \backslash H$, where H is the upper half-plane and Γ a discrete subgroup of $PSL(2, \mathbb{R})$. The hyperbolic metric on M induces the natural scalar product on T^q :

$$(\varphi, \psi) = \int_{\Gamma \backslash H} \varphi(z) \overline{\psi(z)} y^{2q-2} dx dy.$$

Let \mathfrak{H}^q be the L^2 -closure of T^q with respect to this scalar product.

We recall now the link between q -forms and automorphic forms of weight $2q$, also called q -automorphic forms for the following reason. As before, we make use of the uniformization theorem.

Using the notations of [11] and [7], set

$$j_\gamma(z) = \frac{(cz+d)^2}{|cz+d|^2} = \frac{cz+d}{c\bar{z}+d} = \left(\frac{\gamma'z}{|\gamma'z|} \right)^{-1} \quad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma.$$

Let \mathcal{F}_q be the space of all functions $f : H \rightarrow \mathbb{C}$ with

$$f(\gamma z) = j_\gamma(z)^q f(z), \quad \gamma \in \Gamma,$$

and if $\mathcal{D} = \Gamma \backslash H$ is a fundamental domain for Γ , define the Hilbert space $\mathcal{H}_q = \{f \in \mathcal{F}_q, \langle f, f \rangle_{\mathcal{D}} = \int_{\mathcal{D}} |f(z)|^2 d\mu(z) < \infty\}$ with areal measure $d\mu(z) = \frac{dx dy}{y^2}$ and the inner product

$$(8) \quad \langle f, g \rangle = \int_{\mathcal{D}} f(z) \overline{g(z)} d\mu(z).$$

An element in \mathcal{H}_q is called an automorphic form of weight $2q$. \mathcal{H}_q is isometric to \mathfrak{H}^q through the correspondence

$$(9) \quad I : \mathfrak{H}^q \ni f(dz)^q \mapsto y^q f \in \mathcal{H}_q.$$

Maass introduced the differential operators

$$\begin{aligned} L_q &= (\bar{z} - z) \frac{\partial}{\partial \bar{z}} - q & : \mathcal{F}_q &\rightarrow \mathcal{F}_{q-1} \\ K_q &= (z - \bar{z}) \frac{\partial}{\partial z} + q & : \mathcal{F}_q &\rightarrow \mathcal{F}_{q+1} \end{aligned}$$

We also have

$$\begin{aligned} L_q &= -2iy^{1+q} \frac{\partial}{\partial \bar{z}} y^{-q} = \overline{K_{-q}}, \\ K_q &= 2iy^{1-q} \frac{\partial}{\partial z} y^q = \overline{L_{-q}}. \end{aligned}$$

²Some authors refer to them as weight q or $-2q$

We write

$$\begin{aligned} -L_{q+1}K_q &= -\Delta_{2q} + q(q+1), \\ -K_{q-1}L_q &= -\Delta_{2q} + q(q-1) \end{aligned}$$

with

$$\Delta_{2q} = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) - 2iqy \frac{\partial}{\partial x}.$$

These second order differential operators are self-adjoint on \mathcal{H}_q .

Now, the metric and complex structure determine a covariant derivative

$$\nabla = \nabla^q \oplus \nabla_q : \mathcal{E}^{0,0}(M, \omega_M^q) \longrightarrow \mathcal{E}^{1,0}(M, \omega_M^q) \oplus \mathcal{E}^{0,1}(M, \omega_M^q)$$

on the line bundle ω_M^q . With the identifications $\mathcal{E}^{1,0}(M, \omega_M^q) \cong T^{q+1}$ and $\mathcal{E}^{0,1}(M, \omega_M^q) \cong T^{q-1}$, we have

$$\nabla^q : T^q \rightarrow T^{q+1}, \quad \nabla_q : T^q \rightarrow T^{q-1}.$$

Under the correspondence $I_k, I_k : \mathfrak{H}^k \ni f(dz)^k \mapsto y^k f \in \mathcal{H}_k$, $k = q-1, q, q+1$ the operators ∇_q, ∇^q go over to the Maass operators according to the commutative diagram

$$\begin{array}{ccccc} \mathfrak{H}^{q-1} & \xleftarrow{\nabla_q} & \mathfrak{H}^q & \xrightarrow{\nabla_q} & \mathfrak{H}^{q+1} \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{H}_{q-1} & \xleftarrow{L_q} & \mathcal{H}_q & \xrightarrow{K_q} & \mathcal{H}_{q+1} \end{array}$$

and so are given locally by

$$\nabla^q = I_{q+1}^{-1} K_q I_q = 2i\rho^q \partial \rho^{-q} \quad \text{and} \quad \nabla_q = I_{q-1}^{-1} L_q I_q = -2i\rho^{-1} \bar{\partial},$$

where $\partial = \frac{\partial}{\partial z}$ and $\bar{\partial} = \frac{\partial}{\partial \bar{z}}$.

The Laplacians Δ_q^+ and Δ_q^- on T^q are defined by $\Delta_q^+ = -\nabla_{q+1} \nabla^q = -I_q^{-1} L_{q+1} I_{q+1} I_{q+1}^{-1} K_q I_q = I_q^{-1} (-L_{q+1} K_q) I_q$, $\Delta_q^- = -\nabla^{q-1} \nabla_q = I_q^{-1} (-K_{q-1} L_q) I_q$ and then the isometry I conjugates Δ_q^+ with $-\Delta_{2q} + q(q+1)$ and Δ_q^- with $-\Delta_{2q} + q(q-1)$. Thus $\Delta_M = \Delta_0^\pm$ is the Laplacian on functions. The Δ_q^\pm are non-negative self-adjoint operators.

We are first interested in the case $q = 1$. Let Δ_{Diff} be the (positive) Laplacian on 1-forms on a geometrically finite hyperbolic surface, $\Delta_{\text{Diff}} = d\delta + \delta d$, $\delta = -*d*$ with $*$ the Hodge operator. In the following, we write $\Delta_{\text{Diff}} = \Delta$. If ω is a 1-form in the holomorphic cotangent bundle, $\omega = f(z) dz$, then we define its image by the isometry I to be $I(\omega) = I(f dz) = yf(z) = \tilde{f}(z)$. We have $y\Delta(f dz) = -(\Delta_2 \tilde{f}) dz$, in other words, in the preceding notation, $\Delta = \Delta_1^-$.

3.3. Generalized eigenfunctions. We are going to give the spectral expansion in eigenforms of Δ ; we use [7], [24], [1]. With the notation of Section 1, for a finitely generated group of the second kind, for each cusp and for each funnel of the quotient there is a corresponding Eisenstein series, which is what we are going to develop now.

We will denote $R_{s,q}$ the resolvent operator defined for $\text{Re } s > 1/2$, $s \notin [1/2, 1]$ by $R_{s,q} = (\Delta_{2q} + s(1-s))^{-1}$.

Proposition 3.1. *For an integer q , for $\text{Re } s > \delta$, $G_s(z, w, q)$, the kernel of the resolvent $R_{s,q}$, for the self-adjoint operator Δ_{2q} , acting on the Hilbert space \mathcal{H}_q of*

automorphic forms of weight $2q$, is given by the convergent series

$$G_s(z, w, q) = \sum_{\gamma \in \Gamma} j_\gamma(w)^q g_s(z, \gamma w, q)$$

with

$$g_s(z, w, q) = - \left(\frac{w - \bar{z}}{z - \bar{w}} \right)^q \frac{\Gamma(s+q)\Gamma(s-q)}{4\pi\Gamma(2s)} \cosh^{-2s}(d(z, w)/2) F(s+q, s-q, 2s; \cosh^{-2}(d(z, w)/2))$$

and F being Gauss's hypergeometric function.

We collect some properties we can find for example in [24], vol I, p. 100, with [1], Chapters 6 and 7. The resolvent kernel $G_s(z, w, q)$ has a meromorphic continuation to all of \mathbb{C} . Following these references, we define generalized eigenfunctions.

For the funnel case, we identify z' with the standard coordinates (ρ', t') (see (1)) in the funnel F_j , and, with $x' = e^{t'}$, we denote

$$(10) \quad x'^s E_{j,q}^f(s, z, x') = (1-2s) \lim_{\rho' \rightarrow 0} \rho'^{-s} G_s(z, z', q),$$

for $j = 1, \dots, n_f$. In the cusp C_j , with standard coordinates $z' = (\rho', t')$, we set

$$(11) \quad E_{j,q}^c(s, z) = (1-2s) \lim_{\rho' \rightarrow 0} \rho'^{1-s} G_s(z, z', q),$$

for $j = 1, \dots, n_c$.

We will call them respectively (standard) funnel Eisenstein series and cuspidal Eisenstein series. The Poisson kernel is

$$P(z, \zeta) = \text{Im}(z)/|z - \zeta|^2$$

where $z \in H$ and $\zeta \in \mathbb{R}$. For $b \in O(\Gamma) = \mathbb{R} \cup \{\infty\} - L(\Gamma)$, define the Eisenstein series ([24] [3])

$$E_b(z, s, k) = \sum_{\gamma \in \Gamma} j(\gamma, z)^k P(\gamma(z), b)^s (\gamma(z), b)^k,$$

where $j(\gamma, z) = \gamma'(z)/|\gamma'(z)|$ and $(z, b) = (\bar{z} - b)/(z - b)$. It converges uniformly on compact subsets of H if $\text{Re } s > \delta$.

Proposition 3.2. *The series $E_b(z, s, k)$ can be continued to the whole complex plane as a meromorphic function in s .*

One verifies that $-\Delta_{2k} E_b(z, s, k) = s(1-s)E_b(z, s, k)$.

Remark 3.1. *Thus, if $\delta < 1/2$, then $E_b(z, s, k)$ is analytic in a neighbourhood of $\text{Re } s = 1/2$.*

For the standard funnel F_l which corresponds to the region $\text{Re } z \geq 0$ in the model $C_l = \Gamma_l \backslash H$, we have (see [7], p. 200)

$$\begin{aligned} E_{l,q}^f(s, z, x') &= (1-2s) \lim_{z' \rightarrow x'} (\text{Im } z')^{-s} G_s(z, z', q) \\ &= -(1-2s) \frac{4^s}{4\pi} \frac{\Gamma(s+q)\Gamma(s-q)}{\Gamma(2s)} E_{x'}(z, s, q). \end{aligned}$$

In the case of 1-forms we write,

$$\mathcal{E}_{f_l}(s, z, x') = \frac{E_{l,1}^f(s, z, x')}{y} dz.$$

Now we point out the link between cuspidal and classical Eisenstein series.

Remark 3.2. Recall the definition of the classical Eisenstein series. We will assimilate the standard cusp of a parabolic point with the latter. The stabilizer of a cusp \mathfrak{a} is an infinite cyclic group generated by a parabolic motion,

$$\Gamma_{\mathfrak{a}} = \{\gamma \in \Gamma : \gamma \mathfrak{a} = \mathfrak{a}\} = \langle \gamma_{\mathfrak{a}} \rangle ,$$

say. There exists a $\sigma_{\mathfrak{a}} \in SL_2(\mathbb{R})$, called a scaling matrix of the cusp \mathfrak{a} , such that $\sigma_{\mathfrak{a}}\infty = \mathfrak{a}$, $\sigma_{\mathfrak{a}}^{-1}\gamma_{\mathfrak{a}}\sigma_{\mathfrak{a}} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. $\sigma_{\mathfrak{a}}$ is determined up to composition with a translation from the right. The Eisenstein series for the cusp \mathfrak{a} is then defined by

$$E_{\mathfrak{a}}(s, z) = \sum_{\Gamma_{\mathfrak{a}} \backslash \Gamma} y(\sigma_{\mathfrak{a}}^{-1}\gamma z)^s ,$$

where s is a complex variable with $\operatorname{Re} s > \delta$.

Definition 3.1. In a similar way we define the Eisenstein series of weight $2q$ associated to a cusp \mathfrak{a} as the automorphic form of weight $2q$, for $\operatorname{Re} s > \delta$:

$$E_{\mathfrak{a},q}(s, z) = \sum_{\gamma \in \Gamma_{\mathfrak{a}} \backslash \Gamma} y(\sigma_{\mathfrak{a}}^{-1}\gamma z)^s j_{\sigma_{\mathfrak{a}}^{-1}\gamma}(z)^{-q} = \sum_{\gamma \in \Gamma_{\mathfrak{a}} \backslash \Gamma} y(\sigma_{\mathfrak{a}}^{-1}\gamma z)^s \left(\frac{(\sigma_{\mathfrak{a}}^{-1}\gamma)'z}{|(\sigma_{\mathfrak{a}}^{-1}\gamma)'z|} \right)^q .$$

We call a horocyclic Eisenstein series the 1-form corresponding to the Eisenstein series of weight 2 associated to a cusp \mathfrak{a} , $E_{\mathfrak{a},1}$ and defined for $\operatorname{Re} s > 1$ by

$$(12) \quad \mathcal{E}_{\mathfrak{a}}(s, z) = \sum_{\gamma \in \Gamma_{\mathfrak{a}} \backslash \Gamma} y(\sigma_{\mathfrak{a}}^{-1}\gamma z)^{s-1} d(\sigma_{\mathfrak{a}}^{-1}\gamma z) = \frac{E_{\mathfrak{a},1}(s, z)}{y} dz .$$

We now verify that this corresponds to the defining formula (11), that is $E_{\mathfrak{a},q}^c = E_{\mathfrak{a},q}$.

For the standard cusp, we write for $\operatorname{Re} s > \delta$,

$$G_s(z, z', q) = \sum_{\Gamma_{\infty} \backslash \Gamma} \left(\frac{c\bar{z} + d}{cz + d} \right)^q G_s^{\Gamma_{\infty}}(\gamma z, z', q) ,$$

where $G_s^{\Gamma_{\infty}}(z, z', q)$ is the resolvent kernel of the standard cusp for automorphic forms of weight $2q$. We use then [7], p. 155 (38), p. 177 for $\operatorname{Im} z' > \operatorname{Im} \gamma z$, p. 172 (see also [1], pp. 72, 102) to conclude that

$$\lim_{y' \rightarrow \infty} y'^{s-1} G_s(z, z', q) = \sum_{\Gamma_{\infty} \backslash \Gamma} \left(\frac{c\bar{z} + d}{cz + d} \right)^q \frac{(\operatorname{Im} \gamma z)^s}{1-2s} = \frac{1}{1-2s} E_{\infty,q}(s, z) .$$

In particular from the foregoing and the references cited, we have

Lemma 3.1. The funnel and cuspidal Eisenstein series, $E_{j,q}^f(s, z, t')$ and $E_{\mathfrak{a},q}(s, z)$, can be continued to the whole complex plane as a meromorphic function in s .

We recall the property (see for example [7], p. 196, [24], [1], p.94)

Remark 3.3. The possible poles of the resolvent $G_s(z, z', q)$ in $\operatorname{Re} s > 1/2$ are simple and at $s_{1,q}, \dots, s_{N,q} \in (\frac{1}{2}, 1)$ corresponding to the eigenvalues $\lambda_{k,q}$ $1 \leq k \leq N$ of the Laplacian $-\Delta_{2q}$ on S_l .

Recall the decomposition (see Section 1) $M = K \cup_{j=1}^{n_c} C_j \cup_{j=1}^{n_f} F_j$, with the preceding notation and formula(8), we then have (see for example [7], [24])

Proposition 3.3. For $f \in \mathcal{H}_q$,

$$\begin{aligned} f(z) &= \sum_{k=1}^N (f)_{\lambda_{k,q}}(z) + \frac{1}{4\pi} \sum_{j=1}^{n_c} \int_{-\infty}^{+\infty} \langle f, E_{j,q}^c(1/2 + it, \cdot) \rangle E_{j,q}^c(1/2 + it, z) dt + \\ &\quad \frac{1}{4\pi} \sum_{j=1}^{n_f} \int_{-\infty}^{+\infty} \left[\int_1^{e^{lj}} \langle f, E_{j,q}^f(1/2 + it, \cdot, b) \rangle E_{j,q}^f(1/2 + it, z, b) db \right] dt; \end{aligned}$$

where the first sum in the right member is the projection of f on the discrete spectrum.

Let's formulate the spectral decomposition we needed in the case of 1-forms.

Proposition 3.4. For $w = f(z) dz$ square integrable,

$$\begin{aligned} w(z) &= \sum_{i=1}^m (w)_{\lambda_i}(z) + \frac{1}{4\pi} \sum_{j=1}^{n_c} \int_{-\infty}^{+\infty} (w, \mathcal{E}_{c_j}(1/2 + it, \cdot)) \mathcal{E}_{c_j}(1/2 + it, z) dt + \\ &\quad \frac{1}{4\pi} \sum_{j=1}^{n_f} \int_{-\infty}^{+\infty} \left[\int_1^{e^{lj}} (w, \mathcal{E}_{f_j}(1/2 + it, \cdot, b)) \mathcal{E}_{f_j}(1/2 + it, z, b) db \right] dt; \end{aligned}$$

where the first sum in the right member is the projection of w on the discrete spectrum, \mathcal{E}_{c_j} and \mathcal{E}_{f_j} are the Eisenstein series associated to the cusp C_j and the funnel F_j , respectively.

Remark 3.4. One can easily deduce the formula for an arbitrary square integrable 1-form, with the following notations,

$$\begin{aligned} \mathcal{E}_{c_j}(s, z)_+ &= \mathcal{E}_{c_j}(s, z) & \mathcal{E}_{f_k}(s, z, b)_+ &= \mathcal{E}_{f_k}(s, z, b) \\ \mathcal{E}_{c_j}(s, z)_- &= \overline{\mathcal{E}_{c_j}(\bar{s}, z)} & \mathcal{E}_{f_k}(s, z, b)_- &= \overline{\mathcal{E}_{f_k}(\bar{s}, z, b)} \end{aligned}$$

$$\begin{aligned} \Omega(z) &= f dz + g d\bar{z} = \sum_{i=1}^m (\Omega)_{\lambda_i}(z) + \\ &\quad \frac{1}{4\pi} \sum_{j=1}^{n_c} \int_{-\infty}^{+\infty} (\Omega, \mathcal{E}_{c_j}(1/2 + it, \cdot)_+) \mathcal{E}_{c_j}(1/2 + it, z)_+ + (\Omega, \mathcal{E}_{c_j}(1/2 + it, \cdot)_-) \mathcal{E}_{c_j}(1/2 + it, z)_- dt \\ &\quad + \frac{1}{4\pi} \sum_{j=1}^{n_f} \int_{-\infty}^{+\infty} \int_1^{e^{lj}} (\Omega, \mathcal{E}_{f_j}(1/2 + it, \cdot, b)_+) \mathcal{E}_{f_j}(1/2 + it, z, b)_+ db dt \\ &\quad + \frac{1}{4\pi} \sum_{j=1}^{n_f} \int_{-\infty}^{+\infty} \int_1^{e^{lj}} (\Omega, \mathcal{E}_{f_j}(1/2 + it, \cdot, b)_-) \mathcal{E}_{f_j}(1/2 + it, z, b)_- db dt \end{aligned}$$

For simplicity, we will write

$$(13) \quad \begin{aligned} \Omega(z) &= (\Omega)_{\lambda_i}(z) + \frac{1}{4\pi} \int_{-\infty}^{+\infty} (\Omega, \mathcal{E}_{c_j}(1/2 + it, \cdot)_{\pm}) \mathcal{E}_{c_j}(1/2 + it, z)_{\pm} dt + \\ &\quad \frac{1}{4\pi} \int_{-\infty}^{+\infty} \left[\int_1^{e^{lk}} (\Omega, \mathcal{E}_{f_k}(1/2 + it, \cdot, b)_{\pm}) \mathcal{E}_{f_k}(1/2 + it, z, b)_{\pm} db \right] dt. \end{aligned}$$

3.4. **Harmonic dual form.** We are now going to see

Proposition 3.5. *If c is an oriented simple closed geodesic on M , then, for $\text{Re } s > 0$, the hyperbolic Eisenstein series $\Omega_c(s, \cdot)$ are square integrable.*

Proof. We consider a fundamental domain D contained in $\{z, 1 \leq |z| \leq e^l\}$ in which the segment (i, ie^l) represents the geodesic c . We write $C_\lambda = \{z \in D, d(z, c) = \lambda\}$ and $F_\lambda = \{z \in D, d(z, c) \geq \lambda\}$. Without loss of generality we may assume that there is only one funnel on M and no cusps. Let V_λ be the volume of $F_\lambda - F_{\lambda+1}$. Then there exists a constant c_1 such that $V_\lambda \geq c_1(\sinh(\lambda + 1) - \sinh(\lambda))$. For $\text{Re } s = \sigma > 0$, $\|\Omega_c(s, z)\| = \|\Omega(s, z)\| \leq \frac{1}{|k(s)|} \sum_{\Gamma_1 \backslash \Gamma} \frac{1}{(\cosh x_2(\gamma z))^{\sigma+1}}$. Let $\eta(z) = \sum_{\Gamma_1 \backslash \Gamma} \frac{1}{(\cosh x_2(\gamma z))^{\sigma+1}}$. We know from Section 2 that there exists a constant $K > 0$ such that $\forall z \in H, \eta(z) \leq K$.

We have

$$\begin{aligned} \int_D \|\Omega_c(s, z)\|^2 d\mu(z) &\leq \frac{1}{|k(s)|^2} \int_D \eta^2(z) d\mu(z) \\ &\leq \frac{1}{|k(s)|^2} \int_{1 \leq x_1 \leq e^l, -\infty < x_2 < +\infty} \eta(z) \frac{1}{(\cosh x_2(z))^{\sigma+1}} \cosh x_2 dx_1 dx_2 \\ &\leq \frac{K}{|k(s)|^2} \int_{1 \leq x_1 \leq e^l, -\infty < x_2 < +\infty} \frac{1}{(\cosh x_2(z))^{\sigma+1}} \cosh x_2 dx_1 dx_2. \end{aligned}$$

The last integral is $\frac{\Gamma(\frac{1}{2})\Gamma(\frac{\sigma}{2})}{\Gamma(\frac{1}{2} + \frac{\sigma}{2})}(e^l - 1)$ and the result follows. \square

As in [20], we verify that

$$\Delta(\Omega(s, z)) + s(s+1)\Omega(s, z) = s(s+1)\Omega(s+2, z).$$

This formula has the consequence that for fixed s with $\text{Re } s > 0$, the function $\Delta^k(\Omega(s, z))$ is again square integrable for any $k > 0$.

Set $\text{Re } s > 0$. Then with the notation of Remark 3.4, in particular in the last equality (13), we decompose the first sum of the right member as $\Omega_0(z)$ the harmonic part of $\Omega(s, z)$ plus the projection on the non zero discrete spectrum. More precisely

$$\begin{aligned} \Omega(s, z) &= \Omega_0(z) + a_i(s)\varphi_i(z) + \frac{1}{4\pi} \int_{-\infty}^{+\infty} h_{\pm}^{c_j}(s, t)\mathcal{E}_c(1/2 + it, z)_{\pm} dt \\ (14) \quad &+ \frac{1}{4\pi} \int_{-\infty}^{+\infty} \left[\int_1^{e^l k} H_{\pm}^{f_k}(s, t, b)\mathcal{E}_f(1/2 + it, z, b)_{\pm} db \right] dt, \end{aligned}$$

where, $\{\varphi_i\}$ is a complete orthonormal basis of eigenforms of Δ with corresponding positive eigenvalues, $a_i(s) = (\Omega(s, \cdot), \varphi_i)$, $h_{\pm}^{c_j}(s, t) = (\Omega(s, \cdot), \mathcal{E}_{c_j}(1/2 + it, \cdot)_{\pm})$ and $H_{\pm}^{f_k}(s, t, b) = (\Omega(s, \cdot), \mathcal{E}_{f_k}(1/2 + it, \cdot, b)_{\pm})$.

We obtain with H corresponding to any $H_{\pm}^{f_k}$

$$(1/4 + t^2 + s(s+1))H(s, t, b) = s(s+1)H(s+2, t, b).$$

From this we get a continuation of H to the region $\text{Re } s > -1/2$ and we note that for all t and all b we have $H(0, t, b) = 0$.

Moreover, for $\text{Re } s > -1/2$, $\text{Re}(s+2) > 0$ and we may substitute in (14) to obtain a continuation of $\Omega(s, z)$ to $\text{Re } s > -1/2$. In particular $s = 0$ is a regular value. Substituting $s = 0$ into $\Delta\Omega(s, z) + s(s+1)\Omega(s, z) = s(s+1)\Omega(s+2, z)$ we obtain $\Delta\Omega(0, z) = 0$. Moreover for any closed oriented cycle c' , for $\text{Re } s > 0$,

$\int_{c'} \Omega(s, \cdot) = c.c'$ and by analytic continuation, for $\operatorname{Re} s > -1/2$, in particular for $s = 0$. Thus we have proved the following theorem:

Theorem 3.1. $\Omega(s, z)$ has a meromorphic continuation to $\operatorname{Re} s > -1/2$ with $s = 0$ a regular point and $\Omega(0, z)$ is a harmonic form which is dual to c .

Remark 3.5. 1) Another way to see this is:

write $\Omega(s, z) = (\Delta + s(s+1))^{-1}(s(s+1)\Omega(s+2, z))$ and use the meromorphic continuation of the resolvent (see for example [1], [25]).

2) Using methods analogous to those of [20] (see also [18]), we can obtain a complete description of the singularities of the hyperbolic Eisenstein series.

4. THE CASE OF AN INFINITE GEODESIC JOINING TWO POINTS.

Without loss of generality, we suppose that the two cusps p and q are 0 and ∞ and, as the lift of the geodesic, we take the imaginary axis. Let η be the infinite geodesic $]p, q[$. Can we carry out the same construction as Kudla and Millson? As in the finite volume case, the problem reduces to studying the following series for $\operatorname{Re} s > 1$:

$$(15) \quad \hat{\eta}^s(z) = \frac{1}{k(s-1)} \sum_{\gamma \in \Gamma} \gamma * \left[\left(\frac{y}{|z|} \right)^{s-1} \operatorname{Im}(z^{-1} dz) \right] = \operatorname{Im}(\theta^s(z)),$$

where

$$\theta^s(z) = \frac{1}{k(s-1)} \sum_{\gamma \in \Gamma} \gamma * \left[\left(\frac{y}{|z|} \right)^{s-1} \frac{dz}{z} \right],$$

$k(s-1) = \frac{\Gamma(1/2)\Gamma(s/2)}{\Gamma(1/2+s/2)}$, and the notation

$$\operatorname{Im}(\theta^s(z)) = \frac{1}{2ik(s-1)} \left(\sum_{\gamma \in \Gamma} \gamma * \left[\left(\frac{y}{|z|} \right)^{s-1} \frac{dz}{z} \right] - \gamma * \left[\left(\frac{y}{|z|} \right)^{s-1} \frac{d\bar{z}}{\bar{z}} \right] \right).$$

4.1. Some useful estimates. As usual, we can suppose $\Gamma_\infty = \langle z \mapsto z+1 \rangle$ to be the stabilizer of ∞ in Γ and the stabilizer of 0, Γ_0 , is then generated by $z \mapsto \frac{z}{-c_0^2 z + 1}$ (for some non-zero constant c_0).

First of all we note that, unlike in the finite volume case (see Proposition 3.1 and (11)):

Lemma 4.1. *The series $\sum_{\gamma \in \Gamma_\infty \backslash \Gamma} \operatorname{Im}(\gamma z)$ is convergent.*

Another way to see this is ‘by hand’: We know that for $\operatorname{Re} s > \delta$, $\sum_{T \in \Gamma} e^{-sd(i, Tz)}$ converges; moreover there exists a constant $C > 0$ such that $\sum_{\gamma \in \Gamma_\infty \backslash \Gamma} \operatorname{Im}^{\operatorname{Re} s}(\gamma z) \leq C \sum_{T \in \Gamma} e^{-\operatorname{Re} s d(i, Tz)}$, as in our case $\delta < 1$, we have the result.

As $\sum_{\gamma \in \Gamma_\infty \backslash \Gamma} \left| \operatorname{Im}^s(\gamma z) \frac{c\bar{z}+d}{cz+d} \right| = \sum_{\gamma \in \Gamma_\infty \backslash \Gamma} \operatorname{Im}^{\operatorname{Re} s}(\gamma z)$, we also deduce the convergence of the series representing $E_{\infty,1}(1, z)$ (3.2).

Lemma 4.2. *We have the following asymptotic behaviour for $\operatorname{Re} s > \delta$:*

- 1) in a funnel, for all cusps \mathfrak{a} , $E_{\mathfrak{a}}(s, z)$ is square integrable;
- 2) at $\mathfrak{a} = \infty$, $E_\infty(s, z) - y^s = O(y^{1-s})$ and $E_0(s, z) = O(y^{1-s})$;

3) near $\mathfrak{a} = 0$, $E_0(s, z) - y^s/(c_0^2|z|^2)^s = O(y^{1-s}/(c_0^2|z|^2)^{s-1})$ and $E_\infty(s, z) = O(y^{1-s}/(c_0^2|z|^2)^{s-1})$.

Proof. With the notations of 3.1 and taking ρ to be the standard coordinate for the cusp \mathfrak{a} , we re-write the results of [1](p. 110) in the following way:

$$(16) \quad E_{\mathfrak{a}}(s, \cdot) = \rho^{-s}(1 - \chi_0(\rho)) + O(\rho_f^s \rho_c^{s-1}),$$

where we define $\chi_0 \in C_0^\infty(M)$ such that, r being the distance to the compact core for a point in cusp \mathfrak{a} ,

$$\chi_0(\rho) = \chi_0(e^{-r}) = \begin{cases} 1, & r \leq 0 \\ 0, & r \geq 1 \end{cases};$$

and ρ_f (resp. ρ_c) is the standard coordinate in the funnels (resp. cusps). Now Γ contains parabolic elements then $\delta > 1/2$. The area form in standard funnel coordinates is $dA = \rho_f^{-2} d\rho_f dt + O(1)$, so $\rho_f^{1/2}$ is the threshold for L^2 asymptotic behavior in a funnel. Then for $\text{Re } s > \delta > 1/2$ we have 1). The cusp area form is $dA = d\rho_c dt$, so borderline L^2 behavior is $\rho_c^{-1/2}$. For 2) and 3) we can use (16), see also [5] proposition 3.1. \square

4.2. Convergence of the hyperbolic Eisenstein series and its analytic continuation. The computations to prove the convergence of (15) are easily adapted from the finite volume case. For the convenience of the reader, we recall the essential points.

We have $\|\sum_{\gamma \in \Gamma} \gamma * \left(\frac{y}{|z|}\right)^{s-1} \text{Im}(z^{-1} dz)\| \leq \sum_{\gamma \in \Gamma} \left(\frac{y}{|z|}\right)^\sigma (\gamma z)$, where $\sigma = \text{Re } s > 1$ and if we put $\mathcal{S} = \sum_{\gamma \in \Gamma} \left(\frac{y}{|z|}\right)^\sigma (\gamma z)$, we have

$$\mathcal{S} = \sum_{\gamma \in \Gamma_\infty \setminus \Gamma} y^\sigma(\gamma z) \sum_{n \in \mathbb{Z}} \frac{1}{|\gamma z + n|^\sigma} = S_1 + S_2,$$

with

$$S_1 = \sum_{\gamma \in \Gamma_\infty \setminus \Gamma} \frac{y^\sigma(\gamma z)}{|\gamma z|^\sigma} \quad \text{and} \quad S_2 = \sum_{\gamma \in \Gamma_\infty \setminus \Gamma} y^\sigma(\gamma z) \sum_{n \in \mathbb{Z}^*} \frac{1}{|\gamma z + n|^\sigma}.$$

Let S_z be a system of representatives of $\Gamma_\infty \setminus \Gamma$ such that $|\text{Re } \gamma z| \leq 1/2$. Then

$$\|\mathcal{S}\| \leq S_1 + 2 \sum_{\gamma \in S_z} y^\sigma(\gamma z) \sum_{n=1}^{\infty} \frac{1}{(n - 1/2)^\sigma}.$$

We have

$$\begin{aligned} \sum_{\gamma \in \Gamma_\infty \setminus \Gamma} \frac{y^\sigma(\gamma z)}{|\gamma z|^\sigma} &= \sum_{\Gamma_0 \setminus (\Gamma_\infty \setminus \Gamma)} \frac{y^\sigma(\gamma z)}{|\gamma z|^\sigma} \sum_{n \in \mathbb{Z}} \frac{1}{|nc_0^2 \gamma z + 1|^\sigma} \\ &= \sum_{\gamma \in \Gamma_0 \setminus (\Gamma_\infty \setminus \Gamma)} \frac{y^\sigma(\gamma z)}{|\gamma z|^\sigma} + \sum_{\gamma \in \Gamma_0 \setminus (\Gamma_\infty \setminus \Gamma)} \frac{y^\sigma(\gamma z)}{|\gamma z|^\sigma} \sum_{n \in \mathbb{Z}^*} \frac{1}{|nc_0^2|^\sigma [(x(\gamma z) - 1/nc_0^2)^2 + y^2(\gamma z)]^{\sigma/2}}. \end{aligned}$$

For K a compact set in H there exists m in H such that

$$\forall z \in K, \forall \gamma \in \Gamma_0 \setminus \Gamma_\infty \setminus \Gamma, |\gamma z| \geq |m| \text{ and } \text{Im } \gamma z \geq \text{Im } m.$$

So

$$\begin{aligned} \mathcal{S}_1 &\leq \sum_{\gamma \in \Gamma_0 \setminus (\Gamma_\infty \setminus \Gamma)} \frac{y^\sigma(\gamma z)}{|m|^\sigma} + \sum_{\gamma \in \Gamma_0 \setminus (\Gamma_\infty \setminus \Gamma)} \frac{y^\sigma(\gamma z)}{|m|^\sigma} \sum_{n \in \mathbb{Z}^*} \frac{1}{|nc_0^2|^\sigma (\text{Im } m)^\sigma} \\ &\leq \frac{1}{|m|^\sigma} \sum_{\Gamma_\infty \setminus \Gamma} y^\sigma(\gamma z) + 2 \sum_{n \in \mathbb{N}^*} \frac{1}{(nc_0^2)^\sigma} \frac{1}{|m|^\sigma (\text{Im } m)^\sigma} \sum_{\Gamma_\infty \setminus \Gamma} y^\sigma(\gamma z); \end{aligned}$$

and lastly, for all z in K

$$\begin{aligned} \|\mathcal{S}\| &\leq \frac{1}{|m|^\sigma} \sum_{\Gamma_\infty \setminus \Gamma} y^\sigma(\gamma z) + 2 \sum_{n \in \mathbb{N}^*} \frac{1}{(nc_0^2)^\sigma} \frac{1}{|m|^\sigma (\text{Im } m)^\sigma} \sum_{\Gamma_\infty \setminus \Gamma} y^\sigma(\gamma z) + \\ &2 \sum_{\Gamma_\infty \setminus \Gamma} y^\sigma(\gamma z) \sum_{n=1}^{\infty} \frac{1}{(n-1/2)^\sigma}, \end{aligned}$$

with uniform convergence on all compact subsets of H and all compact subsets of $\text{Re } s > 1$.

From this last inequality we conclude that θ^s is square integrable in the funnels as the Eisenstein series \mathcal{E}_∞ . Moreover using the notation in formula (4) we can write $\eta^s(z) = \sum_{\gamma \in \Gamma} \gamma^* \varphi_s$, where $\varphi_s = \frac{1}{k(s-1)} \frac{dx_2}{(\cosh x_2)^s}$ satisfies $\Delta \varphi_s + s(s-1)\varphi_s = s(s-1)\varphi_{s+2}$. To conclude, we have the following theorem:

Theorem 4.1. *For $\text{Re } s > 1$, the Eisenstein series associated to the geodesic $\eta = (p, q)$ converges uniformly on all compact sets. It represents a C^∞ closed form which is dual to η . For $\text{Re } s > 1$ it satisfies the differential functional equation:*

$$\Delta \hat{\eta}^s = s(1-s)[\hat{\eta}^s - \hat{\eta}^{s+2}].$$

Now we want to prove that $\hat{\eta}^s$ has an analytic continuation to $\text{Re } s > 1/2$ and $s = 1$ is a regular value. For this, first of all, we are going to show that $\theta^s(z) - 1/i(\mathcal{E}_\infty(1, z) - \mathcal{E}_0(1, z))$ is square integrable. As we have shown that θ^s is square integrable in the funnels, what we have to do is to investigate the Fourier expansion of θ^s at each inequivalent cusp, i.e. at 0 and ∞ , and to show that $\|\theta^s\|$ is bounded at the cusps. As in the finite volume case, we have ([5]):

Proposition 4.1. *At ∞*

$$\theta^s(z) = \left(\frac{1}{i} + O(1/y)\right) dz,$$

and at 0

$$\theta^s(z) = \left(-\frac{1}{ic_0^2 z^2} + O(1/y)\right) dz.$$

By Proposition 4.1 and Lemma 4.2, we conclude:

Proposition 4.2. *The 1-forms $\theta^s(z) - 1/i(\mathcal{E}_\infty(1, z) - \mathcal{E}_0(1, z))$ and $\hat{\eta}^s(z) + \text{Re}(\mathcal{E}_\infty(1, z) - \mathcal{E}_0(1, z))$ are square integrable.*

Proof. Let's, for example, treat the case at ∞ . Recall that

$$\eta^s(z) = \frac{1}{2ik(s-1)} \left(\sum_{\gamma \in \Gamma} \gamma^* \left[\left(\frac{y}{|z|} \right)^{s-1} \frac{dz}{z} \right] - \gamma^* \left[\left(\frac{y}{|z|} \right)^{s-1} \frac{d\bar{z}}{\bar{z}} \right] \right).$$

The asymptotic behavior of $\frac{1}{k(s-1)} \sum_{\gamma \in \Gamma} \gamma^* \left[\left(\frac{y}{|z|} \right)^{s-1} \frac{d\bar{z}}{z} \right]$ can be deduced from the preceding study. For example at ∞ it behaves like $-\left(\frac{1}{i} + O(1/y)\right) dz$. We conclude that $\eta^s(z) = -\frac{1}{2}(dz + d\bar{z}) + O(1/y dz)$ and the result. \square

Lastly, as in [5]:

Theorem 4.2. *The 1-form $\hat{\eta}^s$ has a meromorphic continuation to $\text{Re } s > 1/2$, with $s = 1$ a regular point and $\hat{\eta}$ is a harmonic form which is dual to η .*

5. DEGENERATIONS.

5.1. Background material and the main results. A family of degenerating geometrically finite hyperbolic surfaces consists of a surface M and a smooth family $(g_t)_{t>0}$ of Riemannian metrics that meet the following assumptions:

- (1) The Riemannian manifold $M_t = (M, g_t)$ is a geometrically finite hyperbolic surface for each t .
- (2) There are finitely many disjoint open subsets $\mathcal{C}_{l,i} \subset M$ that are diffeomorphic to cylinders $\mathbb{R}/\mathbb{Z} \times J_i$ where $J_i \subset \mathbb{R}$ is a connected neighborhood of 0 with the metric $(x, a) \mapsto (l_i(l)^2 + a^2)dx^2 + ((l_i(l)^2 + a^2)^{-1}da^2$ and $l_i(l) \rightarrow 0$ as $l \rightarrow 0$. The curve $c_i = \mathbb{R}/\mathbb{Z} \times \{0\}$ is a closed geodesic of length $l_i(l)$.
- (3) The complement of $(C_1 \cup \dots \cup C_{n_c}) \cup (F_1 \cup \dots \cup F_{n_f}) \cup_i \mathcal{C}_{l,i}$ where we may have some $F_j \subset \mathcal{C}_{l,i}$ is relatively compact.
- (4) On $M_0 := M \setminus \cup_i c_i$, the metrics g_t converge smoothly to a hyperbolic metric g_0 as $l \rightarrow 0$. M_0 is a possibly non connected hyperbolic surface that contains a pair of cusps for each i .

First of all, we recall some material and results. The following lemma can be found, for example, in [1], p. 252-253. The neighbourhood of points within a distance a of a geodesic γ , where $d(z, \gamma)$ is the hyperbolic distance from z to γ ,

$$G_a = \{z \in K, d(z, \gamma) \leq a\},$$

is isometric for small a to a half-collar $[0, a] \times S^1$, $ds^2 = dr^2 + l^2 \cosh^2 r d\theta^2$.

Lemma 5.1. *Suppose that γ is a simple closed geodesic of length $l(\gamma)$ on a geometrically finite hyperbolic surface M . Then γ has a collar neighbourhood of half-width d , such that*

$$\sinh(d) = \frac{1}{\sinh(l(\gamma)/2)}.$$

As a consequence, if η is any other closed geodesic intersecting γ transversally (still assuming γ is simple), then the lengths of the two geodesics satisfy the inequality

$$\sinh(l(\eta)/2) \geq \frac{1}{\sinh(l(\gamma)/2)}.$$

Lemma 5.2. *Let γ be a simple closed geodesic of length l on a complete hyperbolic surface M . If α is a simple closed geodesic that does not intersect γ , then, putting $d(\gamma, \alpha)$ to be the hyperbolic distance of γ to α ,*

$$\cosh d(\gamma, \alpha) \geq \coth(l/2).$$

A standard collar for a geodesic of length l is a cylinder isometric to $\langle z \mapsto e^l z \rangle \backslash \mathcal{C}$ with $\mathcal{C} = \{z = re^{i\theta}, 1 \leq r \leq e^l, l < \theta < \pi - l\} \subset H$ with the restriction of the hyperbolic metric, and $\langle z \mapsto e^l z \rangle$ the cyclic group generated by the transformation $z \mapsto e^l z$. There is a constant k_0 (the short geodesic constant) such that each closed geodesic on M_l of length at most k_0 has a neighbourhood isometric to the standard collar and each cusp for M_l has a neighbourhood isometric to the standard cusp; furthermore, the collars for short geodesics and the cusp regions are all mutually disjoint.

Here we consider a family of surfaces $S_l = \Gamma_l \backslash H$ degenerating to the surface S with only one geodesic c_l being pinched, Γ_l containing the transformation $\sigma_l(z) = e^l z$ corresponding to c_l . Let K_l be S_l minus C_l the standard collar for c_l . There exist homeomorphisms f_l from $S_l \setminus c_l$ to S , with f_l tending to isometries C^2 -uniformly on the compact core $K_l \subset S_l$; define $\pi_l = f_l^{-1}$. Suppose that p is one of the two cusps of S arising from pinching c_l . Let $S_0 = \Gamma \backslash H$ be the component of S containing p and conjugate Γ to represent the cusp by the translation $w \mapsto w + 1$. In the following, $p = \infty$.

Let, for $\operatorname{Re} s > 1$, $\alpha_l(s, z) = \sum_{\gamma \in \langle \sigma_l \rangle \backslash \Gamma_l} \gamma * \left[\left(\frac{y}{|z|} \right)^{s-1} \operatorname{Im}(z^{-1} dz) \right]$ be such that the hyperbolic Eisenstein series $\Omega_{c_l} = \Omega_l$ is related by $\Omega_l(s, z) = \frac{1}{k(s)} \alpha_l(s+1, z)$. Without loss of generality we suppose S_l has only one funnel F_1 . With the notation of the beginning of the paper, $S_l = K \cup (C_1 \cup \dots \cup C_{n_c}) \cup F_1$ and c_l is the one geodesic of the boundary of the compact core K . We consider the specific case of p , the limit of the right side of the c_l -collar, contained in $S_l \setminus F_1$. With the misuse of notation already mentioned

Theorem 5.1. *Let $\operatorname{Re} s > 1$. Then the family of 1-forms $\frac{1}{l^s} \alpha_l(s, \pi_l(\cdot))$ converges uniformly on compact subsets of S_0 to $\operatorname{Im} \mathcal{E}_\infty(s, \cdot)$.*

This is a particular case of Theorem 5.2 below.

The sketch of the proof of this theorem follows those of the finite volume case ([4], [27], see also [8]). To study the right side of the c_l -collar, let $w = \frac{1}{l} \log z$, with the principal branch $z \in H$, and conjugate Γ_l by the map w to obtain $\tilde{\Gamma}_l$ acting on $\mathcal{S}_l = \{w, 0 < \operatorname{Im} w < \pi/l\}$. The hyperbolic metric on \mathcal{S}_l is $ds_l^2 = \left(\frac{l|dw|}{\sin(l \operatorname{Im} w)} \right)^2$, which tends uniformly on compact subsets to $\left(\frac{|dw|}{\operatorname{Im} w} \right)^2$. $\tilde{\Gamma}_l$ is a (non-Möbius) group of deck transformations acting on \mathcal{S}_l ; the quotient $\tilde{\Gamma}_l \backslash \mathcal{S}_l$ is S_l .

In the following if a group G acts on a domain D , we will denote by G_A the stabilizer in G of a subset $A \subset D$.

Let \hat{f}_l be the restriction of f_l to the component $\mathcal{S}_l^{(r)}$ of $S_l \setminus c_l$ containing the right half-collar for c_l . Let F_l be a lift of \hat{f}_l to the universal covers \mathcal{A}_l and H , where \mathcal{A}_l is the simply connected component of $H \setminus \pi_l^{-1}(c_l)$ which contains the standard right collar $\{z = re^{i\theta}, 1 \leq r \leq e^l, l < \theta < \pi/2\}$. $\mathcal{S}_l^{(r)} = (\Gamma_l)_{\mathcal{A}_l} \backslash \mathcal{A}_l$. More precisely [4], p. 350, [27]:

Lemma 5.3. *The simply connected component \mathcal{A}_l contains $\{z = re^{i\theta}, 1 \leq r \leq e^l, l c(l) < \theta < \pi/2\}$ where $c(l) \rightarrow 0, l \rightarrow 0$.*

Start with the standard Γ fundamental domain $\mathcal{F} = \{w \in H, 0 \leq \operatorname{Re} w < 1, \operatorname{Im} w \geq \operatorname{Im} A(w), \forall A \in \Gamma\}$. Set $D_l = F_l^{-1}(\mathcal{F})$, then D_l is a fundamental domain of $\mathcal{S}_l^{(r)}$. Divide the cosets of $\langle \sigma_l \rangle \backslash (\Gamma_l - \langle \sigma_l \rangle)$ into two classes $D = \{[A], A \in \Gamma_l, \inf \operatorname{Re} A(D_l) > 0\}$ and $G = \{[A], A \in \Gamma_l, \sup \operatorname{Re} A(D_l) < 0\}$.

Then \hat{f}_l has a lift \tilde{f}_l , a homeomorphism from a subdomain of \mathcal{S}_l to H : $\tilde{f}_l = F_l \circ w^{-1} : w(\mathcal{A}_l) \rightarrow H$. The homeomorphism \tilde{f}_l induces a group homomorphism $\rho_l : \Gamma \rightarrow \tilde{\Gamma}_l$, $A \mapsto \tilde{f}_l^{-1} A \tilde{f}_l$, $A \in \Gamma$. Now by our normalizations for $\tilde{\Gamma}_l$ and Γ , the translation $w \mapsto w + 1$ corresponds to itself and induces an isomorphism from Γ to $(\tilde{\Gamma}_l)_{w(\mathcal{A}_l)}$. We call $\rho_l(A) \in \tilde{\Gamma}_l$ the element corresponding to $A \in \Gamma$. If we specify the further normalization $\tilde{f}_l(i) = i$, then the lifts \tilde{f}_l are uniquely determined and then we have (see, e.g. [27], p. 107)

Lemma 5.4. *The \tilde{f}_l tend uniformly on compact subsets to the identity, and thus for $A \in \Gamma$, the corresponding elements $\rho_l(A)$ tend uniformly on compact subsets to A .*

For $q \in \mathbb{N}$, we associate to the pinching geodesic c_l , the q -form defined for $\operatorname{Re} s > 1$ by

$$A_{l,q}(s, z) = \sum_{\langle \sigma_l \rangle \backslash \Gamma_l} \left(\frac{\gamma'(z)}{\gamma(z)} \right)^q \sin^{s-q} \theta(\gamma z) dz^q.$$

Remark 5.1. *As suggested by the referee, it would be interesting to have a geometric meaning of $A_{l,q}(s, z)$. As far as I know, in the case of functions, they have been studied in particular in [8], [18], [19]. In the case of 1-forms, the analytic continuation of the hyperbolic Eisenstein series gives a harmonic dual form to the corresponding geodesic and the construction of an explicit geometric basis for the space of holomorphic 1-forms ([20]). In the case of 2-forms (quadratic differentials), they permit to define vector fields on Teichmüller space ([27], Chapter 2). For q -forms ($q > 2$) I could maybe find some answer in [21].*

Divide the cosets $\langle z \mapsto z + 1 \rangle \backslash (\tilde{\Gamma}_l - \langle z \mapsto z + 1 \rangle)$ into two classes, the left and the right: for $\mathcal{F}_l = \tilde{f}_l^{-1}(\mathcal{F})$, $L = wGw^{-1} = \{[A], A \in \tilde{\Gamma}_l, \inf \operatorname{Im} A(\mathcal{F}_l) > \pi/2l\}$ and $R = wDw^{-1} = \{[A], A \in \tilde{\Gamma}_l, \sup \operatorname{Im} A(\mathcal{F}_l) < \pi/2l\}$ (the line $\{\operatorname{Im} w = \pi/2l\}$ is a lift of c_l , and we write $[A]$ for the $\langle z \mapsto z + 1 \rangle$ coset of A). In particular the cosets $\langle z \mapsto z + 1 \rangle \backslash (\Gamma - \langle z \mapsto z + 1 \rangle)$ correspond to the right cosets of $\tilde{\Gamma}_l$: $\{[\rho_l(A)], A \in \Gamma, \langle w \mapsto w + 1 \rangle \subset R\}$. Then we can write, where χ_+ is the characteristic function of $\{\operatorname{Re} z > 0\}$ and χ_- that of $\{\operatorname{Re} z \leq 0\}$,

$$\begin{aligned} A_{l,q}(s, z) &= \sum_{\langle \sigma_l \rangle \backslash \Gamma_l} \left(\frac{\gamma'(z)}{\gamma(z)} \right)^q \sin^{s-q} \theta(\gamma z) dz^q \\ &= y(z)^{s-q} \left(\frac{1}{z^q |z|^{s-q}} \chi_+ + \sum_D \frac{\gamma'(z)^q |\gamma'(z)|^{s-q}}{\gamma(z)^q |\gamma(z)|^{s-q}} \right) dz^q + \\ &\quad y(z)^{s-q} \left(\frac{1}{z^q |z|^{s-q}} \chi_- + \sum_G \frac{\gamma'(z)^q |\gamma'(z)|^{s-q}}{\gamma(z)^q |\gamma(z)|^{s-q}} \right) dz^q \end{aligned}$$

and the q -form on $\mathcal{S}_l^{(r)}$:

$$A_{l,q}^R(s, z) = y(z)^{s-q} \left(\frac{1}{z^q |z|^{s-q}} \chi_+ + \sum_D \frac{\gamma'(z)^q |\gamma'(z)|^{s-q}}{\gamma(z)^q |\gamma(z)|^{s-q}} \right) dz^q$$

$$(17) \quad = l^q (\sin l \operatorname{Im} w)^{s-q} (\chi + \sum_R (\tilde{\gamma}' w)^q |\tilde{\gamma}' w|^{s-q}) dw^q,$$

with $w = \frac{1}{l} \log z$ and χ the characteristic function of $w(\{\operatorname{Re} z > 0\})$.

Theorem 5.2. *Let $S_l = \Gamma_l \backslash H$ be a family of geometrically finite hyperbolic surfaces degenerating to the surface S with only one geodesic c_l being pinched and with S_l having only one funnel F_1 ; Γ_l contains the transformation $\sigma_l(z) = e^l z$ corresponding to c_l and the right half-collar for c_l is in $S_l \setminus F_1$. Let $S_0 = \Gamma \backslash H$ be the component of S containing p , the cusp arising from the right half-collar for c_l , $p = \infty$. Let $w = \frac{1}{l} \log z$ and write again with a little misuse of notation $A_{l,q}(s, w) = a_{l,q}(s, w) dw^q$ and the corresponding q -automorphic form (for $\tilde{\Gamma}_l$) $\hat{a}_{l,q}(s, w) = \operatorname{Im}(w)^q a_{l,q}(s, w)$. Denote by $\hat{a}_{l,q}^R$ the right half of $\hat{a}_{l,q}$. Then the family $(\frac{1}{l^s} \hat{a}_{l,q}^R(s, \pi_l(\cdot)))_l$ converges uniformly on compact subsets of S_0 and on compact subsets of $\operatorname{Re} s > 1$ to the Eisenstein series for weight $2q$:*

$$E_{\infty,q}(s, w) = \sum_{\Gamma_\infty \backslash \Gamma} (\operatorname{Im} \gamma w)^s \left(\frac{c\bar{w} + d}{cw + d} \right)^q.$$

Remark 5.2. *We have the analogous result: if we denote by $\hat{a}_{l,q}^L$ the left half of $\hat{a}_{l,q}$, then $\frac{1}{l^s} \hat{a}_{l,q}^L(s, \pi_l(\cdot))$ converges to $(-1)^q E_{\mathfrak{a},q}(s, \cdot)$, where $E_{\mathfrak{a},q}(s, \cdot)$ is the Eisenstein series of weight $2q$ (see Definition 3.1) associated to \mathfrak{a} , the other cusp arising from the left half-collar for c_l .*

Before proving this theorem we give some complements and its corollaries.

For $\operatorname{Re} s > 1$ let $b_q(s) = e^{i\pi q/2} \int_0^\pi (\sin u)^{s-2} e^{-iqu} du$. Note that $b_1(s) = k(s-1)$. The function b_q has the following properties (see e.g. [18]): $b_q(s+2) = \frac{s(s-1)}{s^2 - q^2} b_q(s)$, b_q admits a meromorphic continuation to all $s \in \mathbb{C}$, more precisely

$$b_q(s) = \pi 2^{-s+2} \frac{\Gamma(s-1)}{\Gamma(\frac{s+q}{2}) \Gamma(\frac{s-q}{2})}.$$

In order to be consistent with the definition of Kudla and Millson's hyperbolic Eisenstein series, we may use the normalized q -automorphic forms

$$(18) \quad \Xi_{l,q}(s, z) = \frac{1}{b_q(s)} A_{l,q}(s, z).$$

We have indeed $\Xi_{l,1}(s, z) = \Theta(s-1, z)$ defined in (7) Section 2.1.

We recall that the series (18) converges absolutely and locally uniformly for any $z \in H$ and $s \in \mathbb{C}$ with $\operatorname{Re} s > 1$, and that it is invariant with respect to Γ .

Let $\tilde{a}_{l,q}(s, z)$ (respectively, $\tilde{f}_{l,q}(s, z)$) be the q -automorphic form associated to $A_{l,q}(s, z)$ (respectively, $\Xi_{l,q}(s, z)$) via the correspondence (9). More precisely and with the notation of Theorem 5.2

$$\begin{aligned} A_{l,q}(s, z) &= \frac{\tilde{a}_{l,q}(s, z)}{y^q} dz^q \\ &= \frac{\hat{a}_{l,q}(s, w)}{w^q} dw^q, \\ \Xi_{l,q}(s, z) &= \frac{\tilde{f}_{l,q}(s, z)}{y^q} dz^q, \quad \hat{f}_{l,q}(s, w) = \frac{1}{b_q(s)} \hat{a}_{l,q}(s, w). \end{aligned}$$

A straightforward computation shows that these series satisfy the functional differential equations

$$\Delta_{2q}\tilde{a}_{l,q}(s, z) + s(1-s)\tilde{a}_{l,q}(s, z) = (s+q)(q-s)\tilde{a}_{l,q}(s+2, z),$$

and

$$(19) \quad \Delta_{2q}\tilde{f}_{l,q}(s, z) + s(1-s)\tilde{f}_{l,q}(s, z) = s(1-s)\tilde{f}_{l,q}(s+2, z);$$

and so, the series $A_{l,q}(s, z)$ satisfies the functional differential equation

$$\Delta_q^\pm A_{l,q}(s, z) - (s \pm q)(1 - s \pm q)A_{l,q}(s, z) = (s+q)(s-q)A_{l,q}(s+2, z),$$

and the series (18)

$$\Delta_q^\pm \Xi_{l,q}(s, z) - (s \pm q)(1 - s \pm q)\Xi_{l,q}(s, z) = s(s-1)\Xi_{l,q}(s+2, z).$$

Proposition 5.1. *The series $A_{l,q}(s, z)$ (as well as $\Xi_{l,q}(s, z)$) admits a meromorphic continuation to all of \mathbb{C} .*

Proof. There are different ways to prove this: one is to use the functional differential equation (18) and to apply the method developed in [20] (see also [18]). More precisely, from (19) we have

$$\tilde{f}_{l,q}(s, z) = s(1-s) \int_{\mathcal{D}_l} G_s^l(z, z', q) \tilde{f}_{l,q}(s+2, z') d\mu(z'),$$

where \mathcal{D}_l is a fundamental domain of S_l and $G_s^l(z, z', q)$ has been defined in proposition 3.1. Another way is to use [7]. We make precise another calculation which we will develop further. We can rewrite $\tilde{a}_{l,q}(s, z)$ as

$$\tilde{a}_{l,q}(s, z) = \sum_{\langle \sigma_l \rangle \setminus \Gamma_l} \left(\frac{c\bar{z} + d}{cz + d} \right)^q \left(\frac{\bar{\gamma}z}{\gamma z} \right)^{q/2} \left(\frac{\text{Im } \gamma z}{|\gamma z|} \right)^s$$

and using the Fourier development of $G_s(z, z', q)$ and the expansion of $\int_1^{e^l} G_s(z, iy', q) d \ln y'$ we obtain the result (see [7] corollary 4.2, p. 188). \square

We recall from Section 3.3 that

Remark 5.3. *The possible poles of the resolvent $G_s^l(z, z', q)$ in $\text{Re } s > 1/2$ are simple, in finite number and at $s_{1,q}^l, \dots, s_{n_l,q}^l \in (\frac{1}{2}, 1)$ corresponding to the eigenvalues $\lambda_{k,q}^l$, $1 \leq k \leq n_l$ of the Laplacian $-\Delta_{2q}$ on S_l .*

In particular, $\tilde{f}_{l,q}(s, z)$ is holomorphic in $\Sigma_l = \{\text{Re } s > 1/2\} \setminus \{s_{1,q}^l, \dots, s_{n_l,q}^l\}$.

We can then refine Theorem 5.2 to the following.

Theorem 5.3. *Let λ_k^q , $1 \leq k \leq n$, be the eigenvalues of the Laplacian $-\Delta_{2q}$ on S_0 such that $0 < \lambda_1^q < \lambda_2^q < \dots < \lambda_n^q < 1/4 \leq \lambda_{n+1}^q$ and the corresponding $s_k^q = \frac{1}{2} + \sqrt{\frac{1}{4} - \lambda_k^q}$ with $\text{Re } s_k^q \geq 1/2$ and let $\Sigma = \{\text{Re } s > 1/2\} \setminus \{s_1^q, \dots, s_n^q\}$.*

(1) *In the case where the geodesic c_l is not separating, without loss of generality we can suppose that the two limiting cusps are represented by ∞ and 0, the family $(\frac{1}{s} \tilde{f}_{l,q}(s, \pi_l(\cdot)))_l$ converges uniformly on compact subsets of S_0 and*

$$\text{on compact subsets of } \Sigma \text{ to } \frac{1}{b_q(s)} E_{\infty,q}(s, \cdot) + \frac{(-1)^q}{b_q(s)} E_{0,q}(s, \cdot).$$

- (2) In the case where c_l is the geodesic boundary of a funnel, the family $(\frac{1}{l^s} \hat{f}_{l,q}(s, \pi_l(\cdot)))_l$ converges uniformly on compact subsets of S_0 and Σ to $\frac{1}{b_q(s)} E_{\infty,q}(s, \cdot)$.

Remark 5.4. In the case c_l is separating and different from the geodesic boundary of a funnel, as in Theorem 5.2 and Remark 5.2, we can define the right and left half of $\frac{1}{l^s} \hat{f}_{l,q} = \frac{1}{l^s} \hat{a}_{l,q}^R + \frac{1}{l^s} \hat{f}_{l,q}^L$ each series in the right member converges to the Eisenstein series of the corresponding cusp.

Remark 5.5. We remark that as σ_l is the geodesic in the funnel, it follows that $D = \langle \sigma_l \rangle \backslash (\Gamma_l - \langle \sigma_l \rangle)$ and $G = \emptyset$.

We have seen that in the infinite volume case, $s = 1$ is a regular value for $E_{\infty,q}(s, z)$ for every $q \in \mathbb{N}$. In the finite volume case, we know that $s = 1$ is a simple pole for $E_{\infty,0}(s, z)$, which is no longer the case for $q \geq 1$:

Remark 5.6. For $q \in \mathbb{N}^*$, $s = 1$ is a regular value of $E_{\infty,q}(s, z)$.

Proof. This comes from the Fourier development of $E_{\infty,q}(s, z)$: we have ([7], p. 175)

$$E_{\infty,q}(s, z) = y^s + \varphi_q(s) y^{1-s} + \sum_{m \neq 0} a_m(y, s)_q e^{2i\pi m x},$$

with $\varphi_q(s) = \frac{\Gamma^2(s)}{\Gamma(s-q)\Gamma(s+q)} e^{-\pi i q} \varphi_0(s)$. The function φ_0 has a simple pole at $s = 1$ (with residue $\text{Vol}(S_0)/\pi$) and $1/\Gamma(s-q)$ has a zero at $s = 1$. \square

We now give the poles of $b_q(s)$:

Lemma 5.5. For an even q , the poles of $b_q(s)$ are simple and at the numbers $s = 1 - 2k$, $k \in \mathbb{N}$.

For an odd q , the poles of $b_q(s)$ are simple and at the numbers $s = -2k$, $k \in \mathbb{N}$.

From the preceding results and Theorem (5.3) we deduce

Corollary 5.1. With the preceding notation,

- (1) Suppose q is odd. If the geodesic c_l is not separating (respectively, the geodesic boundary of a funnel) the family $(\frac{1}{l^s} \hat{a}_{l,q}(s, \pi_l(\cdot)))_l$ converges uniformly on compact subsets of S_0 and on compact subsets of Σ to $E_{\infty,q}(s, \cdot) + (-1)^q E_{0,q}(s, \cdot)$ (respectively, $E_{\infty,q}(s, \cdot)$).
- (2) Suppose q is even, we have the same results as the preceding, on replacing Σ by $\Sigma \setminus \{1\}$.

Remark 5.7. For $q = 1$ we obtain the result announced in the Introduction.

5.2. Proofs of the main theorems and final remarks.

Proof of Theorem 5.2.

It is enough to demonstrate the convergence for β a relatively compact set in the fundamental domain \mathcal{F} . Given $\epsilon > 0$, denote by G_ϵ the set of cosets and representatives for $\langle z \mapsto z + 1 \rangle \backslash \Gamma$ such that $\sup \text{Im } A(\mathcal{F}) < \epsilon$ for $[A] \notin G_\epsilon$ and let R_l be the corresponding cosets of $\langle z \mapsto z + 1 \rangle \backslash \tilde{\Gamma}_l$ with the corresponding representatives.

Lemma 5.6. The set G_ϵ is finite.

Proof. We have to show that the number of representatives for $\langle z \mapsto z + 1 \rangle \backslash \Gamma$ such that $\sup \operatorname{Im} A(\mathcal{F}) \geq \epsilon$ is finite. Take a real a such that the cusp for the parabolic point ∞ , $\{0 \leq \operatorname{Re} z < 1, \operatorname{Im} z \geq a\} \subset \mathcal{F}$. As for all $g \in G_\epsilon$, different from the identity, $g(\mathcal{F})$ and \mathcal{F} are disjoint, we have $\sup \operatorname{Im} g(\mathcal{F}) \geq \epsilon$ and $\sup \operatorname{Im} g(\mathcal{F}) \leq a$. Let \mathcal{S} a system of representatives of $\langle z \mapsto z + 1 \rangle \backslash \Gamma$ such that the disjoint sets $\{[A](\mathcal{F}), [A] \in \mathcal{S}\}$ recover $B = \{0 \leq \operatorname{Re} z < 1\}$. The disjoint sets $\{g(\mathcal{F}), g \in G_\epsilon\}$ recovering the compact set $\{0 \leq \operatorname{Re} z \leq 1, \epsilon \leq \operatorname{Im} z \leq a\}$ of B , the cardinal of G_ϵ must be finite. \square

The cosets G_ϵ of Γ satisfy (modulo the action of $\langle z \mapsto z + 1 \rangle$) $\{0 \leq \operatorname{Re} w < 1, \operatorname{Im} w > \epsilon\} \subset \cup_{A \in G_\epsilon} A(\mathcal{F})$; thus for l sufficiently small the cosets R_l satisfy (modulo the action of $\langle z \mapsto z + 1 \rangle$) $\{0 \leq \operatorname{Re} w < 1, 2\epsilon < \operatorname{Im} w < \pi/2l\} \subset \cup_{A \in R_l} A(\mathcal{F}_l)$ (this is a consequence of the convergence on compact subsets of the f_l and of the fact that \mathcal{F}_l contains the right half-collar for c_l , $\{0 \leq \operatorname{Re} w < 1, c(l) \leq \operatorname{Im} w < \pi/2l\}$, see Lemma 5.3. Now for a right coset $[A] \in R - R_l$, we have that $A(\mathcal{F}_l)$ lies below the c_l geodesic $\{\operatorname{Im} w = \pi/2l\}$ and is disjoint modulo the action $\langle z \mapsto z + 1 \rangle$ from $\{0 \leq \operatorname{Re} w < 1, 2\epsilon < \operatorname{Im} w < \pi/2l\}$, since the latter is covered by the R_l cosets. Thus for $[A] \in R - R_l$, modulo the action of $\langle z \mapsto z + 1 \rangle$, we have $A(\mathcal{F}_l) \subset \{0 \leq \operatorname{Re} w < 1, \operatorname{Im} w < 2\epsilon\}$. For $w \in \beta$, using (17), we write

$$\begin{aligned} \frac{1}{l^s} \hat{a}_{l,q}^R(s, w) &= \frac{1}{l^{s-q}} (\operatorname{Im} w)^q (\sin l \operatorname{Im} w)^{s-q} \left(\chi + \sum_R (\tilde{\gamma}' w)^q |\tilde{\gamma}' w|^{s-q} \right) \\ &= \frac{1}{l^{s-q}} (\operatorname{Im} w)^q (\sin l \operatorname{Im} w)^{s-q} \left(\chi + \sum_{R_l} (\tilde{\gamma}' w)^q |\tilde{\gamma}' w|^{s-q} \right) \\ &+ \frac{1}{l^{s-q}} (\operatorname{Im} w)^q (\sin l \operatorname{Im} w)^{s-q} \left(\sum_{R-R_l} (\tilde{\gamma}' w)^q |\tilde{\gamma}' w|^{s-q} \right). \end{aligned}$$

Now on β for l sufficiently small, χ is identically unity and because of Lemma 5.4, the coset representatives for R_l approximate the coset representatives for G_ϵ . Thus for $\frac{1}{l^s} \hat{a}_{l,q}^R(s, \cdot)$, in the last equality, the first sum is uniformly close to the corresponding terms for $E_{\infty,q}(s, \cdot)$.

The principal problem lies in estimating the second sum: it remains to show that

$$\lim_{l \rightarrow 0} \sum_{R-R_l} |\tilde{\gamma}' w|^\sigma = 0,$$

where $\sigma = \operatorname{Re} s$.

$$\sum_{R-R_l} |\tilde{\gamma}' w|^\sigma = \sum_{w^{-1}(R-R_l)w} |z_l \frac{\gamma' z_l}{\gamma z_l}|^\sigma \leq |z_l|^\sigma \sum_{w^{-1}(R-R_l)w} |\gamma' z_l|^\sigma,$$

where $w = \frac{1}{l} \log z_l$ and $\gamma \mathcal{F}_l \subset w^{-1}(\{0 \leq \operatorname{Re} w < 1, \operatorname{Im} w < 2\epsilon\} \cap \beta)$. We deduce

$$\sum_{R-R_l} |\tilde{\gamma}' w|^\sigma \leq \frac{1}{\sin^\sigma(l \operatorname{Im} w)} \sum_{w^{-1}(R-R_l)w} \operatorname{Im}^\sigma(\gamma z_l).$$

Let $\epsilon_0 \in]0, \sinh^{-1} 1[$ such that for all l sufficiently small and all $w \in \beta$, $B(z_l, \epsilon_0) \subset \mathcal{F}_l$, then $\cup_{\gamma \in w^{-1}(R-R_l)w} B(\gamma z_l, \epsilon_0) \subset \{z, 0 < \arg z < 2\epsilon l, 1 \leq |z| \leq e^l\}$. Moreover there exists a Λ_{ϵ_0} independent of $z_0 \in H$ such that

$$\int_{B(z_0, \epsilon_0)} y^\sigma \frac{dx dy}{y^2} = \Lambda_{\epsilon_0} y(z_0)^\sigma;$$

this is a particular case of the following Proposition 5.2, needed further in a more general case. Hence

$$\sum_{w^{-1}(R-R_l)w} (\operatorname{Im} \gamma z_l)^\sigma = \frac{1}{\Lambda_{\epsilon_0}} \sum_{w^{-1}(R-R_l)w} \int_{B(\gamma z_l, \epsilon_0)} y^\sigma \frac{dx dy}{y^2}.$$

Now (see, e.g. [27], p. 102):

Lemma 5.7. *The multiplicity of the projection map $H \rightarrow \Gamma \backslash H$ restricted to $B(z_0, \eta)$ with $2\eta < c_0$ is at most $M\rho^{-2}(z_0)$, where M is a constant and $\rho(z_0)$ the injectivity radius at z_0 .*

For the convenience of the reader, we give a detailed proof.

Proof. If $B(z_0, \eta) \cap B(\gamma z_0, \eta) \neq \emptyset$, $\gamma \in \Gamma$, then $d(z_0, \gamma z_0) < 2\eta < c_0$ and z_0 is in a cusp region or the collar for a short geodesic. Let $c = c_0/2$. Then $\rho(z_0) < c$. Denote by $m(\eta)$ the multiplicity of the projection restricted to $B(z_0, \eta)$. As $2\eta + \rho(z_0) < 3c$ and the $B(\gamma z_0, \rho(z_0))$ are disjoint, we have

$$m(\eta)\mu(B(z_0, \rho(z_0))) \leq \mu(B(z_0, 3c)).$$

$$\text{Hence } m(\eta) \leq \frac{\cosh 3c - 1}{\cosh \rho(z_0) - 1} \leq 2(\cosh 3c - 1)\rho(z_0)^{-2}. \quad \square$$

Now we have

$$\begin{aligned} \sum_{\gamma \in w^{-1}(R-R_l)w} (\operatorname{Im} \gamma z_l)^\sigma &\leq \frac{1}{\Lambda_{\epsilon_0}} m(\epsilon_0) \int_{\cup_{\gamma \in w^{-1}(R-R_l)w} B(\gamma z_l, \epsilon_0)} y^{\sigma-2} dx dy \\ &\leq \frac{A(c_0)}{\Lambda_{\epsilon_0}} \rho^{-2}(z_l) \int_{\{z, 0 < \arg z < 2\epsilon l, 1 \leq |z| \leq e^l\}} y^{\sigma-2} dx dy, \end{aligned}$$

with $A(c_0) = 2(\operatorname{ch} \frac{3c_0}{2} - 1)$.

Then

$$(20) \quad \sum_{w^{-1}(R-R_l)w} (\operatorname{Im} \gamma z_l)^\sigma \leq \frac{A(c_0)}{\Lambda_{\epsilon_0}} \rho^{-2}(z_l) \frac{e^{l\sigma} - 1}{\sigma} \frac{(2\epsilon l)^{\sigma-1}}{\sigma - 1}.$$

Moreover, as $B(z_l, \epsilon_0) \subset \mathcal{F}_l$, $\rho(z_l) \geq \epsilon_0$ and the conclusion follows.

Proof of Theorem 5.3.

The main point of the proof is to use Vitali's theorem, as in the finite volume case, we follow [4], Proof of Theorem 4.2. More precisely, first we will recall a definition.

Definition 5.1. *A family (f_n) of meromorphic functions on a domain \mathcal{O} in \mathbb{C} is called bounded in \mathcal{O} if*

- (1) $\exists (z_m)_m$ a discrete subset in \mathcal{O} ,
- (2) $\forall K$ compact set of $\mathcal{O} \setminus \{z_m\}$, $\exists n(K)$, $\forall n \geq n(K)$, f_n has no pole in K ,
- (3) $M_K = \sup_{n \geq n(K)} (\sup_{z \in K} |f_n(z)|) < +\infty$.

As before, β will denote a relatively compact set in the fundamental domain \mathcal{F} of S_0 . The family $(\frac{1}{s} \hat{f}_{l,q}(s, \pi(w)))_l$, where $\hat{f}_{l,q}(s, w) = \frac{1}{b_q(s)} \hat{a}_{l,q}(s, w)$, is a family of meromorphic functions on $\Sigma = \{\operatorname{Re} s > 1/2\} \setminus \{s_1^q, \dots, s_n^q\}$. Now, by Theorem (5.2), for every compact $C \subset \{\operatorname{Re} s > 1\}$ this family converges uniformly (a fortiori simply) on $C \times \beta \subset \{\operatorname{Re} s > 1\} \times \mathcal{F}$ to

- (1) $\frac{1}{b_q(s)}(E_{\infty,q}(s, \cdot) + (-1)^q E_{0,q}(s, \cdot))$ in case (1) of Theorem (5.2),
 (2) $\frac{1}{b_q(s)} E_{\infty,q}(s, \cdot)$ in case (2).

We will show that for every compact set K in Σ , there exists $l(K)$ such that for all l less than $l(K)$, $s \mapsto \frac{1}{l^s} \hat{f}_{l,q}(s, \pi(\cdot))$ has no pole on K and $M_K = \sup_{l \leq l(K)} (\sup_{s \in K} \|\frac{1}{l^s} \hat{f}_{l,q}(s, \pi(\cdot))\|_{\infty, \beta}) < +\infty$ —and then we can conclude the result.

The outline of the proof being independent of q , we will omit writing q and only indicate the changes needed in the case of general q if necessary. In particular in this paragraph we can write Δ instead of Δ_{2q} .

As in the finite volume case, we start from the equality

$$\frac{1}{l^s} \tilde{f}_l(s, z) = s(1-s) \int_{\mathcal{D}_l} G_s(z, z') \tilde{f}_l(s+2, z') / l^s d\mu(z');$$

where, to simplify notation, we omit writing $q = 0$; but we can write $G_s^l(z, z')$ to insist on the dependence in l . The definition and properties of $G_s^l(z, z')$ are given in Section 3.3. We start from Remark (5.3), the poles of $G_s^l(z, z')$ in $\text{Re } s > 1/2$ are simple, in finite number and in $(\frac{1}{2}, 1)$, they correspond to the small eigenvalues ($< 1/4$) of the Laplacian on S_l . As l tends to zero, these poles tend to the corresponding poles of the resolvent of the limit surface; in other words the small eigenvalues tend to the small eigenvalues (see [26],[7]). Now take K a compact of Σ , there exists $l(K)$ such that for $l \leq l(K)$, all the finite number of poles of $G_s^l(z, z')$ are in the open set $\Sigma \setminus K$, from the same Remark (5.3), $s \mapsto \frac{1}{l^s} \hat{f}_{l,q}(s, \pi(\cdot))$ has no pole on K . We will suppose C is a compact subset of S_0 , $Y \subset S_0$ is such that $\pi_l(Y)$ is a standard collar, and $\pi_l(Y) \cap \pi_l(C) = \emptyset$, $X = S_0 \setminus Y$, K' a compact subset of $\Sigma' = \Sigma \setminus \{1\}$. Decompose the preceding integral and write

$$\frac{1}{l^s} \tilde{f}_l(s, \pi_l(w)) = I_l + J_l,$$

where

$$I_l = \int_{\pi_l(Y)} G_s(\pi_l(w), z') \tilde{f}_l(s+2, z') / l^s d\mu(z'), \quad J_l = \int_{\pi_l(X)} G_s(\pi_l(w), z') \tilde{f}_l(s+2, z') / l^s d\mu(z').$$

Now the main step is to estimate $G_s(z, z', q)$. We apply [7] Corollary 1.3, p. 151:

Proposition 5.2. *If $\Delta_{2q} f = s(s-1)f$ in some non-Euclidean disc B_r of radius r about $z_0 \in H$, then f has the mean value property:*

$$f(z_0) = \frac{1}{m_q(r, s)} \int_{B_r} f(z) \left(\frac{z - \bar{z}_0}{z_0 - \bar{z}} \right)^q d\mu(z) \text{ where } m_q(r, s) = 2\pi \int_1^r \text{sh } r P_{s,q}(r) \approx \pi r^2 \text{ as } r \rightarrow 0;$$

$P_{s,q}(r) = (1 - \tanh^2 \frac{r}{2})^s F(s-q, s+q, 1; \tanh^2 \frac{r}{2})$ and F is the Gauss's hypergeometric function.

to conclude that

$$(21) \quad \forall w \in C, \forall z' \in \pi_l(Y), \forall s \in K', |G_s(\pi_l(w), z', q)| \leq O(1) \|G_s(\pi_l(w), \cdot, q)\|_{L_2}$$

(see also [4] Lemme 4.4).

The next point is to show that

$$(22) \quad \forall (w, s) \in C \times K', \|G_s(\pi_l(w), \cdot, q)\|_{L_2} = O(1),$$

the constants in question depend only on the compact sets. We can use [26], theorem 15(1), but we give here another presentation using the spectral decomposition.

In effect, we have, using the notation in Section 3.3, in particular Proposition (3.3), where $\{\varphi_n\}$ is a complete orthonormal basis of eigenfunctions of Δ with corresponding eigenvalues $\{\lambda_n\}$ and where we don't note the dependence on l of the eigenfunctions, cuspidal and funnel Eisenstein series:

Lemma 5.8. *For $a > 1$*

$$\begin{aligned} G_s^l(\pi_l(w), z) &= G_a^l(\pi_l(w), z) + \sum_n \left[\frac{1}{s(1-s) - \lambda_n} - \frac{1}{a(1-a) - \lambda_n} \right] \varphi_n(\pi_l(w)) \overline{\varphi_n(z)} + \\ &\frac{1}{4\pi} \sum_j \int_{-\infty}^{+\infty} \left[\frac{1}{s(1-s) - (1/4 + t^2)} - \frac{1}{a(1-a) - (1/4 + t^2)} \right] E_j^c\left(\frac{1}{2} + it, \pi_l(w)\right) \overline{E_j^c\left(\frac{1}{2} + it, z\right)} dt + \\ &\frac{1}{4\pi} \sum_k \int_{-\infty}^{+\infty} \left[\frac{1}{s(1-s) - (1/4 + t^2)} - \frac{1}{a(1-a) - (1/4 + t^2)} \right] \int_1^{e^{tk}} E_k^f\left(\frac{1}{2} + it, \pi_l(w), b\right) \overline{E_k^f\left(\frac{1}{2} + it, z, b\right)} db dt, \end{aligned}$$

all the sums are finite.

Proof. We can use and follow the method in for example [13] (and the references therein), [1], p. 103, to prove it, but not to be too long we can also remark as in [16], p. 87-88, [1], Theorem 6.2, p. 82, that, with E_α corresponding to any E_j^c, E_k^f

$$\Delta E_\alpha\left(\frac{1}{2} + it, \cdot\right) + (1/4 + t^2)E_\alpha\left(\frac{1}{2} + it, \cdot\right) = 0,$$

and $E_\alpha\left(\frac{1}{2} + it, \cdot\right) \in \rho^{-1/2}L^2(S_l)$.

So for $u, \operatorname{Re} u > 1$ ($\operatorname{Re} u - 1/2 > 1/2$)

$$R_u(\Delta + u(1-u))E_\alpha = E_\alpha.$$

We deduce that,

$$\left(\overline{G_u^l(\pi_l(w), \cdot)}, E_\alpha\left(\frac{1}{2} + it, \cdot\right) \right) = \frac{1}{\bar{u}(1-\bar{u}) - (1/4 + t^2)} \overline{E_\alpha\left(\frac{1}{2} + it, \pi_l(w)\right)}$$

and in a similar way as $\varphi_n \in L^2(S_l)$,

$$\frac{1}{u(1-u) - s_n(1-s_n)} \varphi_n(\pi_l(w), b) = \int_{S_l} G_u^l(\pi_l(w), z) \varphi_n(z) d\mu(z).$$

Applying Proposition 3.3 to $\overline{G_s^l(\pi_l(w), \cdot)} - \overline{G_a^l(\pi_l(w), \cdot)}$ by complex conjugation we obtain the lemma for $\operatorname{Re} s > 1$ and by analytic continuation, to $\operatorname{Re} s > 1/2$. \square

From this we now deduce, writing $\lambda_u = u(1-u)$,

Lemma 5.9.

$$\begin{aligned} \|G_s^l(\pi_l(w), \cdot) - G_a^l(\pi_l(w), \cdot)\|_{L^2}^2 &= |\lambda_a - \lambda_s|^2 \sum_n \frac{|\varphi_n(\pi_l(w))|^2}{|\lambda_s - \lambda_n|^2 |\lambda_a - \lambda_n|^2} + \\ &|\lambda_a - \lambda_s|^2 \frac{1}{4\pi} \int_{-\infty}^{+\infty} \frac{|E_j^c(1/2 + it, \pi_l(w))|^2}{|\lambda_s - (1/4 + t^2)|^2 |\lambda_a - (1/4 + t^2)|^2} dt + \\ &|\lambda_a - \lambda_s|^2 \frac{1}{4\pi} \int_{-\infty}^{+\infty} \frac{1}{|\lambda_s - (1/4 + t^2)|^2 |\lambda_a - (1/4 + t^2)|^2} \int_1^{e^{tk}} |E_k^f(1/2 + it, \pi_l(w), b)|^2 db dt. \end{aligned}$$

So there exists a constant M , depending only on the compact set K' of Σ' , such that

$$\|G_s(\pi_l(w), \cdot, l) - G_a(\pi_l(w), \cdot, l)\|_{L^2}^2 \leq \frac{|\lambda_a - \lambda_s|^2}{M} \|G_a(\pi_l(w), \cdot, l)\|_{L^2}^2.$$

The fact that we have for $a > 1$, $\|G_a(\pi_l(w), \cdot, q)\|_{L^2} = O(1)$, the constants in question depend only on the compact sets, is an adaptation of [13] or [14].

Now we are going to prove that $I_l = O(1)$ and $J_l = O(l^2)$.

With the same proof as [4] Lemma 4.3, we have $\int_{\pi_l(Y)} |\tilde{f}_l(s+2, z)| d\mu(z) = O(1)l^{\operatorname{Re} s}$ true for $\operatorname{Re} s > 1/2$.

Recall that $S_l = R \cup C_1 \cup \dots \cup C_{n_c} \cup F_1$ where here, we denote by R the compact core. $J_l = \int_{\pi_l(X) \cap R} G_s(z, z') \tilde{f}_l(s+2, z')/l^s d\mu(z') + \int_{\pi_l(X) \cap (C_1 \cup \dots \cup C_{n_c} \cup F_1)} G_s(z, z') \tilde{f}_l(s+2, z')/l^s d\mu(z')$.

By Theorem 5.2 and (22), $\int_{\pi_l(X) \cap K} G_s(z, z') \tilde{f}_l(s+2, z')/l^s d\mu(z') = O(l^2)$.

For $z' \in C_1 \cup \dots \cup C_{n_c} \cup F_1$ we have the same estimates as in (21). Now let's verify that we have also $\int_{R \cap (C_1 \cup \dots \cup C_{n_c} \cup F_1)} \tilde{f}_l(s+2, z')/l^s d\mu(z') = O(1)$.

We have

$$\begin{aligned} \int_{R \cap (C_1 \cup \dots \cup C_{n_c} \cup F_1)} \tilde{f}_l(s+2, z') d\mu(z') &\leq \frac{2}{|k(s+1)|} \int_{1 \leq r \leq e^r} (\sin \theta)^{\operatorname{Re} s} \frac{dr d\theta}{r} \\ &\leq \frac{2}{|k(s+1)|} l \times l^{\operatorname{Re} s+1} \end{aligned}$$

This allows concluding that $I_l = O(1)$ and $J_l = O(l^2)$. This ends like in [5].

Remark 5.8. 1) We refer also to the result of Fay [6], final remark, p. 201–202.
2) It is possible to establish a link between hyperbolic Eisenstein series and generalized eigenfunctions (I thank F. Naud for this suggestion) as presented in Borthwick ([1], p. 68)

Let $E_{f_l}(s, z, t)$ correspond to the funnel with pinching boundary geodesic. We use the first point of this remark and its notations to write $E_{f_l}(s, z, t) = b^s E^f(s, z, b) = \sum_{n \in \mathbb{Z}} F_{n,0}(z, s) b^{\bar{n}}$, where $b = e^t$ and $\bar{n} = 2i\pi n/l$; and conclude that as $l \rightarrow 0$,

$$\frac{1}{l} \int_0^l E_{f_l}(s, z, t) dt \rightarrow E_\infty(s, z).$$

Application. One of the applications we can think about is in studying the degeneration of the residues of the hyperbolic Eisenstein series. Let us give an example: we look at the case of a degenerating family of compact Riemann surfaces, a non-separating geodesic being pinched, with the family of scalar-valued hyperbolic Eisenstein series degenerating. Remember that we obtained that the series $\frac{1}{l^s} A_l(s, z)$ converges uniformly on compact subsets of S_0 to $E_\infty(s, z) + E_0(s, z)$. The last sum of Eisenstein series has no poles on $[1/2, 1]$ except at $s = 1$ and a finite number of $s_k = \frac{1}{2} + \sqrt{\frac{1}{4} - \lambda_k}$ where the λ_k correspond to the residual spectrum. In other words, if $(\lambda_k(l))$ converges to λ_k , where λ_k is a small cuspidal eigenvalue, then $\operatorname{Res}(\frac{1}{l^s} A_l(s, z)) \rightarrow 0$, otherwise $\operatorname{Res}(\frac{1}{l^s} A_l(s, z)) \rightarrow \operatorname{Res}(E_\infty(s, z) + E_0(s, z))$. There are many obstacles to carrying this calculation further. First of all, we are only dealing with small eigenvalues and so here there is no hope to characterize the embedded eigenvalues through degeneration. Moreover, we need to take into account the multiplicity of an eigenvalue $\lambda_k(l)$. So the easiest result we can obtain is a characterization of the residual spectrum (recall that an eigenvalue in the

residual spectrum is simple) with $\alpha_k(l) = \int_{c_l} \psi_{k,l}(z) d\mu(z)$ where the eigenfunction $\psi_{k,l}$ is associated to the eigenvalue $\lambda_k(l)$; if λ_k is a pole of $E_\infty(s, z) + E_0(s, z)$, then $\alpha_k(l) = O(l^{1/2 + \sqrt{1/4 - \lambda_k(l)}})$, and otherwise $\alpha_k(l) = o(l^{1/2 + \sqrt{1/4 - \lambda_k(l)}})$.

References

- [1] D. Borthwick, *Spectral Theory of Infinite-Area Hyperbolic Surfaces*, volume 256, Progress in Mathematics, Birkhäuser, Boston, MA, 2007.
- [2] G. Carron, L^2 harmonic forms on non-compact Riemannian manifolds, *Surveys in Analysis and Operator Theory* (Canberra, 2001).
- [3] J. Elstrodt, Die Resolvente zum Eigenwertproblem der automorphen Formen in der hyperbolischen Ebene, Teil I. *Math. Ann.* **203** (1973) 295–330, Teil II. *Math. Z.* **132** (1973) 99–134, Teil III. *Math. Ann.* **208** (1974) 99–132.
- [4] T. Falliero, Dégénérescence de séries d’Eisenstein hyperboliques, *Math. Ann.* **339** (2007), 341–375.
- [5] ———, Harmonic differentials and infinite geodesics joining two punctures on a Riemann surface, *Trans. Amer. Math. Soc.* **363** (2011), 3473–3488.
- [6] J. D. Fay, *Theta Functions on Riemann Surfaces*, Lecture Notes in Math., vol. 352, Springer-Verlag, Berlin 1973.
- [7] ———, Fourier coefficients of the resolvent for a Fuchsian group, *J. Reine Angew. Math.* **293** (1977), 143–203.
- [8] D. Garbin, J. Jorgenson and M. Munn, On the appearance of Eisenstein series through degeneration, *Comment. Math. Helv.* **83** (2008), 701–721.
- [9] D. Garbin and A.-M. v. Pippich, On the Behavior of Eisenstein Series Through Elliptic Degeneration, *Commun. Math. Phys.* **292** (2009), 511–528.
- [10] P. Gérardin, *Formes automorphes associées aux cycles géodésiques des surfaces de Riemann hyperboliques*, Lecture Notes in Math., 901, November 1980.
- [11] D. A. Hejhal, *The Selberg Trace Formula for PSL(2,R)*, vol. 1. Lecture Notes in Math., vol. 548, Springer-Verlag, Berlin 1976.
- [12] ———, *The Selberg Trace Formula for PSL(2,R)*, vol. 2. Lecture Notes in Math., vol. 1001, Springer-Verlag, Berlin 1983.
- [13] ——— *Modular Forms. A continuity method for spectral theory on Fuchsian groups*. Chichester, Ellis-Horwood Edition, 1984.
- [14] ———, Regular b-groups, Degenerating Riemann surfaces and Spectral Theory, *Mem. Am. Math. Soc.*, **88** (437), November 1990.
- [15] ———, Kernel functions, Poincaré series, and LVA, *Contemp. Math.* **256** (2000), 173–201.
- [16] H. Iwaniec *Introduction to the Spectral Theory of Automorphic Forms*, Biblioteca de la Revista Matemática Iberoamericana, 1995.
- [17] L. Ji, Spectral degeneration of hyperbolic Riemann surfaces, *J. Differ. Geom.* **38** (1993), 263–313.
- [18] J. Jorgenson, J. Kramer and A.-M. v. Pippich, On the spectral expansion of hyperbolic Eisenstein series, *Math. Ann.* **346** (2010), 931–947.
- [19] J. Jorgenson, A.-M. v. Pippich, L. Smajlović, On the wave representation of hyperbolic, elliptic, and parabolic Eisenstein series, *Adv. Math.* **288** (2016), 887–921.
- [20] J. Kudla and S. Millson, Harmonic differentials and closed geodesics on a Riemann surface, *Invent. Math.* **54** (1979), 193–211.
- [21] A. McIntyre and L.A. Takhtajan, Holomorphic factorization of determinants of Laplacians on Riemann surfaces and a higher genus generalization of Kronecker’s first limit formula, *Geom. Funct. Anal.* **16** (2006), 1291–1323.
- [22] K. Obitsu, The asymptotic Behavior of Eisenstein Series and a Comparison of the Weil–Peterson and the Zograf–Takhtajan Metrics, *Publ. Res. Inst. Math. Sci.* **37** (2001), 459–478.
- [23] ———, Asymptotics of degenerating Eisenstein series, Infinite dimensional Teichmüller spaces and moduli spaces, 115–126, *RIMS Kōkyūroku Bessatsu*, B17, Res. Inst. Math. Sci. (RIMS), Kyoto, 2010.
- [24] S. Patterson, The Laplacian operator on a Riemann surface, I, II, III, *Compositio Math.* **31** (1975), 83–107, **32** (1976), 71–112, **33** (1976), 227–310.

- [25] M. Risager, On the distribution of modular symbols for compact surfaces, *Int. Math. Res. Not.* **41** (2004), 2125–2146.
 - [26] M. Schulze, On the resolvent of the Laplacian on functions for degenerating surfaces of finite geometry, *J. Funct. Anal.* **236** (2006), 120–160.
 - [27] S. A. Wolpert, Spectral limits for hyperbolic surfaces, II, *Invent. Math.* **108** (1992), 91–129.
- Email address:* `therese.falliero@univ-avignon.fr`