



HAL
open science

Nurses' and Midwives' clinical leadership development needs: A mixed methods study

Mary Casey, Martin S Mcnamara, Gerard M. Fealy, Ruth Geraghty, Philomena A Halligan, Margaret Treacy, Michelle M Butler, Maree Johnson

► **To cite this version:**

Mary Casey, Martin S Mcnamara, Gerard M. Fealy, Ruth Geraghty, Philomena A Halligan, et al.. Nurses' and Midwives' clinical leadership development needs: A mixed methods study. *Journal of Advanced Nursing*, 2011, 67 (7), pp.1502. 10.1111/j.1365-2648.2010.05581.x . hal-00614756

HAL Id: hal-00614756

<https://hal.science/hal-00614756>

Submitted on 16 Aug 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Nurses' and Midwives' clinical leadership development needs: A mixed methods study

Journal:	<i>Journal of Advanced Nursing</i>
Manuscript ID:	JAN-2010-0185.R4
Manuscript Type:	Manuscript/Short Report
Keywords:	

SCHOLARONE™
Manuscripts
Copy

Feedback Addressed on Journal Manuscript ID – JAN 2010-0185 R3

11th November, 2010

Editor's & Reviewer's Feedback	Actions Taken
<p>1. Check the entire paper for grammar. For example, this sentence, The roles of clinical leaders have been specified (Cooke, 2001a and 2001b, Davidson et al. 2006), and its impact on care outlined, should be revised to read: ... and their impact on care (you are referring to roles, not role).</p>	<p>The entire paper has been thoroughly review for grammar.</p>
<p>2. The response rate is exceedingly low, which is a major limitation of the study. Moreover, the usable sample of surveys does not meet the required number, so sample size is not adequately justified. This issue is not yet adequately addressed in the text discussion section.</p>	<p>This section now includes a more complete explanation.</p> <p>The low response rate to the national postal questionnaire is a limitation of this study. This can be account for in part in the fact that over five hundred returned questionnaires were from nurses and midwives who were ineligible to participate in the study as they were not employed to work in the public health service, a criterion stipulated by the funding body. Furthermore, the use of telephone or e-mail as a complementary method for enhancing the effectiveness of the Dillman method (Axford 1997) was not possible since the sample was anonymous to the researchers.</p> <p>Comparison of the study response rate with that of other large-scale national postal surveys among nurses and midwives in Ireland indicates that the response rates were very similar to those obtained in the <i>Experiences of Empowerment</i> study (Scott et al. 2003), which and reported a response rate 33.1% from a postal survey of 4,000 nurses and midwives and higher than the response of 20% from a survey of 10,000 nurses obtained in the <i>Report on the Continuing Professional Development of Staff Nurses and Staff Midwives</i> (National Council for the Professional Development of Nursing and Midwifery</p>

	<p>(NCNM) 2004). A high response rate to questionnaires is considered desirable, particularly if the data are to be used for predictive purposes (Meehan <i>et al.</i> 2005).</p> <p>The present study sought data on self-reported need and hence the burden of interpretation with reference to predicting need may be considered lower than that demanded in intervention and interrelationship studies. For the present study the response rate achieved must be considered with reference to sampling frame and the context in which the questionnaire was administered. The sampling frame was the Active Register of the Register of Nurses in Ireland, which does not permit targeted sampling of discrete subgroups such as nurses and midwives or those employed in either public or private sectors. In addition, the survey was administered at a time of uncertainty in Ireland when all public servants experienced large reductions in salaries and this mood may have impacted on data collection.</p>
3. Place the ethical considerations section after the survey instrument section and before the data analysis section.	Done
4. Citations to the literature used for item generation are needed.	Done
5. The survey questionnaire description remains insufficiently clear. What is Scale A and what is Scale B?	Done
6. This sentence is unclear or grammatically incorrect: A scale score being an average of the scores on each item with a higher mean value indicated a higher self-reported development need.	<p>This section now reads</p> <p>Scale A measured the perceived importance of each item in the list of items in relation to effective performance of the clinical leader role in practice setting using a five-point scale from 1 ('not important') to 5 ('very important'). Scale B measured respondents' self-perceived clinical leadership development needs in relation to each item on the same list from 1 ('no need') to 5 ('very high</p>

	need').
7. The description of the factor analysis continues to read as if separate factor analyses were done for each subsection of the questionnaire. A factor analysis that includes all items will yield a certain number of factors from 1 to x. Each item of the questionnaire loads more or less better on a certain factor. Items are retained on the basis of preset criteria for loading and items that load on more than one factor are typically deleted. Thus, it is unclear why you wrote, for example, that a particular subsection of the questionnaire was made up of two components.	Included in this section is the following explanation To further define the traits in each scale, factor analysis (Johnson & Wichern 2002) was conducted on each scale, and where appropriate new subscales defined.
8. It is unclear why the factor analysis results mention the subsections when the data in Table 4 are only for the 5 subscales that also appear in the left column in Table 2.	Table 2 provides details of the results of factor analysis in terms of subscale titles and indicative content. Table 4 indicates the dimensions of clinical leadership in terms of mean, standard deviation and skewness
9. Citations are not consistent with JAN format (should be earliest date of publication first rather than alphabetical order)	Done

Nurses' and Midwives' clinical leadership development needs: A mixed methods study

Aim

This paper is a report of a descriptive study of nurses' and midwives' clinical leadership development needs.

Background

Nurses and midwives are expected to fulfill a leadership role at all levels, yet efforts to strategically support them are often unfocused. An analysis of clinical leadership development needs can provide the foundation for leadership initiatives to support staff.

Method

A mixed methods design was used. A questionnaire was sent to 911 nurses and midwives and 22 focus groups comprising 184 participants were conducted. Data were collected between March and June 2009 across all promotional grades of nurses and midwives in Ireland. Repeated measures ANOVA with Greenhouse-Geisser adjustment was used for post-hoc pair wise comparisons of the subscale dimensions of clinical leadership. ANOVA with Tukey's post-hoc method was used for comparison between grades on each individual subscale. Thematic analysis was undertaken on the focus group data.

Results

Results reveal that needs related to development of the profession were highest for all grades. The staff grade expressed a higher need in relation to 'managing clinical area', 'managing the patient care' and 'skills for clinical leadership' than managers. Qualitative analysis yielded five themes; 'clinical leadership and leaders from a nursing and midwifery perspective'; 'quality service from a nursing and midwifery perspective'; 'clinical leaders' roles and functions'; 'capital and 'competences for clinical leaders and leadership and the context of clinical leadership'.

Conclusion.

Clinical leadership concerns quality, safety and effectiveness. Nurses and midwives are ideally placed to offer the clinical leadership that is required to ensure these patient care outcomes. Development initiatives must address the leader and leadership competencies to support staff.

Keywords: Clinical leaders, leadership development, Needs analysis, Mixed Methods.

SUMMARY STATEMENT

What is already known about this topic

- Leadership is essential for organizational systems and processes to function effectively.
- Clinical leadership in nursing and midwifery is essential for optimizing the environment of care and for improving the effectiveness and outcomes of care.
- Clinical leadership development programmes enhance self confidence, improve care, job satisfaction and enhance leadership skills and capabilities in such areas

1
2
3 as team effectiveness, communications, change management and management of
4
5
6 conflict.

7 8 **What this paper adds**

- 9
- 10 • The staff grade of nurses and midwives have higher clinical leadership
11 development needs than managers in relation to managing the clinical area and
12 patient care.
13
 - 14 • Leadership development need was greater across all grades of nurses and
15 midwives in relation to development of the profession which involves
16 organizational and interdisciplinary interactions with other health care
17 professionals, and influencing clinical decision making and health policy more
18 generally.
19
 - 20 • Clinical leadership in nursing and midwifery is about integrating the clinical
21 leadership role and function into everyday practice to provide a quality service.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

37 **Implications for Practice and/or Policy**

- 38
- 39 • Clinical leadership development must focus on individual and organizational
40 competency development within each individual health care context.
41
 - 42 • Leadership development initiatives are likely to be more successful if the various
43 dimensions of clinical leadership development need and the promotional grade of
44 the individual are taken into account.
45
 - 46 • Clinical leadership development initiatives must take a partnership approach
47 between clinical and academic input for the design and delivery of programmes.
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

From a global perspective, as providers of front-line clinical care, nurses and midwives are ideally placed to offer clinical leadership that is required to ensure quality and optimum patient care. The roles of clinical leaders have been specified (Cook, 2001a and 2001b, Davidson *et al.* 2006), and their impact on care outlined (Dierckx de Casterlé 2008, Hix *et al.* 2009). However, it is the way that these roles and skills are exercised within these roles that determines leadership effectiveness (Gopee & Galloway 2009).

BACKGROUND

Leadership in nursing and midwifery

Leadership in healthcare involves the creation of conditions for success and the achievement of professional and organizational goals (Milward & Bryan 2005). In nursing and midwifery, leadership involves taking responsibility for direct care and for monitoring service more generally (Davidson *et al.* 2006, Carryer *et al.* 2007,). Effective leadership empowers individuals and groups to engage in change (Casey 2006) and improve quality (Alleyne & Jumma 2007, Shaw 2007). Improving nurses' leadership skills enhances the provision of quality healthcare (Borrill *et al.* 2002, Large *et al.* 2005).

The American Association of Colleges of Nursing (AACN) defines a clinical nurse leader as someone who supports and leads innovations that improve outcomes of care, ensures quality and reduces costs, integrates research into practice, is recognised as a leader and an advocate for transforming the health system and implementing best practice (Smith & Dabbs 2007). Clinical leadership enables evidence-based care to improve patient outcomes (Milward & Bryan 2005) by integrating activities and

processes within disciplines and across-service (Tornabeni & Miller 2008). Hence, clinical leadership is essential in complex health systems, in which multiple professional groups, often with complex interdisciplinary social relationships and at times divergent disciplinary goals, work closely together. The challenge for contemporary leadership in healthcare is to maintain a perspective that combines the lofty goals of strategy formulation with the minutiae associated with the day-to-day management of direct care in the clinical context (Casey 2006).

Leadership development takes time (Hartley & Hinksman 2003) and involves sustained interaction between the clinical leader and coworkers (Dierckx de Casterlé *et al.* 2008). Leader development requires human capital, expressed as a range of intrapersonal competences, and social capital which underpins the interpersonal competences required for leadership development as seen in Table 1.

Table 1 Key Aspects of leader and leadership development (Day 2001)

Aspects of leader development (Human Capital)	Aspects of leadership development (Social Capital)
Building intrapersonal competence	Building interpersonal competence
Developing individual knowledge, skills and abilities	Building network relationships through interpersonal exchange
Self awareness (emotional awareness, self confidence)	Social awareness (empathy, service orientation and enabling others)
Self regulation (self control, trustworthiness, ability to cope with change)	Social skills (collaboration, partnership, cooperation, conflict management)
Self motivation (commitment, take initiative, positive outlook)	Membership of influential social networks (Bourdieu 1997, Maton 2005)

Both leader and leadership competency development must be addressed for the integration of the cognitive, affective and behavioural dimensions of the role required to enact leadership. Essentially, leadership development involves building individual and organizational capacity so that members' intrapersonal competences can be effectively

1
2
3 expressed, in the interpersonal domain, in the leadership roles and processes (McCauley
4
5
6 *et al.* 1998).

10 THE STUDY

13 Aim

14
15 The aim of this study was to describe nurses' and midwives' clinical leadership
16
17 development needs.

22 Design

23
24 A concurrent triangulation mixed methods design involving a survey and focus group
25
26 interviews was used.

32 Sample

34 *Survey respondents*

35
36 The sampling frame was 68,000 nurses and midwives (An Bord Altranais 2008) and with
37
38 a precision of $\pm 3\%$ at a 95% confidence interval, it was considered that a sample size of
39
40 1,100 would be required. A simple random sample of 3,000 nurses and midwives
41
42 employed within the Health Service Executive (HSE) in Ireland was selected from the
43
44 Active file of the Register of Nurses. This ensured that all grades and in each of the five
45
46 divisions of the Register, viz. general, children's, psychiatric, intellectual disability, and
47
48 midwife, would be represented. At the closing date for return of questionnaires, a total of
49
50 802 (26.73%) replies were received, 534 (17.8%) of which were returned completed and
51
52 268 (8.9%) were incomplete. Using the Dillman (2000) procedure another circulation of
53
54
55
56
57
58
59
60

1
2
3 questionnaires were sent to the non-respondents which yielded 630 (27.91%) replies, of
4
5 which 377 (16.7%) were completed and 253 (11.2%) incomplete. This procedure
6
7 involves three contacts with the sample: a pre-notification A5 postcard, an initial
8
9 administration of the questionnaire and a second administration to non-respondents of the
10
11 first attempt. Accordingly, a total of 1,432 responses were received, of which 521
12
13 questionnaires were unusable. From the unusable questionnaires, 477 were categorised as
14
15 not employed in the public sector (273) or provided no indication of the sector in which
16
17 employed (194). An additional 29 questionnaires indicated that they were either no
18
19 longer working in nursing or midwifery, were ill or living abroad. A further 25
20
21 questionnaires were returned as undeliverable. This resulted in a valid sample of 2,946
22
23 and a useable sample of 911 questionnaires, representing a response rate of 30.92%.
24
25
26
27
28
29
30
31

32 *Focus group participants*

33
34 A purposive sample of all grades of nurses and midwives from each of the four HSE
35
36 administrative regions was selected to elicit their view on clinical leadership and their
37
38 perceived clinical leadership development needs. Twenty-two focus groups were
39
40 conducted at fourteen different sites which included a mixture of national, regional and
41
42 local tertiary care centres. The mean number of participants per group was 8 and the
43
44 range was 3–12 members. Group homogeneity was achieved by grade and discipline in
45
46 all but one focus group and a total of 184 participants representing all disciplines and
47
48 clinical and promotional grades participated in the focus groups
49
50
51
52
53
54
55
56
57
58
59
60

Data collection

Data collection took place between March and June 2009. Data for the national postal survey were collected using the Clinical Leadership Analysis of Need Questionnaire (CLAN-Q©). Focus group data were collected over the same period. Each focus group was held in private, audio recorded and lasted approximately sixty minutes.

Survey instrument

A 'questionnaire' the Clinical Leadership Analysis of Need Questionnaire '(CLAN-Q©)' was developed to measure the clinical leadership development needs of nurses and midwives. An initial item pool (Bray *et al.* 2003) of over one hundred items that reflected critical tasks associated with effective performance as a clinical leader were generated from the literature (Christian and Norman 1998, Cook 2001a and 2001b, Williams 2004, Large *et al.* 2005, Davidson *et al.* 2006, Dopson and Fitzgerald 2006, Flood 2007, Lunn *et al.* 2008, Sorenson *et al.* 2008, Hix *et al.* 2009) and critically reviewed by the research team acting as content experts. The items were presented in two Likert scales. Scale A measured the perceived importance of each item in the list of items in relation to effective performance of the clinical leader role in practice setting using a five-point scale from 1 ('not important') to 5 ('very important'). Scale B measured respondents' self-perceived clinical leadership development needs in relation to each item on the same list from 1 ('no need') to 5 ('very high need').

A small pilot study was conducted among a convenience sample of one junior and three senior grades of nurses and midwives in different settings and this provided confirmation of the feasibility of the questionnaire (Hallberg 2008). Following pilot

1
2
3 testing it was observed that responses to Scale A in all categories elicited responses that
4 were in the higher end of perceived importance (i.e. important and very important). This
5 indicated a threat to the discriminating power of the items with respect to Scale A.
6
7
8 However, given the absence of any extant list of leadership behaviours and capabilities
9 with which to construct a needs analysis scale, the research team considered that it was
10 necessary to retain Scale A as a self-report mechanism of perceived importance as this
11 provided internal validation of each item with respect to Scale B. This consideration
12 outweighed that limitation, and the fact that 80% of the respondents ranked these items as
13 'very important' lends support for claims of construct validity of the instrument items.
14
15 The CLAN-Q incorporated a demographic questionnaire and took approximately twenty
16 minutes to complete.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 *Validity and reliability of survey instrument*

33
34 Instrument content validation involved domain identification, item generation, and
35 instrument construction (Grant & Davis 1997). Each item was independently examined
36 by four experts from the research team and items were excluded or retained on
37 consensus. Finally, a list of 68 items was retained and compiled into three main category
38 scales based on specific concepts based on their fit with emergent constructs, namely
39 'improving the environment for care delivery' (31 items), 'personal and professional
40 development' (14 items) and 'skills for clinical leadership' (23 items). The multiple
41 iterations of the instrument items and the use of cognitive interviewing principles
42 (Drennan 2003) added rigor to the process of content validation.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

To further define the traits in each scale, factor analysis (Johnson & Wichern 2002) was conducted on each scale, and where appropriate new sub-scales defined. It was determined that the scale 'skills for clinical leadership' consisted a single component. The scale relating to 'improving the environment for care delivery' (31 items) was determined to have two components based on Eigen value greater than 1 from a principal component extraction. Varimax rotation was used to identify which component each item contributed to based on a loading factor greater than 0.5. The two components or subscales were identified as 'managing clinical area' (6 items) and 'managing patient care' (25 items).

A similar analysis of scale 'personal and professional development' (contained 14 items) also identified two components or subscales; 'development of individual' (9 items) and 'development of the profession' (5 items). Based on scale reliability calculations it was noted that the third subscale relating to 'skills for clinical leadership' improved if one item was deleted. The item 'influencing key stakeholders within the profession' was omitted from the survey, leaving 22 items for this third subscale. The results of factor analysis and the indicative content of the final five subscales are listed in Table 2.

Table 2 Results of factor analysis, subscale titles and indicative content of CLAN-Q instrument

Subscales, factor analysis, number of items of category scale 'improving the environment for care delivery'	
Subscale Title	Number of items and indicative content of subscale
Managing clinical area	Contained 6 items related to coordinating care in the work setting, ensuring adequate resources to provide optimum care, identifying priorities for service improvement, and ensuring that team members carry out duties appropriate to their grade.
Eigenvalue 20.53	
% of Variance 66.24	
Cumulative % 66.24	
Managing patient care	Contained 25 items related to protecting dignity and confidentiality of patients, involving patients in their own care, contributing to the
Eigenvalue 1.34	

1
2
3 % of Variance 4.36 development of clinical practice guidelines, and ensuring the outcomes of
4 Cumulative % 70.60 care and other interventions are documented
5
6

7 **Subscales, factor analysis, number of items of category scale ‘personal and professional development**

8 Development of individual Contained 9 items related to demonstrating commitment to lifelong learning,
9 Eigenvalue 8.53 recognising own strengths and weaknesses, coping effectively with pressure,
10 % of Variance 60.93 and acting as a mentor to colleagues
11 Cumulative % 60.93
12

13
14 Development of profession Contained 5 items related to participating in nursing/midwifery forums,
15 Eigenvalue 1.63 networking and understanding the impact of organisational politics on the
16 % of Variance 11.61 work of the profession.
17 Cumulative % 72.54
18
19

20 **Subscales, factor analysis, number of items of category scale ‘skills for clinical leadership’**

21 Skills for clinical leadership Contained 22 items related to effective working relationships with the
22 interdisciplinary team, motivating others, accepting accountability, and
23 initiating change to ensure optimal care
24
25

26
27
28 The reliability of each subscale was examined by calculating a Cronbach’s Alpha
29 coefficient. An acceptable Alpha level is 0.7 or above, and the properties of the five
30 scales were: ‘managing clinical area’ contained 6 items, $\alpha=0.91$, ‘managing patient care’
31 contained 25 items, $\alpha=0.98$; ‘development of individual’ contained 9 items, $\alpha=0.96$;
32
33 ‘development of profession’ contained 5 items $\alpha=0.91$ and ‘skills for clinical leadership’
34 contained 22 items, $\alpha=0.98$.
35
36
37
38
39
40
41
42
43
44

45 **Ethical considerations**

46
47 Permission for the study was received from the university. Return of the CLAN-Q was
48 accepted as consent to participate in the survey and each participant in the focus group
49 gave informed written consent on the basis of confidentiality.
50
51
52
53
54
55
56
57
58
59
60

Data analysis

Data analysis were undertaken using the Statistical Package for the Social Sciences© (SPSS Version 15.0) software. Descriptive variables related to respondents' demographic characteristics were summarised using frequency distributions to describe the sample. As the distribution of scores are centred near the middle of the range, and are essentially symmetric (skew near 0), the scores were assumed to be normally distributed. Based on repeated measures ANOVA with Greenhouse-Geisser adjustment for non spherical data to correct degrees of freedom, post hoc comparisons of subscales were performed using the Bonferroni method for controlling Type 1 error to 0.05. ANOVA with Tukey's post-hoc method was used for comparison between grades on each individual scale.

Dependability and credibility of focus groups

The focus group questions were prepared in advance and structured using the same categories that were developed for the national survey instrument, viz. 'the care environment', 'personal and professional development needs', 'professional skills for leadership'. Dependability was enhanced by the use of a topic guide (Holloway and Wheeler 2002), based on the three categories as the questionnaire and by judicious group moderation and by analysis of the data by two members of the research team which achieved complete agreement on the interpretation and analysis of themes. Data analysis involved data reduction, exploration and synthesis (Attride-Stirling 2001). Credibility of the analysis was enhanced by adherence to systematic analytical process which involved reading each transcript closely and breaking down significant segments into themes which were allocated to free nodes using NVivo 7©. As significant relationships between

1
2
3 the free nodes were identified and further explored in the existing and incoming data, a
4
5 tree node structure was devised. This process allowed relationships between themes to be
6
7 explored, revealing underlying patterns and relationships in the data, thus assisting
8
9 dependability. Dependability was also enhanced using a coding framework based largely
10
11 on the work of Day (2001). This framework was modified to refine the tree node
12
13 structure until the maximum amount of data was accounted for, with minimal overlap and
14
15 redundancy in order to identify and describe the clinical leadership development needs of
16
17 nurses and midwives.
18
19
20
21
22
23

24 RESULTS

25
26
27 The justification for combining quantitative and qualitative methods was to provide a
28
29 more complete account of clinical leadership development needs and a more detailed
30
31 explanation of these perceived needs (Bryman 2006). To this end, the results of the
32
33 survey and focus group are presented separately and ‘the mix’ is in the interpretative
34
35 treatment of the results manifest in the unifying discussion (Plano-Clarke and Creswell
36
37 2008).
38
39
40
41
42

43 Survey sample characteristics

44
45
46 The majority of the sample was female 92% (n=836) and all promotional grades were
47
48 represented. The mean age of the sample was 42 years (SD=10.14, range 22–70). Two-
49
50 thirds (66.3%, n=584) worked in a public or public voluntary hospital setting, 21%
51
52 (n=180) worked in the community or in nursing homes and other non acute care sector.
53
54
55 The majority of respondents (61.5% (n=547) were employed as general nurses and
56
57
58
59
60

midwives accounted for just over 5% (n=46). Details of the grades of each participant are provided in Table 3.

Table 3. Survey results in relation to grade at which currently employed

Grade	Percentage (number)
Staff Nurse	66.6 (n=599)
Clinical Nurse Manager level 1 (CNM1)	6.3 (n=57)
Clinical Nurse Manager level 2 (CNM2)	10.6 (n=95)
Clinical Nurse Manager level 3 (CNM3)	1.1 (n=10)
Clinical Nurse Specialist	5.1 (n=46)
Advanced Nurse Practitioner	0.4 (n=4)
Assistant Director of Nursing	3.1 (n=28)
Director of Nursing	0.8 (n=7)
Staff Midwife	3.9 (n=35)
Clinical Midwife Manager level 1 (CMM1)	0.2 (n=2)
Clinical Midwife Manager level 2 (CMM2)	0.8 (n=7)
Clinical Midwife Manager level 3 (CMM3)	0.4 (n=4)
Clinical Midwife Specialist	0.2 (n=2)
Advanced Midwife Practitioner	0.1 (n=1)
Assistant Director of Midwifery	0.2 (n=2)
Director of Midwifery	0 (n=0)
Missing	1.3 (n=12)
Total	100 (n=911)

The sample was representative when compared with the national Register of Nurses statistics of just under 92,000 qualifications. In the study sample of registrations of 1,306, general nursing was underrepresented 58% (n=755) compared to 63% (n=57,474) in the national Register. Psychiatric Nursing was (9% n=112) approximately 2.1% less representative than the national Register (11% n=9796) and there was slight over representation of midwifery of 17% (n=222) compared to 14% (n=12,988) and, intellectual disability accounted for 7% (n=86) compared to 5% (n=4,233) and public health over representative at 4% (n=47) than the national Register of 3% (n= 2,378).

Dimensions of clinical leadership development

Based on repeated measures ANOVA with Greenhouse-Geisser adjustment for non spherical data, at least one scale was statistically significant from others, $F(3, 2068.7) = 32.2, p < 0.001$. The subscale 'development of profession' ($M=3.22, SD=0.99$) was significantly higher than 'managing clinical area' ($M=3.07, SD=1.00, p < 0.001$), 'skills for clinical leadership' ($M=3.07, SD=1.0, p < 0.001$), 'development of individual' ($M=3.04, SD=1.00, p < 0.001$) and 'managing patient care' ($M=2.92, SD=1.1, p < 0.001$).

Additionally, the subscale 'managing patient care' ($M=2.92, SD=1.1$) was significantly lower than 'managing clinical area' ($M=3.07, SD=1.00, p < 0.001$), 'skills for clinical leadership' ($M=3.07, SD=1.0, p < 0.001$) and 'development of individual' ($M=3.04, SD=1.00, p < 0.001$). The five subscales of dimensions of clinical leadership development need are presented in Table 4.

Table 4 Subscales of clinical leadership development need

Subscale of Clinical Leadership Development Need	Mean	SD	Skewness	n	Missing values*
Managing the clinical area	3.07	1.0	0.19	880	31
Managing patient care	2.92	1.1	0.31	834	77
Development of individual	3.04	1.0	0.24	863	48
Development of profession	3.22	0.99	0.05	872	39
Skills for clinical leadership	3.07	1.0	0.15	834	77

*Non responses

Clinical leadership development and promotional grades

For the purpose of analysing differences by grade, nurses and midwives were grouped together and recoded into four categories; 1) staff (incorporating public health nurse), 2) clinical manager, 3) specialist and advanced practitioner and 4) senior manager and director of nursing/midwifery.

1
2
3 A significant difference among promotional grades was observed in the subscale
4
5 'managing the clinical area' ($F(3,865) = 2.906, p = 0.034$). Pair wise post hoc tests did not
6
7 identify any significant differences. A pooled analysis combining staff and specialists
8
9 views, based on their greater engagement and proximity to clinical care, with those of
10
11 middle and senior management levels revealed a significant difference between frontline
12
13 workers ($M = 3.12, SD = 1.00$) and those more distant from the bedside ($M = 2.89, SD = 0.97,$
14
15 $p = 0.003$), where frontline staff expressed a higher leadership development need.
16
17
18

19
20 A significant difference between grades in the subscale 'managing patient care'
21
22 was also observed ($F(3,819) = 3.41, p = 0.017$). Pair wise comparison indicated a
23
24 significant difference between staff ($M = 2.99, SD = 1.14$) and manager grades ($M = 2.67,$
25
26 $SD = 1.04, p = 0.010$), with staff reporting higher clinical leadership development need.
27
28

29
30 No significant differences between grades for the subscale 'development of the
31
32 individual' ($F(3,847) = 2.43, p = 0.064$) was observed. While the scale comparisons
33
34 indicated that the highest scale across all grades was in relation to 'development of the
35
36 profession', there were no significant differences between grades
37
38 ($F(3,856) = 0.36, p = 0.781$).
39

40
41 There is at least one difference between grades for the 'kills for clinical
42
43 leadership' ($F(3,819) = 3.232, p = 0.022$). Post-hoc comparisons identified that staff grade
44
45 ($M = 3.14, SD = 1.031$) differed from manager grade ($M = 2.85, SD = .94, p = 0.011$), with staff
46
47 expressing a higher clinical leadership development need.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Results: Focus groups
4

5
6 Just over half the participants in the twenty-two focus groups were employed in general
7
8 nursing (52%, n=93), a quarter (24%, n= 44) in midwifery and the remainder in either
9
10 children's nursing (7.7%, n=14), psychiatric nursing (6.6%, n=12) or intellectual
11
12 disability nursing (4.9%, n=9). The majority (92%, n=169) were female and the average
13
14 age was 40 years with a range age of 23-60 years.
15
16

17
18
19
20 *Clinical leadership and views on clinical leaders*
21

22 Focus group participants found it difficult to agree on a definition of clinical leadership,
23
24 but agreed that it is founded on clinical expertise, experience and credibility. Being a
25
26 clinical leader was seen as more challenging for clinical nurse managers and midwifery
27
28 managers (CNM/CMM) due to their position in the organization (middle management)
29
30 their multiple and conflicting responsibilities and poor role definition:
31
32

33
34 We describe ourselves as the meat in the sandwich...we are like the
35 filling. Stuck in the middle...we are the ones that are being pushed
36 from the top down.... (CNMG1)
37
38

39
40 Clinical leaders were seen as the 'guardians' of patient care and expected to advance
41
42 nursing and midwifery practice by advocating for patients' interests and by maintaining
43
44 standards:
45

46
47 I think at the core of it is patient advocacy, patient-centred
48 care...emphasis is on quality and safety ... maintenance of standards is
49 at the core of what we're at. (DoN&M)
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *Quality nursing and midwifery services*
4

5 Participants portrayed a very strong commitment to their role as clinical leaders, patient
6 advocates and protectors of patient dignity and privacy and were uncompromising about
7
8 the difficulties they encounter in so doing:
9
10

11
12 ...you are striving for best practice but the way things are with the
13 patient numbers and the lack of staff it's very hard... to give them a
14 small bit of privacy to see a doctor never mind anything else. (SNG
15 G2)
16
17

18
19
20 *Roles and functions of clinical leaders*
21

22 Nurses and midwives emphasised their coordinating role as integrating diverse
23 disciplinary inputs in the interests of patient care; indeed, they regarded themselves as the
24 main group to whom this responsibility fell:
25
26
27

28
29 ...the whole thing is multi-disciplinary...call in this discipline or that
30 discipline but as nurses we're the ones who seem to pull it all together.
31 (CNM G4)
32
33

34
35 Nurses and midwives were seen as ideally placed to provide clinical leadership but this
36 role potential was frequently underutilized and their contributions unrecognized:
37
38

39
40 nurses are uniquely positioned...the difficulty is the level of
41 understanding within other groups may not value that role or may
42 not understand that role ...we've been struggling for so many years
43 as a profession, articulating what it is that we bring to the table.
44 (DoN & M)
45
46

47
48 All grades were able to provide examples of specific initiatives that had made a
49 difference to patient care. Directors, specialists and some managers emphasised the
50 importance of challenging existing practices and initiating change at organizational level.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1
2
3 patient care. Where these result from the actions or omissions of nursing or midwifery
4
5 colleagues, participants indicated that they would try to deal with concerns at the lowest
6
7 appropriate level and refer upwards according to their professional judgment and
8
9 established policies:
10
11

12 the best way is to talk to the person privately...in a safe, professional
13 way...I think being honest and professional is the right way to
14 approach a colleague ...it depends how serious it was...you try and
15 solve it as local as possible. (SNGG4)
16
17
18
19

20 Dealing with concerns about medical colleagues was considered much more of a
21
22 challenge:
23

24 the majority of your battles when it comes to patients, and they are
25 right royal battles sometimes, is about inappropriate
26 discharging...there is huge social implications. (CNMG1)
27
28
29

30 *resources and competencies for clinical leaders and leadership*

31
32 A lack of personal and professional autonomy was a recurring theme in the data:
33

34 'we' can literally do absolutely nothing because 'we' don't have power
35 ...to do anything....there is a lack of support there...there's no
36 support. (CMMG1)
37
38
39

40
41 Participants indicated a leadership development need at the interpersonal level associated
42
43 with intrapersonal needs expressed through feelings of powerlessness and poor self and
44
45 professional image:
46
47

48 we're looked down on ... if you're down in the pecking order you end
49 up hitting a brick wall...they don't listen to us. (SNID)
50
51
52

53 There were numerous examples of nurses' and midwives' motivation, personal initiative
54
55 and commitment to initiating change and while there was consensus that managing
56
57
58
59
60

1
2
3 change was a vital ingredient of clinical leadership, participants believed that their
4
5 initiative was not effectively harnessed. This again suggested a leadership need at the
6
7 wider organizational level:
8
9

10 one of the big problems in changing policy ...they don't have the
11 time...ward meetings are going out the window...there's nowhere to
12 come together and say, how do we feel things are going, how can we
13 improve? (SMG1)
14
15

16 17 *The context of clinical leadership* 18

19 Whether leaders' human capital, as manifest in their intrapersonal competences,
20 translates into social capital and effective leadership depends on the context in which
21 they operate. Overall, motivation, recognition of contribution, teamwork, mutual respect,
22 organizational support and inclusiveness were identified as the key ingredients of a
23 supportive work context:
24
25
26
27
28
29
30

31 from the time a staff nurse comes to work in a unit... there's a
32 development plan put in place...the hospital is very proactive in
33 allowing staff to participate in these programmes. (CNM G5)
34
35
36

37 DISCUSSION

38 Study limitations

39
40 Leadership is culturally influenced and context-dependent, hence the outcomes of this
41 study are most relevant to how nursing and midwifery is delivered in Ireland, and
42 regulatory frameworks and clinical environments in other countries may elicit different
43 leadership development needs.
44
45
46
47
48
49
50

51 The low response rate to the national postal questionnaire is a limitation of this
52 study. This can be account for in part in the fact that over five hundred returned
53 questionnaires were from nurses and midwives who were ineligible to participate in the
54
55
56
57
58
59
60

1
2
3 study as they were not employed to work in the public health service, a criterion
4 stipulated by the funding body. Furthermore, the use of telephone or e-mail as a
5
6 complementary method for enhancing the effectiveness of the Dillman method (Axford
7
8 1997) was not possible since the sample was anonymous to the researchers. Comparison
9
10 of the study response rate with that of other large-scale national postal surveys among
11
12 nurses and midwives in Ireland indicates that the response rates were very similar to
13
14 those obtained in the *Experiences of Empowerment* study (Scott *et al.* 2003), which and
15
16 reported a response rate 33.1% from a postal survey of 4,000 nurses and midwives and
17
18 higher than the response of 20% from a survey of 10,000 nurses obtained in the *Report*
19
20 *on the Continuing Professional Development of Staff Nurses and Staff Midwives*
21
22 (National Council for the Professional Development of Nursing and Midwifery (NCNM)
23
24 2004). A high response rate to questionnaires is considered desirable, particularly if the
25
26 data are to be used for predictive purposes (Meehan *et al.* 2005).
27
28
29
30
31
32
33

34 The present study sought data on self-reported need and hence the burden of
35
36 interpretation with reference to predicting need may be considered lower than that
37
38 demanded in intervention and interrelationship studies. For the present study the response
39
40 rate achieved must be considered with reference to sampling frame and the context in
41
42 which the questionnaire was administered. The sampling frame was the Active Register
43
44 of the Register of Nurses in Ireland, which does not permit targeted sampling of discrete
45
46 subgroups such as nurses and midwives or those employed in either public or private
47
48 sectors. In addition, the survey was administered at a time of uncertainty in Ireland when
49
50 all public servants experienced large reductions in salaries and this mood may have
51
52 impacted on data collection.
53
54
55
56
57
58
59
60

Defining clinical leadership

There was consensus that clinical leadership is directed at ensuring quality patient services, and nurses and midwives closest to the patient consider themselves well-positioned to make a difference to patient and quality care. The staff grade expressed higher development needs in relation to 'managing the clinical area', 'managing patient care' and 'skills for clinical leadership' which involved activities such as coordinating care, developing effective working relationships with the interdisciplinary team, managing conflict, implementing change, coordinating care and ensuring adequate resources for care. The principal mechanisms through which nurses and midwives impacted on patient care were patient advocacy and safety, maintaining dignity and privacy through the co-ordination of the care. These mechanisms are well supported in the literature where a clinical leader is an expert clinician involved in providing direct clinical care and as a clinician can influence others to achieve optimal care (Cook 2001a, 2001b). While the management of relationships between patients, multidisciplinary team members and the wider organization is widely acknowledged to be a key function of clinical leaders (Milward & Bryan 2005), the ability to articulate concise definitions of clinical leadership and of needs related to its development at the individual, group and wider organizational level varied across the focus groups. It is not surprising therefore that although the subscale 'development of the profession', which included items as networking and organizational politics, did not indicate any significant differences between the grades, it was the highest expressed leadership development need.

1
2
3 Describing and distinguishing need
4

5 The staff grade expressed higher needs in relation to ‘managing the clinical area’,
6
7
8 ‘managing patient care’ and ‘skills for clinical leadership’ and than any manager grade.
9
10 Focus group participants spoke of their commitment to motivating other s to improve
11
12 care and while the acknowledged their responsibility and accountability in providing
13
14 quality care, they also emphasized their lack of self confidence and autonomy and lack of
15
16 control over local resources. Clinicians can exert influence in directing patient care by
17
18 exemplary leadership and by excellence in their practice domain (Davidson *et al.* 2006),
19
20 but their scope of leadership is influenced by their position, status and the everyday
21
22 challenges of limited resources. In contrast, managers are somewhat removed from the
23
24 patient interface but perceive their influence as being greater.
25
26
27
28

29 The subscale ‘development of the profession’ concerning leadership at team,
30
31 interdepartmental and organizational levels of leadership was the highest reported need
32
33 among all grades. Focus group data also confirmed that nurses and midwives struggle to
34
35 get their voices heard at the policy table and feel increasingly excluded within their
36
37 organizations, which is consistent with the evidence from other Irish studies (Tracey
38
39 2003, Hogan 2006, O’Shea 2008). Clinical leadership involves not only taking
40
41 responsibility for direct care and its outcomes but also for monitoring and developing
42
43 services in general (Davidson *et al.* 2006, Carryer *et al.* 2007). As the focus shifts away
44
45 from the bedside and the micro-system in which the nurse or midwife is central (Hix *et*
46
47 *al.* 2009) and outwards to the multidisciplinary team and wider organisation, the focus
48
49 group data suggested that nurses and midwives are much less assured of their roles.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Further examination of these needs is warranted to determine what role the organisation
4 and individual can contribute to meeting this leadership development need deficit.
5
6

7
8 Whether and to what extent nurses and midwives gain recognition and exert
9 influence at the individual, team, department and organizational levels will ultimately
10 impact on the quality of patient services. The need for nurses and midwives to be
11 involved at strategic and operational levels of decision-making is essential to the
12 development of more accountable and integrated health services (O'Shea 2008). That
13 nurses and midwives in management positions should be so constrained in acting as
14 leaders removes a source of role models and mentors for junior colleagues and an
15 incentive to assume such roles. Hence, a distinct leadership development need for nurses
16 and midwives is the ability to better represent their unique contributions to patient
17 outcomes in a credible manner.
18
19
20
21
22
23
24
25
26
27
28
29
30

31
32 Clinical leaders in nursing and midwifery advocate for patients and for their
33 professions (NCNM 2005) and shared decision-making among doctors, nurses and other
34 health professionals is an important condition for such wider advocacy (Office for Health
35 Management 2003). This need is linked to their ability to participate fully in
36 multidisciplinary teams, work in partnership and to drive change. When this need is met
37 it gives rise to effective leadership in the way that the leader is aware of the political
38 nuances that impact on clinical practice and can effectively manage conflict and
39 departmental turfism (Casey 2008). This requires better role differentiation for
40 meaningful integration and highlights the importance of developing both individual
41 leaders and collective leadership (Glatter 2008) thereby building individual and
42 organizational capacity. These competences require capacity-building through
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 interpersonal competences such as communication and decision making, advocacy,
4
5 empowerment, quality, partnership, and political awareness. This finding supports
6
7 leadership development approaches that address the intrapersonal, such as self awareness
8
9 and the interpersonal domains of competence associated with the individual, team,
10
11 departmental and organizational levels (McAuliffe *et al.* 2002, Large *et al.* 2005,
12
13 Sorensen *et al.* 2008, Stanley 2008, Tornabeni & Miller 2008)
14
15
16
17
18
19

20 Clinical leadership development

21
22 Needs in relation to the dimensions of clinical leadership development were successfully
23
24 identified using the ©CLAN-Q. These needs are related to managing clinical area,
25
26 managing patient care, development of the individual, development of the profession and
27
28 skills for clinical leadership. On the evidence in this study *leader* development, in terms
29
30 of self awareness, and *leadership* development (collective organizational context) must
31
32 be considered as part of the spectrum of clinical leadership development. Moreover, there
33
34 is a difference in the degree of need depending on the promotional grades as highlighted
35
36 between staff and manager grades. Any strategy for clinical leadership development
37
38 therefore must take cognisance of the promotional grade and particular context in which
39
40 nurses and midwives operate as well as the overall development needs of nursing and
41
42 midwifery as professional disciplines (Williams 2004, Large *et al.* 2005, Tornabeni &
43
44 Miller 2008, Proctor-Thompson 2008, Sorensen *et al.* 2008, Stanley 2008).
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

CONCLUSION

Leader and leadership competency development must be key outcomes of clinical leadership development initiatives for nursing and midwifery. This requires a focus on the individual, the professions and the organizations in which they operate. Undertaking clinical leadership needs assessment and developing mechanisms to address leadership deficits at individual, team, department and organisational levels are necessary. Clinical leadership development is a shared responsibility and those in leadership positions must work together across professional boundaries to secure the recognition and influence that they need so that clinical leadership capacity can be harnessed, developed and deployed in the interests of patient care.

References

An Bord Altranais (2008) *Register Statistics 2008* An Bord Altranais. Dublin, Available online at: <http://www.nursingboard.ie/en/homepage.aspx> (Retrieved 30 June 2009).

Alleyne J. & Jumaa M.O. (2007) Building the capacity for evidenced-based clinical nursing leadership: the role of executive co-coaching and group clinical supervision for quality patient services. *Journal of Nursing Management* **15**, 230-243.

Attride-Stirling J. (2001) Thematic networks: an analytic tool for qualitative research. *Qualitative Research* **1**(3), 385–405.

1
2
3 Axford R. (1997) Enhancing Dillman's total design method for mailed/ telephone
4 surveys using current technology to maximise cost-benefit ratios *Journal of Sociology*
5
6
7
8 **33**(3), 387–393.
9

10
11
12 Bourdieu P. (1997) The forms of capital. In *Education, Culture, Economy, Society*
13 (Halsey A.H.L., Brown P. & Wells A.S., eds), Oxford University Press, Oxford.
14
15
16
17

18
19
20 Borrill C., West M., Dawson J., Shapiro D., Rees A., Richards A., Garrod S., Carletta J.
21 & Carter A. (2002) *Team Effectiveness in Health Care: Findings from the Health Care*
22 *Team Effectiveness Project*. Report from Aston Centre for Health Service Organization
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Bray C.O., Nash K. & Fromanz R.D. (2003) Validation of measures of middle schoolers' self efficacy for physical and emotional health, and academic tasks. *Research in Nursing and Health* **26**, 376–386.

Bryman A. (2006) Integrating quantitative and qualitative research: How is it done? *Qualitative Research* **6**(1), 97-113.

Carryer J., Gardner G., Dunn S. & Gardner A. (2007) The core role of the nurse practitioner: practice, professionalism and clinical leadership. *Journal of Clinical Nursing* **16**, 1818–1825.

1
2
3 Casey M. (2006) *Developing a Framework for Partnership between Organizations that*
4 *provide Nursing and Midwifery Education*. Unpublished PhD Thesis, School of Business,
5
6 University of Dublin, Trinity College, Dublin.
7
8
9

10
11
12 Casey M. (2008) Partnership: success factors of international relationships. *Journal of*
13 *Nursing Management* **16**, 72 – 83.
14
15
16
17

18
19
20 Christian S. & Norman I. (1998) Clinical Leadership in nursing development units.
21 *Journal of Advanced Nursing* 27, 108-116.
22
23
24

25
26
27 Cook M. (2001a) The renaissance of clinical leadership. *International Nursing Review*
28 48, 38–46.
29
30
31

32
33
34 Cook M. (2001b) The attributes of effective clinical nurse leaders. *Nursing Standard* 15,
35 33–6.
36
37
38

39
40
41 Davidson P., Elliot D. & Daly J. (2006) Clinical Leadership in contemporary clinical
42 practice: implications for nursing in Australia. *Journal of Nursing Management* **14**, 180–
43 187.
44
45
46
47

48
49
50 Day D.V. (2001) Leadership development: A review in context. *Leadership Quarterly*
51 **11**(4), 581–613.
52
53
54
55

1
2
3 Dierckx de Casterlé B., Willemse A., Verschueren M. & Milisen K. (2008) Impact of
4 clinical leadership development on the clinical leader, nursing team and care-giving
5 process: A case study. *Journal of Nursing Management* **16**, 753–763.
6
7

8
9
10
11
12 Dillman D. (2000) *Mail and Internet Surveys: The Tailored Design Approach* (2nd ed.),
13 John Wiley & Sons, New York.
14
15

16
17
18
19
20 Dopson S. and Fitzgerald L. (2006) The role of the middle manager in the
21 implementation of evidenced-based health care. *Journal of Nursing Management* **14**, 43-
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

30 Drennan J. (2003) Cognitive interviewing: verbal data in the development and pre-testing
31 of questionnaires. *Journal of Advanced Nursing* **42**(1), 57-63.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

36 Flood A. (2007) *The Clinical Nurse Manager (2): Unlocking the potential of this Key
37 Role in the Transformation of Irish Health Care*. Unpublished PhD Thesis, University of
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

46 Glatter R. (2008) *Of leadership, management and wisdom: A brief synthesis of selected
47 reports and documents on leadership development* Prepared for the National College for
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Gopee N. & Galloway J. (2009) *Leadership and Management in Healthcare* Sage
4
5
6 Publications Ltd., London.
7

8
9
10 Grant J. & Davis L. (1997) Selection and use of content experts for instrument
11
12 development. *Research in Nursing & Health* **20**, 269–274.
13
14

15
16
17 Hallberg I.R. (2008) Surveys, In (Watson, R. McKenna, H. Cowman, S. and Keady J.
18
19 eds). *Nursing Research: Design and Methods* Edinburgh: Elsevier, 180–197.
20
21

22
23
24 Hartley J. & Hinksman B. (2003), *Leadership Development: A Systematic Review of the*
25
26 *Literature*, NHS Leadership Centre, London.
27
28

29
30
31 Hix C., McKeon L. & Walters S. (2009) Clinical nurse leader impact on microsystems
32
33 outcomes. *The Journal of Nursing Administration* **39**(2), 71–76.
34
35

36
37
38 Hogan J. (2006) *A Grounded Theory of the Role of the Directors of Nursing in Band One*
39
40 *Teaching Hospitals in Ireland*. MSc. (Research), Dublin City University.
41
42

43
44
45 Holloway I. and Wheeler S. (2002) *Research in Nursing* 2nd edn Blackwell Publishing,
46
47 Oxford.
48
49

50
51
52 Hyrkäs K. & Dende D. (2008) Clinical nursing leadership – perspectives on current
53
54 topics. *Journal of Nursing Management* **16**, 495-498.
55
56

1
2
3 Large S., Macleod A., Cunningham G. & Kitson A. (2005) *A Multiple-Case Study*
4
5
6 *Evaluation of the RCN Clinical Leadership Programme in England*. RCN Institute,
7
8 London.

9
10
11
12 Lunn C. MacCurtain S. & McMahon J. (2008) *Clinical Leadership Pilot Evaluation*
13
14
15 *Report*. Nursing and Midwifery Planning and Development Unit. Health Services
16
17 Executive, Dublin.

18
19
20
21 Johnson R.A. & Wichern D.W. (2002). *Applied Multivariate Statistical Analyses* (5th
22
23
24 Edition). Upper Saddle River, Prentice Hall, New Jersey.

25
26
27
28 Jumaa M. & Marrow C. (2008) Nursing leadership development: why should any nurse
29
30
31 be led by you? *Journal of Nursing Management* **16**, 893–897.

32
33
34
35
36 Large S., Macleod A., Cunningham G. & Kitson A. (2005) *A Multiple-Case Study*
37
38
39 *Evaluation of the RCN Clinical Leadership Programme in England*. RCN Institute,
40
41 London.

42
43
44
45 Lunn C., MacCurtain S. & McMahon J. (2008) *Clinical Leadership Pilot Evaluation*
46
47
48 *Report* Nursing and midwifery Planning and Development Unit Health Services
49
50 Executive, Dublin.

1
2
3 Maton K. (2005) A question of autonomy: Bourdieu's field approach and higher
4 education policy. *Journal of Education Policy* **20**(6), 687-704.
5
6
7
8

9
10 Meehan TC, Kemple M, Butler M, Drennan J, Treacy M, Johnson M (2005) *Nursing and*
11 *Midwifery Research Priorities for Ireland* NCNM, Dublin.
12
13
14

15
16
17 McAuliffe E., Coghlan D. & Pathe A. (2002) Organization development and change
18 management in the Irish health boards: Current policies and practices. *Administration*
19 **50**(3), 42-63.
20
21
22
23
24

25
26
27 McCauley C.D., Moxley R.S. & Van Velsor E. (1998) *The Center for Creative*
28 *Leadership Handbook of Leadership Development* Jossey-Bass, San Francisco.
29
30
31

32
33
34 McCauley C.D. (2000) A systemic approach to leadership development. Paper presented
35 at the 15th Annual Conference of the Society for Industrial and Organizational
36 Psychology in D. Day. (2000) Leadership Development: A Review of Context.
37
38
39 *Leadership Quarterly* **11**(4) 581-613.
40
41
42
43
44

45
46 Milward L. & Bryan K. (2005) Clinical leadership in health care: a position statement.
47
48 *Leadership in Health Services* **18**, xiii-4xxv.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 National Council for the Professional Development of Nursing & Midwifery (2004)
4
5 *Report on the Continuing Professional Development of Staff Nurses and Staff Midwives*
6
7
8 NCNM, Dublin.
9

10
11
12 National Council for the Professional Development of Nursing and Midwifery (2005) A
13
14 *Preliminary Evaluation of the Role of the Advanced Nurse Practitioner* NCNM, Dublin.
15
16
17

18
19
20 National Health Service Scotland (2004) *Leadership Development Framework for*
21
22 *Discussion* NHS Scotland.
23
24

25
26
27 Office of Health Management (2003) *Clinicians in Management: A Review of Clinical*
28
29 *Leadership – Discussion Paper 4* Office of Health Management, Dublin.
30
31

32
33
34 O'Shea Y. (2008) *Nursing & Midwifery in Ireland: A Strategy for Professional*
35
36 *Development in a Changing health Service*, Blackhall Publishing, Dublin.
37
38

39
40
41 Plano-Clark V. and Creswell J. (2008) *The Mixed Methods Reader* Sage Publications Ltd,
42
43
44 London.
45

46
47
48 Proctor-Thomson S.B. (2008) *Constellations or stars? What is being developed in*
49
50 *leadership development?* Lancaster University Management School, Centre for
51
52
53 Excellence in Leadership (CEL).
54
55

1
2
3 Scott PA, Matthews A, Corbally M (2003) *Nurses' and Midwives' Understanding and*
4
5
6 *Experiences of Empowerment in Ireland*, Department of Health and Children, Dublin.
7
8
9

10 Shaw S. (2007) *Nursing Leadership* Blackwell Publishing, Oxford.
11

12 Smith D. & Dabbs M. (2007) Transforming the care delivery model in preparation for the
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

clinical nurse leader. *Journal of Nursing Administration* **37**(4), 157–160.

Sorensen R., Iedema R. & Severinsson E. (2008) Beyond profession: nursing leadership
in contemporary healthcare. *Journal of Nursing Management* **16**, 535–544.

Stanley D. (2008) Congruent leadership: values in action. *Journal of Nursing
Management* **16**, 519–524.

Tornabeni J. & Miller J.F. (2008) The power of partnership to shape the future of
nursing: the evolution of the clinical nurse leader. *Journal of Nursing Management* **16**,
608–613.

Tracey C.A. (2003) *Experiences of top-level nurses* Unpublished PhD Thesis, University
of Dublin, Trinity College, Dublin.

Williams S. (2004) *Literature review: Evidence of the contribution leadership
development for professional groups makes in driving their organizations forward*
Henley Management College, Oxford.