



HAL
open science

Prevalence of the variant allele rs61764370 T>G in the 3'UTR of among Dutch , and non-/ breast cancer families

Antoinette Hollestelle, Cory Pelletier, Maartje Hooning, Ellen Crepin, Mieke Schutte, Maxime Look, J. Margriet Collee, Anja Nieuwlaat, Lambert C. J. Dorssers, Caroline Seynaeve, et al.

► **To cite this version:**

Antoinette Hollestelle, Cory Pelletier, Maartje Hooning, Ellen Crepin, Mieke Schutte, et al.. Prevalence of the variant allele rs61764370 T>G in the 3'UTR of among Dutch , and non-/ breast cancer families. *Breast Cancer Research and Treatment*, 2010, 128 (1), pp.79-84. <10.1007/s10549-010-1080-z>. <hal-00615370>

HAL Id: hal-00615370

<https://hal.science/hal-00615370v1>

Submitted on 19 Aug 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Prevalence of the variant allele rs61764370 T>G in the 3'UTR of *KRAS* among Dutch *BRCA1*, *BRCA2* and non-*BRCA1/BRCA2* breast cancer families

Antoinette Hollestelle, Cory Pelletier, Maartje Hooning, Ellen Crepin, Mieke Schutte, Maxime Look, J. Margriet Collee, Anja Nieuwlaat, Lambert C.J. Dorssers, Caroline Seynaeve, Yurii S. Aulchenko, John W.M. Martens, Ans M.W. van den Ouweland, Joanne B. Weidhaas

A. Hollestelle (✉), M. Hooning, E. Crepin, M. Schutte, M. Look, C. Seynaeve, J.W.M. Martens

Department of Medical Oncology, Josephine Nefkens Institute and Daniel den Hoed Cancer Center, Erasmus University Medical Center, P.O. Box 2040, 3000 CA Rotterdam, The Netherlands

e-mail: a.hollestelle@erasmusmc.nl

A. Hollestelle

Cancer Genomics Centre, Rotterdam, The Netherlands

J. M. Collee, A. Nieuwlaat, A.M.W. van den Ouweland

Department of Clinical Genetics, Erasmus University Medical Center, Rotterdam, The Netherlands

L.C.J. Dorssers

Department of Pathology, Josephine Nefkens Institute Erasmus University Medical Center, Rotterdam, The Netherlands

Y. S. Aulchenko

Department of Epidemiology, Erasmus University Medical Center, Rotterdam, The Netherlands

C. Pelletier

Department of Genetics, Yale University, New Haven, USA

C. Pelletier, J. B. Weidhaas (✉)

Department of Therapeutic Radiology, Yale University, 333 Cedar Street, New Haven, CT, 06520, USA.

e-mail: joanne.weidhaas@yale.edu

Keywords *KRAS*-variant; *let-7*; breast cancer susceptibility; association; *BRCA1*

Summary

Recently, a variant allele in the 3'UTR of the *KRAS* gene (rs61764370 T>G) was shown to be associated with an increased risk for developing non-small cell lung cancer, as well as ovarian cancer, and was most enriched in ovarian cancer patients from hereditary breast and ovarian cancer families. This functional variant has been shown to disrupt a *let-7* miRNA binding site leading to increased expression of *KRAS in vitro*. In the current study, we have genotyped this *KRAS*-variant in breast cancer index cases from 268 *BRCA1* families, 89 *BRCA2* families, 685 non-*BRCA1/BRCA2* families and 797 geographically-matched controls. The allele frequency of the *KRAS*-variant was found to be increased among patients with breast cancer from *BRCA1*, but not *BRCA2* or non-*BRCA1/BRCA2* families as compared to controls. Because *BRCA1* carriers mostly develop ER-negative breast cancers, we also examined the variant allele frequency among indexes from non-*BRCA1/BRCA2* families with ER-negative breast cancer. The prevalence of the *KRAS*-variant was, however, not significantly increased as compared to controls, suggesting that the variant allele not just simply associates with ER-negative breast cancer. Subsequent expansion of the number of *BRCA1* carriers with breast cancer by including other family members in addition to the index cases resulted in loss of significance for the association between the variant allele and mutant *BRCA1* breast cancer. In this same cohort, the *KRAS*-variant did not appear to modify breast cancer risk for *BRCA1* carriers. Importantly, results from the current study suggest that *KRAS*-variant frequencies might be increased among *BRCA1* carriers, but solid proof requires confirmation in a larger cohort of *BRCA1* carriers.

Introduction

Breast cancer is the most frequently diagnosed type of cancer in women from western countries and an appreciable fraction is attributable to genetic predisposition to the disease. Disease-causing mutations in the high-risk breast cancer susceptibility genes *BRCA1* and *BRCA2* together explain about 20% of the familial clustering of breast cancer. Population-based studies have estimated the cumulative risk of breast cancer to be 65% and 45% for *BRCA1* and *BRCA2* mutation carriers, respectively [1]. The penetrance of *BRCA1* and *BRCA2* mutations, however, is generally higher in women from families with multiple affected individuals, suggesting the presence of other risk-modifying

factors in these families [2-4]. Indeed, it was shown that low-risk breast cancer susceptibility alleles exist that may act multiplicatively on the breast cancer risk for *BRCA1* and *BRCA2* mutation carriers [5, 6].

Besides *BRCA1* and *BRCA2*, three other high-risk breast cancer susceptibility genes have been recognized: *p53*, *PTEN* and *STK11*. However, these account for only a small fraction (less than 1%) of the familial breast cancer risk and are rarely observed outside the context of the cancer syndromes they cause (e.g. Li-Fraumeni syndrome, Cowden syndrome and Peutz-Jeghers syndrome, respectively). Another 5% of the familial breast cancer risk is explained by mutations in moderate-risk genes *ATM*, *BRIP1*, *CHEK2*, *NBS1* and *PALB2* and low-risk alleles such as those in or near the *FGFR2*, *TNRC9/TOX3* and *MAP3K1* genes [7, 8]. This means that so far three quarters of the familial clustering of breast cancer remains unexplained. Importantly, twin studies suggested that a significant proportion of the remaining familial breast cancer risk is most probably due to genetic factors [9, 10].

MicroRNAs (miRNAs) are a class of small non-coding RNA molecules that negatively regulate gene expression by binding partially complementary sites in the 3' untranslated region (UTR) of their target mRNAs. The importance of miRNAs in cancer is stressed by their widespread deregulation in virtually all cancer types [11, 12]. Recently, a variant allele in the 3'UTR of the *KRAS* gene (rs61764370 T>G) was shown to disrupt a *let-7* miRNA binding site leading to increased expression of *KRAS in vitro* and lower *let-7* miRNA levels *in vivo*. [13]. Consistent with the oncogenic nature of *KRAS*, the variant allele was shown to confer an increased risk for developing non-small cell lung cancer (NSCLC), as well as ovarian cancer ([13] and Ratner *et al.*, in press), although the results in NSCLC were not replicated [14]. In head and neck cancers, the *KRAS*-variant was shown to be a genetic marker of poor outcome as well [15]. Because the frequency of the *KRAS*-variant was found to be increased among unselected ovarian cancer cases as well as within the context of hereditary breast and ovarian cancer (HBOC) families, we wondered whether the variant allele frequency might also be increased in breast cancer families. Therefore, we genotyped the *KRAS*-variant in breast cancer index cases from 268 *BRCA1* families, 89 *BRCA2* families and 685 non-*BRCA1/BRCA2* families and compared its prevalence with 797 geographically-matched controls. We also investigated the association of the *KRAS*-variant with specific characteristics of *BRCA1* families.

Materials and Methods

Study population

Families with clustering of breast cancer were selected from the database of the Rotterdam Family Cancer Clinic at Erasmus University Medical Center - Daniel den Hoed Cancer Center in Rotterdam, representing the Southwestern part of the Netherlands. Selected families included at least two cases of female breast cancer or at least one case of female breast cancer and one case of ovarian cancer in first- or second-degree relatives. For each family, the youngest breast cancer patient who had been tested for *BRCA1* and *BRCA2* was assigned to be the index case. The control population was geographically matched and included 797 females from cystic fibrosis families who were either spouses of individuals at risk of being carrier of a *CFTR* mutation or individuals who were tested negative for a *CFTR* mutation. This unmatched case-control study was approved by the local ethical committee and all individuals gave full informed consent to search for susceptibility alleles.

Genotyping

DNA isolated from peripheral blood was amplified using a custom-made Taqman genotyping assay (Applied Biosystems, Foster City, CA) designed specifically to identify the T or variant G allele of the *KRAS*-variant (rs61764370). Forward primer: 5'-GCCAGGCTGGTCTCGAA-3', reverse primer: 5'-CTGAATAAATGAGTTCTGCAAACAGGTT-3', VIC reporter probe: 5'-CTCAAGTGATTCACCCAC-3' and FAM reporter probe: 5'-CAAGTGATTCACCCAC-3'. Because of the low frequency of homozygotes for the variant allele, patient samples that were either heterozygous (TG) or homozygous (GG) for the variant allele were considered positive for the *KRAS*-variant.

Statistical analyses

Associations between the *KRAS*-variant and *BRCA1*-, *BRCA2*-, or non-*BRCA1/BRCA2*-related familial breast cancer, as well as associations between the *KRAS*-variant and specific characteristics of *BRCA1* families were tested for significance by a χ^2 -test. Case-control odds ratios and their confidence intervals were calculated using Woolf approximations. The difference in mean age of onset between *BRCA1* index cases carrying the variant allele and *BRCA1* index cases not carrying the variant allele was determined by a t-test. The result of the Shapiro-Wilk test showed that the assumption of a normal distribution for the age of onset variable was valid. *KRAS*-variant frequencies

among affected and unaffected *BRCA1* carriers were compared using a Cox proportional hazard model. For this purpose, *BRCA1* carriers were censored at either age of breast cancer diagnosis or age at last follow-up. Only *BRCA1* carriers that were censored at breast cancer diagnosis were considered to be affected. Hazard ratios for the TG and GG genotypes combined were estimated using TT homozygotes as a baseline. *P* values of 0.05 or smaller were considered statistically significant. All statistical analyses were performed with STATA statistical package, release 11.0 (STATA Corp, College Station, TX).

Results

To establish whether the *KRAS*-variant (rs61764370 T>G) is implicated in breast cancer susceptibility, we have evaluated the prevalence of this variant in patients with breast cancer from *BRCA1*, *BRCA2* and non-*BRCA1/BRCA2* families as compared to controls. The variant allele was detected in 108 of 685 (15.8%) of indexes with breast cancer from non-*BRCA1/BRCA2* families and 12 of 89 (13.5%) indexes with breast cancer from *BRCA2* families. These frequencies were not significantly different from the 17.3% (138 of 797) prevalence among the control individuals. In contrast, the variant allele was present in 63 of 268 (23.5%) indexes with breast cancer from *BRCA1* families, which was significantly different from the prevalence in controls (Table 1; *P*=0.025; OR=1.47 (1.05-2.06)). However, after adjustment for multiple testing (Bonferroni correction for 3 tests) this difference was not significant anymore (*P*=0.073).

Because *BRCA1* carriers mostly develop ER-negative breast cancers and *BRCA2* carriers mostly develop ER-positive breast cancers, we wondered whether the *KRAS*-variant was associated with ER-negative breast cancer rather than mutant *BRCA1* breast cancer. We had information on ER status of indexes from 483 non-*BRCA1/BRCA2* families of whom 127 were ER-negative and 356 were ER-positive. The variant allele was present in 18 (14.2%) indexes with ER-negative breast cancer and 63 (17.7%) indexes with ER-positive breast cancer which was not significantly different from the prevalence in the controls (Table 1). These results suggest that the *KRAS*-variant not just simply associates with ER-negative breast cancer.

To evaluate whether the variant allele associates with a particular phenotype in *BRCA1* families, we compared family characteristics of *BRCA1* families with a *KRAS*-variant positive index case to *BRCA1* families with a *KRAS*-variant negative index case (Table 2). There was no evidence

for the variant to be associated with the presence of either one or more ovarian cancer cases or bilateral breast cancer cases in the family. We did find a higher frequency of the *KRAS*-variant in families with 3 or more breast cancer cases younger than 60 years compared with families with 2 or less breast cancer cases younger than 60 years (20.4% versus 27.9%), however this was not statistically significant ($P=0.16$). Additionally, we found no difference in mean age of onset between *BRCA1* index cases carrying the variant allele and *BRCA1* index cases not carrying the variant allele (39.8 versus 39.0 years, respectively; $P=0.55$).

In order to gain more statistical power for the analysis of association between the *KRAS*-variant and mutant *BRCA1* breast cancer, we expanded this analysis by including other family members in addition to the index cases. In total, we had DNA from 390 *BRCA1* carriers with breast cancer available for genotyping among the 268 *BRCA1* families. However, the *KRAS*-variant was present in only 20.3% (79 of 390) of affected *BRCA1* carriers (Table 3), which is lower than the initial 23.5% we found among the index cases (Table 1). Consequently, the prevalence of the *KRAS*-variant among *BRCA1* carriers was not significantly different from the controls ($P=0.22$; OR=1.21 (0.89-1.65)), despite expansion of the number of *BRCA1* carriers with breast cancer.

To determine whether or not the *KRAS*-variant modifies *BRCA1* mutant breast cancer risk, all affected and unaffected *BRCA1* carriers among the 268 *BRCA1* families were genotyped for the variant allele. Mean age at censor was 41.0 years for affected *BRCA1* carriers and 45.5 years for unaffected *BRCA1* carriers. Similar to the frequency in the controls, the *KRAS*-variant was present in 17.3% (45 of 260) unaffected *BRCA1* carriers (Table 3). Importantly, there was no evidence of association between the *KRAS*-variant and breast cancer risk among *BRCA1* carriers (Table 3; $P=0.50$; HR=1.07 (0.88-1.30)).

Discussion

In the current study we have found an increase in the frequency of the *KRAS*-variant (rs61764370 T>G) in breast cancer index cases from *BRCA1* families as compared to geographically-matched controls, whereas the frequency of this variant allele was not significantly different among breast cancer index cases from *BRCA2* or non-*BRCA1/BRCA2* families. However, expansion of the number of *BRCA1* carriers with breast cancer by including other family members in addition to the index cases did not improve significance for the association between the variant allele and mutant *BRCA1* breast

cancer. Additionally, the *KRAS*-variant did not appear to modify breast cancer risk for *BRCA1* carriers. Still the question remains whether the breast cancer risk conferred by the *KRAS*-variant is not too small to be detected in our case-control study. Considering the relatively high prevalence in the general population, the variant allele would not be expected to confer a dramatically increased breast cancer risk. Also, the breast cancer risk conferred by the *KRAS*-variant may be larger in GG homozygotes than TG heterozygotes. Therefore, we feel that this trend towards association between the *KRAS*-variant and mutant *BRCA1* breast cancer requires validation in a larger cohort of *BRCA1* carriers, providing more statistical power and enabling the analysis of TG heterozygotes and GG homozygotes separately.

Interestingly, stratification of breast cancer patients from non-*BRCA1/BRCA2* families according to their tumor's ER-status did not reveal an increased *KRAS*-variant frequency among ER-negative breast cancers, suggesting that the variant is not just simply associated with ER-negative breast cancer. This may seem unexpected as the *KRAS*-variant has been shown to lead to increased *KRAS* expression *in vitro* and hyperactivation of the MAPK pathway has been shown to downregulate ER α expression *in vitro* [13, 16]. However, one may wonder whether the level of increased *KRAS* expression as a result of the *KRAS*-variant is similar to that of an oncogenic *RAS* or *RAF* mutation. This may thus be a matter of dosage and therefore MAPK pathway activation should be assessed in tumors from patients that carry the *KRAS*-variant. Unfortunately, due to missing ERBB2 status of the vast majority of non-*BRCA1/BRCA2* cases we have not been able to assess variant allele frequencies in index cases with triple (ER, PR, and ERBB2) negative breast cancers, preventing us to study whether the *KRAS*-variant frequency is increased specifically in *BRCA1* carriers or more generally in patients with triple negative or basal-like breast cancer.

If the *KRAS*-variant were to be specifically associated with mutant *BRCA1* breast cancer, this would suggest a biological interaction between *BRCA1* and *KRAS* or between *BRCA1* and *let-7* miRNA. Many human cancers, including breast cancers, display altered expression of the *let-7* miRNA family and *let-7* has been shown to act as a tumor suppressor regulating the expression of *RAS*, *HMGA2*, *LIN28* and *PEBP1* [11, 17, 18]. Recently, *let-7* was shown to regulate stem cell-like properties in breast tumor initiating cells as downregulation of *let-7* increased proliferation and self-renewal, whereas expression of *let-7* enhanced differentiation [19]. Because the *KRAS*-variant allele was associated with lower *let-7* levels in NSCLC patients as compared to patients without the variant

allele, the *KRAS*-variant may therefore act in tumorigenesis through maintaining a proliferative cell state [13]. Interestingly, knock-down of *BRCA1* in primary breast epithelial cells was shown to increase the number of cells displaying a stem/progenitor phenotype [20]. The presence of the *KRAS*-variant in the germline of a *BRCA1* carrier may thus further facilitate this stem cell phenotype imposed by *BRCA1*. If the *KRAS*-variant would eventually be conclusively associated with mutant *BRCA1* breast cancer, it might be interesting to study this hypothesis in future research.

miRNAs make up one of the most abundant classes of regulatory genes, with 30% of the human genes harboring miRNA target sites [21, 22]. Thus far, many miRNAs have already been implicated in cancer as either oncogenes or tumor suppressor genes [11, 17]. This makes both the miRNA genes themselves as well as the miRNA binding sites in the 3'UTR of well-known cancer genes attractive candidate breast cancer susceptibility alleles. Although several key targets of many miRNAs have been identified, also many target genes remain elusive. A significant portion of these miRNA target genes include well-known breast cancer oncogenes and tumor suppressor genes, such as *ERBB2*, *CCND1*, *MYC* and *PTEN*, but also breast cancer susceptibility genes such as *ATM* [17, 23]. The *KRAS*-variant was the first single nucleotide variant in a miRNA binding site shown to be implicated in cancer risk, but many more are likely to be discovered. Screening of miRNA genes and their target sequences in the 3'UTR of genes known to be implicated in breast cancer may thus provide a productive strategy for defining part of the excess familial breast cancer risk.

Acknowledgements

We wish to thank all members of the breast cancer families that have participated in this study. We also thank Renée Foekens, Bahar Öztürk, Marc Kribbe and Rahena van Kampen-Binda for assistance with collecting DNA samples. Financial support was provided by the Netherlands Genomics Initiative (NGI)/Netherlands Organization for Scientific Research (NWO), a K08 grant from the NIH [CA124484] and a CTSA Grant, Number UL1 RR024139 from the National Center for Research Resources (NCR), a component of the National Institutes of Health (NIH), and NIH roadmap for Medical Research.

Conflict of interest statement

No authors have a conflict of interest with the content of this manuscript.

References

- 1 Antoniou A, Pharoah PD, Narod S, Risch HA, Eyfjord JE, Hopper JL, Loman N, Olsson H, Johannsson O, Borg A, Pasini B, Radice P, Manoukian S, Eccles DM, Tang N, Olah E, Anton-Culver H, Warner E, Lubinski J, Gronwald J, Gorski B, Tulinius H, Thorlacius S, Eerola H, Nevanlinna H, Syrjakoski K, Kallioniemi OP, Thompson D, Evans C, Peto J, Lalloo F, Evans DG, Easton DF (2003) Average risks of breast and ovarian cancer associated with BRCA1 or BRCA2 mutations detected in case Series unselected for family history: a combined analysis of 22 studies. *Am J Hum Genet* 72:1117-1130.
- 2 Ford D, Easton DF, Bishop DT, Narod SA, Goldgar DE (1994) Risks of cancer in BRCA1-mutation carriers. Breast Cancer Linkage Consortium. *Lancet* 343:692-695.
- 3 Ford D, Easton DF, Stratton M, Narod S, Goldgar D, Devilee P, Bishop DT, Weber B, Lenoir G, Chang-Claude J, Sobol H, Teare MD, Struewing J, Arason A, Scherneck S, Peto J, Rebbeck TR, Tonin P, Neuhausen S, Barkardottir R, Eyfjord J, Lynch H, Ponder BA, Gayther SA, Zelada-Hedman M, et al. (1998) Genetic heterogeneity and penetrance analysis of the BRCA1 and BRCA2 genes in breast cancer families. The Breast Cancer Linkage Consortium. *Am J Hum Genet* 62:676-689.
- 4 Struewing JP, Hartge P, Wacholder S, Baker SM, Berlin M, McAdams M, Timmerman MM, Brody LC, Tucker MA (1997) The risk of cancer associated with specific mutations of BRCA1 and BRCA2 among Ashkenazi Jews. *N Engl J Med* 336:1401-1408.
- 5 Antoniou AC, Sinilnikova OM, Simard J, Leone M, Dumont M, Neuhausen SL, Struewing JP, Stoppa-Lyonnet D, Barjhoux L, Hughes DJ, Coupier I, Belotti M, Lasset C, Bonadona V, Bignon YJ, Genetic Modifiers of Cancer Risk in BMCS, Rebbeck TR, Wagner T, Lynch HT, Domchek SM, Nathanson KL, Garber JE, Weitzel J, Narod SA, Tomlinson G, Olopade OI, Godwin A, Isaacs C, Jakubowska A, Lubinski J, Gronwald J, Gorski B, Byrski T, Huzarski T, Peock S, Cook M, Baynes C, Murray A, Rogers M, Daly PA, Dorkins H, Epidemiological Study of BaBMC, Schmutzler RK, Versmold B, Engel C, Meindl A, Arnold N, Niederacher D, Deissler H, German Consortium for Hereditary Breast and Ovarian C, Spurdle AB, Chen X, Waddell N, Cloonan N, Kathleen Cuningham Consortium for Research into Familial Breast C, Kirchhoff T, Offit K, Friedman E, Kaufmann B, Laitman Y, Galore G, Rennert G, Lejbkovicz F, Raskin L,

- Andrulis IL, Ilyushik E, Ozcelik H, Devilee P, Vreeswijk MP, Greene MH, Prindiville SA, Osorio A, Benitez J, Zikan M, Szabo CI, Kilpivaara O, Nevanlinna H, Hamann U, Durocher F, Arason A, Couch FJ, Easton DF, Chenevix-Trench G, Consortium of Investigators of Modifiers of B (2007) RAD51 135G-->C modifies breast cancer risk among BRCA2 mutation carriers: results from a combined analysis of 19 studies. *Am J Hum Genet* 81:1186-1200.
- 6 Antoniou AC, Spurdle AB, Sinilnikova OM, Healey S, Pooley KA, Schmutzler RK, Vermold B, Engel C, Meindl A, Arnold N, Hofmann W, Sutter C, Niederacher D, Deissler H, Caldes T, Kampjarvi K, Nevanlinna H, Simard J, Beesley J, Chen X, Kathleen Cuningham Consortium for Research into Familial Breast C, Neuhausen SL, Rebbeck TR, Wagner T, Lynch HT, Isaacs C, Weitzel J, Ganz PA, Daly MB, Tomlinson G, Olopade OI, Blum JL, Couch FJ, Peterlongo P, Manoukian S, Barile M, Radice P, Szabo CI, Pereira LH, Greene MH, Rennert G, Lejbkowitz F, Barnett-Griness O, Andrulis IL, Ozcelik H, Ocggn, Gerdes AM, Caligo MA, Laitman Y, Kaufman B, Milgrom R, Friedman E, Swedish BaBsc, Domchek SM, Nathanson KL, Osorio A, Llort G, Milne RL, Benitez J, Hamann U, Hogervorst FB, Manders P, Ligtenberg MJ, van den Ouweland AM, collaborators D-H, Peock S, Cook M, Platte R, Evans DG, Eeles R, Pichert G, Chu C, Eccles D, Davidson R, Douglas F, Embrace, Godwin AK, Barjhoux L, Mazoyer S, Sobol H, Bourdon V, Eisinger F, Chompret A, Capoulade C, Bressac-de Paillerets B, Lenoir GM, Gauthier-Villars M, Houdayer C, Stoppa-Lyonnet D, Gemo, Chenevix-Trench G, Easton DF, Cimba (2008) Common breast cancer-predisposition alleles are associated with breast cancer risk in BRCA1 and BRCA2 mutation carriers. *Am J Hum Genet* 82:937-948.
- 7 Stratton MR, Rahman N (2008) The emerging landscape of breast cancer susceptibility. *Nat Genet* 40:17-22.
- 8 Hollestelle A, Wasielewski M, Martens JWM, Schutte M (2010) Discovering moderate-risk breast cancer susceptibility genes. *Current Opinion in Genetics & Development* 20:268-276.
- 9 Ahlbom A, Lichtenstein P, Malmstrom H, Feychting M, Hemminki K, Pedersen NL (1997) Cancer in twins: genetic and nongenetic familial risk factors. *J Natl Cancer Inst* 89:287-293.
- 10 Mack TM, Hamilton AS, Press MF, Diep A, Rappaport EB (2002) Heritable breast cancer in twins. *Br J Cancer* 87:294-300.
- 11 Esquela-Kerscher A, Slack FJ (2006) Oncomirs - microRNAs with a role in cancer. *Nat Rev Cancer* 6:259-269.

- 12 Trang P, Weidhaas JB, Slack FJ (2008) MicroRNAs as potential cancer therapeutics. *Oncogene* 27 Suppl 2:S52-57.
- 13 Chin LJ, Ratner E, Leng S, Zhai R, Nallur S, Babar I, Muller RU, Straka E, Su L, Burki EA, Crowell RE, Patel R, Kulkarni T, Homer R, Zelterman D, Kidd KK, Zhu Y, Christiani DC, Belinsky SA, Slack FJ, Weidhaas JB (2008) A SNP in a let-7 microRNA complementary site in the KRAS 3' untranslated region increases non-small cell lung cancer risk. *Cancer Res* 68:8535-8540.
- 14 Nelson HH, Christensen BC, Plaza SL, Wiencke JK, Marsit CJ, Kelsey KT (2010) KRAS mutation, KRAS-LCS6 polymorphism, and non-small cell lung cancer. *Lung Cancer* 69:51-53.
- 15 Christensen BC, Moyer BJ, Avissar M, Ouellet LG, Plaza SL, McClean MD, Marsit CJ, Kelsey KT (2009) A let-7 microRNA-binding site polymorphism in the KRAS 3' UTR is associated with reduced survival in oral cancers. *Carcinogenesis* 30:1003-1007.
- 16 Oh AS, Lorant LA, Holloway JN, Miller DL, Kern FG, El-Ashry D (2001) Hyperactivation of MAPK induces loss of ERalpha expression in breast cancer cells. *Mol Endocrinol* 15:1344-1359.
- 17 O'Day E, Lal A MicroRNAs and their target gene networks in breast cancer. *Breast Cancer Res* 12:201.
- 18 Johnson SM, Grosshans H, Shingara J, Byrom M, Jarvis R, Cheng A, Labourier E, Reinert KL, Brown D, Slack FJ (2005) RAS is regulated by the let-7 microRNA family. *Cell* 120:635-647.
- 19 Yu F, Yao H, Zhu P, Zhang X, Pan Q, Gong C, Huang Y, Hu X, Su F, Lieberman J, Song E (2007) let-7 regulates self renewal and tumorigenicity of breast cancer cells. *Cell* 131:1109-1123.
- 20 Liu S, Ginestier C, Charafe-Jauffret E, Foco H, Kleer CG, Merajver SD, Dontu G, Wicha MS (2008) BRCA1 regulates human mammary stem/progenitor cell fate. *Proc Natl Acad Sci U S A* 105:1680-1685.
- 21 Lewis BP, Burge CB, Bartel DP (2005) Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 120:15-20.
- 22 Rajewsky N (2006) microRNA target predictions in animals. *Nat Genet* 38 Suppl:S8-13.
- 23 Hu H, Du L, Nagabayashi G, Seeger RC, Gatti RA ATM is down-regulated by N-Myc-regulated microRNA-421. *Proc Natl Acad Sci U S A* 107:1506-1511.

Table 1 *KRAS* 3'UTR SNP genotype frequencies among indexes from *BRCA1*, *BRCA2* and non-*BRCA1/BRCA2* families

Population	Genotype	Frequency	OR	95% CI	P value
Controls	GG+TG	138/797 (17.3%)	1.00		
	TT	659/797 (82.7%)			
<i>BRCA1</i> carriers	GG+TG	63/268 (23.5%)	1.47	1.05-2.06	0.025*
	TT	205/268 (76.5%)			
<i>BRCA2</i> carriers	GG+TG	12/89 (13.5%)	0.74	0.39-1.40	0.36
	TT	77/89 (86.5%)			
Non- <i>BRCA1/BRCA2</i> carriers	GG+TG	108/685 (15.8%)	0.89	0.68-1.18	0.42
	TT	577/685 (84.2%)			
ER-positive	GG+TG	63/356 (17.7%)	1.03	0.74-1.43	0.87
	TT	293/356 (82.3%)			
ER-negative	GG+TG	18/127 (14.2%)	0.79	0.46-1.34	0.38
	TT	109/127 (85.8%)			

Differences in *KRAS*-variant allele frequencies between cases and controls were tested for significance by a χ^2 -test. Case-control odds ratios and their confidence intervals were calculated using Woolf approximations.

*After adjustment for multiple testing (Bonferroni correction for 3 tests) the *P*-value between *BRCA1* carriers and controls was not statistically significant anymore (*P*=0.073).

Table 2 Association of the *KRAS*-variant with specific characteristics of *BRCA1* families

Family characteristic	Frequency of the GG+TG genotype	OR	95% CI	P value
No OvC in family	38/152 (25.0%)	0.89	0.49-1.63	0.71
One or more OvC in family	22/96 (22.9%)			
No Bil BrC in family	36/147 (24.5%)	0.90	0.50-1.63	0.73
One or more Bil BrC in family	24/106 (22.6%)			
Less than 3 BrC < 60 years in family	29/142 (20.4%)	1.51	0.84-2.70	0.16
3 or more BrC < 60 years in family	31/111 (27.9%)			

Association of the *KRAS*-variant with specific characteristics of *BRCA1* families was tested for significance by a χ^2 -test. Case-control odds ratios and their confidence intervals were calculated using Woolf approximations.

OvC, ovarian cancer; Bil, bilateral; BrC, breast cancer.

Table 3 *KRAS* 3'UTR SNP genotype frequencies among affected and unaffected *BRCA1* carriers

Genotype	Frequency in unaffected carriers	Frequency in carriers with breast cancer	HR	95% CI	P value
TT	215/260 (82.7%)	311/390 (79.7%)	1.00	0.88-1.30	0.50
GG+TG	45/260 (17.3%)	79/390 (20.3%)	1.07		

Differences in *KRAS*-variant allele frequencies between affected and unaffected *BRCA1* carriers were tested for significance by a Cox proportional hazards model. Mean age at censor was 41.0 years for affected *BRCA1* carriers and 45.5 years for unaffected *BRCA1* carriers. Hazard ratios for the TG+GG genotype were estimated using TT homozygotes as a baseline.