



HAL
open science

Continuity in the Hurst parameter of the law of the Wiener integral with respect to the fractional Brownian motion

Maria Jolis, Noèlia Viles

► **To cite this version:**

Maria Jolis, Noèlia Viles. Continuity in the Hurst parameter of the law of the Wiener integral with respect to the fractional Brownian motion. *Statistics and Probability Letters*, 2010, 80 (7-8), pp.566. <10.1016/j.spl.2009.12.011>. <hal-00616893>

HAL Id: hal-00616893

<https://hal.science/hal-00616893v1>

Submitted on 25 Aug 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Accepted Manuscript

Continuity in the Hurst parameter of the law of the Wiener integral with respect to the fractional Brownian motion

Maria Jolis, Noèlia Viles


PII: S0167-7152(09)00469-6
DOI: 10.1016/j.spl.2009.12.011
Reference: STAPRO 5594

To appear in: *Statistics and Probability Letters*

Received date: 16 September 2009
Revised date: 9 December 2009
Accepted date: 10 December 2009

Please cite this article as: Jolis, M., Viles, N., Continuity in the Hurst parameter of the law of the Wiener integral with respect to the fractional Brownian motion. *Statistics and Probability Letters* (2009), doi:10.1016/j.spl.2009.12.011

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Continuity in the Hurst parameter of the law of the Wiener integral with respect to the fractional Brownian motion

Maria Jolis

Noèlia Viles

*Departament de Matemàtiques, Universitat Autònoma de Barcelona
08193-Bellaterra, Barcelona, Spain.*

Abstract

We prove the convergence in law, in the space of continuous functions $\mathcal{C}([0, T])$, of the Wiener integral of a deterministic function f with respect to the fractional Brownian motion with Hurst parameter H to the Wiener integral of f with respect to the fractional Brownian motion with parameter H_0 , when H tends to $H_0 \in (0, 1/2]$.

Key words: Fractional Brownian motion, Wiener integral, Convergence in law .
2000 MSC: 60B12, 60H05, 60G15.

1. Introduction

Consider the laws in $\mathcal{C}([0, T])$ of the family of processes $\{B^H, H \in (0, 1)\}$, where each $B^H = \{B_t^H, t \in [0, T]\}$ is a fractional Brownian motion with Hurst parameter H . It is easily seen that these laws converge weakly to that of B^{H_0} , when H tends to $H_0 \in (0, 1)$ (for a proof of this result, see the Introduction of Jolis and Viles (2007b)).

It is an interesting question to study if some important functionals of the fractional Brownian motion conserve this property. That is, we ask if the law of the functional of B^H remains near to that of the same functional of B^{H_0} , when H is near to H_0 . It is worth to mention that this kind of results justify the use of $B^{\hat{H}}$ as a model in applied situations where the true value of the Hurst parameter is unknown and \hat{H} is some estimation of it.

In the previous works Jolis and Viles (2007a) and Jolis and Viles (2007b), we have considered this problem for the functionals given by the multiple fractional integrals with $H \in (1/2, 1)$ and the local time, respectively.

In Jolis and Viles (2007a), as a first and easy step to prove the weak convergence of the multiple Itô-Wiener and Stratonovich integrals, we have proved the convergence in law, in $\mathcal{C}([0, T])$, of the family of Wiener integrals of a deterministic function with respect to B^H with $H \in (\frac{1}{2}, 1)$, when $H \rightarrow H_0 \in [\frac{1}{2}, 1)$.

In this note we study the weak convergence of the Wiener integrals when $H \in (0, \frac{1}{2})$. We do not consider the problem of the convergence of multiple integrals, because the domains and the norms involved in the multiple integrals are much more complicated in this case (see Bardina and Jolis (2006), for instance).

We will prove in our main result (see Theorem 3.5) that for f belonging to the domain of the Wiener integral with respect to $B^{H'}$, with $0 < H' < H_0$, the Wiener integral of f with respect to B^H , with $H \in [H', 1/2]$ converges in law to the Wiener integral of f with respect to B^{H_0} .

We have organized the paper as follows. In Section 2 we give some preliminaries about the Wiener integral with respect to the fractional Brownian motion with Hurst parameter $H < 1/2$, following Bardina and Jolis (2006). Using the characterization of the domain of the integral given in this last paper, we prove in Section 3 the result about the convergence in law of the Wiener integral of a function with respect to B^H , in $\mathcal{C}([0, T])$, when $H \rightarrow H_0$.

Email addresses: mjolis@mat.uab.cat (Maria Jolis), nviles@mat.uab.cat (Noèlia Viles)

2. Preliminaries

Let $B^H = \{B_t^H, t \in [0, T]\}$ be a fractional Brownian motion with Hurst parameter $H \in (0, 1)$. That is, B^H is a centered Gaussian process with covariance function given by

$$R_H(s, t) = E[B_s^H B_t^H] = \frac{1}{2}(t^{2H} + s^{2H} - |t - s|^{2H}).$$

Next, we introduce the Wiener integral with respect to B^H and, from now on, we only consider the case $H < 1/2$, although H_0 can be also equal to $1/2$. Consider the space of simple functions \mathcal{S} in $[0, T]$ of the form

$$f = \sum_{i=1}^N f_i \mathbf{1}_{[a_i, b_i)}, \quad \text{where } [a_i, b_i) \subset [0, T].$$

The Wiener integral of f as above with respect to the fractional Brownian motion B^H is defined in the natural way as

$$I_1^H(f) = \sum_{i=1}^N f_i (B_{b_i}^H - B_{a_i}^H).$$

One can easily see that $I_1^H(f)$ does not depend on the particular representation of f as a simple function and that I_1^H is a linear map from \mathcal{S} into a subspace of $L^2(\Omega)$.

For any $f, g \in \mathcal{S}$ one define the following scalar product

$$\Psi_H(f, g) = E[I_1^H(f) I_1^H(g)].$$

The Wiener integral can be extended in a standard way to the Hilbert space \mathcal{L}^H that is the completion of \mathcal{S} with respect to the inner product Ψ_H . This completion is called *the domain of the Wiener integral*. This space is well-known and can be characterized in terms of fractional derivatives (see, for instance, Alòs et al. (2001), Decreusefond and Üstünel (1999) or Pipiras and Taqqu (2001)).

Nevertheless, along this paper we will use another characterization of the domain proved in Bardina and Jolis (2006) because it is related in a more direct form with the usual fractional Sobolev spaces. This characterization is based in the following result:

Lemma 2.1. (Lemma 2.1., Bardina and Jolis (2006)) For all $f, g \in \mathcal{S}$, we have that

$$\begin{aligned} \Psi_H(f, g) &= \frac{1}{2} H(1 - 2H) \int_0^T \int_0^T \frac{(f(x) - f(y))(g(x) - g(y))}{|x - y|^{2-2H}} dx dy \\ &\quad + H \int_0^T f(x)g(x) \left[\frac{1}{x^{1-2H}} + \frac{1}{(T-x)^{1-2H}} \right] dx. \end{aligned}$$

It can be given also a compact form of the above equality as follows (see Bardina and Jolis (2006)):

$$\Psi_H(f, g) = \frac{1}{2} H(1 - 2H) \iint_{\mathbb{R}^2} \frac{(\bar{f}(x) - \bar{f}(y))(\bar{g}(x) - \bar{g}(y))}{|x - y|^{2-2H}} dx dy, \quad (1)$$

where

$$\bar{f}(x) = \begin{cases} f(x), & \text{if } x \in [0, T] \\ 0, & \text{otherwise.} \end{cases}$$

For all measurable $f : [0, T] \rightarrow \mathbb{R}$, we introduce the following, possibly infinite, quantity:

$$\begin{aligned} \|f\|_H &= \left[\frac{1}{2} H(1 - 2H) \int_0^T \int_0^T \frac{(f(x) - f(y))^2}{|x - y|^{2-2H}} dx dy \right. \\ &\quad \left. + H \int_0^T f(x)^2 \left(\frac{1}{x^{1-2H}} + \frac{1}{(T-x)^{1-2H}} \right) dx \right]^{1/2}. \end{aligned} \quad (2)$$

In the following theorem (see Theorem 2.5, Bardina and Jolis (2006)) the characterization of the space \mathcal{L}^H is given.

Theorem 2.2. (Theorem 2.5, Bardina and Jolis (2006)) The domain of the Wiener integral I_1^H is given by

$$\mathcal{L}^H = \{f \in L^2([0, T]) : \|f\|_H < +\infty\}.$$

If we provide this space with the scalar product

$$\begin{aligned} \Psi_H(f, g) &= \frac{1}{2}H(1 - 2H) \int_0^T \int_0^T \frac{(f(x) - f(y))(g(x) - g(y))}{|x - y|^{2-2H}} dx dy \\ &\quad + H \int_0^T f(x)g(x) \left[\frac{1}{x^{1-2H}} + \frac{1}{(T-x)^{1-2H}} \right] dx, \end{aligned}$$

then the Wiener integral I_1^H is an isometry between \mathcal{L}^H and a closed subspace of $L^2(\Omega)$.

We recall also here the definition of the Sobolev spaces of fractionary order on an interval $[0, T]$. Let $\alpha \in \mathbb{R}$ and $p \in (1, +\infty)$, for any measurable $f : [0, T] \rightarrow \mathbb{R}$ define the following (possibly infinite) quantity

$$\|f\|_{\alpha,p} = \left(\iint_{[0,T]^2} \frac{|f(x) - f(y)|^p}{|x - y|^{1+\alpha p}} dx dy \right)^{1/p}.$$

For $\alpha > 0$, the Sobolev space $W^{\alpha,p}([0, T])$ is defined as

$$W^{\alpha,p}([0, T]) = \{f : \|f\|_{L^p([0,T])} + \|f\|_{\alpha,p} < +\infty\}.$$

This space provided with the norm $\|f\|_{L^p([0,T])} + \|f\|_{\alpha,p}$ is a Banach space.

When $p = 2$, $W^{\alpha,2}([0, T])$ is a Hilbert space with scalar product defined by

$$\langle f, g \rangle_{W^{\alpha,2}([0,T])} = \iint_{[0,T]^2} \frac{(f(x) - f(y))(g(x) - g(y))}{|x - y|^{1+2\alpha}} dx dy + \langle f, g \rangle_{L^2([0,T])}.$$

In the following section we will work with the fractionary Sobolev spaces $W^{1/2-H,2}([0, T])$ of order $\alpha = \frac{1}{2} - H$, that is,

$$W^{1/2-H,2}([0, T]) = \left\{ f \in L^2([0, T]) : \iint_{[0,T]^2} \frac{(f(x) - f(y))^2}{|x - y|^{2-2H}} dx dy < +\infty \right\}.$$

Remark 2.3. Owing Theorem 2.2, we can give the following characterization of the domain of the Wiener integral with respect to the fractional Brownian motion when $H \in (0, 1/2)$:

$$\mathcal{L}^H = \left\{ f \in W^{1/2-H,2}([0, T]) : \int_0^T f(x)^2 \left(\frac{1}{x^{1-2H}} + \frac{1}{(T-x)^{1-2H}} \right) dx < +\infty \right\}$$

and, using the compact expression of Ψ_H given in (1), we have also this other characterization of the space \mathcal{L}_T^H :

$$\mathcal{L}^H = \{f \in L^2([0, T]) : \bar{f} \in W^{1/2-H,2}(\mathbb{R})\},$$

where $W^{1/2-H,2}(\mathbb{R})$ is the Sobolev space defined by

$$W^{1/2-H,2}(\mathbb{R}) = \left\{ f \in L^2(\mathbb{R}) : \iint_{\mathbb{R}^2} \frac{(f(x) - f(y))^2}{|x - y|^{2-2H}} dx dy < +\infty \right\}.$$

We will need also the following Sobolev Embedding Theorem (see Theorem 5.4, Adams (1975)).

Theorem 2.4. (Sobolev Embedding Theorem) Suppose that $0 < \alpha < \frac{1}{p}$ with $p \in (1, +\infty)$. Then,

$$W^{\alpha,p}([0, T]) \hookrightarrow L^q([0, T]),$$

where $p \leq q \leq \frac{p}{1-\alpha p}$.

Applying the above Sobolev Embedding Theorem with $p = 2$, $\alpha = \frac{1}{2} - H$ and $q = \frac{p}{1-\alpha p} = \frac{1}{H}$, we obtain that

$$W^{1/2-H,2}([0, T]) \hookrightarrow L^{1/H}([0, T]),$$

and, as a consequence,

$$\mathcal{L}^H \subset L^{1/H}([0, T]).$$

3. Convergence in law of the Wiener integral

3.1. Results in order to obtain Tightness

Along this section, we will denote by

$$\mathcal{I}_1^H(f) = \{\mathcal{I}_1^H(f \mathbf{1}_{[0,t]}), t \in [0, T]\},$$

the Wiener integral of f with respect to B^H as a process.

Next, we state some technical results in order to prove the tightness of the family of laws of $\{\mathcal{I}_1^H(f)\}_H$ for H belonging to a neighborhood of H_0 . The first one, is a very simple inequality.

Lemma 3.1. *Let $T, k > 0$ positive real numbers, then for any $x \in \mathbb{R}$ such that $|x| \leq T$ and $0 < \alpha_1 < \alpha_2 \leq k$, there exists a constant $C_{T,k}$ (only depending on T and k) for which we have*

$$|x|^{\alpha_2} \leq C_{T,k} |x|^{\alpha_1}.$$

The following proposition is the main result of this section since it provides an estimation of the second order moment of the increments $\mathcal{I}_1^H(f)$ that will give the tightness of the family of laws of the Wiener integral processes for H in a neighborhood of H_0 .

Proposition 3.2. *Let $f \in \mathcal{L}^H$ with $H' \in (0, \frac{1}{2})$. Then, for $H_1 \in (H', \frac{1}{2})$, there exists a positive constant $C = C_{H', H_1, f, T}$, only depending on H', H_1, f and T , such that*

$$\sup_{H \in [H_1, \frac{1}{2})} E|\mathcal{I}_1^H(f \mathbf{1}_{[s,t]})|^2 \leq C(t-s)^{2(H_1-H')}, \quad (3)$$

for all $0 \leq s \leq t \leq T$.

Proof. Let $H \in [H_1, \frac{1}{2})$. In order to simplify the notations, and taking into account that we are only interested on the possible dependence on H of the constants, from now on we will use C to denote all the constants in this proof, except those that depend on H .

Using expression (2) we have that for all $s \leq t$

$$\begin{aligned} E|\mathcal{I}_1^H(f \mathbf{1}_{[s,t]})|^2 &= \|f \mathbf{1}_{[s,t]}\|_H^2 \\ &= \frac{1}{2}H(1-2H) \left(\int_s^t \int_s^t \frac{(f(x)-f(y))^2}{|x-y|^{2-2H}} dx dy + \int_s^t \int_{[0,T] \setminus [s,t]} \frac{f(y)^2}{|x-y|^{2-2H}} dx dy \right. \\ &\quad \left. + \int_{[0,T] \setminus [s,t]} \int_s^t \frac{f(x)^2}{|x-y|^{2-2H}} dx dy \right) + H \int_s^t f(x)^2 \left[\frac{1}{x^{1-2H}} + \frac{1}{(T-x)^{1-2H}} \right] dx. \end{aligned} \quad (4)$$

Applying Hölder's inequality with measure $\mu(dx, dy) = (f(x) - f(y))^2 dx dy$, taking $p = \frac{1-H'}{1-H} > 1$ and q its conjugate, we majorize the integral of the first term of (4) in the following way:

$$\begin{aligned} \int_s^t \int_s^t \frac{(f(x)-f(y))^2}{|x-y|^{2-2H}} dx dy &\leq \left(\int_s^t \int_s^t \frac{(f(x)-f(y))^2}{|x-y|^{2-2H'}} dx dy \right)^{1-H/1-H'} \\ &\quad \times \left(\int_s^t \int_s^t (f(x)-f(y))^2 dx dy \right)^{H-H'/1-H'}. \end{aligned}$$

Since $H \geq H_1$, we have that

$$\left(\int_s^t \int_s^t \frac{(f(x)-f(y))^2}{|x-y|^{2-2H'}} dx dy \right)^{1-H/1-H'} \leq \max(1, \|f\|_{W^{1/2-H', 2}([0,T])}^2)^{\frac{(1-H_1)}{1-H'}}.$$

Since, due to the Sobolev Embedding Theorem, $f \in L^{1/H'}([0, T])$, and applying again Hölder's inequality, with $p = \frac{1}{2H'}$, $q = \frac{1}{1-2H'}$, we obtain

$$\begin{aligned} \left(\int_s^t \int_s^t f(x)^2 dx dy \right)^{\frac{H-H'}{1-H'}} &\leq \left(\int_s^t \left(\int_s^t |f(x)|^{1/H'} dx \right)^{2H'} (t-s)^{1-2H'} dy \right)^{\frac{H-H'}{1-H'}} \\ &= \left(\int_s^t |f(x)|^{1/H'} dx \right)^{2H' \left(\frac{H-H'}{1-H'} \right)} (t-s)^{2(H-H')} \\ &\leq C(t-s)^{2(H-H')}. \end{aligned}$$

In this way, we have an upper bound for the first summand of (4) of the form $C(t-s)^{2(H-H')}$.

The remaining terms can be handled easily. In fact, the second and third summands of (4) coincide. Computing the integral with respect to x of the second term and making some simple majorizations,

$$\begin{aligned} & \int_s^t \int_{[0,T] \setminus [s,t]} \frac{f(y)^2}{|x-y|^{2-2H}} dx dy \\ &= \frac{1}{1-2H} \left[\int_s^t f(y)^2 ((y-s)^{2H-1} - y^{2H-1}) dy + \int_s^t f(y)^2 ((t-y)^{2H-1} - (T-y)^{2H-1}) dy \right] \\ &\leq \frac{1}{1-2H} \left(\int_s^t f(y)^2 (y-s)^{2H-1} dy + \int_s^t f(y)^2 (t-y)^{2H-1} dy \right). \end{aligned}$$

Using again that $f \in L^{1/H'}([0, T])$, applying Hölder's inequality to each summand of the right-hand side of the last inequality and using that the exponent $\frac{2H-1}{1-2H} > -1$, due to the fact that $H \geq H_1 > H'$, we have that

•

$$\begin{aligned} \int_s^t f(y)^2 (y-s)^{2H-1} dy &\leq \left(\int_s^t |f(y)|^{1/H'} dy \right)^{2H'} \left(\int_s^t (y-s)^{2H-1/1-2H'} dy \right)^{1-2H'} \\ &\leq C \left(\frac{1-2H'}{2(H-H')} \right)^{1-2H'} (t-s)^{2(H-H')}. \end{aligned}$$

•

$$\begin{aligned} \int_s^t f(y)^2 (t-y)^{2H-1} dy &\leq \left(\int_s^t |f(y)|^{1/H'} dy \right)^{2H'} \left(\int_s^t (t-y)^{2H-1/1-2H'} dy \right)^{1-2H'} \\ &\leq C \left(\frac{1-2H'}{2(H-H')} \right)^{1-2H'} (t-s)^{2(H-H')}. \end{aligned}$$

So, taking into account these inequalities and using that $H - H' > H_1 - H' > 0$, we obtain that

$$\frac{1}{2} H(1-2H) \int_{[0,T] \setminus [s,t]} \int_s^t \frac{f(x)^2}{|x-y|^{2-2H}} dx dy \leq C(t-s)^{2(H-H')}.$$

In a similar way we deal with the fourth summand of (4). Indeed, first using that $f \in L^{1/H'}([0, T])$ and then applying in a convenient form Hölder's inequality, we obtain that

$$\begin{aligned} & \int_s^t f(x)^2 \left[\frac{1}{x^{1-2H}} + \frac{1}{(T-x)^{1-2H}} \right] dx \\ &\leq \left(\int_s^t |f(x)|^{1/H'} dx \right)^{2H'} \left(\int_s^t x^{2H-1/1-2H'} dx \right)^{1-2H'} \\ &\quad + \left(\int_s^t |f(x)|^{1/H'} dx \right)^{2H'} \left(\int_s^t (T-x)^{2H-1/1-2H'} dx \right)^{1-2H'} \\ &\leq C [t^{2(H-H')} - s^{2(H-H')} + (T-s)^{2(H-H')} - (T-t)^{2(H-H')}]. \end{aligned} \tag{5}$$

Since for $0 < \alpha < 1$,

$$x^\alpha - y^\alpha \leq (x-y)^\alpha,$$

we have that:

- $t^{2(H-H')} - s^{2(H-H')} \leq (t-s)^{2(H-H')}.$
- $(T-s)^{2(H-H')} - (T-t)^{2(H-H')} \leq (t-s)^{2(H-H')}.$

Therefore, the term (5) can be bounded by

$$C(t-s)^{2(H-H')}.$$

Finally, using the above inequalities and Lemma 3.1, we deduce that

$$\sup_{H \in [H_1, \frac{1}{2})} E|I_1^H(f \mathbf{1}_{[s,t]})|^2 \leq C(t-s)^{2(H-H')} \leq C(t-s)^{2(H_1-H')},$$

with C a positive constant. □

3.2. Results for the convergence of the finite-dimensional distributions

An important tool in the proof of the convergence of the finite-dimensional distributions is the following general lemma, that we state in our particular setting.

Lemma 3.3. *Let $(G, \|\cdot\|)$ be a normed space, and let $\{\mathcal{J}^H\}_{H \in V_0}$ (where V_0 is an interval that contains H_0) be a family of linear maps defined on G and taking values in $(L^0(\Omega))^m$, the space of m -dimensional finite a.s. random vectors. Denote by $|\cdot|$ the euclidian norm in \mathbb{R}^m . Suppose that there exists a positive constant C such that, for any $f \in G$,*

$$(C) \quad \sup_{H \in V_0} E|\mathcal{J}^H(f)| \leq C\|f\|.$$

Suppose also that, for some dense subset $D \subset G$, we have that

$$\mathcal{J}^H(f) \xrightarrow{\mathcal{L}} \mathcal{J}^{H_0}(f), \quad \text{for all } f \in D, \text{ when } H \rightarrow H_0.$$

Then, $\mathcal{J}^H(f) \xrightarrow{\mathcal{L}} \mathcal{J}^{H_0}(f)$, when $H \rightarrow H_0$, for all $f \in G$.

Next, we will prove this lemma; though it can be found around in the literature, we have not been able to find an explicit proof.

Proof. Let $h \in \mathcal{C}^1(\mathbb{R}^m)$ having bounded partial derivatives. We want to prove that for any $f \in G$

$$|E[h(\mathcal{J}^H(f))] - E[h(\mathcal{J}^{H_0}(f))]|$$

tends to zero when $H \rightarrow H_0$.

Since $h \in \mathcal{C}^1(\mathbb{R}^m)$ and it has bounded partial derivatives, there exists a constant C_h such that for all $x, y \in \mathbb{R}^m$

$$|h(x) - h(y)| < C_h|x - y|.$$

As $D \subset E$ is dense in E we have that for any $f \in E$ and any $\varepsilon > 0$ there exists a function $g \in D$ such that $\|f - g\| < \frac{\varepsilon}{3C_h}$, where C is the constant that appears in the hypothesis (C). We will write

$$|E[h(\mathcal{J}^H(f))] - E[h(\mathcal{J}^{H_0}(f))]| \leq A_1 + A_2 + A_3,$$

where

$$A_1 = C_h E(|\mathcal{J}^H(f) - \mathcal{J}^H(g)|),$$

$$A_2 = |E[h(\mathcal{J}^H(g))] - E[h(\mathcal{J}^{H_0}(g))]|,$$

and

$$A_3 = C_h E(|\mathcal{J}^{H_0}(g) - \mathcal{J}^{H_0}(f)|).$$

By condition (C), the term A_1 can be bounded in the following way

$$\begin{aligned} A_1 &\leq C_h \sup_{H \in V_0} E(|\mathcal{J}^H(f) - \mathcal{J}^H(g)|) \leq C_h \sup_{H \in V_0} E(|\mathcal{J}^H(f - g)|) \\ &\leq C_h C \|f - g\| < \frac{\varepsilon}{3}. \end{aligned}$$

Similarly, we can show that the term A_3 is bounded by $\frac{\varepsilon}{3}$.

Finally, using that for $g \in D$ we have the convergence in law

$$\mathcal{J}^H(g) \xrightarrow{\mathcal{L}} \mathcal{J}^{H_0}(g),$$

we get that the term A_2 can be also bounded by $\frac{\varepsilon}{3}$ if H is sufficiently close to H_0 . □

To check that our family of processes satisfies the condition (C) of the above lemma we need the inequality stated in the next lemma.

Lemma 3.4. *For any $f \in \mathcal{L}^{H'}$ and $H \in [H', \frac{1}{2}]$, there exists a constant $C_{T,H'}$ (only depending on T and H') such that*

$$\|f\|_H^2 \leq C_{T,H'} \|f\|_{H'}^2.$$

Proof. We distinguish the two cases $0 < H < \frac{1}{2}$ and $H = \frac{1}{2}$.

1. Case $H \in (0, \frac{1}{2})$:

As $0 < 1 - 2H \leq 1 - 2H'$ for $H \in [H', \frac{1}{2}]$, and applying Lemma 3.1 we obtain that:

- $|x - y|^{2-2H} \geq C_{T,H'} |x - y|^{2-2H'}$,
- $x^{1-2H} \geq C_{T,H'} x^{1-2H'}$,
- $(T - x)^{1-2H} \geq C_{T,H'} (T - x)^{1-2H'}$.

Applying these inequalities to each summand of the expression of $\|f\|_H$ we have that:

$$\begin{aligned} \|f\|_H^2 &= \frac{1}{2} H(1 - 2H) \int_0^T \int_0^T \frac{(f(x) - f(y))^2}{|x - y|^{2-2H}} dx dy + H \int_0^T f^2(x) \left[\frac{1}{x^{1-2H}} + \frac{1}{(T - x)^{1-2H}} \right] dx \\ &\leq C_{T,H'} \int_0^T \int_0^T \frac{(f(x) - f(y))^2}{|x - y|^{2-2H'}} dx dy + \frac{1}{2} C_{T,H'} \int_0^T f^2(x) \left[\frac{1}{x^{1-2H'}} + \frac{1}{(T - x)^{1-2H'}} \right] dx \end{aligned}$$

From here, we deduce that

$$\|f\|_H^2 \leq C_{H',T} \|f\|_{H'}^2.$$

2. Case $H = \frac{1}{2}$:

Given that $x = \frac{T}{2}$ is the minimum of the function

$$g(x) = \frac{1}{x^{1-2H'}} + \frac{1}{(T - x)^{1-2H'}}, \quad x \in [0, T],$$

we have that

$$\int_0^T f^2(x) \left[\frac{2}{\left(\frac{T}{2}\right)^{1-2H'}} \right] dx \leq \int_0^T f^2(x) \left[\frac{1}{x^{1-2H'}} + \frac{1}{(T - x)^{1-2H'}} \right] dx$$

or equivalently,

$$\int_0^T f^2(x) dx \leq \frac{1}{2} \left(\frac{T}{2}\right)^{1-2H'} \int_0^T f^2(x) \left[\frac{1}{x^{1-2H'}} + \frac{1}{(T - x)^{1-2H'}} \right] dx$$

So, taking into account this last inequality, we obtain easily the bound

$$\|f\|_{1/2}^2 \leq C_{T,H'} \|f\|_{H'}^2.$$

□

3.3. Main result

We finally prove the convergence in law in $\mathcal{C}([0, T])$ of the processes $\{\mathcal{I}_1^H(f)\}_H$, when $H \rightarrow H_0 \in (0, \frac{1}{2}]$, for all $f \in \mathcal{L}^{H'}$, for some $H' < H_0$.

Theorem 3.5. *Let $f \in \mathcal{L}^{H'}$ with $H' < H_0$ and $H_0 \in (0, \frac{1}{2}]$. Then, the family of processes $\{\mathcal{I}_1^H(f)\}_{H \in (H', \frac{1}{2}]}$ converges in law to $\mathcal{I}_1^{H_0}(f)$, in the space $\mathcal{C}([0, T])$, when $H \rightarrow H_0$.*

Proof. The first step is to prove the existence of a continuous version of the Wiener integral $\mathcal{I}_1^H(f)$. Taking into account that $\mathcal{I}_1^H(f)$ is a centered Gaussian process, we only need to apply Proposition 3.2 and Kolmogorov's continuity criterium.

Using again that the processes $\mathcal{I}_1^H(f)$ are centered and Gaussian, by Proposition 3.2 and Billingsley's criterium (see (Billingsley, 1968, Theorem 12.3)) we obtain the tightness of the laws of the family $\{\mathcal{I}_1^H(f), H \in [H_1, \frac{1}{2}]\}$ in $\mathcal{C}([0, T])$, for all $0 < H' < H_1 < \frac{1}{2}$.

It only remains to show the convergence of the finite-dimensional distributions of the processes $\mathcal{I}_1^H(f)$. We prove this by applying Lemma 3.3 taking $G = \mathcal{L}_T^{H'}$ endowed with the norm $\|\cdot\|_{H'}$, and

$$\begin{aligned} \mathcal{J}^H : G &\longrightarrow (L^0(\Omega))^m \\ f &\longmapsto (I_1^H(f\mathbf{1}_{[0,t_1]}), \dots, I_1^H(f\mathbf{1}_{[0,t_m]})). \end{aligned}$$

By Lemma 3.4, the linear maps \mathcal{J}^H satisfy condition (C) of Lemma 3.3. Moreover, for $f \in \mathcal{S}$ (that is a dense subspace of G), the convergence in law of $\mathcal{I}_1^H(f)$ to $\mathcal{I}_1^{H_0}(f)$ is obtained from the fact that the integrals $I_1^H(f\mathbf{1}_{[0,t_j]})$ are linear combinations of increments of B^H , and B^H converges in law to B^{H_0} , as $H \rightarrow H_0$. \square

Acknowledgement

The authors are supported by the grant MEC-FEDER Ref. MTM2006-06427 from the Dirección General de Investigación, Ministerio de Educación y Ciencia, Spain.

The authors would like to thank an anonymous referee for helpful suggestions.

References

References

- Adams, R. A. *Sobolev spaces*. Academic Press, New York-London, 1975. Pure and Applied Mathematics, Vol. 65.
- Alòs, E., Mazet, O., Nualart, D. Stochastic calculus with respect to Gaussian processes. *Ann. Probab.*, 29, no. 2, 766–801, 2001.
- Billingsley, P. *Convergence of probability measures*. John Wiley & Sons Inc., New York, 1968.
- Bardina, X., Jolis, M. Multiple fractional integral with Hurst parameter less than $\frac{1}{2}$. *Stochastic Process. Appl.*, 116, no. 3, 463–479, 2006.
- Decreusefond, L., Üstünel, A. S. Stochastic analysis of the fractional Brownian motion. *Potential Anal.*, 10, no. 2, 177–214, 1999.
- Jolis, M. On the Wiener integral with respect to the fractional Brownian motion on an interval. *J. Math. Anal. Appl.* 330, no. 2, 1115–1127, 2007.
- Jolis, M., Viles, N. Continuity with respect to the Hurst parameter of the laws of the multiple fractional integrals. *Stochastic Process. Appl.* 117, no. 9, 1189–1207, 2007.
- Jolis, M., Viles, N. Continuity in law with respect to the Hurst parameter of the local time of the fractional Brownian motion. *J. Theoret. Probab.* 20, no. 2, 133–152, 2007.
- Pipiras, V., Taqqu, M. S. Are classes of deterministic integrands for fractional Brownian motion on an interval complete? *Bernoulli*, 7, no. 6, 873–897, 2001.
- Rudin, W. *Real and complex analysis*. McGraw-Hill Book Co., New York, third edition, 1987.
- Samorodnitsky, G., Taqqu, M.S. *Stable non-Gaussian random processes*. Stochastic Modeling. Chapman & Hall, New York, 1994. Stochastic models with infinite variance.