

Inflammation in Transgenic Mouse Models of Neurodegenerative Disorders

Claudia Schwab, Andis Klegeris, Patrick L. Mcgeer

▶ To cite this version:

Claudia Schwab, Andis Klegeris, Patrick L. Mcgeer. Inflammation in Transgenic Mouse Models of Neurodegenerative Disorders. Biochimica et Biophysica Acta - Molecular Basis of Disease, 2010, 1802 (10), pp.889. 10.1016/j.bbadis.2009.10.013 . hal-00616908

HAL Id: hal-00616908 https://hal.science/hal-00616908

Submitted on 25 Aug 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Inflammation in Transgenic Mouse Models of Neurodegenerative Disorders

Claudia Schwab, Andis Klegeris, Patrick L. McGeer

PII: DOI: Reference: S0925-4439(09)00249-X doi:10.1016/j.bbadis.2009.10.013 BBADIS 63029

BBA - Molecular Basis of Disease

To appear in:

Received date: Revised date: Accepted date:

7 July 2009 9 October 2009 23 October 2009



Please cite this article as: Claudia Schwab, Andis Klegeris, Patrick L. McGeer, Inflammation in Transgenic Mouse Models of Neurodegenerative Disorders, *BBA - Molecular Basis of Disease* (2009), doi:10.1016/j.bbadis.2009.10.013

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Inflammation in transgenic mouse models

INFLAMMATION IN TRANSGENIC MOUSE MODELS OF NEURODEGENERATIVE DISORDERS

Claudia Schwab (a,*), Andis Klegeris (b) and Patrick. L. McGeer (a)

(a) Department of Psychiatry, Kinsmen Laboratory of Neurological Research, University of British Columbia, 2255 Wesbrook Mall, Vancouver, BC V6T 1Z3, Canada
(b) Laboratory of Cellular and Molecular Pharmacology, Department of Biology, I.K. Barber School of Arts and Sciences, University of British Columbia Okanagan, 3333 University Way, Kelowna, BC V1V 1V7, Canada

* Corresponding author:

Department of Psychiatry, Kinsmen Laboratory of Neurological Research, University of British Columbia, 2255 Wesbrook Mall, Vancouver, BC V6T 1Z3, Canada E-mail: claudia.schwab@ubc.ca, Fax : 1-604-822-7086

- 1. INTRODUCTION
- 2. TRANSGENIC MOUSE MODELS OF ALZHEIMER'S DISEASE
- 2.1. Complement activation in AD mouse models
- 2.2. Microglial activation in AD mouse models
- 2.3. Cytokines in AD mouse models
- 2.4. Efficiency of NSAIDs in AD mouse models
- 3. TRANSGENIC MOUSE MODELS OF PARKINSON'S DISEASE
- 3.1 Alpha-synuclein transgenic mouse models
- 3.2. DJ-1, Parkin and PINK transgenic mouse models
- 4. TRANSGENIC MOUSE MODELS OF HUNTINGTON'S DISEASE
- 5. SUMMARY

2

Inflammation in transgenic mouse models

Keywords: Transgenic mice, Alzheimer's disease, Parkinson's disease, Huntington's disease, immune system, microglia, cytokines, complement system

Abstract:

Much evidence is available that inflammation contributes to the development of neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease and Huntington's disease. Our review investigates how well current mouse models reflect this aspect of the pathogenesis.

Transgenic models of AD have been available for several years and are the most extensively studied. Modulation of cytokine levels, activation of microglia and, to a lesser extent, activation of the complement system have been reported. Mouse models of PD and HD so far show less evidence for the involvement of inflammation. An increasing number of transgenic mouse strains is being created to model human neurodegenerative diseases. A perfect model should reflect all aspects of a disease. It is important to evaluate continuously the models for their match with the human disease and reevaluate them in light of new findings in human patients.

Although none of the transgenic mouse models recapitulates all aspects of the human disorder they represent, all models have provided valuable information on basic molecular pathways. In particular, the mouse models of Alzheimer disease have also led to the development of new therapeutic strategies such as vaccination and modulation of microglial activity.

Inflammation in transgenic mouse models

1. INTRODUCTION

The perfect animal model of a human disorder should reflect (i) the symptoms, (ii) the lesions and (iii) the causes [1]. Here we will review how transgenic mouse models meet these criteria with special consideration of the inflammation hypothesis. We discuss the role and modulation of inflammatory processes in these models.

The inflammation hypothesis proposes that a chronic inflammatory response occurs in neurodegenerative disorders such as Alzheimer's disease (AD), Parkinson's disease (PD) and Huntington's disease (HD), and that this response exacerbates the characteristic neurodegeneration. Several findings support the hypothesis for AD and to some extent for PD and HD: (i) markers of inflammation are elevated in postmortem brain tissue (gliosis, proinflammatory cytokines, complement activation); (ii) epidemiological studies indicate that long-term use of non-steroidal anti-inflammatory drugs (NSAIDs) lowers the risk of disease; (iii) DNA polymorphism of some cytokines and acute phase proteins modulates the risk; and (iv) animal and cell culture models show that modulation of inflammation is effective in curbing the disease process (reviewed in [2-4]). When involvement of inflammation in these disorders was first discovered, it was viewed solely as a detrimental factor. More recently it has been suggested that certain aspects of the immune response can play a positive role in regeneration and clearance of pathological aggregates, creating great interest in the possible modulation of immune responses in order to obtain beneficial effects [5-7].

2. TRANSGENIC MOUSE MODELS OF ALZHEIMER'S DISEASE

In AD, a disturbance of beta amyloid protein (Aβ) metabolism leads to the formation of soluble oligomers and larger, insoluble aggregates. It is now thought that the oligomers themselves may be neurotoxic, causing synapse loss and neuronal degeneration, accompanied by aberrant tau phosphorylation and formation of neurofibrillary tangles (NFT). Transgenic mouse models of AD have been created by introducing to the mouse genome mutated forms of the human amyloid precursor protein (APP) or the presenilin 1 (PS1) gene found in familial AD cases (sporadic AD is not connected with such mutations). Introduction of wild-type forms of APP and PS1 as found in normals and non-familial AD cases does not result in mouse pathology. The mice expressing

Inflammation in transgenic mouse models

pathological human mutant proteins readily develop $A\beta$ deposits, show behavioral deficits, and display some aberrant tau-phosphorylation, even though very little neuronal death occurs (for recent reviews see: [1, 8, 9]). Neurofibrillary tangles (one of the major pathological hallmarks of AD) are not found in these mouse strains. Different transgenic models with additional pathological tau mutations have been created to mimic this aspect of AD [10]. Even though these tau mutations cause other tauopathies, they are not associated with AD.

2.1. Complement activation in AD mouse models

The complement system is a major component of the body's immune defense. It is an ancient and highly conserved system which is a key component of the innate defense against microbes and other infections. It is also a key component of the adaptive immune system clearing antigen-antibody complexes. In addition, it is involved in clearance of damaged tissue and necrotic cells of the body itself. The complement pathways are cascades of proteins which are sequentially cleaved and terminally assembled. They can be activated via the classical pathway initiated by C1q binding, the alternative pathway, starting with activation of C3, or the lectin pathway. Full complement activation results in formation of the membrane attack complex (MAC or C5b-9) consisting of the assembly of the proteins C5b through C9. MAC inserts into membranes and lyses cells. The complement system is a potent mechanism and is tightly regulated by inhibitors such as C1 inhibitor, factor H and C4b-binding protein. Activation of the complement system results in opsonization of targets for phagocytosis, production of pro-inflammatory anaphylatoxins (C5a, C3a) and in lysis of cells. For a review of the complement system see [11].

Complement activation in AD was first described more than two decades ago (for review see [12, 13]). This activation likely occurs as an attempt to clear toxic protein aggregates and debris of dying cells. Complement activation attracts microglia, promotes secretion of inflammatory cytokines and induces neuronal damage via MAC and free radicals, contributing to the pathogenesis of AD [3]. It has been demonstrated that $A\beta$ is a powerful activator of complement. It binds to C1q [14, 15], initiating the classical pathway of complement activation. A β also activates the alternative pathway by

4

Inflammation in transgenic mouse models

formation of complexes with activation products of C3 [16]. In AD complement is activated fully [17, 18] and early [19] in the disease process. With the exception of C4b binding protein [20], expression of complement inhibitors is deficient and the complement system activation is not adequately controlled [19, 21]. Markers of complement activation have also been detected in several AD mouse models. In a triple mutation model of cerebral amyloid angiopathy (SwDI; Dutch E693Q, Iowa D694N and Swedish K670N/M671L mutations), Fan et al. [22] described that mRNAs of the early complement components C1q and C3 were upregulated. C1q, C3, and C4 protein levels were also elevated in Western blots. In immunostaining, the complement proteins colocalized with fibrillary A β deposits and microglia. Matsuoka and colleagues showed that mouse C1q strongly colocalized with fibrillary A β , and was also upregulated in some plaque-associated microglia in PS/APP mice [23]. Failure to detect robust C1q binding to plaques in another report [24] may be due to differences between transgenic strains or the anti-C1q antibodies used. A more recent study found C1q deposition to increase with age in Tg2576 mice [25].

Clq knock-out mice (Q-/-) are not able to activate the classical complement pathway. Crossing this strain with Tg2576 or APP/PS1 mice reduced neuropathology as demonstrated by a decreased loss of the presynaptic marker synaptophysin and diminished glia activation, while the plaque load remained unchanged [26]. In the double-transgenic strain (APPQ-/-), Zhou and colleagues reported reduced C4 expression and upregulated C3 expression [25], suggesting that components of the alternative complement pathway (C3b, C3a, or C5a) could have a neuroprotective function. However, a recent study showed that C5a may actually play a detrimental role [27]. Additional evidence for a neuroprotective function of C3 of the alternative pathway was provided by Maier et al. [28] who found that C3 deficiency increased Aβ plaque formation and neuronal loss in APP mice which were crossed with C3 knock-out mice. The phenotype of microglia was switched towards an increase in CD45 expressing cells. Overexpression of the C3 inhibitor, soluble complement receptor-related protein y (sCrry), also caused a 2-3 fold increase in A β deposition and increased neuropathology in APP mice [29]. A different approach to complement modulation was used by Pillay and colleagues by employing Vaccinia virus complement control protein (VCP), which

5

Inflammation in transgenic mouse models

inhibits both the classical and alternate complement pathways. It improved memory performance in APP/PS1 mice [30].

Although an activation of the complement system occurs in mouse models of AD, several studies have pointed out significant differences between the human and mouse systems (for reviews see [31, 32]). It has been show that mouse C1q binds to A β [23, 26], but as Webster and colleagues demonstrated binding was less efficient compared to the human form, resulting in a lower activation of the mouse complement complex in transgenic models compared with humans [33]. Immunostaining of plaque-associated complement proteins was generally weaker in mice compared to AD [24](see figure 1). By using erythrocyte lysis assays Ong et al. [34] showed that mice have lower serum complement protein levels compared to humans and other species. Ebanks and colleagues found that mouse complement C4 lacked C5 convertase activity, preventing complement activation via the classical pathway [35].

Taken together, these studies show that the complement system is activated in transgenic models of AD (for summary see table 1). Complement protein expression is increased and associated with plaques and glial cells. While C1q deficiency reduces neuronal degeneration and improves learning, inhibition of the alternative pathway (C3) increases A β pathology and neuronal loss. Important differences between AD and mouse models are the lower expression levels of the complement proteins, and possibly reduced binding of C1q to A β as well as lack of late-stage complement activation [17]. These mouse models also fail to display extensive tau pathology and widespread neuronal death. Tau aggregates are, in addition to A β aggregates, possible activators of the complement system. Several approaches of modulating the complement system in transgenic mice have been shown to reduce neuropathology and/or A β load. It remains to be determined if these interventions can also be effective in human AD, particularly in early stages, before full-blown complement activation exacerbates the severe degeneration in end stage disease.

2.2. Microglial activation in AD mouse models

There is a plethora of evidence that activated microglia contribute to neurodegeneration in AD. Microglial cells are the immune effector cells in the CNS and are part of the

Inflammation in transgenic mouse models

mononuclear phagocytic system, expressing characteristic markers of phagocytes and immune cells, such as complement components and their receptors, major histocompatibility complex (MHC) glycoproteins, and scavenger receptors (for review see [36, 37]). Once activated, microglia phagocytose targets, release cytokines and, if fully activated, discharge toxic free radicals [38-40]. In neurodegenerative disorders, such as AD, microglia fail to clear pathological protein aggregates and cellular debris and become chronically activated, continuously releasing neurotoxic substances and contributing to the pathogenesis [41]. It was long thought that microglia exist in either an activated or a resting state. More recently it has been proposed that microglia can be activated into either anti-inflammatory (alternative, adaptive) or pro-inflammatory (classical, chronic) states, having either beneficial (A β clearance, release of neurotrophic factors) or detrimental effects (release of pro-inflammatory cytokines, free radicals) [7, 42].

Microglia accumulate around A\beta deposits and are activated in mouse models In transgenic mouse models, activated microglial cells accumulate in the vicinity of A β plaques, similarly to AD (figure 1). This has been shown in various strains including Tg2576 (Swedish), APP 22 (Swedish/London), APP 23 (Swedish), TgCRND8 (Indiana/Swedish) [43-46]. While in other mouse models microglia are mainly associated with compact (dense core) plaques [47], in TgCRND8 mice, which develop diffuse and compact plaques earlier than other strains (at 9-10 weeks), microglia and astroglia were associated early on with both types of plaques [46]. Early activation of microglia stained with HLA-DR and CD11b was also reported in the PS19 model of tauopathy which carry the human tau P301S mutation. Microglial activation coincided with synaptic pathology, followed later by fibrillary tau pathology and astrogliosis [48]. Although Howlett and colleagues found no up-regulation of microglial CD11b in TASTPM mice (double APP/PS1 transgenic), expression of Iba-1 was greatly increased [9, 49], suggesting a change in activation state.

Recent advances in imaging have made it possible to observe individual cells in vivo over time. Using in vivo multiphoton microscopy Meyer-Luehmann and colleagues [50] demonstrated that plaques form within days in both Tg2576 and PS1/APP mice and that microglial cells aggregate around these newly formed plaques within 1 to 2 days,

7

Inflammation in transgenic mouse models

followed by dysmorphic changes to neurites. Despite the close association with microglia, the plaques were not cleared, but possibly restricted from growing larger. In a different in-vivo imaging study, it was observed that additional microglial cells continue to migrate to plaques, that their volume predicts changes in plaque size, and that the microglia closest to plaques phagocytose more actively [51].

Microglia in SwDI mice were found to express complement proteins C1q, C3, C4 [22]. Microglia expressing complement receptor-3 (CR3, CD11) and astroglia were detected in APP/PS1 (Tg2576/5.1PS1) transgenic mice as soon as the first A β deposits formed at 6 months, but MHC-II expression was not found until 12 months of age, coinciding with stabilization of fibrillary A β and suggesting a more severe state of inflammation and activation [52].

Modulation of states of microglial activity in transgenic mice

Several recent studies have focused on the regulation of microglial activity via their receptors. CD40-CD40 ligand (CD40L) suppression shifts microglia towards increased phagocytic activity ("anti-inflammatory state"). Tan and colleagues crossed Tg2576 mice with CD40L deficient mice and reported decreased plaque loads and microgliosis [53, 54]. Interruption of CD40-CD40L interaction in vaccinated PSAPP and Tg2576 mice increased A β clearance via increasing the phagocytotic activity of microglia, suggesting that CD40–CD40L acts as a molecular switch between pro- and anti-inflammatory states of microglial activation[55]. Increased microglial phagocytosis was also shown in Tg2576 and PSAPP mice after infusion with human umbilical cord blood cells, via suppression of CD40-CD40L interaction [56]. CD40L or CD40 deficiency also reduced pathological tau phosphorylation in Tg2576 mice [57].

Impairment of microglia recruitment to plaques by inducing a deficiency of CCR2, a microglial chemokine receptor [58], or the innate immune receptor toll-like receptor 2 (TLR2, [59]) increased A β aggregation and accelerated memory impairment, indicating that microglial chemotaxis is important in controlling A β deposition.

Liver x receptor (LXR) plays a dual role in controlling cholesterol homeostasis, while inhibiting inflammation. Loss of LXR in APP/PS1 mice leads to increased plaque load. In cell culture, LXR activation reduced the microglial inflammatory response to fibrillary Aβ and enhanced microglial phagocytosis [60]. Treatment of Tg2576 mice with the LXR

8

Inflammation in transgenic mouse models

agonist GW3965, reduced A β deposits by half, and was accompanied by reduction in the number of CD45+ microglia as well as interleukin (IL)-6 mRNA expression [61]. The authors suggest that LXR acts via increased expression of lipidated ApoE. Microglial activity can also be altered by modification of cytokine levels. Treatment of APP/PS1 mice with macrophage colony-stimulating factor (M-CSF) increased the number of microglial cells, as well as their phagocytosis of A β , and reduced the number of A β deposits [62]. Nichol and colleagues found that exercise induced a shift from proinflammatory to anti-inflammatory microglial activation in Tg2576 mice. Exercise resulted in reduced expression of the pro-inflammatory cytokines IL-1ß and tumor necrosis factor (TNF)- α , increased MHC-II levels (indicating a shift to an antigenpresenting state), and induced a trend towards reduction in levels of aggregated A β [63]. As reviewed earlier in this article, there is good evidence for an activation of the complement system in AD mouse models. Alterations in the level of complement components also modulate microglial activation. Microglial accumulation, C4 levels and neurodegeneration were reduced in C1q deficient Tg2576 and APP/PS1 mice, while C3 levels were increased and A β load remained unchanged [25, 26]. Extrinsic C1g can shift microglia into a pro-inflammatory activation state, as demonstrated recently in wildtype rat cell culture [64]. In a different set of experiments Maier et al. [28] showed that reducing levels of C3 led to increased microglia/macrophage CD45 levels while F4/80 and CD68 showed a trend toward reduction, suggesting a switch in phenotype of

microglia.

Butovsky and colleagues showed that T-cell-based immunization of APP/PS1 mice with glatimer acetate caused a phenotype switch in microglia inducing CD11c and insulin-like growth factor 1 expression as well as increased clearance of A β . This was also accompanied by increased neurogenesis [65].

Intracerebral injections of lipopolysaccharide (LPS) induced neuroinflammation associated with clearance of A β and microglial activation. Dexamethasone inhibited microglial activation and A β clearance [66].

Minocycline is a tetracycline with anti-inflammatory properties and reduces microglial activation. Although minocycline treatment did not alter A β deposition in SwDI mice, it reduced the number of MHC-II microglia and IL-6 expression as well as improving

Inflammation in transgenic mouse models

learning performance [67]. The studies of minocycline treatment and C1q deficiency show that the harmful effects of microglia can be ameliorated in mouse models independently of A β levels.

A β itself can influence the balance between pro- and anti-inflammatory activation. It drives microglia towards a pro-inflammatory state. While microglia of wild-type mice subjected to traumatic brain injury predominantly produce anti-inflammatory factors, microglia in APPswe mice have increased expression of pro-inflammatory genes [68]. Jimenez and colleagues [69] reported that in 6-month-old APP/PS1 mice, microglia are TNF- α negative and phagocytose A β in accordance with an anti-inflammatory phenotype, while microglia of 18-month old mice have switched to a pro-inflammatory phenotype with increased TNF- α and inducible nitric oxide synthase (iNOS) expression. The authors suggest that soluble oligomeric A β , which had increased 10-fold by 18 months, may be responsible for this switch. This phenotypic switch may be the murine equivalent of 'microglial senescence' leading to deficiency in phagocytosis and immune defense in aged human brains [70] or 'inflammaging' [71], a chronic up-regulation of the pro-inflammatory state associated with aging.

Resident brain microglia vs. peripheral, bone-marrow derived cells

An interesting question is whether infiltrating peripheral macrophages enter the brain and whether they take part in clearance of A β . It has been shown that in whole-body irradiated mice, peripheral macrophages cross the blood-brain barrier and differentiate into CD11c positive microglia. Such infiltrated cells are more efficient as phagocytes than the resident cells [72, 73]. Considerable infiltration of peripheral macrophages and accumulation near vessels and plaques was induced when transforming growth factor (TGF)- β -Smad2/3 signaling was interrupted in immune cells of Tg2576 mice, accompanied with a substantial reduction of A β deposits and improved learning [74]. The potential of bone-marrow derived cells for A β clearance was also demonstrated by Butovsky and colleagues [75], who specifically ablated these cells by irradiation in APP/ PS1 chimeric mice. The number of CD11c+ cells in brain was reduced and A β deposition increased. It is still a point of debate as to whether peripheral recruitment can occur without damage of the blood-brain-barrier, since much of the data are derived from

Inflammation in transgenic mouse models

experiments using whole-body irradiation [72, 75]. A different approach was used by Ajami and colleagues who created parabiotic mice with peripheral blood chimerism in which one of the chimeric partners expressed GFP in hematopoietic cells. When microgliosis was induced in the GFP-negative partner, none of the microglial cells was GFP-positive, suggesting that the microgliosis resulted from expansion of the local brain microglial pool [76].

Microglia are involved in vaccination-mediated clearance of AB

Both active and passive A β based vaccination protocols have been used successfully in AD mouse models to reduce the A β burden and to improve behavioral functions (for reviews see [77-79]). Encouraged by such positive results, a clinical trial in human patients was initiated, but was hampered by unwanted side effects, such as meningoencephalitis, stroke and even death [80, 81]. Subsequent clinical trials are aimed at avoidance of inflammation, complement activation and inappropriate T-cell responses. Several distinct mechanisms for vaccination results have been proposed but we include only studies with relevance to microglial activation and phagocytosis in this review. It has been proposed that one of the effects of A β vaccination is a shift in the state of microglia towards a phenotype with increased phagocytotic activity and limited production of inflammatory cytokines [82]. Wilcock and colleagues reported activation of CD45+ microglia after intracranial injection of anti-AB antibodies [83]. Antiinflammatory drugs reduced microglial activation and inhibited fibrillary AB clearance. This suggests that microglial activation is necessary for clearance of compact deposits. The same authors showed also that microglial activation (upregulated CD45) was transient after repeated vaccination of APP/PS1 mice with A β [84]. Using in vivo two-photon microscopy imaging in PDAPP mice with GFP labeled microglia, Koenigsknecht-Talboo and colleagues [85] showed that microglia migrate to A β plaques within 3 days after systemic administration of an anti-A β antibody (m3D6). Microglia did not react to a different antibody which recognizes only soluble A β (mHJ5) or to Fab fragments of m3D6. This indicates that antibodies against different AB epitopes have differing effects on microglia activation. Similar effects were also shown by Bard et

11

Inflammation in transgenic mouse models

al. and Gray et al. [86, 87] who found that antibodies against the A β N-terminus activated phagocytosis, while C-terminal antibodies acted by peripheral sequestration of A β . A β DNA vaccines were used as an alternate approach for immunotherapy and they have been shown to work effectively in mouse models. Okura and colleagues [88] reported that increased phagocytotic activity and numbers of microglia were the main drivers of A β clearance in their model (APP23), while inflammatory cytokine TNF- α expression remained unchanged. Mouri and coworkers used A β cDNA in an oral vaccination paradigm in Tg2576 mice and reported reduction in A β burden without lymphocytic infiltration [89].

Although most studies show involvement of microglial activation in vaccination-induced A β clearance, others suggest that microglia have only a limited role [90]. The number of microglial cells associated with plaques significantly increased when the anti-A β antibody 10D5 was directly applied into the brain of APP/PS1 mice and was further increased by IFN- γ . Inhibition of microglia with anti-Mac1-saporin or minocycline reduced the number of microglia. Effects of microglial inhibition or activation on A β clearance, in addition to 10D5 effects, were limited.

In summary (see also table 1), these studies demonstrate that in transgenic AD mouse models, microglia are activated, accumulate around A β deposits, and are involved in their clearance (although one study only shows a limited role [90]). It will be very interesting to see if further studies on modulating microglial activity towards alternate, antiinflammatory states can lead to an improved clearance of plaques and/or reduced neurodegeneration. It will also be important to search for possible differences between mouse and human microglial activation, such as NOS production and microglial senescence. In contrast to human microglia which do not produce measurable amounts of NOS, murine microglia produce large quantities when subjected to the same type of stimulation [91].

2.3. Cytokines in AD mouse models

Up- or down-regulation of the expression of various pro- and anti-inflammatory cytokines has been widely reported in AD patients (for reviews see: [7, 92, 93]). Similar

Inflammation in transgenic mouse models

changes in cytokine expression have been shown in transgenic mouse models of AD. The cytokines IFN- γ , IL-12, TNF- α , IL-6, IL-1 β , IL-1 α , TGF- β , and IL-10, monocyte chemoattractant protein (MCP)-1 and granulocyte-macrophage (GM)-CSF were reported to be up-regulated in various transgenic mouse models (Tg2576, TgAPPsw, PS1/APPsw, 3xTg-AD), while IL-4 was found to be down-regulated [94-98]. In addition, experimental manipulation of the cytokine system has been shown to affect the AD type pathology in these models. Disruption of the soluble TNF- α signaling with dominant negative TNF inhibitors (DN-TNFs) in the 3xTg-AD (APP/PS1/tau) model prevents intraneuronal A β accumulation [99]. Deletion of the TNF- α receptor TNFR1 in APP23 mice resulted in reduced plaque formation and limited neuronal loss [100]. Exercise was shown to reduce TNF- α and IL-1 β in Tg2576 mice to wild-type levels, accompanied by an increase in IFN- γ , macrophage inflammatory protein (MIP)-1 α , CD40, MHC-II, CD11c and reduced A β levels [63].

Suppression of IFN-y signaling in Tg2576 mice via IFN-y receptor 1 knockout reduced A β plaque load and gliosis [101]. The same authors showed that in glial cell cultures TNF- α and BACE1 expression was reduced in mutants, suggesting that this pathway is involved in Abeta formation. Microglial production of TNF- α , IL-1 β , IL-10 and IL-17 is up-regulated in APPswe mice, but this up-regulation was blocked by knock-out of the TLR-4 gene [102]. Shaftel and colleagues [103] demonstrated that in APPswe/PS1 mice, local overexpression of IL-1 β resulted in an inflammatory response and led to a reduction in A β deposition, possibly through enhanced microglial A β degradation. A moderate increase in astroglial TGF-B1 expression resulted in a reduction of AB deposition in hAPP mice (K670M/N671L and V717F) [104]. These findings seem inconsistent with the reported increased AB production in TGF-B1 transgenic mice (not transgenic for APP) [105]. Over-expression of CCL2 (MCP-1) in double transgenic Tg2576 mice increased glial activation and A β deposition [106], while gene delivery of a dominantnegative CCL2 mutant reduced gliosis and AB deposition in APP/PS1 mice [107]. In summary, inflammatory cytokines levels in transgenic mouse models reflect similar changes in AD. Their modulation regulates microglia activity in these models. These

Inflammation in transgenic mouse models

studies are summarized in table 1. It will be interesting to see if this can be translated into a useful therapeutic strategy in humans.

2.4. Efficiency of NSAIDs in AD mouse models

Evidence of chronic inflammation in the brain of AD patients, including upregulation of inflammatory cytokines, over-activation of microglia, and neurotoxic effects from the activated glia, suggest that NSAIDs may be useful in the treatment and prevention of AD. In addition to the pathological evidence for activation of the immune system, the majority of epidemiological studies have demonstrated that long-term exposure to certain NSAIDs reduce the risk of AD in humans. The large Rotterdam study demonstrated that long-term use of ibuprofen (two years and longer) reduced the relative risk of developing AD to 0.2 (95% CI 0.05–0.83) [108]. Incidence of AD was also reduced in users of NSAIDs (ibuprofen, naproxen and indomethacin) in the Baltimore longitudinal study of aging [109]. These and other epidemiological studies are reviewed in [110]. Although some newer studies found no benefits of NSAIDs, another recent study using records of the Veterans Affairs Health Care System showed that long-term NSAID use protected against AD [111].

With such epidemiological rationale, several trials in transgenic mouse models of AD have been conducted to assess the effectiveness of NSAIDs. Ibuprofen, the most commonly used NSAID in humans, was successfully utilized in the majority of transgenic studies. Feeding 10 month-old Tg2576 mice with chow mixed with ibuprofen for 6 months reduced the number and area of A β plaques, as well as IL-1 β and glial fibrillary acidic protein levels [112]. Yan et al. [113] reported a similar reduction of plaque load (60%) in Tg2576 when ibuprofen treatment was started at 11 months of age. But they found no effect of pioglitazone (a peroxisome proliferator-activated receptors (PPAR) γ agonist). In double-transgenic mice (APP/PS1), which develop plaques at an earlier age, Jantzen et al. [114] found that a nitric oxide-releasing derivative of flurbiprofen (NCX-2216) reduced the A β load dramatically, accompanied by an increase in the number of MHC-II expressing microglia. Ibuprofen reduced the A β load moderately, and celecoxib (a cyclooxygenase (COX)-2-selective NSAID) had no effect.

Inflammation in transgenic mouse models

A different nitric oxide-releasing derivative of flurbiprofen (HCT 1026) also reduced the $A\beta$ load and microglial activation near plaques [115].

R-flurbiprofen, which is not a COX inhibitor, was found to be effective in improving spatial learning deficits in Tg2576 if administered to younger mice [116]. If the drug was used after plaque development ensued, the plaque load was reduced but the learning deficit was not ameliorated. When treating 10 month-old APP V1711 transgenic mice with pioglitazone (at a higher dose than [113]) and ibuprofen for only 7 days, Heneka et al. [117] found a reduced number of activated microglia and reactive astrocytes. decreased expression of COX-2, iNOS and BACE1, and reduction of AB deposits and soluble A β 42. Also, in the Tg2576 mouse model, Morihara et al.[118] reported that ibuprofen reduced IL-1 β and the pro-amyloidogenic α 1-antichymotrypsin (ACT) expression. They suggested that ibuprofen reduced the A β burden via this pathway. In a recent study on triple transgenic mice (APPswe, PS1M146V, tauP301L), ibuprofen reduced intraneuronal oligometric A β , cognitive deficits, and hyperphosphorylated tau immunoreactivity [119]. But when ibuprofen was given to 3 month-old Tg2576 mice for a very short duration (4 days), no effect on A β levels was observed [120]. Indomethacin and a lecithin derivative (DP-155) were shown to be effective in reducing A β levels in Tg2576 mice [121-123]. Drugs which inhibit COX-2 did not reduce Aβ burden or even increased A β levels in mouse models [114, 122, 124].

The proposed mechanisms of action for NSAIDs include reduction of inflammation via COX inhibition [125, 126] or inhibition of cytokine expression (IL-1 β , α 1-ACT, [118]). In addition, certain NSAIDs have been shown at high concentration to reduce A β production by modulation of γ -secretase activity [127]. Some also act as agonists of PPAR γ and by modulation of Rho-GTPase (reviewed in [128]).

Despite the epidemiological evidence and successful trials with NSAIDs in transgenic mice, the outcomes of clinical trials in humans have been disappointing and have, in some cases, been accompanied by cardiovascular or gastrointestinal side effects. Several COX-2-selective NSAIDs (Nimesulide, Celecoxib, Rofecoxib) and non-selective NSAIDs (indomethacin, diclofenac, naproxen, ibuprofen) have been used in trials, but benefits were either very small or absent (for review see [110, 128]).

Inflammation in transgenic mouse models

3. TRANSGENIC MOUSE MODELS OF PARKINSON'S DISEASE

The 1988 report describing reactive microglia within the substantia nigra of PD patients was the first line of evidence to support the involvement of the immune system in PD [129]. Since then there have been numerous studies showing that inflammatory processes play an important role in the pathogenesis of PD, and several recent reviews have summarized the current state of our understanding of neuroinflammatory processes in PD [4, 130-132]. Similarly to AD, the evidence stems from observations in human postmortem tissue of reactive microglia and astrocytes in the affected areas. Such findings were supported by genetic studies identifying correlations between polymorphisms of genes associated with inflammatory responses and the risk of PD. They were further supported by epidemiological data showing protective effects of chronic use of anti-inflammatory drugs and by functional animal studies showing that activation of the inflammatory response in the central nervous system may lead to selective loss of dopaminergic neurons and that anti-inflammatory treatments ameliorate neuronal and behavioural deficits observed in these animals.

Additional support for the inflammatory hypothesis in the case of PD comes from the investigation of an accidental human model of the disease. In 1983 Langston et al. reported that drug addicts who consumed a synthetic product containing the previously undiscovered neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) developed symptoms that were very similar to idiopathic PD [133]. A dissimilarity was noted in the pathological examination at autopsy since these patients did not develop Lewy bodies. Nevertheless, there was a selective loss of dopaminergic neurons and continued presence of activated microglia. In one case, this sign of inflammation was seen as late as 16 years after the exposure to the agent [134]. These observations were supported by later studies on MPTP-treated monkeys that showed both microglial and astroglial reactions up to 14 years after the experimental treatment [135, 136]. The presence of reactive immune cells years after the initial toxic insult indicates that inflammatory processes in these cases, and possibly also in idiopathic PD, may evolve into self-sustaining mechanisms that are damaging to selective groups of neurons. A number of functional animal models of PD that use various toxins (e.g. 6-OH-dopamine,

Inflammation in transgenic mouse models

MPTP, rotenone) or exogenous immune mediators (e.g. LPS) have demonstrated robust activation of neuroimmune mechanisms that are accompanied by neuronal death in the substantia nigra. Furthermore, several of these models rely on immune activation of glial cells as the trigger of selective loss of dopaminergic neurons (reviewed in [41, 131, 137]). In transgenic animal models of PD, the identification of inflammatory mechanisms has proven to be more challenging. In similarity to AD, transgenic animal models of PD have been created based on knowledge of gene mutations associated with the human disease. They include α-synuclein (Asyn), DJ-1, LRRK2, Parkin, PINK1 and UCH-L1 [138-141]. Thus far transgenic PD models have been created in three different species: mice, as well as Drosophila melanogaster and Caenorhabditis elegans. Only in the latter two species has the incorporation of Asyn mutations resulted in a consistent loss of dopaminergic neurons. This illustrates that genetic manipulation alone may not be sufficient to trigger the pathogenesis of PD and that additional environmental factors are needed [137]. Even though the genetically modified animals display certain pathophysiological and behavioural characteristics of PD, there have been very few reports describing neuroimmune responses associated with the genetic manipulations. A scarcity of pertinent information does not rule out the possibility that immune responses are evoked by genetic manipulations. Future experiments in transgenic animals may establish such correlations. Below we review the evidence obtained thus far from the various groups of transgenic animal models of PD.

3.1 Alpha-synuclein transgenic mouse models

This is the most studied group of animals, perhaps due to the fact that alpha-synuclein (Asyn, PARK 1) was the first gene to be linked to PD. Asyn has also become the most studied PD-associated protein. Three different mutations (A30P, A53T and E46K) as well as genomic duplication and triplication of a locus containing the Asyn gene have been linked to familial forms of PD [142-145]. The first Asyn transgenic mice were created by Masliah and colleagues by expressing wild-type human Asyn in neuronal cells under the regulatory control of the platelet-derived growth factor–b (PDGF-b) promoter [146]. This resulted in progressive accumulation of Asyn and ubiquitin-immunoreactive inclusions in neurons in the neocortex, hippocampus and substantia nigra. Although loss

Inflammation in transgenic mouse models

of dopaminergic terminals in the basal ganglia and motor impairments were observed, there was no substantial decrease in the number of dopaminergic neurons and no glial activation was reported [146]. Since then, pan-neuronal expression of Asyn has been achieved by using PDGF-b, Thy1 or prion protein promoters. In contrast the tyrosine hydroxylase (TH) promoter has been used to restrict Asyn expression to catecholaminergic neurons.

Even though some of the transgenic models mimic certain stages or features of human PD, none of them display the full pathological cascade of PD (reviewed in [147, 148]). While neurodegeneration is absent in some cases, it is non-specific or of late onset in others. Similarly, the evidence of immune system involvement has not been consistent and depends on the type of Asyn and the promoter used.

Thy1 is another promoter that has been used in addition to PDGF-b to achieve broad expression of Asyn in mouse neurons. The Thy1 transgenic mouse lines have a wide range of phenotypes with different patterns of Asyn expression [147]. Activated microglia associated with Asyn inclusions have been observed in these mice [138]. However this upregulation was not confirmed in the same mice after paraquat injection, which may have been due to the fact that microglial activation occurs concomitant to neuronal death and is not detectable 2 weeks after the onset of neuronal degeneration [149]. Lewy-like pathological changes in neurons were reported in mice expressing the Thy1-driven A53T mutant form of human Asyn. Astrocytic gliosis and microglial activation accompanied motor neuron pathology, which was also seen in other affected brain regions [150].

Giasson and colleagues reported that pan-neuronal overexpression of A53T Asyn leads to significant astrogliosis in the spinal cord, which was interpreted as being consistent with neuronal injury. These mice displayed motor impairments although no significant loss of motor neurons was observed [151]. A more severe phenotype of A53T Asyn mice was described by Lee et al. [152] who reported adult-onset neurodegeneration leading to progressive motor dysfunction and eventual death of the animals. A prominent astroglial reaction was observed in areas affected by neurodegenerative changes. Interestingly, mice expressing the wild-type or A30P variant of Asyn did not develop such pathology [152]. The above two studies used the mouse prion protein promoter which generally

Inflammation in transgenic mouse models

leads to high levels of Asyn protein expression. Despite this fact, not all mice expressing A53T even under the control of the prion promoter displayed a glial reaction. Negative results have also been reported [153].

High levels of A30P Asyn expression in the Tg5093 mouse line was associated with a progressive motor disorder characterized by rigidity, dystonia, gait impairment, and tremor. Even though these mice did not exhibit specific deterioration of the nigrostriatal dopaminergic neurons, they did display neuropathological features including age-related central nervous system astrocytosis and microgliosis. Gliosis was evident in the hippocampal formation at the age of 7–8 months in heterozygote mice and at 4 months in the homozygote mice [154].

In a more recent study, Su et al. (2008) characterized inflammatory responses in mice expressing human wild-type Asyn under control of the rat TH promoter [155]. Asyn expressing (SynWT+/+) animals exhibited a significant increase in the number of activated microglia in the substantia nigra at one month compared to age-matched control animals. They also had significantly higher levels of expression of the pro-inflammatory cytokine tumor necrosis factor (TNF)- α . Similarly, expression of truncated human Asyn(1–120), driven by the rat TH promoter, led to the formation of pathological inclusions in the substantia nigra; even though no significant cell loss was observed, an increase in microglial cells was evident [156]. A lack of microgliosis has also been reported in mice expressing a truncated (1-130) form of Asyn with the A53T mutation [157].

Transgenic mice with oligodendrocyte-specific expression of wild-type Asyn were created to model multiple system atrophy (MSA) since Asyn-positive glial inclusions are a major histopathological hallmark of idiopathic MSA [158]. These mice showed early and progressive microglial activation in the substantia nigra pars compacta (SNc), which was associated with increased expression of iNOS and loss of dopaminergic neurons. Suppression of microglial activation by minocycline protected dopaminergic SNc neurons [158]. Transgenic mice overexpressing human Asyn have also been used to perform functional animal experiments by using various toxins and immune mediators to trigger neurodegeneration. For example, Gao et al. (2008) studied the effects of bacterial LPS injection in mice that expressed human wild-type or mutant A53T Asyn in the

Inflammation in transgenic mouse models

absence of murine Asyn (Asyn-null background). Stereotaxic injection of LPS into the substantia nigra caused a similar inflammatory reaction in all animals studied. However, loss of dopaminergic neurons and aggregates of Asyn were observed in mice overexpressing wild-type and mutant Asyn but not in Asyn-null control mice [159]. Furthermore, ex vivo glial cultures or mixed neuron/glial cultures from transgenic animals overexpressing or lacking Asyn have been used to demonstrate differences in microglial phenotype and their ability to produce pro-inflammatory and toxic species [159, 160]. Thus microglia from Asyn knock-out animals displayed a reactive phenotype and increased cytokine secretion, but impaired phagocytic ability [160]. This indicates that Asyn plays a major role in regulating the microglial activation state. Human Asyn overexpressing mice were also used to study the effects of immunization with Asyn. Since immunization has been shown to reduce the amyloid burden and reduce the pathology in a number of AD transgenic mouse lines, Masliah et al. (2005) vaccinated human Asyn overexpressing transgenic mice with human Asyn. In mice that produced high affinity antibodies against Asyn there was a decreased accumulation of Asyn aggregates and reduced neurodegeneration [161]. Even though vaccination did not induce a significant glial reaction, these experiments clearly demonstrate involvement of the immune system in regulating Asyn aggregation and its associated neurodegenerative pathology.

In summary, due to the various Asyn forms, as well as different promoters and strains of animal used, there have been significant inconsistencies between various studies. It appears that glial reactions typically accompany neuronal cell death, if and when it occurs. However, several reports have described glial activation in the absence of significant neuronal death, which may indicate that Asyn is a signaling molecule released not only from dying neurons, but also from neurons that are functionally compromised or are just simply overexpressing Asyn. Support for this hypothesis comes from a study demonstrating that neurons overexpressing Asyn secrete high concentrations of Asyn which are capable of activating microglia [155], and from in vitro observations that extracellular Asyn can be recycled by neurons and is an effective pro-inflammatory stimulant of both microglia and astrocytes [162-166]. Alternatively, the formation of

Inflammation in transgenic mouse models

neuronal inclusions, the development of neuronal deficits, and neuroimmune responses, may not be linked in a causal chain.

3.2. DJ-1, Parkin and PINK transgenic mouse models

Mutations in DJ-1, PARK and PINK genes produce rare inherited forms of PD. DJ-1 at the PARK7 locus has been linked to multiple cellular processes including oxidative stress responses. At least three different lines of DJ-1 knockout mice have been created, which show decreased motor functions without clear loss of dopaminergic neurons [139, 167-169]. None of these studies reported inflammatory responses in DJ-1 knockout mice. PINK1 (PARK6) is believed to play an important role in mitochondrial function, while Parkin (PARK2) is important in protein targeting for proteosomal degradation. PINK1defficient and Parkin (PARK2) knockout mice do not exhibit loss of dopaminergic neurons and alterations in inflammatory markers have not been reported [139, 170-178]. Recently, Frank-Cannon et al. (2008) reported that long-term intraperitoneal administration of LPS induced neuroinflammation, which was accompanied by fine motor deficits and loss of dopaminergic neurons in Parkin-deficient (parkin -/-) mice, but not in wild-type animals [170]. Meanwhile Thomas et al. 2007 reported that nigrostriatal dopaminergic neurons in Parkin knockout mice did not show increased susceptibility to the parkinsonian neurotoxin, MPTP, which was administered in various doses [179]. There are a number of other transgenic mouse models that resemble human PD that still have not been characterized with regard to immune system involvement. One example are MitoPark mice, which were created by conditional knockout of the gene for mitochondrial transcription factor A (Tfam) in dopaminergic neurons. These mice have reduced mitochondrial DNA (mtDNA) expression and respiratory chain deficiency in midbrain DA neurons. This leads to adult onset formation of intraneuronal Asyn negative inclusions, dopaminergic cell death, and associated progressive impairment of motor function [180].

Other transgenic mouse models involve genes that do not have identified mutations associated with PD, but which nevertheless may be involved in the pathogenesis of this disease. Thus, the naturally occurring murine weaver mutant (gene symbol, wv) carries a mutation in the gene encoding the G-protein which inwardly rectifies the potassium

21

Inflammation in transgenic mouse models

channel Girk2. It has been reported that in these mice the degeneration of midbrain dopaminergic neurons, unlike other types of neurons, is mediated via neuroinflammatory mechanisms [181].

Mallajosyula et al. (2008) reported the creation of inducible monoamine oxidase B (MAO-B) transgenic mice. Astrocytic MAO-B levels were specifically induced in adult animals, which caused selective and progressive loss of dopaminergic neurons in the substantia nigra. Since local microglial activation was observed within the substantia nigra in addition to global astrogliosis, these mice may provide a new model for the investigation of the neuroinflammatory features of PD [182].

Transgenic mice deficient in various key signaling molecules have been widely used to investigate the role of specific molecular pathways in neuroinflammatory processes associated with PD pathology. Examples include mice deficient in MAPK-activated protein kinase 2 (MK2) [183], complement C3 [184], protease-activated receptor 1 (PAR1) [185], IL-1 α/β [186], MCP-1/CC chemokine ligand 2 (Mcp-1/Ccl2) [187], (TNF)- α receptors R1 and R2 [188] and prostaglandin E₂ receptor subtype 2 (EP2) [189]. Transgenic models of PD have also been created by using invertebrates, namely C. elegans and Drosophila. While these species may offer certain advantages, such as the absence of endogenous Asyn which allows for the study of this protein in naïve organisms [190], the obvious disadvantage in using any fly or worm model is the potential risk that a critical factor in the pathology of the diseases is specific to mammals in general or humans in particular [191]. Immune reactions may be an example of a mammal-specific mechanism, which could explain why we were unable to identify any publications describing glial or other immune activation in the two invertebrate models of PD.

4. TRANSGENIC MOUSE MODELS OF HUNTINGTON'S DISEASE

Huntington's disease is a hereditary polyglutamine disorder caused by extension of the triplet repeat region of the huntingtin (htt) gene. HD is characterized by severe atrophy of the striatum, where up to 95% of the medium spiny neurons degenerate, as well as by the presence of intracellular and intranuclear htt inclusions, and by reactive gliosis.

Inflammation in transgenic mouse models

Evidence for involvement of inflammation in the pathogenesis of HD includes reports on activation of the complement system, increased microglial activity and up-regulation of inflammatory cytokines. Singhrao and colleagues found complement activation and increased mRNA levels for complement proteins and regulators (C1q C chain, C1r, C4, C3, C1 inhibitor, clusterin, membrane cofactor protein, decay accelerating factor, CD59) in HD brain tissue [192]. Increased protein expression of complement components C7 and C9, complement inhibitor clusterin and acute phase protein α -2-macroglobulin was shown in HD plasma and cerebrospinal fluid [193]. Activated microglia expressing HLA-DR were reported in affected areas in HD [194]. These microglial cells were found in close association with degenerating neurons [195]. The peripheral benzodiazepine receptor is selectively expressed by brain microglia. Increased binding of its ligand PK11195 has been described in postmortem HD putamen and frontal cortex [196]. Using PK11195 for positron emission tomography (PET) imaging, increased microglial activation was demonstrated in HD patients and in presymptomatic carriers of the mutated htt gene, indicating that microglial activation is an early event in HD pathogenesis [197, 198]. Microglial activation correlated inversely with raclopride binding, a marker of striatal cell function [199].

Data on microglial activation in HD mouse models is relatively sparse. In both HD patients and R6/2 mice, expression of ferritin in microglia is increased and the microglial cells appear dystrophic [200]. But unlike the human HD cases, microglial size and density is reduced with age in R6/2 mice [201].

Upregulation of inflammatory and anti-inflammatory cytokines has been reported in the plasma (IL-6, IL-8, IL-4, IL-10, TNF- α) and striatum (IL-6, IL-8 and TNF- α) of HD patients [193, 202]. Interestingly, IL-6 plasma levels were found to be increased in presymptomatic HD mutation carriers 16 years prior to the predicted onset of the disease, suggesting that inflammatory changes commence very early in the disease process [202]. HD monocytes showed increased IL-6 expression when activated by LPS and IFN- γ . R6/2 microglia and YAC128 macrophages were similarly hyperactivated and protein expression of IL-6, IL-10, IL-1 β or IL-12p70 was increased in R6/2, Hdh 150Q/150Q, and to a lesser extent in YAC128 mouse models. Less severe and more localized pathology was also observed in the YAC128 model (for review see [203]). In contrast to

Inflammation in transgenic mouse models

these studies, Godavarthi and colleagues found no differences in cytokine expression (MCP-1, IL-6, IL-10, TNF- α , interferon- γ , IL-12) and microglia numbers in 12 week old R6/2 mice, at an age when these mice show severe symptoms [204]. The anti-inflammatory tetracycline minocycline has been shown to improve behavioral and neuropathological deficits in R6/2 mice [205, 206] and was also protective in an excitotoxic rat model of HD [207]. In a different excitotoxic model of HD (3-nitropropionic acid) treatment with C5a receptor antagonists reduced neuropathology, behavioural and motor deficits [208].

Abnormalities in the kynurenic acid pathway are an interesting aspect of HD pathology. Quinolinic acid and 3-hydroxykynurenine are neurotoxic metabolites that are elevated in HD and in the R6/2, Hdh 150Q/150Q and YAC128 mouse models. The kynurenine pathway is mainly localized in glial cells, and it is possible that activation of microglia and astroglia may result in production of these excitotoxic metabolites before morphological markers of glial activation are apparent [209-211]. The data on activation of the inflammatory system in transgenic HD mouse models is sparse and contradictory, possibly due to considerable differences in the pathology between different transgenic strains, but also due to the paucity of studies. However, data in HD patients shows that the inflammatory system is activated very early and thus may be an important therapeutic target.

5. SUMMARY

As stated in the introduction, the perfect animal model of a human disease should reflect the symptoms, lesions and causes of the disease [1]. Our review explored how transgenic mouse models of AD, PD and HD matched these requirements in the particular area of neuroinflammation. Possible mechanisms involving inflammatory responses in transgenic mouse models are shown in figure 2.

Mouse models of AD have been available for a number of years and have been studied widely (for recent reviews comparing mouse models and AD in general see: [1, 8, 9]). These transgenic mice develop symptoms comparable to AD (memory and behavioral deficits). They build up A β deposits, one of the hallmarks of AD, but lack two others; NFTs and overt neuronal death. Since APP and PS1 mutations do not cause sporadic AD

Inflammation in transgenic mouse models

(the majority of AD cases), models with these mutations reflect the cause of disease only for the subset of familial cases. Evidence is readily available that A β deposition in AD mouse models is accompanied by glial accumulation and activation, and, to variable degrees, modulation of the levels of inflammatory cytokines and complement activation. This inflammatory response is incomplete and less severe compared to AD. There are several possible explanations: (i) neurodegenerative diseases in humans take many years or even decades to develop; mice may simply not get old enough to develop full-blown inflammation; (ii) in most models human genes are introduced and human proteins are expressed. Interactions with the mouse proteome may not reflect the situation in humans. (iii) Different mechanisms of repair and compensation may exist in long-lived organisms such as humans, and interspecies differences in the complement system have been reported [34, 35].

Similarly to AD, only a fraction of PD cases is caused by mutations. These mutations have been used to create mouse models. Compared with AD transgenic models, the available PD models show less behavioral deficits and pathological changes. Only little evidence for activation of the immune system is available. This may in part be due to the fact that these models have been developed recently and only a few studies have been conducted to investigate the involvement of inflammation in the disease process. Since HD is a hereditary disorder, models created using genes of extended triplet repeats mirror the cause of the disease in humans. Introduction of segments or the complete HD gene causes a range of pathology from wide-spread, non-specific degeneration to mild disease [203]. Again only little and conflicting information on the involvement of inflammation in these mice can be found in the literature.

The seeming lack of inflammatory changes in PD and HD mouse models may be due to (i) the fact that the aggregated proteins in AD are deposited extracellularly, while in HD (and in some models of PD), pathological proteins aggregate intracellularly; (ii) that inflammatory processes may play a lesser role in the mouse models and/or in human PD and HD; or (iii) the current lack of studies focused on the involvement of inflammation in these models.

In conclusion, transgenic models have been proven to be useful tools in the exploration of molecular mechanisms of neurodegenerative disorders and in the discovery of potential

Inflammation in transgenic mouse models

therapeutic strategies. Particularly in AD these transgenic models represent an improvement over previous toxic or lesion models. Especially interesting are data in the field of in-vivo multi photon imaging, differential microglial activation and the promising results of vaccination studies. The latter example also demonstrates the importance of careful scrutiny when translating results from mouse models into clinical application. Like most models, the available transgenic mouse strains are imperfect models of human neurodegenerative diseases. However, further studies will lead to a continuously improving match with human disorders for the development of useful therapeutic strategies. Careful comparison with data from human pathology and studies on other organisms, such as dogs or non-human primates [212] will be the most useful additions to mouse models. The recent report describing creation of the first transgenic nonhuman primate line [213] will hopefully encourage further efforts into the development of new transgenic animal models of neurodegenerative disease that more closely resemble human pathology, including neuroimmune responses.

Acknowledgements:

This research was supported by the Jack Brown and Family AD Research Foundation, the Pacific Alzheimer Research Foundation and the Natural Sciences and Engineering Research Council of Canada.

References:

- [1] C. Duyckaerts, M.C. Potier, B. Delatour, Alzheimer disease models and human neuropathology: similarities and differences, Acta Neuropathol 115 (2008) 5-38.
- [2] A.J. Rozemuller, W.A. van Gool, P. Eikelenboom, The neuroinflammatory response in plaques and amyloid angiopathy in Alzheimer's disease: therapeutic implications, Curr Drug Targets CNS Neurol Disord 4 (2005) 223-233.
- [3] E.G. McGeer, A. Klegeris, P.L. McGeer, Inflammation, the complement system and the diseases of aging, Neurobiol Aging 26 Suppl 1 (2005) 94-97.
- [4] A. Klegeris, E.G. McGeer, P.L. McGeer, Therapeutic approaches to inflammation in neurodegenerative disease, Curr Opin Neurol 20 (2007) 351-357.
- [5] I. Glezer, A.R. Simard, S. Rivest, Neuroprotective role of the innate immune system by microglia, Neuroscience 147 (2007) 867-883.

- [6] M. Griffiths, J.W. Neal, P. Gasque, Innate immunity and protective neuroinflammation: new emphasis on the role of neuroimmune regulatory proteins, Int Rev Neurobiol 82 (2007) 29-55.
- [7] C. Schwab, P.L. McGeer, Inflammatory aspects of Alzheimer disease and other neurodegenerative disorders, J Alzheimers Dis 13 (2008) 359-369.
- [8] D.A. Morrissette, A. Parachikova, K.N. Green, F.M. LaFerla, Relevance of transgenic mouse models to human Alzheimer disease, J Biol Chem 284 (2009) 6033-6037.
- [9] D.R. Howlett, J.C. Richardson, The pathology of APP transgenic mice: a model of Alzheimer's disease or simply overexpression of APP?, Histol Histopathol 24 (2009) 83-100.
- [10] S. Oddo, A. Caccamo, J.D. Shepherd, M.P. Murphy, T.E. Golde, R. Kayed, R. Metherate, M.P. Mattson, Y. Akbari, F.M. LaFerla, Triple-transgenic model of Alzheimer's disease with plaques and tangles: intracellular Abeta and synaptic dysfunction, Neuron 39 (2003) 409-421.
- [11] A.P. Sjoberg, L.A. Trouw, A.M. Blom, Complement activation and inhibition: a delicate balance, Trends Immunol 30 (2009) 83-90.
- [12] P.L. McGeer, E.G. McGeer, Complement proteins and complement inhibitors in Alzheimer's disease, Res Immunol 143 (1992) 621-624.
- [13] J. Rogers, J. Schultz, L. Brachova, L.F. Lue, S. Webster, B. Bradt, N.R. Cooper, D.E. Moss, Complement activation and beta-amyloid-mediated neurotoxicity in Alzheimer's disease, Res Immunol 143 (1992) 624-630.
- [14] H. Jiang, D. Burdick, C.G. Glabe, C.W. Cotman, A.J. Tenner, beta-Amyloid activates complement by binding to a specific region of the collagen-like domain of the C1q A chain, J Immunol 152 (1994) 5050-5059.
- [15] P. Tacnet-Delorme, S. Chevallier, G.J. Arlaud, Beta-amyloid fibrils activate the C1 complex of complement under physiological conditions: evidence for a binding site for A beta on the C1q globular regions, J Immunol 167 (2001) 6374-6381.
- [16] B.M. Bradt, W.P. Kolb, N.R. Cooper, Complement-dependent proinflammatory properties of the Alzheimer's disease beta-peptide, J Exp Med 188 (1998) 431-438.
- [17] D.A. Loeffler, D.M. Camp, D.A. Bennett, Plaque complement activation and cognitive loss in Alzheimer's disease, J Neuroinflammation 5 (2008) 9.
- [18] K. Yasojima, C. Schwab, E.G. McGeer, P.L. McGeer, Up-regulated production and activation of the complement system in Alzheimer's disease brain, Am J Pathol 154 (1999) 927-936.
- [19] H. Zanjani, C.E. Finch, C. Kemper, J. Atkinson, D. McKeel, J.C. Morris, J.L. Price, Complement activation in very early Alzheimer disease, Alzheimer Dis Assoc Disord 19 (2005) 55-66.
- [20] L.A. Trouw, H.M. Nielsen, L. Minthon, E. Londos, G. Landberg, R. Veerhuis, S. Janciauskiene, A.M. Blom, C4b-binding protein in Alzheimer's disease: binding to Abeta1-42 and to dead cells, Mol Immunol 45 (2008) 3649-3660.
- [21] K. Yasojima, E.G. McGeer, P.L. McGeer, Complement regulators C1 inhibitor and CD59 do not significantly inhibit complement activation in Alzheimer disease, Brain Res 833 (1999) 297-301.

- [22] R. Fan, K. DeFilippis, W.E. Van Nostrand, Induction of complement proteins in a mouse model for cerebral microvascular A beta deposition, J Neuroinflammation 4 (2007) 22.
- [23] Y. Matsuoka, M. Picciano, B. Malester, J. LaFrancois, C. Zehr, J.M. Daeschner, J.A. Olschowka, M.I. Fonseca, M.K. O'Banion, A.J. Tenner, C.A. Lemere, K. Duff, Inflammatory responses to amyloidosis in a transgenic mouse model of Alzheimer's disease, Am J Pathol 158 (2001) 1345-1354.
- [24] C. Schwab, M. Hosokawa, P.L. McGeer, Transgenic mice overexpressing amyloid beta protein are an incomplete model of Alzheimer disease, Exp Neurol 188 (2004) 52-64.
- [25] J. Zhou, M.I. Fonseca, K. Pisalyaput, A.J. Tenner, Complement C3 and C4 expression in C1q sufficient and deficient mouse models of Alzheimer's disease, J Neurochem 106 (2008) 2080-2092.
- [26] M.I. Fonseca, J. Zhou, M. Botto, A.J. Tenner, Absence of C1q leads to less neuropathology in transgenic mouse models of Alzheimer's disease, J Neurosci 24 (2004) 6457-6465.
- [27] M.I. Fonseca, R.R. Ager, S.H. Chu, O. Yazan, S.D. Sanderson, F.M. LaFerla, S.M. Taylor, T.M. Woodruff, A.J. Tenner, Treatment with a C5aR antagonist decreases pathology and enhances behavioral performance in murine models of Alzheimer's disease, J Immunol 183 (2009) 1375-1383.
- [28] M. Maier, Y. Peng, L. Jiang, T.J. Seabrook, M.C. Carroll, C.A. Lemere, Complement C3 deficiency leads to accelerated amyloid beta plaque deposition and neurodegeneration and modulation of the microglia/macrophage phenotype in amyloid precursor protein transgenic mice, J Neurosci 28 (2008) 6333-6341.
- [29] T. Wyss-Coray, F. Yan, A.H. Lin, J.D. Lambris, J.J. Alexander, R.J. Quigg, E. Masliah, Prominent neurodegeneration and increased plaque formation in complement-inhibited Alzheimer's mice, Proc Natl Acad Sci U S A 99 (2002) 10837-10842.
- [30] N.S. Pillay, L.A. Kellaway, G.J. Kotwal, Early detection of memory deficits and memory improvement with vaccinia virus complement control protein in an Alzheimer's disease model, Behav Brain Res 192 (2008) 173-177.
- [31] D.A. Loeffler, Using animal models to determine the significance of complement activation in Alzheimer's disease, J Neuroinflammation 1 (2004) 18.
- [32] A.J. Tenner, M.I. Fonseca, The double-edged flower: roles of complement protein C1q in neurodegenerative diseases, Adv Exp Med Biol 586 (2006) 153-176.
- [33] S.D. Webster, A.J. Tenner, T.L. Poulos, D.H. Cribbs, The mouse C1q A-chain sequence alters beta-amyloid-induced complement activation, Neurobiol Aging 20 (1999) 297-304.
- [34] G.L. Ong, M.J. Mattes, Mouse strains with typical mammalian levels of complement activity, J Immunol Methods 125 (1989) 147-158.
- [35] R.O. Ebanks, D.E. Isenman, Mouse complement component C4 is devoid of classical pathway C5 convertase subunit activity, Mol Immunol 33 (1996) 297-309.
- [36] E.G. McGeer, P.L. McGeer, Brain inflammation in Alzheimer disease and the therapeutic implications, Curr Pharm Des 5 (1999) 821-836.

- [37] W.J. Streit, R.E. Mrak, W.S. Griffin, Microglia and neuroinflammation: a pathological perspective, J Neuroinflammation 1 (2004) 14.
- [38] D.W. Dickson, S.C. Lee, L.A. Mattiace, S.H. Yen, C. Brosnan, Microglia and cytokines in neurological disease, with special reference to AIDS and Alzheimer's disease, Glia 7 (1993) 75-83.
- [39] A. Klegeris, P.L. McGeer, A. Klegeris, P.L. McGeer, beta-amyloid protein enhances macrophage production of oxygen free radicals and glutamate, Journal of Neuroscience Research 49 (1997) 229-235.
- [40] S.L. Yates, L.H. Burgess, J. Kocsis-Angle, J.M. Antal, M.D. Dority, P.B. Embury, A.M. Piotrkowski, K.R. Brunden, Amyloid beta and amylin fibrils induce increases in proinflammatory cytokine and chemokine production by THP-1 cells and murine microglia, J Neurochem 74 (2000) 1017-1025.
- [41] M.L. Block, L. Zecca, J.S. Hong, Microglia-mediated neurotoxicity: uncovering the molecular mechanisms, Nat Rev Neurosci 8 (2007) 57-69.
- [42] U.K. Hanisch, H. Kettenmann, Microglia: active sensor and versatile effector cells in the normal and pathologic brain, Nat Neurosci 10 (2007) 1387-1394.
- [43] S.A. Frautschy, F. Yang, M. Irrizarry, B. Hyman, T.C. Saido, K. Hsiao, G.M. Cole, Microglial response to amyloid plaques in APPsw transgenic mice, Am J Pathol 152 (1998) 307-317.
- [44] C. Sturchler-Pierrat, D. Abramowski, M. Duke, K.H. Wiederhold, C. Mistl, S. Rothacher, B. Ledermann, K. Burki, P. Frey, P.A. Paganetti, C. Waridel, M.E. Calhoun, M. Jucker, A. Probst, M. Staufenbiel, B. Sommer, Two amyloid precursor protein transgenic mouse models with Alzheimer disease-like pathology, Proc Natl Acad Sci U S A 94 (1997) 13287-13292.
- [45] B.J. Hock, Jr., B.T. Lamb, Transgenic mouse models of Alzheimer's disease, Trends Genet 17 (2001) S7-12.
- [46] S. Dudal, P. Krzywkowski, J. Paquette, C. Morissette, D. Lacombe, P. Tremblay, F. Gervais, Inflammation occurs early during the Abeta deposition process in TgCRND8 mice, Neurobiol Aging 25 (2004) 861-871.
- [47] D.W. Dickson, Microglia in Alzheimer's disease and transgenic models. How close the fit?, Am J Pathol 154 (1999) 1627-1631.
- [48] Y. Yoshiyama, M. Higuchi, B. Zhang, S.M. Huang, N. Iwata, T.C. Saido, J. Maeda, T. Suhara, J.Q. Trojanowski, V.M. Lee, Synapse loss and microglial activation precede tangles in a P301S tauopathy mouse model, Neuron 53 (2007) 337-351.
- [49] D.R. Howlett, K. Bowler, P.E. Soden, D. Riddell, J.B. Davis, J.C. Richardson, S.A. Burbidge, M.I. Gonzalez, E.A. Irving, A. Lawman, G. Miglio, E.L. Dawson, E.R. Howlett, I. Hussain, Abeta deposition and related pathology in an APP x PS1 transgenic mouse model of Alzheimer's disease, Histol Histopathol 23 (2008) 67-76.
- [50] M. Meyer-Luehmann, T.L. Spires-Jones, C. Prada, M. Garcia-Alloza, A. de Calignon, A. Rozkalne, J. Koenigsknecht-Talboo, D.M. Holtzman, B.J. Bacskai, B.T. Hyman, Rapid appearance and local toxicity of amyloid-beta plaques in a mouse model of Alzheimer's disease, Nature 451 (2008) 720-724.

- [51] T. Bolmont, F. Haiss, D. Eicke, R. Radde, C.A. Mathis, W.E. Klunk, S. Kohsaka, M. Jucker, M.E. Calhoun, Dynamics of the microglial/amyloid interaction indicate a role in plaque maintenance, J Neurosci 28 (2008) 4283-4292.
- [52] M.N. Gordon, L.A. Holcomb, P.T. Jantzen, G. DiCarlo, D. Wilcock, K.W. Boyett, K. Connor, J. Melachrino, J.P. O'Callaghan, D. Morgan, Time course of the development of Alzheimer-like pathology in the doubly transgenic PS1+APP mouse, Exp Neurol 173 (2002) 183-195.
- [53] J. Tan, T. Town, F. Crawford, T. Mori, A. DelleDonne, R. Crescentini, D. Obregon, R.A. Flavell, M.J. Mullan, Role of CD40 ligand in amyloidosis in transgenic Alzheimer's mice, Nat Neurosci 5 (2002) 1288-1293.
- [54] K.P. Townsend, T. Town, T. Mori, L.F. Lue, D. Shytle, P.R. Sanberg, D. Morgan, F. Fernandez, R.A. Flavell, J. Tan, CD40 signaling regulates innate and adaptive activation of microglia in response to amyloid beta-peptide, Eur J Immunol 35 (2005) 901-910.
- [55] D. Obregon, H. Hou, Y. Bai, W.V. Nikolic, T. Mori, D. Luo, J. Zeng, J. Ehrhart, F. Fernandez, D. Morgan, B. Giunta, T. Town, J. Tan, CD40L disruption enhances Abeta vaccine-mediated reduction of cerebral amyloidosis while minimizing cerebral amyloid angiopathy and inflammation, Neurobiol Dis 29 (2008) 336-353.
- [56] W.V. Nikolic, H. Hou, T. Town, Y. Zhu, B. Giunta, C.D. Sanberg, J. Zeng, D. Luo, J. Ehrhart, T. Mori, P.R. Sanberg, J. Tan, Peripherally administered human umbilical cord blood cells reduce parenchymal and vascular beta-amyloid deposits in Alzheimer mice, Stem Cells Dev 17 (2008) 423-439.
- [57] V. Laporte, G. Ait-Ghezala, C.H. Volmar, C. Ganey, N. Ganey, M. Wood, M. Mullan, CD40 ligation mediates plaque-associated tau phosphorylation in betaamyloid overproducing mice, Brain Res 1231 (2008) 132-142.
- [58] J. El Khoury, M. Toft, S.E. Hickman, T.K. Means, K. Terada, C. Geula, A.D. Luster, Ccr2 deficiency impairs microglial accumulation and accelerates progression of Alzheimer-like disease, Nat Med 13 (2007) 432-438.
- [59] K.L. Richard, M. Filali, P. Prefontaine, S. Rivest, Toll-like receptor 2 acts as a natural innate immune receptor to clear amyloid beta 1-42 and delay the cognitive decline in a mouse model of Alzheimer's disease, J Neurosci 28 (2008) 5784-5793.
- [60] N. Zelcer, N. Khanlou, R. Clare, Q. Jiang, E.G. Reed-Geaghan, G.E. Landreth, H.V. Vinters, P. Tontonoz, Attenuation of neuroinflammation and Alzheimer's disease pathology by liver x receptors, Proc Natl Acad Sci U S A 104 (2007) 10601-10606.
- [61] Q. Jiang, C.Y. Lee, S. Mandrekar, B. Wilkinson, P. Cramer, N. Zelcer, K. Mann,
 B. Lamb, T.M. Willson, J.L. Collins, J.C. Richardson, J.D. Smith, T.A. Comery,
 D. Riddell, D.M. Holtzman, P. Tontonoz, G.E. Landreth, ApoE promotes the
 proteolytic degradation of Abeta, Neuron 58 (2008) 681-693.
- [62] V. Boissonneault, M. Filali, M. Lessard, J. Relton, G. Wong, S. Rivest, Powerful beneficial effects of macrophage colony-stimulating factor on beta-amyloid deposition and cognitive impairment in Alzheimer's disease, Brain 132 (2009) 1078-1092.

- [63] K.E. Nichol, W.W. Poon, A.I. Parachikova, D.H. Cribbs, C.G. Glabe, C.W. Cotman, Exercise alters the immune profile in Tg2576 Alzheimer mice toward a response coincident with improved cognitive performance and decreased amyloid, J Neuroinflammation 5 (2008) 13.
- [64] K. Farber, G. Cheung, D. Mitchell, R. Wallis, E. Weihe, W. Schwaeble, H. Kettenmann, C1q, the recognition subcomponent of the classical pathway of complement, drives microglial activation, J Neurosci Res 87 (2009) 644-652.
- [65] O. Butovsky, M. Koronyo-Hamaoui, G. Kunis, E. Ophir, G. Landa, H. Cohen, M. Schwartz, Glatiramer acetate fights against Alzheimer's disease by inducing dendritic-like microglia expressing insulin-like growth factor 1, Proc Natl Acad Sci U S A 103 (2006) 11784-11789.
- [66] D.L. Herber, M. Mercer, L.M. Roth, K. Symmonds, J. Maloney, N. Wilson, M.J. Freeman, D. Morgan, M.N. Gordon, Microglial activation is required for Abeta clearance after intracranial injection of lipopolysaccharide in APP transgenic mice, J Neuroimmune Pharmacol 2 (2007) 222-231.
- [67] R. Fan, F. Xu, M.L. Previti, J. Davis, A.M. Grande, J.K. Robinson, W.E. Van Nostrand, Minocycline reduces microglial activation and improves behavioral deficits in a transgenic model of cerebral microvascular amyloid, J Neurosci 27 (2007) 3057-3063.
- [68] F.C. Crawford, M. Wood, S. Ferguson, V.S. Mathura, B. Faza, S. Wilson, T. Fan, B. O'Steen, G. Ait-Ghezala, R. Hayes, M.J. Mullan, Genomic analysis of response to traumatic brain injury in a mouse model of Alzheimer's disease (APPsw), Brain Res 1185 (2007) 45-58.
- [69] S. Jimenez, D. Baglietto-Vargas, C. Caballero, I. Moreno-Gonzalez, M. Torres, R. Sanchez-Varo, D. Ruano, M. Vizuete, A. Gutierrez, J. Vitorica, Inflammatory response in the hippocampus of PS1M146L/APP751SL mouse model of Alzheimer's disease: age-dependent switch in the microglial phenotype from alternative to classic, J Neurosci 28 (2008) 11650-11661.
- [70] W.J. Streit, Microglial senescence: does the brain's immune system have an expiration date?, Trends Neurosci 29 (2006) 506-510.
- [71] B. Giunta, F. Fernandez, W.V. Nikolic, D. Obregon, E. Rrapo, T. Town, J. Tan, Inflammaging as a prodrome to Alzheimer's disease, J Neuroinflammation 5 (2008) 51.
- [72] A.R. Simard, S. Rivest, Bone marrow stem cells have the ability to populate the entire central nervous system into fully differentiated parenchymal microglia, Faseb J 18 (2004) 998-1000.
- [73] A.R. Simard, D. Soulet, G. Gowing, J.P. Julien, S. Rivest, Bone marrow-derived microglia play a critical role in restricting senile plaque formation in Alzheimer's disease, Neuron 49 (2006) 489-502.
- [74] T. Town, Y. Laouar, C. Pittenger, T. Mori, C.A. Szekely, J. Tan, R.S. Duman, R.A. Flavell, Blocking TGF-beta-Smad2/3 innate immune signaling mitigates Alzheimer-like pathology, Nat Med 14 (2008) 681-687.
- [75] O. Butovsky, G. Kunis, M. Koronyo-Hamaoui, M. Schwartz, Selective ablation of bone marrow-derived dendritic cells increases amyloid plaques in a mouse Alzheimer's disease model, Eur J Neurosci 26 (2007) 413-416.

- [76] B. Ajami, J.L. Bennett, C. Krieger, W. Tetzlaff, F.M. Rossi, Local self-renewal can sustain CNS microglia maintenance and function throughout adult life, Nat Neurosci 10 (2007) 1538-1543.
- [77] C.A. Lemere, E.T. Spooner, J.F. Leverone, C. Mori, M. Iglesias, J.K. Bloom, T.J. Seabrook, Amyloid-beta immunization in Alzheimer's disease transgenic mouse models and wildtype mice, Neurochem Res 28 (2003) 1017-1027.
- [78] D.M. Wilcock, C.A. Colton, Anti-amyloid-beta immunotherapy in Alzheimer's disease: relevance of transgenic mouse studies to clinical trials, J Alzheimers Dis 15 (2008) 555-569.
- [79] D. Morgan, The role of microglia in antibody-mediated clearance of amyloid-beta from the brain, CNS Neurol Disord Drug Targets 8 (2009) 7-15.
- [80] S. Gilman, M. Koller, R.S. Black, L. Jenkins, S.G. Griffith, N.C. Fox, L. Eisner, L. Kirby, M.B. Rovira, F. Forette, J.M. Orgogozo, Clinical effects of Abeta immunization (AN1792) in patients with AD in an interrupted trial, Neurology 64 (2005) 1553-1562.
- [81] J.A. Nicoll, D. Wilkinson, C. Holmes, P. Steart, H. Markham, R.O. Weller, Neuropathology of human Alzheimer disease after immunization with amyloidbeta peptide: a case report, Nat Med 9 (2003) 448-452.
- [82] D. Morgan, Modulation of microglial activation state following passive immunization in amyloid depositing transgenic mice, Neurochem Int 49 (2006) 190-194.
- [83] D.M. Wilcock, S.K. Munireddy, A. Rosenthal, K.E. Ugen, M.N. Gordon, D. Morgan, Microglial activation facilitates Abeta plaque removal following intracranial anti-Abeta antibody administration, Neurobiol Dis 15 (2004) 11-20.
- [84] D.M. Wilcock, M.N. Gordon, K.E. Ugen, P.E. Gottschall, G. DiCarlo, C. Dickey, K.W. Boyett, P.T. Jantzen, K.E. Connor, J. Melachrino, J. Hardy, D. Morgan, Number of Abeta inoculations in APP+PS1 transgenic mice influences antibody titers, microglial activation, and congophilic plaque levels, DNA Cell Biol 20 (2001) 731-736.
- [85] J. Koenigsknecht-Talboo, M. Meyer-Luehmann, M. Parsadanian, M. Garcia-Alloza, M.B. Finn, B.T. Hyman, B.J. Bacskai, D.M. Holtzman, Rapid microglial response around amyloid pathology after systemic anti-Abeta antibody administration in PDAPP mice, J Neurosci 28 (2008) 14156-14164.
- [86] F. Bard, R. Barbour, C. Cannon, R. Carretto, M. Fox, D. Games, T. Guido, K. Hoenow, K. Hu, K. Johnson-Wood, K. Khan, D. Kholodenko, C. Lee, M. Lee, R. Motter, M. Nguyen, A. Reed, D. Schenk, P. Tang, N. Vasquez, P. Seubert, T. Yednock, Epitope and isotype specificities of antibodies to beta -amyloid peptide for protection against Alzheimer's disease-like neuropathology, Proc Natl Acad Sci U S A 100 (2003) 2023-2028.
- [87] A.J. Gray, G. Sakaguchi, C. Shiratori, A.G. Becker, J. LaFrancois, P.S. Aisen, K. Duff, Y. Matsuoka, Antibody against C-terminal Abeta selectively elevates plasma Abeta, Neuroreport 18 (2007) 293-296.
- [88] Y. Okura, K. Kohyama, I.K. Park, Y. Matsumoto, Nonviral DNA vaccination augments microglial phagocytosis of beta-amyloid deposits as a major clearance pathway in an Alzheimer disease mouse model, J Neuropathol Exp Neurol 67 (2008) 1063-1071.

Inflammation in transgenic mouse models

[89]

A. Mouri, Y. Noda, H. Hara, H. Mizoguchi, T. Tabira, T. Nabeshima, Oral vaccination with a viral vector containing Abeta cDNA attenuates age-related Abeta accumulation and memory deficits without causing inflammation in a

mouse Alzheimer model, Faseb J 21 (2007) 2135-2148.
[90] M. Garcia-Alloza, B.J. Ferrara, S.A. Dodwell, G.A. Hickey, B.T. Hyman, B.J. Bacskai, A limited role for microglia in antibody mediated plaque clearance in

- Bacskai, A limited role for microglia in antibody mediated plaque clearance in APP mice, Neurobiol Dis 28 (2007) 286-292.
- [91] C. Colton, S. Wilt, D. Gilbert, O. Chernyshev, J. Snell, M. Dubois-Dalcq, Species differences in the generation of reactive oxygen species by microglia, Mol Chem Neuropathol 28 (1996) 15-20.
- [92] H. Akiyama, S. Barger, S. Barnum, B. Bradt, J. Bauer, G.M. Cole, N.R. Cooper, P. Eikelenboom, M. Emmerling, B.L. Fiebich, C.E. Finch, S. Frautschy, W.S. Griffin, H. Hampel, M. Hull, G. Landreth, L. Lue, R. Mrak, I.R. Mackenzie, P.L. McGeer, M.K. O'Banion, J. Pachter, G. Pasinetti, C. Plata-Salaman, J. Rogers, R. Rydel, Y. Shen, W. Streit, R. Strohmeyer, I. Tooyoma, F.L. Van Muiswinkel, R. Veerhuis, D. Walker, S. Webster, B. Wegrzyniak, G. Wenk, T. Wyss-Coray, Inflammation and Alzheimer's disease, Neurobiol Aging 21 (2000) 383-421.
- [93] R.E. Mrak, J.G. Sheng, W.S. Griffin, Glial cytokines in Alzheimer's disease: review and pathogenic implications, Hum Pathol 26 (1995) 816-823.
- [94] W.C. Benzing, J.R. Wujek, E.K. Ward, D. Shaffer, K.H. Ashe, S.G. Younkin, K.R. Brunden, Evidence for glial-mediated inflammation in aged APP(SW) transgenic mice, Neurobiol Aging 20 (1999) 581-589.
- [95] J. Apelt, R. Schliebs, Beta-amyloid-induced glial expression of both pro- and antiinflammatory cytokines in cerebral cortex of aged transgenic Tg2576 mice with Alzheimer plaque pathology, Brain Res 894 (2001) 21-30.
- [96] N. Abbas, I. Bednar, E. Mix, S. Marie, D. Paterson, A. Ljungberg, C. Morris, B. Winblad, A. Nordberg, J. Zhu, Up-regulation of the inflammatory cytokines IFN-gamma and IL-12 and down-regulation of IL-4 in cerebral cortex regions of APP(SWE) transgenic mice, J Neuroimmunol 126 (2002) 50-57.
- [97] N.S. Patel, D. Paris, V. Mathura, A.N. Quadros, F.C. Crawford, M.J. Mullan, Inflammatory cytokine levels correlate with amyloid load in transgenic mouse models of Alzheimer's disease, J Neuroinflammation 2 (2005) 9.
- [98] M.C. Janelsins, M.A. Mastrangelo, S. Oddo, F.M. LaFerla, H.J. Federoff, W.J. Bowers, Early correlation of microglial activation with enhanced tumor necrosis factor-alpha and monocyte chemoattractant protein-1 expression specifically within the entorhinal cortex of triple transgenic Alzheimer's disease mice, J Neuroinflammation 2 (2005) 23.
- [99] F.E. McAlpine, J.K. Lee, A.S. Harms, K.A. Ruhn, M. Blurton-Jones, J. Hong, P. Das, T.E. Golde, F.M. LaFerla, S. Oddo, A. Blesch, M.G. Tansey, Inhibition of soluble TNF signaling in a mouse model of Alzheimer's disease prevents pre-plaque amyloid-associated neuropathology, Neurobiol Dis 34 (2009) 163-177.
- [100] P. He, Z. Zhong, K. Lindholm, L. Berning, W. Lee, C. Lemere, M. Staufenbiel, R. Li, Y. Shen, Deletion of tumor necrosis factor death receptor inhibits amyloid beta generation and prevents learning and memory deficits in Alzheimer's mice, J Cell Biol 178 (2007) 829-841.

- [101] M. Yamamoto, T. Kiyota, M. Horiba, J.L. Buescher, S.M. Walsh, H.E. Gendelman, T. Ikezu, Interferon-gamma and tumor necrosis factor-alpha regulate amyloid-beta plaque deposition and beta-secretase expression in Swedish mutant APP transgenic mice, Am J Pathol 170 (2007) 680-692.
- [102] J.J. Jin, H.D. Kim, J.A. Maxwell, L. Li, K. Fukuchi, Toll-like receptor 4dependent upregulation of cytokines in a transgenic mouse model of Alzheimer's disease, J Neuroinflammation 5 (2008) 23.
- [103] S.S. Shaftel, S. Kyrkanides, J.A. Olschowka, J.N. Miller, R.E. Johnson, M.K. O'Banion, Sustained hippocampal IL-1 beta overexpression mediates chronic neuroinflammation and ameliorates Alzheimer plaque pathology, J Clin Invest 117 (2007) 1595-1604.
- [104] T. Wyss-Coray, C. Lin, F. Yan, G.Q. Yu, M. Rohde, L. McConlogue, E. Masliah, L. Mucke, TGF-beta1 promotes microglial amyloid-beta clearance and reduces plaque burden in transgenic mice, Nat Med 7 (2001) 612-618.
- [105] S. Lesne, F. Docagne, C. Gabriel, G. Liot, D.K. Lahiri, L. Buee, L. Plawinski, A. Delacourte, E.T. MacKenzie, A. Buisson, D. Vivien, Transforming growth factorbeta 1 potentiates amyloid-beta generation in astrocytes and in transgenic mice, J Biol Chem 278 (2003) 18408-18418.
- [106] M. Yamamoto, M. Horiba, J.L. Buescher, D. Huang, H.E. Gendelman, R.M. Ransohoff, T. Ikezu, Overexpression of monocyte chemotactic protein-1/CCL2 in beta-amyloid precursor protein transgenic mice show accelerated diffuse betaamyloid deposition, Am J Pathol 166 (2005) 1475-1485.
- [107] T. Kiyota, M. Yamamoto, B. Schroder, M.T. Jacobsen, R.J. Swan, M.P. Lambert, W.L. Klein, H.E. Gendelman, R.M. Ransohoff, T. Ikezu, AAV1/2-mediated CNS gene delivery of dominant-negative CCL2 mutant suppresses gliosis, betaamyloidosis, and learning impairment of APP/PS1 mice, Mol Ther 17 (2009) 803-809.
- [108] B.A. in't Veld, A. Ruitenberg, A. Hofman, B.H. Stricker, M.M. Breteler, Antihypertensive drugs and incidence of dementia: the Rotterdam Study, Neurobiol Aging 22 (2001) 407-412.
- [109] W.F. Stewart, C. Kawas, M. Corrada, E.J. Metter, Risk of Alzheimer's disease and duration of NSAID use, Neurology 48 (1997) 626-632.
- [110] P.L. McGeer, E.G. McGeer, NSAIDs and Alzheimer disease: epidemiological, animal model and clinical studies, Neurobiol Aging 28 (2007) 639-647.
- [111] S.C. Vlad, D.R. Miller, N.W. Kowall, D.T. Felson, Protective effects of NSAIDs on the development of Alzheimer disease, Neurology 70 (2008) 1672-1677.
- [112] G.P. Lim, F. Yang, T. Chu, P. Chen, W. Beech, B. Teter, T. Tran, O. Ubeda, K.H. Ashe, S.A. Frautschy, G.M. Cole, Ibuprofen suppresses plaque pathology and inflammation in a mouse model for Alzheimer's disease, J Neurosci 20 (2000) 5709-5714.
- [113] Q. Yan, J. Zhang, H. Liu, S. Babu-Khan, R. Vassar, A.L. Biere, M. Citron, G. Landreth, Anti-inflammatory drug therapy alters beta-amyloid processing and deposition in an animal model of Alzheimer's disease, J Neurosci 23 (2003) 7504-7509.
- [114] P.T. Jantzen, K.E. Connor, G. DiCarlo, G.L. Wenk, J.L. Wallace, A.M. Rojiani, D. Coppola, D. Morgan, M.N. Gordon, Microglial activation and beta -amyloid

Inflammation in transgenic mouse models

deposit reduction caused by a nitric oxide-releasing nonsteroidal antiinflammatory drug in amyloid precursor protein plus presenilin-1 transgenic mice, J Neurosci 22 (2002) 2246-2254.

- [115] T. van Groen, I. Kadish, Transgenic AD model mice, effects of potential anti-AD treatments on inflammation and pathology, Brain Res Brain Res Rev 48 (2005) 370-378.
- [116] T. Kukar, S. Prescott, J.L. Eriksen, V. Holloway, M.P. Murphy, E.H. Koo, T.E. Golde, M.M. Nicolle, Chronic administration of R-flurbiprofen attenuates learning impairments in transgenic amyloid precursor protein mice, BMC Neurosci 8 (2007) 54.
- [117] M.T. Heneka, M. Sastre, L. Dumitrescu-Ozimek, A. Hanke, I. Dewachter, C. Kuiperi, K. O'Banion, T. Klockgether, F. Van Leuven, G.E. Landreth, Acute treatment with the PPARgamma agonist pioglitazone and ibuprofen reduces glial inflammation and Abeta1-42 levels in APPV717I transgenic mice, Brain 128 (2005) 1442-1453.
- [118] T. Morihara, B. Teter, F. Yang, G.P. Lim, S. Boudinot, F.D. Boudinot, S.A. Frautschy, G.M. Cole, Ibuprofen suppresses interleukin-1beta induction of proamyloidogenic alpha1-antichymotrypsin to ameliorate beta-amyloid (Abeta) pathology in Alzheimer's models, Neuropsychopharmacology 30 (2005) 1111-1120.
- [119] A.C. McKee, I. Carreras, L. Hossain, H. Ryu, W.L. Klein, S. Oddo, F.M. LaFerla, B.G. Jenkins, N.W. Kowall, A. Dedeoglu, Ibuprofen reduces Abeta, hyperphosphorylated tau and memory deficits in Alzheimer mice, Brain Res 1207 (2008) 225-236.
- [120] T.A. Lanz, G.J. Fici, K.M. Merchant, Lack of specific amyloid-beta(1-42) suppression by nonsteroidal anti-inflammatory drugs in young, plaque-free Tg2576 mice and in guinea pig neuronal cultures, J Pharmacol Exp Ther 312 (2005) 399-406.
- [121] J. Quinn, T. Montine, J. Morrow, W.R. Woodward, D. Kulhanek, F. Eckenstein, Inflammation and cerebral amyloidosis are disconnected in an animal model of Alzheimer's disease, J Neuroimmunol 137 (2003) 32-41.
- [122] S. Sung, H. Yang, K. Uryu, E.B. Lee, L. Zhao, D. Shineman, J.Q. Trojanowski, V.M. Lee, D. Pratico, Modulation of nuclear factor-kappa B activity by indomethacin influences A beta levels but not A beta precursor protein metabolism in a model of Alzheimer's disease, Am J Pathol 165 (2004) 2197-2206.
- [123] E. Dvir, A. Elman, D. Simmons, I. Shapiro, R. Duvdevani, A. Dahan, A. Hoffman, J.E. Friedman, DP-155, a lecithin derivative of indomethacin, is a novel nonsteroidal antiinflammatory drug for analgesia and Alzheimer's disease therapy, CNS Drug Rev 13 (2007) 260-277.
- [124] T. Kukar, M.P. Murphy, J.L. Eriksen, S.A. Sagi, S. Weggen, T.E. Smith, T. Ladd, M.A. Khan, R. Kache, J. Beard, M. Dodson, S. Merit, V.V. Ozols, P.Z. Anastasiadis, P. Das, A. Fauq, E.H. Koo, T.E. Golde, Diverse compounds mimic Alzheimer disease-causing mutations by augmenting Abeta42 production, Nat Med 11 (2005) 545-550.

- [125] J.J. Hoozemans, J.M. Rozemuller, E.S. van Haastert, R. Veerhuis, P. Eikelenboom, Cyclooxygenase-1 and -2 in the different stages of Alzheimer's disease pathology, Curr Pharm Des 14 (2008) 1419-1427.
- [126] J.R. Vane, R.M. Botting, Mechanism of action of antiinflammatory drugs, Int J Tissue React 20 (1998) 3-15.
- [127] S. Weggen, J.L. Eriksen, P. Das, S.A. Sagi, R. Wang, C.U. Pietrzik, K.A. Findlay, T.E. Smith, M.P. Murphy, T. Bulter, D.E. Kang, N. Marquez-Sterling, T.E. Golde, E.H. Koo, A subset of NSAIDs lower amyloidogenic Abeta42 independently of cyclooxygenase activity, Nature 414 (2001) 212-216.
- [128] A. Lleo, E. Galea, M. Sastre, Molecular targets of non-steroidal anti-inflammatory drugs in neurodegenerative diseases, Cell Mol Life Sci 64 (2007) 1403-1418.
- [129] P.L. McGeer, S. Itagaki, B.E. Boyes, E.G. McGeer, Reactive microglia are positive for HLA-DR in the substantia nigra of Parkinson's and Alzheimer's disease brains, Neurology 38 (1988) 1285-1291.
- [130] E.C. Hirsch, S. Hunot, Neuroinflammation in Parkinson's disease: a target for neuroprotection?, Lancet Neurol 8 (2009) 382-397.
- [131] P.S. Whitton, Inflammation as a causative factor in the aetiology of Parkinson's disease, Br J Pharmacol 150 (2007) 963-976.
- [132] P.L. McGeer, E.G. McGeer, Glial reactions in Parkinson's disease, Mov Disord 23 (2008) 474-483.
- [133] J.W. Langston, P. Ballard, J.W. Tetrud, I. Irwin, Chronic Parkinsonism in humans due to a product of meperidine-analog synthesis, Science 219 (1983) 979-980.
- [134] J.W. Langston, L.S. Forno, J. Tetrud, A.G. Reeves, J.A. Kaplan, D. Karluk, Evidence of active nerve cell degeneration in the substantia nigra of humans years after 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine exposure, Ann. Neurol. 46 (1999) 598-605.
- [135] P.L. McGeer, C. Schwab, A. Parent, D. Doudet, Presence of reactive microglia in monkey substantia nigra years after 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine administration, Ann. Neurol. 54 (2003) 599-604.
- [136] J. Miklossy, D.D. Doudet, C. Schwab, S. Yu, E.G. McGeer, P.L. McGeer, Role of ICAM-1 in persisting inflammation in Parkinson disease and MPTP monkeys, Exp. Neurol. 197 (2006) 275-283.
- [137] P. Jenner, Functional models of Parkinson's disease: a valuable tool in the development of novel therapies, Annals of Neurology 64 (2008).
- [138] S.M. Fleming, P.O. Fernagut, M.F. Chesselet, Genetic mouse models of parkinsonism: strengths and limitations, NeuroRx 2 (2005) 495-503.
- [139] B.K. Harvey, Y. Wang, B.J. Hoffer, Transgenic rodent models of Parkinson's disease, Acta Neurochir Suppl 101 (2008) 89-92.
- [140] M. Terzioglu, D. Galter, Parkinson's disease: genetic versus toxin-induced rodent models, Febs J 275 (2008) 1384-1391.
- [141] O. von Bohlen und Halbach, Modeling neurodegenerative diseases in vivo review, Neurodegenerative Diseases 2 (2005) 313-320.
- [142] M.C. Chartier-Harlin, J. Kachergus, C. Roumier, V. Mouroux, X. Douay, S. Lincoln, C. Levecque, L. Larvor, J. Andrieux, M. Hulihan, N. Waucquier, L. Defebvre, P. Amouyel, M. Farrer, A. Destee, Alpha-synuclein locus duplication as a cause of familial Parkinson's disease, Lancet 364 (2004) 1167-1169.

- [143] P. Ibanez, S. Lesage, S. Janin, E. Lohmann, F. Durif, A. Destee, A.M. Bonnet, C. Brefel-Courbon, S. Heath, D. Zelenika, Y. Agid, A. Durr, A. Brice, Alpha-synuclein gene rearrangements in dominantly inherited parkinsonism: frequency, phenotype, and mechanisms, Arch Neurol 66 (2009) 102-108.
- [144] P. Ibanez, A.M. Bonnet, B. Debarges, E. Lohmann, F. Tison, P. Pollak, Y. Agid, A. Durr, A. Brice, Causal relation between alpha-synuclein gene duplication and familial Parkinson's disease, Lancet 364 (2004) 1169-1171.
- [145] A.B. Singleton, M. Farrer, J. Johnson, A. Singleton, S. Hague, J. Kachergus, M. Hulihan, T. Peuralinna, A. Dutra, R. Nussbaum, S. Lincoln, A. Crawley, M. Hanson, D. Maraganore, C. Adler, M.R. Cookson, M. Muenter, M. Baptista, D. Miller, J. Blancato, J. Hardy, K. Gwinn-Hardy, alpha-Synuclein locus triplication causes Parkinson's disease, Science 302 (2003) 841.
- [146] E. Masliah, E. Rockenstein, I. Veinbergs, M. Mallory, M. Hashimoto, A. Takeda, Y. Sagara, A. Sisk, L. Mucke, Dopaminergic loss and inclusion body formation in alpha-synuclein mice: implications for neurodegenerative disorders, Science 287 (2000) 1265-1269.
- [147] M.F. Chesselet, In vivo alpha-synuclein overexpression in rodents: a useful model of Parkinson's disease?, Experimental Neurology 209 (2008) 22-27.
- [148] M.F. Chesselet, S. Fleming, F. Mortazavi, B. Meurers, Strengths and limitations of genetic mouse models of Parkinson's disease, Parkinsonism & Related Disorders 14 (2008).
- [149] P.O. Fernagut, C.B. Hutson, S.M. Fleming, N.A. Tetreaut, J. Salcedo, E. Masliah, M.F. Chesselet, Behavioral and histopathological consequences of paraquat intoxication in mice: effects of alpha-synuclein over-expression, Synapse 61 (2007) 991-1001.
- [150] H. van der Putten, K.H. Wiederhold, A. Probst, S. Barbieri, C. Mistl, S. Danner, S. Kauffmann, K. Hofele, W.P. Spooren, M.A. Ruegg, S. Lin, P. Caroni, B. Sommer, M. Tolnay, G. Bilbe, Neuropathology in mice expressing human alphasynuclein, Journal of Neuroscience 20 (2000) 6021-6029.
- [151] B.I. Giasson, J.E. Duda, S.M. Quinn, B. Zhang, J.Q. Trojanowski, V.M. Lee, Neuronal alpha-synucleinopathy with severe movement disorder in mice expressing A53T human alpha-synuclein, Neuron 34 (2002) 521-533.
- [152] M.K. Lee, W. Stirling, Y. Xu, X. Xu, D. Qui, A.S. Mandir, T.M. Dawson, N.G. Copeland, N.A. Jenkins, D.L. Price, Human alpha-synuclein-harboring familial Parkinson's disease-linked Ala-53 --> Thr mutation causes neurodegenerative disease with alpha-synuclein aggregation in transgenic mice, Proceedings of the National Academy of Sciences of the United States of America 99 (2002) 8968-8973.
- [153] S. Gispert, D. Del Turco, L. Garrett, A. Chen, D.J. Bernard, J. Hamm-Clement, H.W. Korf, T. Deller, H. Braak, G. Auburger, R.L. Nussbaum, Transgenic mice expressing mutant A53T human alpha-synuclein show neuronal dysfunction in the absence of aggregate formation, Molecular & Cellular Neurosciences 24 (2003) 419-429.
- [154] T. Gomez-Isla, M.C. Irizarry, A. Mariash, B. Cheung, O. Soto, S. Schrump, J. Sondel, L. Kotilinek, J. Day, M.A. Schwarzschild, J.H. Cha, K. Newell, D.W. Miller, K. Ueda, A.B. Young, B.T. Hyman, K.H. Ashe, Motor dysfunction and

Inflammation in transgenic mouse models

gliosis with preserved dopaminergic markers in human alpha-synuclein A30P transgenic mice, Neurobiology of Aging 24 (2003) 245-258.

- [155] X. Su, K.A. Maguire-Zeiss, R. Giuliano, L. Prifti, K. Venkatesh, H.J. Federoff, Synuclein activates microglia in a model of Parkinson's disease, Neurobiol Aging 29 (2008) 1690-1701.
- [156] G.K. Tofaris, P. Garcia Reitbock, T. Humby, S.L. Lambourne, M. O'Connell, B. Ghetti, H. Gossage, P.C. Emson, L.S. Wilkinson, M. Goedert, M.G. Spillantini, Pathological changes in dopaminergic nerve cells of the substantia nigra and olfactory bulb in mice transgenic for truncated human alpha-synuclein(1-120): implications for Lewy body disorders, Journal of Neuroscience 26 (2006) 3942-3950.
- [157] M. Wakamatsu, A. Ishii, S. Iwata, J. Sakagami, Y. Ukai, M. Ono, D. Kanbe, S. Muramatsu, K. Kobayashi, T. Iwatsubo, M. Yoshimoto, Selective loss of nigral dopamine neurons induced by overexpression of truncated human alpha-synuclein in mice, Neurobiology of Aging 29 (2008) 574-585.
- [158] N. Stefanova, M. Reindl, M. Neumann, P.J. Kahle, W. Poewe, G.K. Wenning, Microglial activation mediates neurodegeneration related to oligodendroglial alpha-synucleinopathy: implications for multiple system atrophy, Movement Disorders 22 (2007) 2196-2203.
- [159] H.M. Gao, P.T. Kotzbauer, K. Uryu, S. Leight, J.Q. Trojanowski, V.M. Lee, Neuroinflammation and oxidation/nitration of alpha-synuclein linked to dopaminergic neurodegeneration, J Neurosci 28 (2008) 7687-7698.
- [160] S.A. Austin, A.M. Floden, E.J. Murphy, C.K. Combs, Alpha-synuclein expression modulates microglial activation phenotype, Journal of Neuroscience 26 (2006) 10558-10563.
- [161] E. Masliah, E. Rockenstein, A. Adame, M. Alford, L. Crews, M. Hashimoto, P. Seubert, M. Lee, J. Goldstein, T. Chilcote, D. Games, D. Schenk, Effects of alpha-synuclein immunization in a mouse model of Parkinson's disease, Neuron 46 (2005) 857-868.
- [162] A. Klegeris, B.I. Giasson, H. Zhang, J. Maguire, S. Pelech, P.L. McGeer, Alphasynuclein and its disease-causing mutants induce ICAM-1 and IL-6 in human astrocytes and astrocytoma cells, Faseb J 20 (2006) 2000-2008.
- [163] A. Klegeris, S. Pelech, B.I. Giasson, J. Maguire, H. Zhang, E.G. McGeer, P.L. McGeer, Alpha-synuclein activates stress signaling protein kinases in THP-1 cells and microglia, Neurobiol Aging 29 (2008) 739-752.
- [164] J. Liu, J.P. Zhang, M. Shi, T. Quinn, J. Bradner, R. Beyer, S. Chen, J. Zhang, Rab11a and HSP90 regulate recycling of extracellular alpha-synuclein, Journal of Neuroscience 29 (2009) 1480-1485.
- [165] W. Zhang, S. Dallas, D. Zhang, J.P. Guo, H. Pang, B. Wilson, D.S. Miller, B. Chen, W. Zhang, P.L. McGeer, J.S. Hong, J. Zhang, Microglial PHOX and Mac-1 are essential to the enhanced dopaminergic neurodegeneration elicited by A30P and A53T mutant alpha-synuclein, Glia 55 (2007) 1178-1188.
- [166] W. Zhang, T. Wang, Z. Pei, D.S. Miller, X. Wu, M.L. Block, B. Wilson, W. Zhang, Y. Zhou, J.S. Hong, J. Zhang, Aggregated alpha-synuclein activates microglia: a process leading to disease progression in Parkinson's disease, Faseb J 19 (2005) 533-542.

- [167] L. Chen, B. Cagniard, T. Mathews, S. Jones, H.C. Koh, Y. Ding, P.M. Carvey, Z. Ling, U.J. Kang, X. Zhuang, Age-dependent motor deficits and dopaminergic dysfunction in DJ-1 null mice, Journal of Biological Chemistry 280 (2005) 21418-21426.
- [168] M.S. Goldberg, A. Pisani, M. Haburcak, T.A. Vortherms, T. Kitada, C. Costa, Y. Tong, G. Martella, A. Tscherter, A. Martins, G. Bernardi, B.L. Roth, E.N. Pothos, P. Calabresi, J. Shen, Nigrostriatal dopaminergic deficits and hypokinesia caused by inactivation of the familial Parkinsonism-linked gene DJ-1.[see comment], Neuron 45 (2005) 489-496.
- [169] R.H. Kim, P.D. Smith, H. Aleyasin, S. Hayley, M.P. Mount, S. Pownall, A. Wakeham, A.J. You-Ten, S.K. Kalia, P. Horne, D. Westaway, A.M. Lozano, H. Anisman, D.S. Park, T.W. Mak, Hypersensitivity of DJ-1-deficient mice to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyrindine (MPTP) and oxidative stress, Proceedings of the National Academy of Sciences of the United States of America 102 (2005) 5215-5220.
- [170] T.C. Frank-Cannon, T. Tran, K.A. Ruhn, T.N. Martinez, J. Hong, M. Marvin, M. Hartley, I. Trevino, D.E. O'Brien, B. Casey, M.S. Goldberg, M.G. Tansey, Parkin deficiency increases vulnerability to inflammation-related nigral degeneration, J Neurosci 28 (2008) 10825-10834.
- [171] M.S. Goldberg, S.M. Fleming, J.J. Palacino, C. Cepeda, H.A. Lam, A. Bhatnagar, E.G. Meloni, N. Wu, L.C. Ackerson, G.J. Klapstein, M. Gajendiran, B.L. Roth, M.F. Chesselet, N.T. Maidment, M.S. Levine, J. Shen, Parkin-deficient mice exhibit nigrostriatal deficits but not loss of dopaminergic neurons, J Biol Chem 278 (2003) 43628-43635.
- [172] T. Kitada, A. Pisani, D.R. Porter, H. Yamaguchi, A. Tscherter, G. Martella, P. Bonsi, C. Zhang, E.N. Pothos, J. Shen, Impaired dopamine release and synaptic plasticity in the striatum of PINK1-deficient mice, Proc Natl Acad Sci U S A 104 (2007) 11441-11446.
- [173] H. Zhou, B.H. Falkenburger, J.B. Schulz, K. Tieu, Z. Xu, X.G. Xia, Silencing of the Pink1 gene expression by conditional RNAi does not induce dopaminergic neuron death in mice, International journal of biological sciences 3 (2007) 242-250.
- [174] J.M. Itier, P. Ibanez, M.A. Mena, N. Abbas, C. Cohen-Salmon, G.A. Bohme, M. Laville, J. Pratt, O. Corti, L. Pradier, G. Ret, C. Joubert, M. Periquet, F. Araujo, J. Negroni, M.J. Casarejos, S. Canals, R. Solano, A. Serrano, E. Gallego, M. Sanchez, P. Denefle, J. Benavides, G. Tremp, T.A. Rooney, A. Brice, J. Garcia de Yebenes, Parkin gene inactivation alters behaviour and dopamine neurotransmission in the mouse, Hum Mol Genet 12 (2003) 2277-2291.
- [175] R. Von Coelln, B. Thomas, J.M. Savitt, K.L. Lim, M. Sasaki, E.J. Hess, V.L. Dawson, T.M. Dawson, Loss of locus coeruleus neurons and reduced startle in parkin null mice, Proceedings of the National Academy of Sciences of the United States of America 101 (2004) 10744-10749.
- [176] J.J. Palacino, D. Sagi, M.S. Goldberg, S. Krauss, C. Motz, M. Wacker, J. Klose, J. Shen, Mitochondrial dysfunction and oxidative damage in parkin-deficient mice, J Biol Chem 279 (2004) 18614-18622.

- [177] F.A. Perez, R.D. Palmiter, Parkin-deficient mice are not a robust model of parkinsonism, Proc Natl Acad Sci U S A 102 (2005) 2174-2179.
- [178] S. Sato, T. Chiba, S. Nishiyama, T. Kakiuchi, H. Tsukada, T. Hatano, T. Fukuda, Y. Yasoshima, N. Kai, K. Kobayashi, Y. Mizuno, K. Tanaka, N. Hattori, Decline of striatal dopamine release in parkin-deficient mice shown by ex vivo autoradiography, J Neurosci Res 84 (2006) 1350-1357.
- [179] B. Thomas, R. von Coelln, A.S. Mandir, D.B. Trinkaus, M.H. Farah, K. Leong Lim, N.Y. Calingasan, M. Flint Beal, V.L. Dawson, T.M. Dawson, MPTP and DSP-4 susceptibility of substantia nigra and locus coeruleus catecholaminergic neurons in mice is independent of parkin activity, Neurobiology of Disease 26 (2007) 312-322.
- [180] M.I. Ekstrand, M. Terzioglu, D. Galter, S. Zhu, C. Hofstetter, E. Lindqvist, S. Thams, A. Bergstrand, F.S. Hansson, A. Trifunovic, B. Hoffer, S. Cullheim, A.H. Mohammed, L. Olson, N.G. Larsson, Progressive parkinsonism in mice with respiratory-chain-deficient dopamine neurons, Proc Natl Acad Sci U S A 104 (2007) 1325-1330.
- [181] J. Peng, L. Xie, F.F. Stevenson, S. Melov, D.A. Di Monte, J.K. Andersen, Nigrostriatal dopaminergic neurodegeneration in the weaver mouse is mediated via neuroinflammation and alleviated by minocycline administration, Journal of Neuroscience 26 (2006) 11644-11651.
- [182] J.K. Mallajosyula, D. Kaur, S.J. Chinta, S. Rajagopalan, A. Rane, D.G. Nicholls, D.A. Di Monte, H. Macarthur, J.K. Andersen, MAO-B elevation in mouse brain astrocytes results in Parkinson's pathology, PLoS ONE [Electronic Resource] 3 (2008).
- [183] T. Thomas, M. Timmer, K. Cesnulevicius, E. Hitti, A. Kotlyarov, M. Gaestel, MAPKAP kinase 2-deficiency prevents neurons from cell death by reducing neuroinflammation--relevance in a mouse model of Parkinson's disease, J Neurochem 105 (2008) 2039-2052.
- [184] Y. Liang, S. Li, Q. Guo, Y. Zhang, C. Wen, Q. Zou, B. Su, Complement 3deficient mice are not protected against MPTP-induced dopaminergic neurotoxicity, Brain Res 1178 (2007) 132-140.
- [185] C.E. Hamill, W.M. Caudle, J.R. Richardson, H. Yuan, K.D. Pennell, J.G. Greene, G.W. Miller, S.F. Traynelis, Exacerbation of dopaminergic terminal damage in a mouse model of Parkinson's disease by the G-protein-coupled receptor proteaseactivated receptor 1, Molecular pharmacology 72 (2007) 653-664.
- [186] A. Vroon, B. Drukarch, J.G. Bol, P. Cras, J.J. Breve, S.M. Allan, J.K. Relton, P.V. Hoogland, A.M. Van Dam, Neuroinflammation in Parkinson's patients and MPTP-treated mice is not restricted to the nigrostriatal system: microgliosis and differential expression of interleukin-1 receptors in the olfactory bulb, Exp Gerontol 42 (2007) 762-771.
- [187] R. Pattarini, R.J. Smeyne, J.I. Morgan, Temporal mRNA profiles of inflammatory mediators in the murine 1-methyl-4-phenyl-1,2,3,6-tetrahydropyrimidine model of Parkinson's disease, Neuroscience 145 (2007) 654-668.
- [188] L. Qin, X. Wu, M.L. Block, Y. Liu, G.R. Breese, J.S. Hong, D.J. Knapp, F.T. Crews, Systemic LPS causes chronic neuroinflammation and progressive neurodegeneration, Glia 55 (2007) 453-462.

- 41
- [189] J. Jin, F.S. Shie, J. Liu, Y. Wang, J. Davis, A.M. Schantz, K.S. Montine, T.J. Montine, J. Zhang, Prostaglandin E2 receptor subtype 2 (EP2) regulates microglial activation and associated neurotoxicity induced by aggregated alphasynuclein, J Neuroinflammation 4 (2007) 2.
- [190] W. Springer, P.J. Kahle, Mechanisms and models of alpha-synuclein-related neurodegeneration, Current neurology and neuroscience reports 6 (2006) 432-436.
- [191] C.D. Link, Transgenic invertebrate models of age-associated neurodegenerative diseases, Mechanisms of ageing and development 122 (2001) 1639-1649.
- [192] S.K. Singhrao, J.W. Neal, B.P. Morgan, P. Gasque, Increased complement biosynthesis by microglia and complement activation on neurons in Huntington's disease, Exp Neurol 159 (1999) 362-376.
- [193] A. Dalrymple, E.J. Wild, R. Joubert, K. Sathasivam, M. Bjorkqvist, A. Petersen, G.S. Jackson, J.D. Isaacs, M. Kristiansen, G.P. Bates, B.R. Leavitt, G. Keir, M. Ward, S.J. Tabrizi, Proteomic Profiling of Plasma in Huntington's Disease Reveals Neuroinflammatory Activation and Biomarker Candidates, Journal of Proteome Research 6 (2007) 2833-2840.
- [194] P.L. McGeer, S. Itagaki, E.G. McGeer, Expression of the histocompatibility glycoprotein HLA-DR in neurological disease, Acta Neuropathol 76 (1988) 550-557.
- [195] E. Sapp, K.B. Kegel, N. Aronin, T. Hashikawa, Y. Uchiyama, K. Tohyama, P.G. Bhide, J.P. Vonsattel, M. DiFiglia, Early and progressive accumulation of reactive microglia in the Huntington disease brain, J Neuropathol Exp Neurol 60 (2001) 161-172.
- [196] K. Messmer, G.P. Reynolds, Increased peripheral benzodiazepine binding sites in the brain of patients with Huntington's disease, Neurosci Lett 241 (1998) 53-56.
- [197] Y.F. Tai, N. Pavese, A. Gerhard, S.J. Tabrizi, R.A. Barker, D.J. Brooks, P. Piccini, Microglial activation in presymptomatic Huntington's disease gene carriers, Brain 130 (2007) 1759-1766.
- [198] Y.F. Tai, N. Pavese, A. Gerhard, S.J. Tabrizi, R.A. Barker, D.J. Brooks, P. Piccini, Imaging microglial activation in Huntington's disease, Brain Res Bull 72 (2007) 148-151.
- [199] N. Pavese, A. Gerhard, Y.F. Tai, A.K. Ho, F. Turkheimer, R.A. Barker, D.J. Brooks, P. Piccini, Microglial activation correlates with severity in Huntington disease: a clinical and PET study, Neurology 66 (2006) 1638-1643.
- [200] D.A. Simmons, M. Casale, B. Alcon, N. Pham, N. Narayan, G. Lynch, Ferritin accumulation in dystrophic microglia is an early event in the development of Huntington's disease, Glia 55 (2007) 1074-1084.
- [201] L. Ma, A.J. Morton, L.F. Nicholson, Microglia density decreases with age in a mouse model of Huntington's disease, Glia 43 (2003) 274-280.
- [202] M. Bjorkqvist, E.J. Wild, J. Thiele, A. Silvestroni, R. Andre, N. Lahiri, E. Raibon, R.V. Lee, C.L. Benn, D. Soulet, A. Magnusson, B. Woodman, C. Landles, M.A. Pouladi, M.R. Hayden, A. Khalili-Shirazi, M.W. Lowdell, P. Brundin, G.P. Bates, B.R. Leavitt, T. Moller, S.J. Tabrizi, A novel pathogenic pathway of immune activation detectable before clinical onset in Huntington's disease, J Exp Med 205 (2008) 1869-1877.

- 42
- [203] M.S. Levine, C. Cepeda, M.A. Hickey, S.M. Fleming, M.-F. Chesselet, Genetic mouse models of Huntington's and Parkinson's diseases: illuminating but imperfect, Trends in Neurosciences 27 (2004) 691-697.
- [204] S.K. Godavarthi, D. Narender, A. Mishra, A. Goswami, S.N. Rao, N. Nukina, N.R. Jana, Induction of chemokines, MCP-1, and KC in the mutant huntingtin expressing neuronal cells because of proteasomal dysfunction, J Neurochem 108 (2009) 787-795.
- [205] E.C. Stack, K.M. Smith, H. Ryu, K. Cormier, M. Chen, S.W. Hagerty, S.J. Del Signore, M.E. Cudkowicz, R.M. Friedlander, R.J. Ferrante, Combination therapy using minocycline and coenzyme Q10 in R6/2 transgenic Huntington's disease mice, Biochim Biophys Acta 1762 (2006) 373-380.
- [206] X. Wang, S. Zhu, M. Drozda, W. Zhang, I.G. Stavrovskaya, E. Cattaneo, R.J. Ferrante, B.S. Kristal, R.M. Friedlander, Minocycline inhibits caspaseindependent and -dependent mitochondrial cell death pathways in models of Huntington's disease, Proc Natl Acad Sci U S A 100 (2003) 10483-10487.
- [207] J.K. Ryu, H.B. Choi, J.G. McLarnon, Combined minocycline plus pyruvate treatment enhances effects of each agent to inhibit inflammation, oxidative damage, and neuronal loss in an excitotoxic animal model of Huntington's disease, Neuroscience 141 (2006) 1835-1848.
- [208] T.M. Woodruff, J.W. Crane, L.M. Proctor, K.M. Buller, A.B. Shek, K. de Vos, S. Pollitt, H.M. Williams, I.A. Shiels, P.N. Monk, S.M. Taylor, Therapeutic activity of C5a receptor antagonists in a rat model of neurodegeneration, Faseb J 20 (2006) 1407-1417.
- [209] P. Guidetti, G.P. Bates, R.K. Graham, M.R. Hayden, B.R. Leavitt, M.E. MacDonald, E.J. Slow, V.C. Wheeler, B. Woodman, R. Schwarcz, Elevated brain 3-hydroxykynurenine and quinolinate levels in Huntington disease mice, Neurobiol Dis 23 (2006) 190-197.
- [210] P. Guidetti, P.H. Reddy, D.A. Tagle, R. Schwarcz, Early kynurenergic impairment in Huntington's disease and in a transgenic animal model, Neurosci Lett 283 (2000) 233-235.
- [211] N. Stoy, G.M. Mackay, C.M. Forrest, J. Christofides, M. Egerton, T.W. Stone, L.G. Darlington, Tryptophan metabolism and oxidative stress in patients with Huntington's disease, J Neurochem 93 (2005) 611-623.
- [212] C.A. Lemere, M. Maier, L. Jiang, Y. Peng, T.J. Seabrook, Amyloid-beta immunotherapy for the prevention and treatment of Alzheimer disease: lessons from mice, monkeys, and humans, Rejuvenation Res 9 (2006) 77-84.
- [213] E. Sasaki, H. Suemizu, A. Shimada, K. Hanazawa, R. Oiwa, M. Kamioka, I. Tomioka, Y. Sotomaru, R. Hirakawa, T. Eto, S. Shiozawa, T. Maeda, M. Ito, R. Ito, C. Kito, C. Yagihashi, K. Kawai, H. Miyoshi, Y. Tanioka, N. Tamaoki, S. Habu, H. Okano, T. Nomura, Generation of transgenic non-human primates with germline transmission, Nature 459 (2009) 523-528.

Inflammation in transgenic mouse models

Legend to figure 1:

Comparison of immunostaining and histology staining of the cortex of APP23 mice (A, C, E, G, I and K) and of a human AD case (B, D, F, H, K and L). (A) and (B) compare Abeta deposits. (C) and (D) compare Bielschowsky histology staining demonstrating only few dystrophic neurites associated with Abeta plaques in APP23 mice but dense fibrillary changes in the AD case. (E) and (F) compare immunostaining for microglial cells. The mouse tissue is stained with an antibody against Iba1 and the human tissue with an antibody against HLA-DR. (G) and (H) compare GFAP immunostaining for astrocytes (G is counterstained with Cresyl Violet, labeling all nuclei). (I) and (J) compare immunostaining of C1q of complement (note that C1q association with plaques in tg strains has been reported by [23, 25]), and (K) and (L) compare immunostaining for C3 of complement. Calibration bar: 50 µm.

Legend to figure 2:

Possible mechanism of inflammatory response in mouse models of AD and interactions between A β , microglia and the complement system.



Inflammation in transgenic mouse models

Table 1: Summary of studies on inflammation in mouse models of AD

Model	Experimental Intervention	Inflammatory Response	References
		COMPLEMENT	
SwDI		UP mRNA C1q, C3; UP Western C1q, C3, C4; colocal. complement w. fib Aβ and microglia	[22]
PS/APP		Colocal Claw AB/MG	[23]
Tg2576		UP C1g deposition w age	[25]
Tg2576	x C1q KO	DN glia activation, DN neurodegn, unchanged plaque load	[26]
Tg2576	x Clq KO	DN C4, UP C3	[25]
APP22	x C3 KO	UP plaque load, neuronal loss MG express CD45	[28]
APP22	x sCrry TG	UP A β deposition, UP neurodegn	[29]
APP/PS1	Vaccinia virus complement control protein	UP memory	[30]
Tg2576, 3xTg-AD	C5a receptor antagonist	DN A β deposition, neurodegn, gliosis, phospho tau, UP behavior	[27]
		MICROGLIA	
Tg2576		MG activation	[43]
APP 22		MG activation	[45]
APP 23		MG activation	[44]
TgCRND8		Early MG association with diffuse and compact $A\beta$	[46]
APP/PS1		UP Iba-1, unchanged CD11b in MG	[9, 49]
PS19 (tau)		MG activation	[48]
APP/PS1 and Tg2576	X	MG aggregate near plaques and phagocytose	[50, 51]
SwDI		MG express C1q, C3, C4	[22]
APP/PS1	G	MG express CR3 (CD11) at 6 mo and MHC-II at 12 mo	[52]
Tg2576	x CD40L deficient strain	DN microgliosis and Aβ load	[53, 54]
PSAPP, Tg2576	x CD40 or neutralizing CD40L antibody, Active Vacc.	Up Aβ clearance, UP MG phagocytosis	[55]
PSAPP and Tg2576	infusion with human umbilical cord blood cells	UP MG phagocytosis via suppression of CD40- CD40L interaction	[56]
Tg2576	x CD40L/CD40 KO	DN tau phosphorylation	[57]
Tg2576	x CCR2 KO	DN MG recruitment to plaques, UP A β aggregation, memory impairment	[58]
APP/PS1	x TLR2 KO	DN MG recruitment to plaques, UP A β aggregation, memory impairment	[59]
APP/PS1	x LXR KO	UP Abeta load	[60]
Tg2576	LXR agonist	DN Aβ deposits, DN CD45+ MG, DN IL-6 mRNA	[61]
APP/PS1	M-CFS	UP number of MG, UP A β phagocytosis, DN A β deposits	[62]
Tg2576	Exercise	DN IL-1 β A β , TNF- α , UP MHC-II	[63]
Tg2576	x Clq KO	DN MG activation, unchanged plaque load	[26], [25]

Inflammation in transgenic mouse models

APP	x C3 KO	UP MG CD45, DN MG F4/80 and CD68	[28]
APP/PS1	Glatimer acetate	UP MG CD11c, UP IGF, Aβ clearance	[65]
	vaccination		
Tg2576	LPS, intracerebral	UP MG activation, UP A β clearance	[66]
	Dexamethasone	Blocks activation	
SwDI	Minocycline	DN MHC-II MG, DN IL-6, UP memory, no change	[67]
		to Aβ deposition	
APPswe	Aged mice	UP MG expression of pro-inflammatory genes	[68]
	UP Abeta		5 (0]
APP/PS1	Aged mice	UP MG TNF- α , iNOS	[69]
	UP Abeta		[02]
APP/PS1	Pass. Vaccine	Activation CD45+ MG, clearance of AB	[83]
PDAPP ADD22	Pass. Vaccine	MG migrate to plaques	
APP23, $T_{\alpha}2576$	DNA vaccines	UP phagocytosis, number of MG, DN AB burden	[88, 89]
1 g2.5 / 0 A PD/PS1	Pass Vaccine	LID number of MG, but no abango in AB algorange	[00]
AII/ISI		CVTOKINES	
Τσ2576		UP IFN-v: II -12 and II -4 absent	[96]
Tg2576		$\begin{array}{c} \text{IP II} - 18 \text{ TGF} & \text{II} - 10 \text{ II} & 6 \end{array}$	[95]
Tg2576		UP II -1 & TGE- & II -6	[94]
Tg2576		UP TNE- α II -6 II -12p40 II -1 β II -1 α GM-	[97]
PS1/APPsw		CSF	
3xTg-AD		UP TNF- α MCP-1	[98]
3xTg-AD	dominant negative	DN AB deposition	[99]
8	TNF inhibitor		
APP23	x TNF-αR KO	DN A β deposition, neuronal loss	[100]
Tg2576	Exercise	DN TNF- α , IL-1 β , A β	[63]
-		UP IFN-γ, MIP-1 α, CD40, MHC-II, CD11c	
Tg2576	x IFN-γR 1 KO	DN TNF- α , BACE1, A β plaques	[101]
APPswe	x TLR-4 KO	DN TNF- α , IL-1 β , IL-10 and IL-17	[102]
APPswe/PS1	UP IL-1β	DN Αβ	[103]
hAPP	x TGF-β1 (AG)	DN Αβ	[104]
TGF-β1		UP A β (murine)	[105]
Tg2576	x CCL2 (MCP-1)	UP glial activation, $A\beta$ deposition	[107]
APP/PS1	AAV - dominant-	DN glial activation, $A\beta$ deposition	[107]
	negative CCL2		. –

AG: astroglia; DN: downregulation, reduction; MG: microglia; MIP: macrophage inflammatory protein; PassVaccine: passive vaccination; UP: upregulation, increase, improvement; x crossbreeding.

Inflammation in transgenic mouse models



46

Inflammation in transgenic mouse models



47