



Poly(ADP-ribose) polymerase-1 (PARP-1) gene deficiency alleviates diabetic kidney disease

Hanna Shevalye, Yury Maksimchyk, Pierre Watcho, Irina G. Obrosova

► To cite this version:

Hanna Shevalye, Yury Maksimchyk, Pierre Watcho, Irina G. Obrosova. Poly(ADP-ribose) polymerase-1 (PARP-1) gene deficiency alleviates diabetic kidney disease. *Biochimica et Biophysica Acta - Molecular Basis of Disease*, 2010, 1802 (11), pp.1020. 10.1016/j.bbadis.2010.07.004 . hal-00623313

HAL Id: hal-00623313

<https://hal.science/hal-00623313>

Submitted on 14 Sep 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript


Poly(ADP-ribose) polymerase-1 (PARP-1) gene deficiency alleviates diabetic kidney disease

Hanna Shevalye, Yury Maksimchyk, Pierre Watcho, Irina G. Obrosova

PII: S0925-4439(10)00131-6
DOI: doi: [10.1016/j.bbadis.2010.07.004](https://doi.org/10.1016/j.bbadis.2010.07.004)
Reference: BBADIS 63129

To appear in: *BBA - Molecular Basis of Disease*

Received date: 16 April 2010
Revised date: 1 July 2010
Accepted date: 5 July 2010



Please cite this article as: Hanna Shevalye, Yury Maksimchyk, Pierre Watcho, Irina G. Obrosova, Poly(ADP-ribose) polymerase-1 (PARP-1) gene deficiency alleviates diabetic kidney disease, *BBA - Molecular Basis of Disease* (2010), doi: [10.1016/j.bbadis.2010.07.004](https://doi.org/10.1016/j.bbadis.2010.07.004)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**POLY(ADP-RIBOSE) POLYMERASE-1 (PARP-1) GENE DEFICIENCY
ALLEVIATES DIABETIC KIDNEY DISEASE**

Hanna Shevalye, Yury Maksimchyk, Pierre Watcho, and Irina G.Obrosova

Pennington Biomedical Research Center,
Louisiana State University System, Baton Rouge, LA

Article type: Original article

Address for correspondence and reprint request:

Irina G.Obrosova, Ph.D.

Pennington Biomedical Research Center

Louisiana State University System

6400 Perkins Road

Baton Rouge, LA 70808

Tel. (225) 763-0276

FAX. (225) 763-0274

E-mail: obrosoig@pbrc.edu

ABSTRACT

Poly(ADP-ribose)polymerase (PARP) inhibitors prevent or alleviate diabetic nephropathy. This study evaluated the role for PARP-1 in diabetic kidney disease using the PARP-1-deficient mouse. PARP-1^{-/-} and the wild-type (129S1/SvImJ) mice were made diabetic with streptozotocin, and were maintained for 12 weeks. Final blood glucose concentrations were increased ~3.7-fold in both diabetic groups. PARP-1 protein expression (Western blot analysis) in the renal cortex was similar in non-diabetic and diabetic wild-type mice (100% and 107%) whereas all knockouts were PARP-1-negative. PARP-1 gene deficiency reduced urinary albumin (ELISA) and protein excretion, prevented diabetes-induced kidney hypertrophy, and decreased mesangial expansion and collagen deposition (both assessed by histochemistry) as well as fibronectin expression. Renal podocyte loss (immunohistochemistry) and nitrotyrosine and transforming growth factor- β_1 accumulations (both by ELISA) were slightly lower in diabetic PARP-1^{-/-} mice, but the differences with diabetic wild-type group did not achieve statistical significance. In conclusion, PARP-1^{-/-} gene deficiency alleviates although does not completely prevent diabetic kidney disease.

Key words: Albuminuria, collagen, diabetic nephropathy, mesangial expansion, poly(ADP-ribose) polymerase-1, poly(ADP-ribose)polymerase-1 deficient mouse, renal podocyte

1. INTRODUCTION

Diabetic nephropathy is one of the most severe complications of diabetes mellitus which develops in 30% to 40% of patients with both Type 1 and Type 2 diabetes mellitus [1,2], and is responsible for at least ~35% of all new cases of end-stage renal disease in the United States [3]. Despite a significant breakthrough in prevention and treatment of diabetic kidney disease in the last decade, due to development of angiotensin-converting enzyme inhibitors and angiotensin receptor blockers, there is still a vital need to discover and target novel pathophysiologic pathways. Recent studies in animal models identified NAD(P)H-oxidase [4], 12/15-lipoxygenase [5], AMP-activated protein kinase [6], fatty acid imbalances [7], and a number of other compelling drug targets for prevention and treatment of diabetic nephropathy.

Poly(ADP-ribose) polymerase-1 (PARP-1) a highly conserved protein of 116 kDa, catalyzes the cleavage of NAD^+ into nicotinamide and ADP-ribose [8]. PARP-1 belongs to a 18 member superfamily of PARP enzymes synthesizing poly(ADP-ribose) polymer which covalently attaches to acceptor proteins i.e., histones, DNA repair enzymes, transcription factors and PARPs themselves [8,9]. PARP-1 has been localized in cell nucleus [8] and mitochondria [10], whereas poly(ADP-ribosyl)ated proteins can also be found in other cell compartments because poly(ADP-ribosyl)ation occurs during protein (e.g., glyceraldehyde 3-phosphate dehydrogenase [11]) trafficking through the nucleus. PARP-1 activation is triggered by reactive oxygen and nitrogen-induced DNA single-strand breakage [8,9,12,13], and by several other factors [14,15], and affects multiple metabolic pathways, transcriptional regulation, and gene expression [8,9,16]. The list of PARP-1-regulated genes includes *interleukin (IL)-6*, *IL-1 β* , *intercellular adhesion molecule-1*, *vesicular adhesion molecule-1*,

c-myc, P- and E-selectins, granulocyte-macrophage colony-stimulating factor, NF- κ B1, NF- κ B2, c-rel, Ikk- α , Ikk- β , Ikk- γ , Ikk-i, I- κ B α , inducible nitric oxide synthase, interferon regulatory factor 1, interferon- β , MIF, Gro-1, IL-1 α , IL-2, IL-4, IL-5, IL-10, IL-12A, IL-12B, IL-16, IL-17, IL-18, LT- β , transforming growth factor (TGF)- α , TGF- β 1, TGF- β 2, TGF- β 3, tumor necrosis factor (TNF)- α , TNF- β , monocyte chemoattractant protein-1, NIK, endothelin-1 and endothelin receptors [16,17]. PARP-1 activation participates in apoptosis by stabilizing p53, by mediating the translocation of apoptosis-inducing factor from mitochondria to the nucleus, or by inhibiting early activation of DNases [8,9,18,19]. Evidence for the important role for PARP and, in particular, the PARP-1 isoform, activations in many major diseases has been obtained in experimental studies with PARP inhibitors and PARP-1-deficient mice and clinical studies and trials (reviewed in [8-10,20-22]). Recently, the PARP-1 isoform has been implicated in the pathogenesis of diabetes mellitus [8,9,23] and diabetic complications including endothelial dysfunction [24] and peripheral neuropathy [25,26]. The present study evaluated the role for PARP-1 in diabetic kidney disease using the PARP-1-deficient mouse and the multiple-dose streptozotocin model of diabetes.

2. MATERIALS AND METHODS

2.1. Reagents

Unless otherwise stated, all chemicals were of reagent-grade quality, and were purchased from Sigma Chemical Co., St. Louis, MO. Rabbit polyclonal anti-PARP-1 antibody was obtained from Enzo Life Sciences International, Inc., Plymouth Meeting, PA. Mouse monoclonal anti-poly(ADP-ribose) antibody was purchased from Trevigen, Inc., Gaithersburg, MD, rabbit polyclonal anti-Wilms tumor gene product-1 (WT1) antibody

from Santa Cruz Biotechnology, Inc., Santa Cruz, CA, and mouse monoclonal anti-fibronectin antibody from BD Transduction Laboratories, Lexington, KY. Other reagents for immunohistochemistry have been purchased from Vector Laboratories, Inc., Burlingame, CA.

2.2. Animals

The experiments were performed in accordance with regulations specified by the National Institutes of Health “Principles of Laboratory Animal Care, 1985 Revised Version” and Pennington Biomedical Research Center Protocol for Animal Studies. Male PARP-/- (129S-Parp1tm1Zqw/J) and the corresponding wild-type (129S1/SvImJ) mice were fed a standard rat chow (PMI Nutrition Int., Brentwood, MO) and had access to water ad libitum. The multiple low-dose streptozotocin (STZ)-diabetes was induced as described [25]. Blood samples for glucose measurements were taken from the tail vein ~48 h after the STZ injection and the day prior to the study termination. All mice with blood glucose levels ≥ 13.8 mM were considered diabetic. The duration of experiment was 12 weeks. At the end of the study, mice were placed in individual metabolic cages (Lab Products, Inc., Seaford, DE) and urine collected for 48 h. Urine specimen were centrifuged at 12,000 g (4°C, 10 min) and frozen for subsequent assessment of albumin by ELISA.

2.3. Anesthesia, euthanasia, and tissue sampling

The animals were sedated by CO₂, and immediately sacrificed by cervical dislocation. Kidneys were weighed. One kidney was immediately frozen in liquid nitrogen for subsequent Western blot analyses of PARP-1, poly(ADP-ribosyl)ated proteins, and fibronectin, and ELISA measurements of transforming growth factor-- β_1 (TGF- β_1), and nitrotyrosine (NT). The second kidney was fixed in normal buffered 4% formalin for further

assessment of collagen deposition, podocyte counts, and periodic acid-Schiff (PAS)-positive substance accumulation.

2.4. Specific methods

2.4.1. Urinary albumin, protein, creatinine, and renal TGF- β_1 , and NT

Urinary albumin was assessed by ELISA (AssayMax mouse albumin ELISA kit, Assaypro, St. Charles, MO). Urinary protein was measured with the bicinchoninic acid protein assay (Pierce Biotechnology, Rockford, IL). Urinary creatinine was measured spectrophotometrically, using Creatinine Parameter assay kit (R&D Systems, Minneapolis, MN). For measurements of TGF- β_1 and NT concentrations, renal cortex samples were homogenized on ice in RIPA buffer (1:10 w/v) containing 50 mM Tris-HCl, pH 7.2; 150 mM NaCl; 0.1% sodium dodecyl sulfate; 1% NP-40; 5 mM EDTA; 1 mM EGTA; 1% sodium deoxycholate and the protease/phosphatase inhibitors leupeptin (10 μ g/ml), aprotinin (20 μ g/ml), benzamidine (10 mM), phenylmethylsulfonyl fluoride (1 mM), sodium orthovanadate (1 mM). Homogenates were sonicated (3 x 5 s) and centrifuged at 14,000g (4°C, 20 min). TGF- β_1 and NT concentrations were measured with the Quantikine mouse/rat/porcine/canine TGF- β_1 kit (R&D Systems, Minneapolis, MN) and the OxiSelect Nitrotyrosine ELISA kit (Cell Biolabs, San Diego, CA). All the assays were performed in accordance with the manufacturers' instructions. Protein was measured with the bicinchoninic acid protein assay (Pierce Biotechnology, Rockford, IL).

2.4.2. Western blot analysis

Western blot analyses of PARP-1, poly(ADP-ribosyl)ated proteins, and fibronectin were performed as described previously [27,28]. Protein bands were visualized with the Amersham ECL Western blotting detection reagents and analysis system (GE Healthcare,

Buckinghamshire, UK). Membranes were then stripped and reprobed with β -actin antibody to verify equal protein loading. All poly(ADP-ribosyl)ated protein bands were used for calculations. The data were quantified by densitometry (Quantity One 4.5.0 software, Bio-Rad Laboratories, Richmond, CA), and the density of non-specific bands (obtained without primary antibody) subtracted from the results.

2.4.3. Immunohistochemical studies

All renal sections were processed by a single investigator and evaluated blindly. Collagen and PAS-positive substance stainings were performed by conventional histochemical methods. At least, ten fields of each section (~50 glomeruli) were examined to select one representative image. Color intensity of renal sections stained for PAS-positive substances was calculated using the ImageJ 1.43q software (National Institutes of Health, Bethesda, MD). 25-40 glomeruli were evaluated in a blinded fashion for each animal. 4- μ m renal sections were stained for collagen with Masson's trichrome as described [29], after which the percentage of areas positively stained for collagen was quantified in 10 randomly taken microphotographs of each section using the Threshold Colour plug-in of ImageJ 1.43q program, and the average per mouse was calculated. Podocyte nuclei were detected in 3- μ m sections with an anti-WT1 antibody and the ABC staining kit, and visualized with the DAB detection kit (both kits from Vector Laboratories Inc., Burlingame, CA). Podocyte numbers were counted per glomerular section, and 20-25 glomeruli were examined for each animal. Low power observations of renal sections were made using a Olympus BX41 microscope. Color images were captured with a MiniVID digital camera (LW Scientific, Inc, Lawrenceville, GA) at 1280 x 1024 resolution.

2.5. Statistical analysis

The results are expressed as Mean $\pm$ SEM. Data were subjected to equality of variance F test, and then to log transformation, if necessary, before one-way analysis of variance. Where overall significance ($p < 0.05$) was attained, individual between group comparisons were made using the Student-Newman-Keuls multiple range test. Significance was defined at $p \leq 0.05$. When between-group variance differences could not be normalized by log transformation (datasets for body weights and plasma glucose), the data were analyzed by the nonparametric Kruskal-Wallis one-way analysis of variance, followed by the Bonferroni/Dunn or Fisher's PLSD tests for multiple comparisons.

3. RESULTS

3.1. Effect of PARP-1 gene deficiency on body weights and glycemia.

The initial (prior to STZ administration) body weights were similar in both groups of wild-type mice as well as in both groups of PARP-1^{-/-} mice (Table 1). Weight gain was reduced in non-diabetic and diabetic PARP-1^{-/-} mice (19% and -16% vs 29% and 6% in non-diabetic and diabetic wild-type mice, respectively). PARP-1 gene deficiency did not affect the development of diabetes induced by multiple low-dose STZ administration. Initial blood glucose concentrations were 118% and 109% higher in diabetic wild-type and diabetic PARP-1^{-/-} mice, respectively, than in the non-diabetic controls. The progression of diabetic hyperglycemia was also similar in the two diabetic groups. Final blood glucose concentrations were increased by 269% and by 267% in diabetic wild-type and diabetic PARP-1^{-/-} mice compared with the corresponding controls. PARP-1 gene deficiency did not affect glycemia in non-diabetic mice.

3.2. Effect of PARP-1 gene deficiency on renal cortex poly(ADP-ribosyl)ation.

PARP-1 expression in renal cortex was 7% higher in diabetic wild-type mice, but the difference with the corresponding non-diabetic group did not achieve statistical significance ($p = 0.20$, Fig.1, A and B). Poly(ADP-ribosyl)ated proteins were clearly identifiable in all experimental groups (Fig.2, A and B). Renal cortex poly(ADP-ribosyl)ation was increased in both diabetic wild-type and diabetic PARP-1^{-/-} mice, compared with the corresponding controls (by 18% and by 14%, $p < 0.01$ and < 0.05 , respectively). No statistically significant differences in poly(ADP-ribosyl)ated protein abundance were found between diabetic wild-type and diabetic PARP-1^{-/-} mice ($p = 0.133$). Poly(ADP-ribosyl)ated protein level was 5% lower in non-diabetic PARP-1^{-/-} mice than in the corresponding wild-type mice, but the difference between the two groups did not achieve statistical significance ($p = 0.41$).

3.3. Effect of PARP-1 gene deficiency on water consumption, urination volume, and urinary albumin, protein, and creatinine excretions.

No statistically significant differences in either water consumption or urination volumes were found between non-diabetic wild-type and PARP-1^{-/-} mice (Table 2). Diabetes was associated with 6.2-fold and 40-fold increases in water consumption and urination volumes in wild-type mice. Interestingly and surprisingly, despite similar levels of hyperglycemia in diabetic wild-type and diabetic PARP-1^{-/-} mice, the latter displayed less dramatic elevation in both variables (2.7-fold and 8.5-fold increases compared with the corresponding non-diabetic group, $p < 0.01$ for both comparisons). Urinary albumin, protein, and creatinine excretions were increased 21-, 14-, and 12.5-fold, and 13-, 7-, and 12-fold, respectively, in diabetic wild-type and diabetic PARP-1^{-/-} mice, compared with the corresponding non-diabetic groups. Urinary albumin/creatinine and protein/creatinine ratios were increased 1.78- and 3-fold in diabetic wild-type mice, and 1.33- and 1.38-fold in

diabetic PARP-/- mice. The differences between the two diabetic groups for all the variables, except creatinine excretion, were of statistical significance ($p < 0.05$). PARP-1 gene deficiency did not affect urinary albumin, creatinine, and protein excretions, or urinary albumin/creatinine and protein/creatinine ratios in non-diabetic mice.

3.4. Effect of PARP-1 gene deficiency on diabetes-associated kidney hypertrophy, mesangial expansion, fibronectin expression, and podocyte loss.

PARP-1 gene deficiency did not affect kidney weight in non-diabetic mice (Fig. 3, A). Whereas diabetic wild-type mice displayed clearly manifest kidney hypertrophy (22% increase in kidney weight, compared with the non-diabetic group, $p < 0.01$), diabetic PARP-1-/- mice maintained the kidney weights indistinguishable from those in the corresponding non-diabetic controls. Because of a reduced weight gain, kidney-to-body weight ratios were increased in both diabetic groups; however, this increase was ~10% greater in diabetic wild-type than in diabetic PARP-1-/- mice ($p < 0.01$, Fig.3, B). PAS-positive substance staining intensity was increased by 52% in diabetic wild-type mice compared with the non-diabetic controls (Fig.4, A and B), indicative of a mesangial expansion, and this increase was blunted in diabetic PARP-/- mice (23%, $p < 0.01$ vs non-diabetic PARP-1-/- mice, and < 0.05 vs diabetic wild-type group). PARP-1 gene deficiency did not affect PAS-positive substance staining intensity in non-diabetic mice. Renal fibronectin expression was increased by 42% in diabetic wild-type mice, and by 20% in diabetic PARP-/- mice, compared with the corresponding non-diabetic groups ($p < 0.01$ for all comparisons, Fig.5, A and B). Both diabetic wild-type and diabetic PARP-1-/- mice displayed the loss of glomerular podocytes (to 75% and 83% vs corresponding non-diabetic groups, $p < 0.01$ for both comparisons, Fig.6, A and B). Although podocyte loss was 8% smaller in diabetic PARP-1-/- mice than in

the diabetic wild-type group, the difference did not achieve statistical significance ($p = 0.36$). Non-diabetic PARP-1^{-/-} mice contained 6% less podocytes in their glomeruli than non-diabetic wild-type mice ($p = 0.068$).

3.5 Effect of PARP-1 gene deficiency on diabetes-induced renal TGF- β accumulation, nitrosative stress, and glomerular collagen deposition.

Both diabetic wild-type and diabetic PARP-1^{-/-} mice displayed clearly manifest renal TGF- β accumulation (to 160% and 141% vs corresponding non-diabetic groups, $p < 0.01$ for both comparisons, Fig.7, A). PARP-1 gene deficiency was associated with an ~15% reduction in TGF- β concentration in non-diabetic mice, but the differences between the two non-diabetic and two diabetic groups did not achieve statistical significance ($p = 0.44$ and $p = 0.27$, respectively). Renal cortex NT concentrations were dramatically increased in both diabetic wild-type and diabetic PARP-1^{-/-} mice (to 282% and 258% compared with the corresponding non-diabetic groups, $p < 0.01$ for both comparisons, Fig.7, B). Collagen deposition in renal glomeruli was increased by 243 % and 137 % in diabetic wild-type and diabetic PARP-1^{-/-} mice vs corresponding non-diabetic groups ($p < 0.01$ for both comparisons, Fig.8, A and B). PARP-1^{-/-} gene deficiency was associated with a 26% reduction in diabetes-induced collagen deposition ($p < 0.01$ vs diabetic wild-type group), but did not affect this variable in the non-diabetic mice.

4. DISCUSSION

The present study revealed that PARP-1 gene deficiency was associated with alleviation of several manifestations of diabetic kidney disease i.e., albuminuria and proteinuria, kidney hypertrophy, glomerular mesangial expansion, fibronectin overexpression, and collagen deposition.

The effects of PARP-1 inhibitors (INO-1001, PJ34, 1,5-isoquinolinediol, and GPI-15,427) in previous reports [27,30,31] and PARP-1 gene deficiency in the present study on diabetic nephropathy are unidirectional, i.e., both PARP inhibition and PARP-1 gene deficiency are nephroprotective. However, whereas PARP inhibitors completely or essentially prevented albuminuria, renal mesangial expansion, and renal collagen deposition in STZ-diabetic rats [27], PARP-1 gene deficiency reduced, but did not completely abrogate, these changes in the STZ-diabetic mouse model. Furthermore, whereas PARP inhibition attenuated podocyte loss, TGF- β accumulation, and nitrosative stress [27], the effects of PARP-1 gene deficiency on those variables were non-significant. Both PARP in general and the PARP-1 isoform activations are known to contribute to multiple pathological conditions via two mechanisms. One of them, NAD⁺ depletion, affects the rate of the NAD-dependent glyceraldehyde 3-phosphate dehydrogenase reaction, which slows down glycolysis, electron transport, and ATP formation, eventually leading to functional impairment or cell death [8,9,20,21]. Another one, poly(ADP-ribosyl)ation, leads to changes in transcriptional regulation and gene expression [8,9]. In the present study, renal cortex poly(ADP-ribosyl)ation was clearly identifiable in both non-diabetic and diabetic PARP-1-deficient mice. Furthermore, poly(ADP-ribosyl)ation was increased in both groups of diabetic mice with a small and statistically non-significant differences between diabetic wild-type and diabetic PARP-1^{-/-} mice. The latter suggests that not only PARP-1, but also other PARPs, provide an important contribution to diabetes-induced poly(ADP-ribosyl)ation in the renal cortex. It is quite plausible that activation of PARP isoforms other than PARP-1 also contributes to NAD⁺ depletion and concomitant metabolic changes. A much more profound nephroprotective effect of PARP inhibition compared with PARP-1 gene deficiency may be

related to the lack of isoform specificity of existing PARP inhibitors. GPI-15,427, employed in our kidney studies in STZ-diabetic rats [27], as well as several other PARP inhibitors inhibit not only PARP-1, but also PARP-2 [32-34, and W.Xu, personal communication]. Other PARP inhibitors i.e., INO-1001, PJ34, and 1,5-isoquinolinediol, employed for diabetic nephropathy-related studies [27,30], have never been tested for their ability to inhibit PARP enzymes other than PARP-1. Thus, the present findings put forward the possibility of a contribution of other PARPs to diabetic nephropathy. Recently, selective PARP-2 inhibitors and PARP-2 deficient mice became available for exploring human disease [35-37]. Activation of PARP-2 was identified as an important player in colitis [38], post-ischemic brain injury [39] and cancer [40]. The roles for PARP-2 and other, different from PARP-1, PARP isoforms in diabetic complications have never been elucidated.

Another possible explanation for different efficacies of PARP inhibition (STZ-rat study [27]) and PARP-1/- gene deficiency in preventing biochemical and morphological manifestations of diabetic nephropathy lies in diverse susceptibility of rats and mice to toxic effects of STZ. STZ-diabetic rat is considered a good model for diabetic nephropathy-related studies because all manifestations of both early (albuminuria, mesangial expansion, podocyte loss) and advanced (glomerulosclerosis, tubulo-interstitial fibrosis) kidney disease are amenable to a variety of pharmacological interventions [7,27,41-44], which means that they do not develop because of STZ nephrotoxicity. In contrast, no good STZ-diabetic mouse model for diabetic nephropathy-related studies has been identified so far [45,46]. In the present study, diabetic 129S1/SvImJ mice developed robust albuminuria, kidney hypertrophy, and mesangial expansion, and modest podocyte loss. Further studies in this

model are needed to determine its susceptibility to STZ nephrotoxicity, and suitability for drug discovery in diabetic nephropathy.

In conclusion, PARP-1 activation is responsible for kidney hypertrophy and contributes to albuminuria, proteinuria, mesangial expansion, and collagen deposition in Type 1 diabetic kidney disease. However, the nephroprotective effect of PARP-1 gene deficiency was less pronounced than that of non-selective PARP inhibitors in previous studies. The current findings provide rationale for 1) development and further studies of PARP inhibitors and PARP inhibitor-containing combination therapies, for prevention and treatment of diabetic nephropathy; and 2) evaluation of the role of PARP isoforms, other than PARP-1, in the pathogenesis of this devastating complications of diabetes mellitus.

ACKNOWLEDGMENTS

The study was supported by the Juvenile Diabetes Research Foundation International Grant 1-2005-223, National Institutes of Health Grants DK070720, DK074517, and DK077141, and American Diabetes Association Grant (all to I.G.O.).

REFERENCES

- [1] M.E. Molitch, R.A. DeFronzo, M.J. Franz, W.F. Keane, C.E. Mogensen, H.H. Parving, M.W. Steffes; American Diabetes Association, Nephropathy in diabetes. *Diabetes Care* 27 (2004) S79-S83.
- [2] G.F. Strippoli, M. Craig, J.J. Deeks, F.P. Schena, J.C. Craig, Effects of angiotensin converting enzyme inhibitors and angiotensin II receptor antagonists on mortality and renal outcomes in diabetic nephropathy: systematic review, *BMJ* 329 (2004) 828.
- [3] R. Nelson, W. Knowler, D. Pettitt, P. Bennett, Kidney diseases in diabetes, in: National Institutes of Diabetes and Digestive and Kidney Diseases (Ed), *Diabetes in America*, NIH Publication No.95-1468, Bethesda, MD, 1995, pp. 349-385.
- [4] A.A. Eid, Y. Gorin, B.M. Fagg, R. Maalouf, J.L. Barnes, K. Block, H.E. Abboud, Mechanisms of podocyte injury in diabetes: role of cytochrome P450 and NADPH oxidases, *Diabetes* 58 (2009) 1201-1211.
- [5] H. Yuan, L. Lanting, Z.G. Xu, S.L. Li, P. Swiderski, S. Putta, M. Jonnalagadda, M. Kato, R. Natarajan, Effects of cholesterol-tagged small interfering RNAs targeting 12/15-lipoxygenase on parameters of diabetic nephropathy in a mouse model of type 1 diabetes, *Am. J. Physiol. Renal. Physiol.* 295 (2008) F605-F617.
- [6] M.J. Lee, D. Feliers, M.M. Mariappan, K. Sataranatarajan, L. Mahimainathan, N. Musi, M. Foretz, B. Viollet, J.M. Weinberg, G.G. Choudhury, B.S. Kasinath, A role for AMP-activated protein kinase in diabetes-induced renal hypertrophy, *Am. J. Physiol. Renal. Physiol.* 292 (2007) F617-F627.

- [7] J.H. Garman, S. Mulroney, M. Manigrasso, E. Flynn, C. Maric, Omega-3 fatty acid rich diet prevents diabetic renal disease, *Am. J. Physiol. Renal. Physiol.* 296 (2009) F306-F316.
- [8] P. Jagtap, C. Szabó, Poly(ADP-ribose) polymerase and the therapeutic effects of its inhibitors, *Nat. Rev. Drug Discov.* 4 (2005) 421-440.
- [9] L. Virág, Structure and function of poly(ADP-ribose) polymerase-1: role in oxidative stress-related pathologies, *Curr. Vasc. Pharmacol.* 3 (2005) 209-214.
- [10] M.N. Rossi, M. Carbone, C. Mostocotto, C. Mancone, M. Tripodi, R. Maione, P. Amati, Mitochondrial localization of PARP-1 requires interaction with mitofilin and is involved in the maintenance of mitochondrial DNA integrity, *J. Biol. Chem.* 284 (2009) 31616-31624.
- [11] X. Du, T. Matsumura, D. Edelstein, L. Rossetti, Z. Zsengellér, C. Szabó, M. Brownlee, Inhibition of GAPDH activity by poly(ADP-ribose) polymerase activates three major pathways of hyperglycemic damage in endothelial cells, *J. Clin. Invest.* 112 (2003) 1049-1057.
- [12] P. Pacher, J.S. Beckman, L. Liaudet, Nitric oxide and peroxynitrite in health and disease, *Physiol. Rev.* 87 (2007) 315-424.
- [13] C. Szabó, H. Ischiropoulos, R. Radi, Peroxynitrite: biochemistry, pathophysiology and development of therapeutics, *Nat. Rev. Drug Discov.* 6 (2007) 662-680.
- [14] T.M. Kauppinen, W.Y. Chan, S.W. Suh, A.K. Wiggins, E.J. Huang, R.A. Swanson, Direct phosphorylation and regulation of poly(ADP-ribose) polymerase-1 by extracellular signal-regulated kinases 1/2, *Proc. Natl. Acad. Sci. U S A* 103 (2006) 7136-1741.

- [15] S. Homburg, L. Visochek, N. Moran, F. Dantzer, E. Priel, E. Asculai, D. Schwartz, V. Rotter, N. Dekel, M. Cohen-Armon, A fast signal-induced activation of poly(ADP-ribose) polymerase: a novel downstream target of phospholipase C, *J. Cell Biol.* 150 (2000) 293-307.
- [16] H.C. Ha, L.D. Hester, S.H. Snyder, Poly(ADP-ribose) polymerase-1 dependence of stress-induced transcription factors and associated gene expression in glia, *Proc. Natl. Acad. Sci. U S A* 99 (2002) 3270-3275.
- [17] A.G. Minchenko, M.J. Stevens, L. White, O.I. Abatan, K. Komjáti, P. Pacher, C. Szabó, I.G. Obrosova, Diabetes-induced overexpression of endothelin-1 and endothelin receptors in the rat renal cortex is mediated via poly(ADP-ribose) polymerase activation, *FASEB J.* 17 (2003) 1514-1516.
- [18] S.W. Yu, H. Wang, M.F. Poitras, C. Coombs, W.J. Bowers, H.J. Federoff, G.G. Poirier, T.M. Dawson, V.L. Dawson, Mediation of poly(ADP-ribose) polymerase-1-dependent cell death by apoptosis-inducing factor, *Science* 297 (2002) 259-263.
- [19] S.W. Yu, S.A. Andrabi, H. Wang, N.S. Kim, G.G. Poirier, T.M. Dawson, V.L. Dawson, Apoptosis-inducing factor mediates poly(ADP-ribose) (PAR) polymer-induced cell death, *Proc. Natl. Acad. Sci. U S A* 103 (2006) 18314-18319.
- [20] P. Pacher, C. Szabó, Role of the peroxynitrite-poly(ADP-ribose) polymerase pathway in human disease, *Am. J. Pathol.* 173 (2008) 2-13.
- [21] P. Pacher, C. Szabó, Role of poly(ADP-ribose) polymerase 1 (PARP-1) in cardiovascular diseases: the therapeutic potential of PARP inhibitors, *Cardiovasc. Drug Rev.* 25 (2007) 235-260.

- [22] S. Love, R. Barber, G.K. Wilcock, Neuronal death in brain infarcts in man, *Neuropathol. Appl. Neurobiol.* 26 (2000) 55-66.
- [23] V. Burkart, Z.Q. Wang, J. Radons, B. Heller, Z. Herceg, L. Stingl, E.F. Wagner, H. Kolb, Mice lacking the poly(ADP-ribose) polymerase gene are resistant to pancreatic beta-cell destruction and diabetes development induced by streptozocin, *Nat. Med.* 5 (1999) 314-319.
- [24] F. Garcia Soriano, L. Virág, P. Jagtap, E. Szabó, J.G. Mabley, L. Liaudet, A. Marton, D.G. Hoyt, K.G. Murthy, A.L. Salzman, G.J. Southan, C. Szabó, Diabetic endothelial dysfunction: the role of poly(ADP-ribose) polymerase activation, *Nat. Med.* 7 (2001) 108-113.
- [25] I.G. Obrosova, F. Li, O.I. Abatan, M.A. Forsell, K. Komjáti, P. Pacher, C. Szabó, M.J. Stevens, Role of poly(ADP-ribose) polymerase activation in diabetic neuropathy, *Diabetes* 53 (2004) 711-720.
- [26] I.G. Obrosova, W. Xu, V.V. Lyzogubov, O. Ilnytska, N. Mashtalir, I. Vareniuk, I.A. Pavlov, J. Zhang, B. Slusher, V.R. Drel, PARP inhibition or gene deficiency counteracts intraepidermal nerve fiber loss and neuropathic pain in advanced diabetic neuropathy, *Free Radic. Biol. Med.* 44 (2008) 972-981.
- [27] V.R. Drel, W. Xu, J. Zhang, I.A. Pavlov, H. Shevalye, B. Slusher, I.G. Obrosova, Poly(Adenosine 5'-diphosphate-ribose) polymerase inhibition counteracts multiple manifestations of experimental type 1 diabetic nephropathy, *Endocrinology* 150 (2009) 5273-5283.

- [28] V.R. Drel, W. Xu, J. Zhang, P.F. Kador, T.K.Ali, J. Shin, U. Julius, B. Slusher, A.B. El-Remessy, I.G. Obrosova, Poly(ADP-ribose)polymerase inhibition counteracts cataract formation and early retinal changes in streptozotocin-diabetic rats. *Invest. Ophthalmol. Vis. Sci.* 50 (2009) 1778-1790.
- [29] J.L. Figarola, S. Loera, Y. Weng, N. Shanmugam, R. Natarajan, S. Rahbar, LR-90 prevents dyslipidaemia and diabetic nephropathy in the Zucker diabetic fatty rat, *Diabetologia* 51 (2008) 882-891.
- [30] C. Szabó, A. Biser, R. Benko, E. Böttinger, K. Suszták, Poly(ADP-ribose) polymerase inhibitors ameliorate nephropathy of type 2 diabetic Leprdb/db mice, *Diabetes* 55 (2006) 3004-3012.
- [31] H. Shevalye, R. Stavniichuk, W. Xu, J. Zhang, S. Lupachyk, Y. Maksimchyk, V.R. Drel, E.Z. Floyd, B. Slusher, I.G. Obrosova, Poly(ADP-ribose) polymerase (PARP) inhibition counteracts multiple manifestations of kidney disease in long-term streptozotocin-diabetic rat model, *Biochem. Pharmacol.* 79 (2010) 1007-1014.
- [32] T.D. Penning, G.D. Zhu, V.B. Gandhi, J. Gong, X. Liu, Y. Shi, V. Klinghofer, E.F. Johnson, C.K. Donawho, D.J. Frost, V. Bontcheva-Diaz, J.J. Bouska, D.J. Osterling, A.M. Olson, K.C. Marsh, Y. Luo, V.L. Giranda, Discovery of the Poly(ADP-ribose) polymerase (PARP) inhibitor 2-[(R)-2-methylpyrrolidin-2-yl]-1H-benzimidazole-4-carboxamide (ABT-888) for the treatment of cancer, *J. Med. Chem.* 52 (2009) 514-523.
- [33] T. Eltze, R. Boer, T. Wagner, S. Weinbrenner, M.C. McDonald, C. Thiemermann, A. Bürkle, T. Klein, Imidazoquinolinone, imidazopyridine, and isoquinolindione derivatives as novel and potent inhibitors of the poly(ADP-ribose) polymerase (PARP): a comparison with standard PARP inhibitors, *Mol. Pharmacol.* 74 (2008) 1587-1598.

- [34] K.A. Menear, C. Adcock, R. Boulter, X.L. Cockcroft, L. Copsey, A. Cranston, K.J. Dillon, J. Drzewiecki, S. Garman, S. Gomez, H. Javaid, F. Kerrigan, C. Knights, A. Lau, V.M. Loh Jr., I.T. Matthews, S. Moore, M.J. O'Connor, G.C. Smith, N.M. Martin, 4-[3-(4-cyclopropanecarbonylpiperazine-1-carbonyl)-4-fluorobenzyl]-2H-phthalazin-1-one: a novel bioavailable inhibitor of poly(ADP-ribose) polymerase-1, *J. Med. Chem.* 51 (2008) 6581-6591.
- [35] F. Moroni, L. Formentini, E. Gerace, E. Camaioni, D.E. Pellegrini-Giampietro, A. Chiarugi, R. Pellicciari, Selective PARP-2 inhibitors increase apoptosis in hippocampal slices but protect cortical cells in models of post-ischaemic brain damage, *Br. J. Pharmacol.* 157 (2009) 854-862.
- [36] R. Pellicciari, E. Camaioni, G. Costantino, L. Formentini, P. Sabbatini, F. Venturoni, G. Eren, D. Bellocchi, A. Chiarugi, F. Moroni, On the way to selective PARP-2 inhibitors. Design, synthesis, and preliminary evaluation of a series of isoquinolinone derivatives, *ChemMedChem* 3 (2008) 914-923.
- [37] A. Huber, P. Bai, J.M. de Murcia, G. de Murcia, PARP-1, PARP-2 and ATM in the DNA damage response: functional synergy in mouse development, *DNA Repair (Amst)* 3 (2004) 1103-1108.
- [38] I. Popoff, H. Jijon, B. Monia, M. Tavernini, M. Ma, R. McKay, K. Madsen, Antisense oligonucleotides to poly(ADP-ribose) polymerase-2 ameliorate colitis in interleukin-10-deficient mice, *J. Pharmacol. Exp. Ther.* 303 (2002) 1145-1154.
- [39] J. Kofler, T. Otsuka, Z. Zhang, R. Noppens, M.R. Grafe, D.W. Koh, V.L. Dawson, J.M. de Murcia, P.D. Hurn, R.J. Traystman, Differential effect of PARP-2 deletion on brain

injury after focal and global cerebral ischemia, *J. Cereb. Blood Flow Metab.* 26 (2006) 135-141.

[40] J. Yélamos, V. Schreiber, F. Dantzer, Toward specific functions of poly(ADP-ribose) polymerase-2, *Trends Mol. Med.* 14 (2008) 169-178.

[41] H. Trachtman, S. Futterweit, J. Maesaka, C. Ma, E. Valderrama, A. Fuchs, A.A. Tarectecan, P.S. Rao, J.A. Sturman, T.H. Boles, .M.-X. Fu, J. Baynes, Taurine ameliorates chronic streptozocin-induced diabetic nephropathy in rats, *Am. J. Physiol.* 269 (1995) F429-F438.

[42] M.F. Melhem, P.A. Craven, J. Liachenko, F.R. DeRubertis, Alpha-lipoic acid attenuates hyperglycemia and prevents glomerular mesangial matrix expansion in diabetes, *J. Am. Soc. Nephrol.* 13 (2002) 108-116.

[43] J.M. Forbes, V. Thallas, M.C. Thomas, H.W. Founds, W.C. Burns, G. Jerums, M.E. Cooper, The breakdown of preexisting advanced glycation end products is associated with reduced renal fibrosis in experimental diabetes, *FASEB J.* 17 (2003) 1762-1764.

[44] J.J. Wang, S.X. Zhang, R. Mott, R.R. Knapp, W. Cao, K. Lau, J.X. Ma, Salutary effect of pigment epithelium-derived factor in diabetic nephropathy: evidence for antifibrogenic activities, *Diabetes* 55 (2006) 1678-1685.

[45] M.D. Breyer, E. Böttinger, F.C. Brosius 3rd, T.M. Coffman, R.C. Harris, C.W. Heilig, K. Sharma; AMDCC, Mouse models of diabetic nephropathy, *J. Am. Soc. Nephrol.* 16 (2005) 27-45.

[46] F.C. Brosius 3rd, C.E. Alpers, E.P. Bottinger, M.D. Breyer, T.M. Coffman, S.B. Gurley, R.C. Harris, M. Kakoki, M. Kretzler, E.H. Leiter, M. Levi, R.A. McIndoe, K. Sharma, O. Smithies, K. Susztak, N. Takahashi, T. Takahashi; Animal Models of Diabetic

Complications Consortium, Mouse models of diabetic nephropathy, J. Am. Soc. Nephrol. 20 (2009) 2503-2512.

ACCEPTED MANUSCRIPT

Table 1.

Initial and final body weights and blood glucose concentrations in control and diabetic wild-type and poly(ADP-ribose) polymerase-1-deficient mice.

Variable	Body weight (g)		Blood glucose (mmol/l)	
	Initial	Final	Initial	Final
PARP-1+/+, C	24.2 ± 0.4	31.2 ± 1.0	6.7 ± 0.1	7.1 ± 0.3
PARP-1+/+, D	24.3 ± 0.5	25.6 ± 0.5**	14.6 ± 0.6**	26.2 ± 0.9**
PARP-1-/-, C	29.9 ± 1.0	35.5 ± 2.0	6.6 ± 0.2	7.3 ± 0.3
PARP-1-/-, D	30.6 ± 1.1	25.8 ± 0.7**	13.8 ± 0.6**	26.8 ± 1.7**

Initial and final blood glucose concentrations were measured after confirmation of diabetes (after 7-8 daily consecutive streptozotocin injections [33]) and at the 12-wk time point, respectively. Data are expressed as Means ± SEM. n = 10-15 per group. C – control mice; D – diabetic mice; PARP-1 – poly(ADP-ribose) polymerase-1. ** - p < 0.01 vs corresponding non-diabetic groups.

Table 2.

Water intake, urination volume, urinary albumin, total protein and creatinine excretions, albumin-to-creatinine and protein-to-creatinine ratios in control and diabetic wild-type and poly(ADP-ribose) polymerase-1-deficient mice.

Variable	Group	PARP-1+/+, C	PARP-1+/+, D	PARP-1-/-, C	PARP-1-/-, D
Water intake (ml)		4.6 ± 1.5	28.5 ± 5.2**	6.1 ± 1.7	16.3 ± 6.0**,##
Urination volume (ml)		0.53 ± 0.14	20.8 ± 4.4**	1.26 ± 0.40	10.7 ± 3.2**,##
Albuminuria (µg 48 h ⁻¹)		14.8 ± 4.2	314 ± 45**	15.5 ± 6.9	200 ± 53**,#
Proteinuria (mg 48 h ⁻¹)		22.0 ± 5.0	317 ± 41**	28.5 ± 7.1	203 ± 51**,#
Creatinine excretion (mg/48 h ⁻¹)		255 ± 44	3,187 ± 473**	287 ± 38	3,517 ± 570**
UA/CR (µg/mg)		0.09 ± 0.01	0.16 ± 0.02**	0.09 ± 0.01	0.12 ± 0.01**,#
UP/CR (mg/mg)		0.05 ± 0.01	0.15 ± 0.02**	0.08 ± 0.01	0.11 ± 0.01**,#

Data are expressed as Means ± SEM. n = 7-15 per group. C – control mice; D – diabetic mice; PARP-1 poly(ADP-ribose) polymerase-1; UA/CR – urinary albumin-to-creatinine ratio; UP/CR – urinary protein-to-creatinine ratio. ** - p < 0.01 vs corresponding non-diabetic groups; #,## - p < 0.05 and < 0.01 vs diabetic wild-type mice.

FIGURE LEGENDS

Fig.1. A) Representative Western blot analysis of renal cortex poly(ADP-ribose) polymerase-1 and B) poly(ADP-ribose)polymerase-1 content (densitometry), in control and diabetic wild-type and poly(ADP-ribose) polymerase-1-deficient mice. C – control; D – diabetic, PARP-1 – poly(ADP-ribose) polymerase-1. Mean $\pm$ SEM, n = 8-10 per group.

Fig.2. A) Representative Western blot analysis of renal cortex poly(ADP-ribosyl)ated proteins and B) poly(ADP-ribosyl)ated protein content (densitometry), in control and diabetic wild-type and poly(ADP-ribose)polymerase-1-deficient mice. C – control; D – diabetic, PARP-1 – poly(ADP-ribose) polymerase-1. Mean $\pm$ SEM, n = 9-12 per group. *, ** - p < 0.05 and < 0.01 vs corresponding non-diabetic groups.

Fig.3. A) Kidney weights and B) kidney weight-to-body weight ratios, in control and diabetic wild-type and poly(ADP-ribose)polymerase-1-deficient mice. C – control; D – diabetic, PARP-1 – poly(ADP-ribose) polymerase-1. Mean $\pm$ SEM, n = 10-15 per group. ** - p < 0.01 vs corresponding non-diabetic groups; ^{##} - p < 0.01 vs diabetic wild-type mice.

Fig.4. A) Representative microphotographs and B) color intensities of PAS-positive substance stainings in the renal cortex in control and diabetic wild-type and poly(ADP-ribose)polymerase-1-deficient mice. C – control; D – diabetic, PARP-1 – poly(ADP-ribose) polymerase-1. Magnification x 400. Mean $\pm$ SEM, n = 10 per group. ** - p < 0.01 vs corresponding non-diabetic groups; [#] - p < 0.05 vs diabetic wild-type mice.

Fig.5. A) Representative Western blot analysis of renal cortex fibronectin and B) fibronectin content (densitometry), in control and diabetic wild-type and poly(ADP-ribose)polymerase-1-deficient mice. C – control; D – diabetic, PARP-1 – poly(ADP-ribose) polymerase-1.

Mean $\pm$ SEM, n = 10-13 per group. *, ** - p < 0.05 and < 0.01 vs corresponding non-diabetic groups; ### - p < 0.01 vs diabetic wild-type mice.

Fig.6. A) Representative microphotographs of glomerular podocyte immunostaining and B) podocyte counts, in control and diabetic wild-type and poly(ADP-ribose)polymerase-1-deficient mice. C – control; D – diabetic, PARP-1 – poly(ADP-ribose) polymerase-1. Magnification x 400. Mean $\pm$ SEM, n = 10 per group. ** - p < 0.01 vs corresponding non-diabetic groups.

Fig.7. A) Transforming growth factor- β and B) nitrotyrosine concentrations, in the renal cortex in control and diabetic wild-type and poly(ADP-ribose)polymerase-1-deficient mice. C – control; D – diabetic, PARP-1 – poly(ADP-ribose) polymerase-1. Mean $\pm$ SEM, n = 9-14 per group. ** - p < 0.01 vs corresponding non-diabetic groups.

Fig.8. A) Representative microphotographs and B) percentage of positively stained for collagen area in the renal cortex in control and diabetic wild-type and poly(ADP-ribose)polymerase-1-deficient mice. C – control; D – diabetic, PARP-1 – poly(ADP-ribose) polymerase-1. Magnification x 200. Mean $\pm$ SEM, n = 8-10 per group. ** - p < 0.01 vs corresponding non-diabetic groups; ### - p < 0.01 vs diabetic wild-type mice.




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8

Research highlights

- PARP-1 activation is responsible for kidney hypertrophy and contributes to albuminuria, proteinuria, renal mesangial expansion, fibronectin overexpression, and collagen deposition in Type 1 diabetic kidney disease
- The nephroprotective effect of PARP-1 gene deficiency was less pronounced than that of non-selective PARP inhibitors in previous studies which suggests that the PARP isoform other than PARP-1 are also involved in the pathogenesis of diabetic nephropathy
- The findings provide rationale for development and further studies of PARP inhibitors and PARP inhibitor-containing combination therapies, for prevention and treatment of diabetic kidney disease