



HAL
open science

Errata to: On Kazhdan-Lusztig cells in type B

Cédric Bonnafé

► **To cite this version:**

Cédric Bonnafé. Errata to: On Kazhdan-Lusztig cells in type B. *Journal of Algebraic Combinatorics*, 2012, 35, pp.515-517. 10.1007/s10801-012-0351-7 . hal-00627160

HAL Id: hal-00627160

<https://hal.science/hal-00627160>

Submitted on 28 Sep 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ERRATA TO: ON KAZHDAN-LUSZTIG CELLS IN TYPE B

CÉDRIC BONNAFÉ

ABSTRACT. Based on a combinatorial result about the domino insertion algorithm [6], which appears to be false, the proof of our main result in [2] is not complete. We explain here what can be saved from [2]: more precisely, we explain what is the scope of validity of [2, Theorem 1.5].

In [2], we have found, using brute force computations, some (not all) Kazhdan-Lusztig relations (let us call them the *elementary relations*) between very particular elements of a Weyl group of type B . This shows in particular that the equivalence classes generated by the elementary relations are contained in Kazhdan-Lusztig cells.

It was announced in [6, Theorems 1.2 and 1.3] that the elementary relations generate the equivalence classes defined by the domino insertion algorithm (let us call them the *combinatorial cells*). As a consequence, we “deduced” that the combinatorial cells are contained in the Kazhdan-Lusztig cells [2, Theorem 1.5], thus confirming conjectures of Geck, Iancu, Lam and the author [3, Conjectures A and B]. However, as it was explained in a revised version of [6] (see [7]), the equivalence classes generated by the elementary relations are in general strictly contained in the combinatorial cells. This has no consequence on most of the intermediate results in [2], but changes the scope of validity of [2, Theorem 1.5]. Indeed, for some special cases of the parameters, T. Pietraho [5] has found that the elementary relations generate the combinatorial cells. So part of [2, Theorem 1.5] can be saved: the aim of this note is to explain precisely what is proved and what remains to be proved.

REMARK - The fact that [6, Theorems 1.2 and 1.3] is false does not imply that the result stated in [2, Theorem 1.5] is also false: it just means that its proof is

Date: September 28, 2011.

1991 Mathematics Subject Classification. According to the 2000 classification: Primary 20C08; Secondary 20C15.

The author is partly supported by the ANR (Project No JC07-192339).

not complete and we still expect the statement to be correct (as conjectured in [3, Conjectures A and B]).

1. PROVED AND UNPROVED RESULTS FROM [2]

Unproved results. We keep the notation of [2]. First of all, the proof of the Theorem stated in the introduction of [2], so its statement remains a conjecture (and similarly for the Corollary stated at the end of this introduction). Also, [2, Theorem 1.5(a)] is still a conjecture. However, [2, Theorem 1.5(b)] is still correct: its proof must only be adapted, using Pietraho's results [5].

Theorem 1. *Let $r \geq 0$ and assume that $b = ra > 0$. Let $? \in \{L, R, LR\}$ and $x, y \in W_n$ be such that $x \approx_?^r y$. Then $x \sim_? y$.*

The proof of Theorem 1 will be given in the next section. It must also be noted that [2, Theorem 1.5] is also valid if $b > (n - 1)a$ (see [4, Theorem 7.7] and [1, Corollaries 3.6 and 5.2]).

Proved results. Apart from the above mentioned results, all other intermediate results (about computations of Kazhdan-Lusztig polynomials, structure constants, elementary relations) are correct.

2. PROOF OF THEOREM 1

In [2, §7.1], we have introduced, following [6], three elementary relations \smile_1 , \smile_2^r and \smile_3^r : for adapting our argument to the setting of [5], we shall need to introduce another relation, which is slightly stronger than \smile_3^r .

Definition 2. *If w and w' are two elements of W_n , we shall write $w \frown_3^r w'$ whenever $w' = tw$ and $|w(1)| > |w(2)| > \dots > |w(r+2)|$. If $r \geq n-1$, then, by convention, the relation \frown_3^r never occurs.*

Using this definition, Pietraho's Theorem [5, Theorem 3.11] can be stated as follows:

Pietraho's Theorem. *The relation \approx_R^r is the equivalence relation generated by \smile_1 , \smile_2^r and \frown_3^{r-1} .*

It is easy to check that, if $w \frown_3^r w'$, then $w \smile_3^r w'$. Therefore, Theorem 1 follows from [2, Lemmas 7.1, 7.2 and 7.3] and the argument in [2, §7.2].

REFERENCES

- [1] C. BONNAFÉ, Two-sided cells in type B (asymptotic case), *J. Algebra* **304** (2006), 216–236.
- [2] C. BONNAFÉ, On Kazhdan-Lusztig cells in type B , *Journal of Alg. Comb.* **31** (2010), 53–82.
- [3] C. BONNAFÉ, M. GECK, L. IANCU & T. LAM, On domino insertion and Kazhdan-Lusztig cells in type B_n , *Representation theory of algebraic groups and quantum groups*, 33-54, *Progress in Mathematics* **284**, Birkhäuser/Springer, New York, 2010.
- [4] C. BONNAFÉ & L. IANCU, Left cells in type B_n with unequal parameters, *Represent. Theory* **7** (2003), 587–609.
- [5] T. PIETRAHO, Knuth relations for the hyperoctahedral groups, *Journal of Alg. Comb.* **29** (2009), 509-535.
- [6] M. TASKIN, Plactic relations for r -domino tableaux, preprint (2008), available at [arXiv:0803.1148v1](https://arxiv.org/abs/0803.1148v1).
- [7] M. TASKIN, Plactic relations for r -domino tableaux (revised version), preprint (2011), available at [arXiv:0803.1148v2](https://arxiv.org/abs/0803.1148v2).

INSTITUT DE MATHÉMATIQUES ET DE MODÉLISATION DE MONTPELLIER (CNRS: UMR 5149), UNIVERSITÉ MONTPELLIER 2, CASE COURRIER 051, PLACE EUGÈNE BATAILLON, 34095 MONTPELLIER CEDEX, FRANCE

E-mail address: `cedric.bonnafe@math.univ-montp2.fr`