



**HAL**  
open science

# Circle Diffeomorphisms: Quasi-reducibility and Commuting Diffeomorphisms

Mostapha Benhenda

► **To cite this version:**

Mostapha Benhenda. Circle Diffeomorphisms: Quasi-reducibility and Commuting Diffeomorphisms. 2011. hal-00628298v2

**HAL Id: hal-00628298**

**<https://hal.science/hal-00628298v2>**

Preprint submitted on 27 Feb 2012 (v2), last revised 28 Feb 2012 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Circle Diffeomorphisms: Quasi-reducibility and Commuting Diffeomorphisms

Mostapha Benhenda\*

February 27, 2012

## Abstract

In this article, we show two related results on circle diffeomorphisms. The first result is on quasi-reducibility: for a Baire-dense set of  $\alpha$ , for any diffeomorphism  $f$  of rotation number  $\alpha$ , it is possible to accumulate  $R_\alpha$  with a sequence  $h_n f h_n^{-1}$ ,  $h_n$  being a diffeomorphism. The second result is: for a Baire-dense set of  $\alpha$ , given two commuting diffeomorphisms  $f$  and  $g$ , such that  $f$  has  $\alpha$  for rotation number, it is possible to approach each of them by commuting diffeomorphisms  $f_n$  and  $g_n$  that are differentiably conjugated to rotations.

In particular, it implies that for  $\alpha$  in this Baire-dense set, and if  $\beta$  is an irrational number such that  $(\alpha, \beta)$  are not simultaneously Diophantine, the set of commuting diffeomorphisms  $(f, g)$  with singular conjugacy, and with rotation numbers  $(\alpha, \beta)$  respectively, is  $C^\infty$ -dense in the set of commuting diffeomorphisms with rotation numbers  $(\alpha, \beta)$ .

## 1 Introduction

It is well-known that there are circle diffeomorphisms with Liouville rotation numbers (i.e. non-Diophantine) that are not smoothly conjugated to rotations [1, 7, 8, 9]. A natural question arises, namely, the problem of smooth quasi-reducibility: *given a smooth diffeomorphism  $f$  of rotation number  $\alpha$ , is it possible to accumulate  $R_\alpha$  in the  $C^\infty$ -norm, with a sequence  $h_n^{-1} f h_n$ ,  $h_n$  being a smooth diffeomorphism?* In this case, we say that  $f$  is smoothly *quasi-reducible* to  $R_\alpha$ . Quasi-reducibility is a question that has been studied by Herman [7, pp.93-99], who showed that for any  $C^2$ -diffeomorphism  $f$  of irrational rotation number  $\alpha$ , it is possible to accumulate  $R_\alpha$  in the  $C^{1+bv}$ -norm, with a sequence  $h_n^{-1} f h_n$ ,  $h_n$  being a  $C^2$ -diffeomorphism (i.e. the total variation of  $D(h_n^{-1} f h_n - R_\alpha)$  converges towards zero). Quasi-reducibility is also related to a problem solved by Yoccoz [10], who showed that it is possible to accumulate a smooth diffeomorphism  $f$  in the  $C^\infty$ -norm with a sequence  $h_n R_\alpha h_n^{-1}$ ,  $h_n$  being a smooth diffeomorphism. However, these two problems are not the same, and the method used by Yoccoz does not directly yield our result. In our case, we determine a Baire-dense set of rotation numbers  $\alpha$  such that for any smooth diffeomorphism  $f$  of rotation number  $\alpha$ ,  $f$  is smoothly quasi-reducible.

---

\*Laboratoire d'Analyse, Geometrie et Applications, Paris 13 University, 99 Avenue J.B. Clement, 93430 Villetaneuse, France. Contact: mostaphabenhenda@gmail.com. I would like to thank the two anonymous referees for their remarks, which helped improving the article.

Connected to the problem of quasi-reducibility is the following question, raised by Mather: *given two commuting  $C^\infty$ -diffeomorphisms  $f$  and  $g$ , is it possible to approach each of them in the  $C^\infty$ -norm by commuting smooth diffeomorphisms that are smoothly conjugated to rotations?* In this paper, we determine a Baire-dense set of rotation numbers  $\alpha$  such that if  $f$  and  $g$  are commuting  $C^\infty$ -diffeomorphisms, with  $f$  of rotation number  $\alpha$ , then  $f$  and  $g$  are accumulated in the  $C^\infty$  norm by commuting  $C^\infty$ -diffeomorphisms that are  $C^\infty$ -conjugated to a rotation. This result is related to a theorem of Fayad and Khanin [6]. They showed that if  $(\alpha, \alpha')$  are simultaneously Diophantine (i.e. there is  $C_d > 0, \beta \geq 0$  such that for any  $p, p' \in \mathbb{Z}$ , any  $q \geq 1$ ,  $\max(|\alpha - p/q|, |\alpha' - p'/q|) \geq C_d/q^{2+\beta}$ ). This set includes some pairs  $(\alpha, \alpha')$  with  $\alpha$  and  $\alpha'$  Liouvillean), and if  $f$  and  $g$  are commuting  $C^\infty$ -diffeomorphisms, with  $f$  and  $g$  of rotation numbers  $\alpha$  and  $\alpha'$  respectively, then  $f$  and  $g$  are smoothly linearizable. Fayad and Khanin's result implies our result of quasi-reducibility in the particular case when the rotation numbers of  $f$  and  $g$  are simultaneously Diophantine. However, in general, our result is not implied by theirs. Indeed, our result holds for a set  $(\alpha, \alpha')$  that is Baire-dense in  $\mathbb{R}^2$  (because  $\alpha$  belongs to a Baire-dense set of  $\mathbb{R}$  and  $\alpha'$  is arbitrary), whereas the set of simultaneously Diophantine numbers is not Baire-dense.<sup>1</sup>

Moreover, for Diophantine rotation numbers, which are of full Lebesgue measure, the question of quasi-reducibility and Mather's problem are trivial, because in this case, the diffeomorphism  $f$  is smoothly conjugated to a rotation. Therefore, these two questions remain open for a meagre set of rotation numbers of zero Lebesgue measure.

In order to derive our results, we use estimates of the conjugacy to rotations of diffeomorphisms having rotation numbers of Diophantine constant type. These estimates were obtained in [2].

The circle is denoted  $\mathbb{T}^1$ . For  $r \in \mathbb{R}_+ \cup \{+\infty\}$ , we work in the universal cover  $D^r(\mathbb{T}^1)$ , which is the group of diffeomorphisms  $f$  of class  $C^r$  of the real line such that  $f - Id$  is  $\mathbb{Z}$ -periodic. For  $\alpha \in \mathbb{R}$ , we denote  $R_\alpha \in D^\infty(\mathbb{T}^1)$  the map  $x \mapsto x + \alpha$ .

Let  $f \in D^0(\mathbb{T}^1)$  be a homeomorphism and  $x \in \mathbb{R}$ . The sequence  $((f^n(x) - x)/n)_{n \geq 1}$  admits a limit independent of  $x$ , denoted  $\rho(f)$ . This limit is called the *rotation number* of  $f$ . This is a real number invariant by conjugacy.

**Theorem 1.1.** *There is a Baire-dense set  $A_1 \subset \mathbb{R}$  such that for any  $f \in D^\infty(\mathbb{T}^1)$  of rotation number  $\alpha \in A_1$ , there is a sequence  $h_n \in D^\infty(\mathbb{T}^1)$  such that  $h_n^{-1} f h_n \rightarrow R_\alpha$  in the  $C^\infty$ -topology.*

**Theorem 1.2.** *There is a Baire-dense set  $A_2 \subset \mathbb{R}$  such that for any  $f \in D^\infty(\mathbb{T}^1)$  of rotation number  $\alpha \in A_2$  and any  $g$  of class  $C^\infty$  with  $fg = gf$ ,  $f$  and  $g$  are accumulated in the  $C^\infty$ -topology by commuting  $C^\infty$ -diffeomorphisms that are  $C^\infty$ -conjugated to rotations.*

<sup>1</sup>The complementary in  $\mathbb{R}^2$  of simultaneously Diophantine numbers (noted  $SD^c$ ) is Baire-dense. Indeed, we have:

$$SD^c = \bigcap_{k \in \mathbb{N}^*} \bigcap_{n \in \mathbb{N}^*} \bigcup_{q \geq n} (A_{q,k} \times A_{q,k})$$

with:

$$A_{q,k} = \left\{ \alpha \in \mathbb{R} \mid \text{there is an integer } p \in \mathbb{Z}, \left| \alpha - \frac{p}{q} \right| < \frac{1}{q^k} \right\}.$$

$A_{q,k}$  is open (and so is  $A_{q,k} \times A_{q,k}$ ), and for any integer  $n$ ,  $\bigcup_{q \geq n} (A_{q,k} \times A_{q,k})$  is dense, because it contains all pairs of rational numbers (if  $\alpha = p_1/q_1$  and  $\alpha' = p_2/q_2$ , then  $(\alpha, \alpha') \in (A_{j_1 q_2, k} \times A_{j_1 q_2, k})$  for any  $j, k \in \mathbb{N}^*$ ). Therefore,  $SD^c$  is Baire-dense.

*Remark 1.3.* The proof of theorem 1.1 also gives that  $h_n R_\alpha h_n^{-1} \rightarrow f$  in the  $C^\infty$ -topology if  $\alpha \in A_1$ .

*Remark 1.4.* Combined with [6, p. 965], theorem 1.2 implies that if  $\alpha \in A_2$ , and  $(\alpha, \beta)$  are not simultaneously Diophantine, then  $S_{\alpha, \beta}$ , the set of couples  $(f, g)$  of smooth commuting circle diffeomorphisms with singular conjugacies to  $R_\alpha$  and  $R_\beta$  respectively, is  $C^\infty$ -dense in  $F_{\alpha, \beta}$ , the set of couples  $(f, g)$  of smooth commuting circle diffeomorphisms with rotation numbers  $\alpha$  and  $\beta$  respectively.

Indeed, our result shows that  $O_{\alpha, \beta}$  is  $C^\infty$ -dense in  $F_{\alpha, \beta}$ . Moreover, in [6, p. 965], for  $(\alpha, \beta)$  not simultaneously Diophantine, Fayad and Khanin described the construction of a couple of  $(f, g)$  of smooth commuting circle diffeomorphisms with singular conjugacies to  $R_\alpha$  and  $R_\beta$  respectively. This construction relies on the method of successive conjugacies, which can be made  $C^\infty$ -dense in  $O_{\alpha, \beta}$  [5].

## 2 Preliminaries

### 2.1 Basic properties

When the rotation number  $\alpha$  of  $f$  is irrational, and if  $f$  is of class  $C^2$ , Denjoy showed that  $f$  is topologically conjugated to  $R_\alpha$ . However, this conjugacy is not always differentiable. It depends on the Diophantine properties of the rotation number  $\alpha$ .

Let  $\alpha = a_0 + 1/(a_1 + 1/(a_2 + \dots))$  be the development of  $\alpha \in \mathbb{R}$  in continued fraction (see [4]). It is denoted  $\alpha = [a_0, a_1, a_2, \dots]$ . Let  $p_{-2} = q_{-1} = 0$ ,  $p_{-1} = q_{-2} = 1$ . For  $n \geq 0$ , we define integers  $p_n$  and  $q_n$  by:

$$p_n = a_n p_{n-1} + p_{n-2}$$

$$q_n = a_n q_{n-1} + q_{n-2}$$

We have  $q_0 = 1$ ,  $q_n \geq 1$  for  $n \geq 1$ . The rationals  $p_n/q_n$  are called the convergents of  $\alpha$ . Remember that  $q_{n+2} \geq 2q_n$ , for  $n \geq -1$ .

For any real number  $\beta \geq 0$ ,  $\alpha \in \mathbb{R} - \mathbb{Q}$  is Diophantine of order  $\beta$  and constant  $C_d$  (a set denoted  $DC(C_d, \beta)$ ) if there is a constant  $C_d > 0$  such that for any  $p/q \in \mathbb{Q}$ , we have:

$$\left| \alpha - \frac{p}{q} \right| > \frac{C_d}{q^{2+\beta}}$$

Each of the following relations characterizes  $DC(C_d, \beta)$ :

1.  $|\alpha - p_n/q_n| > C_d/q_n^{2+\beta}$  for any  $n \geq 0$
2.  $a_{n+1} < \frac{1}{C_d} q_n^\beta$  for any  $n \geq 0$
3.  $q_{n+1} < \frac{1}{C_d} q_n^{1+\beta}$  for any  $n \geq 0$
4.  $a_{n+1} > C_d \alpha_n^{1+\beta}$  for any  $n \geq 0$

$DC(C_d, 0)$  is the set of irrational numbers of *constant type*  $C_d$ . The first derivative of  $f \in D^1(\mathbb{T}^1)$  is denoted  $Df$ .

## 2.2 Some useful lemmas

For any  $n$  integer, let  $\alpha_n = [a_0, \dots, a_n, 1, \dots]$ .

Let  $V_\alpha(n) = \max_{0 \leq i \leq n} a_i$ . Observe that  $\alpha_n \in DC(1/V_\alpha(n), 0)$ . We will need the lemma:

**Lemma 2.1.** *Let  $\alpha$  be an irrational number,  $q_n$  its convergents and  $\alpha_n = [a_0, \dots, a_n, 1, \dots]$ . We have:*

$$|\alpha_n - \alpha| \leq \frac{2}{q_n^2} \leq \frac{4}{2^n}$$

*Proof.* Let  $\tilde{\alpha}_n = [a_0, \dots, a_n, 0, \dots]$ . By induction, we can show that  $\tilde{\alpha}_n = p_n/q_n$ . Moreover,  $\tilde{\alpha}_n$  is also the  $n^{\text{th}}$  convergent of  $\alpha_n$ . Therefore, by the best rational approximation theorem,  $|\alpha - p_n/q_n| \leq 1/q_n^2$  and  $|\alpha_n - p_n/q_n| \leq 1/q_n^2$ . Moreover, since  $q_{n+2} \geq q_n$ , then  $q_n \geq (\sqrt{2})^{n-1}$ . □

We need the lemma:

**Lemma 2.2.** *Let  $\phi : \mathbb{N} \rightarrow \mathbb{R}_+$  be such that  $\phi(n) \rightarrow_{n \rightarrow +\infty} +\infty$ . Let*

$$A = \{\alpha \in \mathbb{R} / V_\alpha(n) < \phi(n) \text{ for an infinity of } n\}$$

*Then  $A$  is Baire-dense.*

*Proof.* First, we show that for any positive integers  $n$  and  $i$ ,

$A_{i,n} = \{\alpha \text{ such that } a_i < \phi(n)\}$  is open. Let  $u(x) = \lfloor x \rfloor$ ,  $v(x) = \frac{1}{x}$  and  $w(x) = v(x) - u(v(x))$ . We have:  $a_{k+1} = v(w^k(x)) - w^{k+1}(x)$ . Since  $v$  is continuous and  $u$  is upper semi-continuous and non-negative, then  $w$  is lower semi-continuous. Moreover,  $w$  is non-negative. Therefore,  $w^k$  and  $w^{k+1}$  are also lower semi-continuous and non-negative. Since  $v$  is decreasing, then  $v \circ w^k - w^{k+1}$  is upper semi-continuous. We conclude that  $A_{i,n}$  is open.

Moreover, for any  $p \geq 0$ ,

$$\bigcup_{n \geq p} \bigcap_{i \leq n} A_{i,n}$$

is dense. Indeed, since  $\phi(n) \rightarrow +\infty$ , then it contains all numbers of constant type, which are dense. This set is also open and therefore,

$$A = \bigcap_{p \geq 0} \bigcup_{n \geq p} \bigcap_{i \leq n} A_{i,n}$$

is Baire-dense. □

## 2.3 Notations

- For any real numbers  $a$  and  $b$ ,  $a \vee b$  denotes  $\max(a, b)$ .
- For  $\phi$  a real  $\mathbb{Z}$ -periodic  $C^r$  function,  $0 \leq r < +\infty$ , we define:

$$\|\phi\|_r = \max_{0 \leq j \leq r} \max_{x \in \mathbb{R}} |D^j \phi(x)|$$

Note that for  $f, g \in D^r(\mathbb{T}^1)$ ,  $f - g$  is  $\mathbb{Z}$ -periodic, and for  $1 \leq j \leq r$ ,  $D^j f$  is  $\mathbb{Z}$ -periodic. For  $f \in D^r(\mathbb{T}^1)$ , we also define:

$$\|f\|_r = \max\left(\|f - id\|_0, \max_{1 \leq j \leq r} \|D^j f\|_0\right)$$

Note that the notation  $\|f\|_r$  is not a norm when  $f \in D^r(\mathbb{T}^1)$ , since  $D^r(\mathbb{T}^1)$  is not a vector space.

- In all the paper,  $C$  denotes a constant depending on  $u$ .  $W(f)$  denotes the total variation of  $\log Df$ , and  $Sf$  denotes the Schwartzian derivative of  $f$ .

## 2.4 Estimates of the conjugacy [2]

The following theorem gives an estimate of the linearization of a diffeomorphism having a rotation numbers of Diophantine constant type. This estimate, obtained in the previous chapter, is necessary to derive our results.

**Theorem 2.3.** *Let  $l \geq 3$  be an integer and  $\eta > 0$ . Let  $f \in D^l(\mathbb{T}^1)$  be of rotation number  $\alpha$ , such that  $\alpha$  is of constant type  $C_d$ . There exists a diffeomorphism  $h \in D^{l-1-\eta}(\mathbb{T}^1)$  conjugating  $f$  to  $R_\alpha$ , and a function  $B$  of  $C_d, l, \eta, W(f), \|Sf\|_{l-3}$ , which satisfy the estimate:*

$$\max\left(\frac{1}{\min Dh}, \|h\|_{l-1-\eta}\right) \leq B(C_d, l, \eta, W(f), \|Sf\|_{l-3}) \quad (1)$$

In particular, we remark that if  $f_n$  is a sequence of diffeomorphisms of rotation number  $\alpha_n$ , if the sequences  $W(f_n)$  and  $\|Sf_n\|_{l-3}$  are bounded (this will hold in our case, because we will take  $f_n = \lambda_n + f$  for a properly chosen  $\lambda_n \in \mathbb{R}$ ), if  $V_\alpha(n) \rightarrow +\infty$  and if  $h_n$  is the conjugacy to a rotation associated with  $f_n$ , then there is a real function  $E(V_\alpha(n))$  such that, for  $n$  sufficiently large, we have:

$$\max\left(\frac{1}{\min Dh_n}, \|h_n\|_{l-1-\eta}\right) \leq E(V_\alpha(n))$$

## 3 Quasi-Reducibility

**Theorem 3.1.** *Let  $l \geq 3$  be an integer,  $f \in D^l(\mathbb{T}^1)$  be of rotation number  $\alpha \in \mathbb{T}^1$ . Let  $\eta > 0$  be a real number. There exists a numerical sequence  $F(n)$ , going to  $+\infty$  as  $n \rightarrow +\infty$ , such that, if*

$$\liminf \frac{V_\alpha(n)}{F(n)} = 0$$

*then there is a sequence  $h_n$  of class  $C^{l-1-\eta}$  such that  $h_n^{-1} f h_n \rightarrow R_\alpha$  in the  $C^{l-2-\eta}$ -topology.*

By applying lemma 2.2, we obtain the corollary:

**Corollary 3.2.** *There is a Baire-dense set  $A_1 \subset \mathbb{R}$  such that if  $l \geq 3$  is an integer,  $f \in D^l(\mathbb{T}^1)$  of rotation number  $\alpha \in A_1$  and  $\eta > 0$ , then  $f$  is  $C^{l-2-\eta}$ -quasi-reducible: there is a sequence  $h_n \in D^{l-1-\eta}(\mathbb{T}^1)$  such that  $h_n^{-1} f h_n \rightarrow R_\alpha$  in the  $C^{l-2-\eta}$ -topology.*

*Proof of theorem 1.1.* We let  $\eta = l/3$  in corollary 3.2. Since  $f$  is smooth, then there is a sequence  $(h_{n,l})_{n \geq 0} \in D^\infty(\mathbb{T}^1)$  such that, for any integer  $l \geq 3$  fixed,

$$\|h_{n,l}^{-1}fh_{n,l} - R_\alpha\|_{2(\frac{l}{3}-1)} \rightarrow_{n \rightarrow +\infty} 0$$

In particular, there is  $n(l)$  such that:

$$\|h_{n(l),l}^{-1}fh_{n(l),l} - R_\alpha\|_{2(\frac{l}{3}-1)} \leq \frac{1}{l}$$

Let  $h_l = h_{n(l),l}$ . Let  $\epsilon > 0$ , and let  $k > 0$  be an integer. There is  $l_0 \geq 0$  such that for any  $l \geq l_0$ , we have:  $\epsilon \geq 1/l$ ,  $k \leq 2(\frac{l}{3} - 1)$  and:

$$\|h_l^{-1}fh_l - R_\alpha\|_k \leq \|h_l^{-1}fh_l - R_\alpha\|_{2(\frac{l}{3}-1)} \leq \frac{1}{l} \leq \epsilon$$

Therefore,  $h_l^{-1}fh_l \rightarrow_{l \rightarrow +\infty} R_\alpha$  in the  $C^k$ -topology, for any  $k$ , and therefore, this convergence holds in the  $C^\infty$ -topology.

□

### 3.1 The one-parameter family $R_\lambda f$

To prove theorem 3.1, we need to consider the one-parameter family  $R_\lambda f = f + \lambda$  [7, p.31]. We have the lemma:

**Lemma 3.3.** *Let  $l \geq 3$  be an integer,  $f \in D^l(\mathbb{T}^1)$ ,  $0 < \eta \leq l - 3$ ,  $\alpha = \rho(f)$ . Let  $\tilde{\alpha}$  be an irrational number of constant type. There exists  $\lambda_0 \in \mathbb{R}$  and a  $C^{l-1-\eta}$ -diffeomorphism  $h$  such that  $h^{-1}R_{\lambda_0}fh = R_{\tilde{\alpha}}$ . Moreover,*

$$\frac{|\lambda_0|}{\min Dh} \geq |\tilde{\alpha} - \alpha| \geq \frac{|\lambda_0|}{\|Dh\|_0}$$

*Proof.* Let  $\mu(\lambda) = \rho(R_\lambda f)$ .  $\mu$  is continuous, non-decreasing and  $\mu(\mathbb{R}) = \mathbb{R}$  (see [7, p. 31]). Therefore, there exists  $\lambda_0 \in \mathbb{R}$  such that  $\tilde{\alpha} = \rho(R_{\lambda_0}f)$ . Since  $\tilde{\alpha}$  is of constant type, there exists a  $C^{l-1-\eta}$ -diffeomorphism  $h$  such that  $h^{-1}R_{\lambda_0}fh = R_{\tilde{\alpha}}$  and that satisfies estimate (1) of theorem 2.3. By the mean value theorem, for any  $x$ , there is  $c(x)$  such that:

$$\tilde{\alpha} + x - h^{-1}fh(x) = R_{\tilde{\alpha}}(x) - h^{-1}fh(x) = h^{-1}R_{\lambda_0}fh(x) - h^{-1}fh(x) = D(h^{-1})(c(x))\lambda_0$$

By integrating this equation on an invariant measure of  $h^{-1}fh$ , we get lemma 3.3. Note that since  $h \in D^1(\mathbb{T}^1)$ , then  $Dh(x) > 0$  for any  $x$ , and  $\min Dh > 0$ .

□

### 3.2 The speed of approximation of $R_\alpha$

The proof of theorem 3.1 is also based on the lemma:

**Lemma 3.4.** *Let  $l \geq 3$  be an integer,  $f \in D^l(\mathbb{T}^1)$ ,  $0 < \eta \leq l - 3$ ,  $\alpha = \rho(f)$ . Let  $\tilde{\alpha}$  be an irrational number of constant type, and let  $\lambda_0 \in \mathbb{R}$  and  $h$  the  $C^{l-1-\eta}$ -diffeomorphism be given by lemma 3.3. Recall that  $C$  denotes a constant that only depends on  $u$ ,  $0 \leq u \leq l - 2 - \eta$ . We have the estimate:*

$$\|h^{-1}fh - R_\alpha\|_u \leq C\|f\|_u^C \|h\|_{u+1}^C \frac{1}{(\min Dh)^C} |\tilde{\alpha} - \alpha|$$

Before proving lemma 3.4, we show how theorem 3.1 is derived from it.

*proof of theorem 3.1.* If  $\alpha$  is of constant type, then  $f$  is reducible and there is nothing to prove. Therefore, we can suppose that  $V_\alpha(n) \rightarrow_{n \rightarrow +\infty} +\infty$ . By applying theorem 2.3, there exists a real function  $\tilde{F}$  strictly increasing with  $V_\alpha(n)$ , such that, for  $\alpha_n$  and for its associated diffeomorphism  $h_n$  given by lemma 3.3, we have, for  $n$  sufficiently large:

$$\|h_n^{-1} f h_n - R_\alpha\|_{l-2-\eta} \leq \exp(\tilde{F}(V_\alpha(n))) |\alpha_n - \alpha|$$

Let  $F(n) = \tilde{F}^{-1}(n^{1/2})$ . By extracting, we can suppose that  $\lim_{n \rightarrow +\infty} \frac{V_\alpha(n)}{F(n)} = 0$ . Therefore,  $V_\alpha(n) \leq F(n)$  for  $n$  sufficiently large and therefore,

$$\tilde{F}(V_\alpha(n)) \leq n^{1/2}$$

We get, for  $n$  sufficiently large,

$$\|h_n^{-1} f h_n - R_\alpha\|_{l-2-\eta} \leq e^{-\frac{n \log 2}{4}} \rightarrow_{n \rightarrow +\infty} 0$$

Hence theorem 3.1. □

Now, we show lemma 3.4:

*proof of lemma 3.4.* We need the Faa-di-Bruno formula (see e.g. [3]):

**Lemma 3.5.** *For every integer  $u \geq 0$  and functions  $\phi$  and  $\psi$  of class  $C^u$ , we have:*

$$D^u [\phi(\psi(x))] = \sum_{j=0}^u D^j \phi(\psi(x)) B_{u,j} (D\psi(x), D^2\psi(x), \dots, D^{(u-j+1)}\psi(x))$$

The  $B_{u,j}$  are the Bell polynomials, defined by  $B_{u,0} = 1$  and, for  $j \geq 1$ :

$$B_{u,j}(x_1, x_2, \dots, x_{u-j+1}) = \sum \frac{u!}{l_1! l_2! \dots l_{u-j+1}!} \left(\frac{x_1}{1!}\right)^{l_1} \left(\frac{x_2}{2!}\right)^{l_2} \dots \left(\frac{x_{u-j+1}}{(u-j+1)!}\right)^{l_{u-j+1}}$$

The sum extends over all sequences  $l_1, l_2, l_3, \dots, l_{u-j+1}$  of non-negative integers such that:  $l_1 + l_2 + \dots = j$  and  $l_1 + 2l_2 + 3l_3 + \dots = u$ .

Therefore, for any  $x$ , we have the estimate:

$$\left| B_{u,j} (D\psi(x), D^2\psi(x), \dots, D^{(u-j+1)}\psi(x)) \right| \leq C (1 \vee \|\psi\|_u^j) \quad (2)$$

Combining this estimate with lemma 3.5, we obtain the corollary:

**Corollary 3.6.** *For every integer  $u \geq 0$  and functions  $\phi$  and  $\psi$  of class  $C^u$ , we have:*

$$\|\phi \circ \psi\|_u \leq C \max_{0 \leq j \leq u} \|D^j \phi \circ \psi\|_0 (1 \vee \|\psi\|_u^u)$$



We apply this corollary to estimate  $\|h^{-1}\|_u$ . We let  $\phi(x) = 1/x$  and  $\psi = Dh \circ h^{-1}$ . We observe that  $D(h^{-1}) = \frac{1}{Dh \circ h^{-1}} = \phi \circ \psi$ . Since there is  $x_0$  such that  $Dh(x_0) = 1$ , then  $\|Dh\|_0 \geq 1$  (and we also have  $1 \geq \min Dh > 0$ ). Therefore, we get:

$$\|D(h^{-1})\|_u \leq C \max_{0 \leq j \leq u} \frac{1}{\|(Dh \circ h^{-1})^{j+1}\|_0} \|Dh \circ h^{-1}\|_u^C$$

By corollary 3.6, we also have:

$$\|Dh \circ h^{-1}\|_u \leq C \|Dh\|_u \|h^{-1}\|_u^C$$

By combining these two estimations, we get:

$$\|D(h^{-1})\|_u \leq C \frac{1}{(\min Dh)^C} \|Dh\|_u^C \|h^{-1}\|_u^C$$

We iterate this estimate to estimate  $\|h^{-1}\|_u$ , for  $u \geq 1$ . We get:

$$\|h^{-1}\|_{u+1} \leq C \frac{1}{(\min Dh)^C} \|h\|_{u+1}^C \|h^{-1}\|_1^C \quad (3)$$

Now, we estimate the  $C^u$ -distance of  $h^{-1}fh$  to  $R_\alpha$ . Let  $\tilde{\alpha}, \lambda_0$  be as in lemma 3.3. We have:

$$h^{-1}fh - R_\alpha = h^{-1}fh - h^{-1}R_{\lambda_0}fh + R_{\tilde{\alpha}} - R_\alpha$$

Therefore,

$$\|h^{-1}fh - R_\alpha\|_u \leq \|h^{-1}fh - h^{-1}R_{\lambda_0}fh\|_u + |\tilde{\alpha} - \alpha| \quad (4)$$

On the other hand, by the Faa-di-Bruno formula, we have:

$$D^u [h^{-1}fh - h^{-1}R_{\lambda_0}fh](x) = \sum_{j=0}^u B_{u,j} (D(fh)(x), \dots, D^{u-j+1}(fh)(x)) \\ [D^j(h^{-1})(fh(x)) - D^j(h^{-1})(fh(x) + \lambda_0)]$$

Since  $|D^j(h^{-1})(fh(x)) - D^j(h^{-1})(fh(x) + \lambda_0)| \leq \|D^{j+1}(h^{-1})\|_0 |\lambda_0|$ , then by applying estimate (2), we get:

$$\|h^{-1}fh - h^{-1}R_{\lambda_0}fh\|_u \leq C \|f \circ h\|_u^C \|h^{-1}\|_{u+1} |\lambda_0|$$

By applying corollary 3.6, we get:

$$\|h^{-1}fh - h^{-1}R_{\lambda_0}fh\|_u \leq C \|f\|_u^C \|h\|_u^C \|h^{-1}\|_{u+1} |\lambda_0|$$

By applying (3), we obtain:

$$\|h^{-1}fh - h^{-1}R_{\lambda_0}fh\|_u \leq C \|f\|_u^C \|h\|_u^C \frac{1}{(\min Dh)^C} \|h\|_{u+1}^C \|h^{-1}\|_1^C |\tilde{\alpha} - \alpha| \|Dh\|_0$$

$$\|h^{-1}fh - h^{-1}R_{\lambda_0}fh\|_u \leq C \|f\|_u^C \|h\|_{u+1}^C \frac{|\tilde{\alpha} - \alpha|}{(\min Dh)^C}$$

By estimate (4), we obtain:

$$\|h^{-1}fh - R_\alpha\|_u \leq C \|f\|_u^C \|h\|_{u+1}^C \frac{1}{(\min Dh)^C} |\tilde{\alpha} - \alpha| \quad (5)$$

Hence lemma 3.4.  $\square$

## 4 Application to commuting diffeomorphisms

**Theorem 4.1.** *There exists a numerical sequence  $G(n)$ , going to  $+\infty$  as  $n \rightarrow +\infty$ , such that, for any  $l \geq 3$  an integer,  $f \in D^l(\mathbb{T}^1)$  of rotation number  $\alpha \in \mathbb{R}$ ,  $\eta > 0$  and  $g$  of class  $C^l$  such that  $fg = gf$ , if*

$$\liminf \frac{V_\alpha(n)}{G(n)} = 0$$

*then there exists two sequences of diffeomorphisms  $f_n$  and  $g_n$  that are  $C^{l-1-\eta}$ -conjugated to rotations, such that  $f_n g_n = g_n f_n$ , and with  $f_n$  and  $g_n$  converging respectively towards  $f$  and  $g$  in the  $C^{l-2-\eta}$ -norm.*

**Corollary 4.2.** *There is a Baire-dense set  $A_2 \subset \mathbb{R}$  such that if  $l \geq 3$  is an integer,  $f \in D^l(\mathbb{T}^1)$  has a rotation number  $\alpha \in A_2$ ,  $g$  is of class  $C^l$  such that  $fg = gf$  and  $\eta \in \mathbb{R}_+$ , then there exists two sequences of diffeomorphisms  $f_n$  and  $g_n$  that are  $C^{l-1-\eta}$ -conjugated to rotations, such that  $f_n g_n = g_n f_n$  and with  $f_n$  and  $g_n$  converging respectively towards  $f$  and  $g$  in the  $C^{l-2-\eta}$ -norm.*

We derive theorem 1.2 from corollary 4.2 by following the same argument as in the proof of theorem 1.1.

### 4.1 The speed of approximation of $g$ by a linearizable and commuting diffeomorphism

To prove theorem 4.1, we consider  $(h_n)_{n \geq 0}$ , the sequence of conjugating diffeomorphisms constructed in the proof of theorem 3.1,  $(\lambda_n)_{n \geq 0}$  the associated sequence of real numbers such that  $f_n = R_{\lambda_n} f = h_n R_{\alpha_n} h_n^{-1}$ . We also consider  $g'_n = h_n^{-1} g h_n$  and  $g_n = h_n R_{g'_n(0)} h_n^{-1}$ . The diffeomorphisms  $f_n$  and  $g_n$  commute, and  $f_n \rightarrow f$  in the  $C^{l-2-\eta}$ -norm. To prove theorem 4.1, it suffices to show that  $g_n \rightarrow g$  in the  $C^{l-2-\eta}$ -norm. This convergence is based on the lemma:

**Lemma 4.3.** *Let  $l \geq 3$  be an integer,  $f \in D^l(\mathbb{T}^1)$  of rotation number  $\alpha \in \mathbb{R}$ ,  $\eta > 0$ ,  $0 \leq u \leq l - 2 - \eta$ , and  $g \in D^l(\mathbb{T}^1)$  be such that  $fg = gf$ . Let  $(q_r)_{r \geq 0}$  be the sequence of denominators of the convergents of  $\alpha$ , and let  $r \geq 0$  be an integer. Let  $\tilde{\alpha}$  be an irrational number of constant type,  $\lambda_0 \in \mathbb{R}$  the associated number and  $h$  the associated  $C^{l-1-\eta}$  diffeomorphism given by lemma 3.3. Let  $f' = h^{-1} f h$  and  $g' = h^{-1} g h$ . We have the estimate:*

$$\|g - h R_{g'(0)} h^{-1}\|_u \leq C \|h\|_{u+1}^C \|f'\|_u^C \|g'\|_{u+1}^C \left( \frac{1}{q_r} + |\tilde{\alpha} - \alpha| \left( \frac{(C \|h\|_{u+1} \|f'\|_{u+1})^{C q_r}}{(\min Dh)^C} \right) \right)$$

*Proof of theorem 4.1.* Assuming lemma 4.3, we show theorem 4.1.

Let  $\tilde{\alpha} = \alpha_n$  and  $h_n$  be the associated diffeomorphism given by lemma 3.3. Since  $V_\alpha(n) \rightarrow +\infty$ , by applying the estimate for the conjugacy  $h_n$ , there exists  $\tilde{G}(x)$  strictly increasing with  $x$  such that, for  $n$  sufficiently large:

$$\|g - h_n R_{g'_n(0)} h_n^{-1}\|_{l-2-\eta} \leq e^{C \tilde{G}(V_\alpha(n) q_r)} \left( \frac{1}{q_r} + \frac{e^{C \tilde{G}(V_\alpha(n) q_r)}}{2^n} \right)$$

Moreover, since  $q_n = a_n q_{n-1} + q_{n-2}$ , and  $q_{n-2} \leq q_{n-1}$ , then

$$(\sqrt{2})^{n-1} \leq q_n \leq \prod_{k=1}^n (a_k + 1) \quad (6)$$

Therefore, we get:

$$\|g - h_n R_{g'_n(0)} h_n^{-1}\|_{l-2-\eta} \leq e^{C\tilde{G}(V_\alpha(n)) - \frac{1}{2}(r-1)\log 2} + e^{C\tilde{G}(V_\alpha(n)) + C\tilde{G}(V_\alpha(n))(V_\alpha(r)+1)^{r-n}\log 2} \quad (7)$$

Let  $G(n) = \tilde{G}^{-1}((\log n)^{1/2})$ . By extracting in the sequence  $V_\alpha(n)/G(n)$ , we can suppose that:

$$\frac{V_\alpha(n)}{G(n)} \rightarrow 0$$

Therefore, for  $n$  sufficiently large, we have:

$$\tilde{G}(V_\alpha(n)) \leq (\log n)^{1/2}$$

Moreover, for  $n$  sufficiently large, we can take an integer  $r_n$  such that:

$$(\log n)^{3/4} \leq r_n \leq (\log n)^{7/8}$$

we get:

$$(V_\alpha(r_n) + 1)^{r_n} = e^{r_n \log(V_\alpha(r_n)+1)} \leq e^{(\log n)^{15/16}}$$

The first term in estimate (7) tends towards 0. Moreover, since, for  $n$  sufficiently large,

$$(\log n)^{1/2} e^{(\log n)^{15/16}} \leq \frac{n}{2} \log 2$$

then the second term also tends towards 0. Hence theorem 4.1.  $\square$

## 4.2 Higher-order analogous of the mean value theorem

*Proof of lemma 4.3.* We need two higher-order analogous of the mean value theorem. The first one is:

**Lemma 4.4.** *Let  $u \geq 0$ ,  $s, t \in D^u(\mathbb{T}^1)$ . Let  $\delta \in \mathbb{R}$ . We have:*

$$\|st - R_\delta t\|_u \leq C \|s\|_{u+1} \|s - R_\delta\|_u \|t\|_u^u$$

Observe the presence of the term  $\|s\|_{u+1}$ , which is absent in the mean value formula. This is because of the estimate (2) on the Bell polynomial, in the Faa-di-Bruno formula.

*Proof.* If  $u = 0$ , the estimate is trivial. We suppose  $u \geq 1$ . For any  $x \in \mathbb{R}$ , the Faa-di-Bruno formula gives:

$$D^u(st)(x) - D^u(R_\delta t)(x) = \sum_{j=0}^u \left( (D^j s)(t(x)) - (D^j R_\delta)(t(x)) \right) B_{u,j} \left( Dt(x), \dots, D^{u-j+1} t(x) \right)$$

Therefore, by estimate (2), and since  $\|t\|_u \geq 1$ ,

$$|D^u(st)(x) - D^u(R_\delta t)(x)| \leq C\|s\|_{u+1}\|s - R_\delta\|_u\|t\|_u^u$$

Hence lemma 4.4. □

The second higher-order analogous of the mean value theorem is:

**Lemma 4.5.** *Let  $u \geq 0$ ,  $s \in D^{u+1}(\mathbb{T}^1)$ ,  $t \in D^u(\mathbb{T}^1)$ ,  $\delta \in \mathbb{R}$ . We have:*

$$\|st - sR_\delta\|_u \leq C\|s\|_{u+1}\|t\|_u^u\|t - R_\delta\|_u$$

Observe the presence of the term  $\|t\|_u$ , which is absent in the mean value formula. As in lemma 4.4, this is because of an estimate on the Bell polynomial, in the Faa-di-Bruno formula.

*Proof.* If  $u = 0$ , the estimate holds. We suppose  $u \geq 1$ . We use the following lemma:

**Lemma 4.6.** *Let  $u \geq 1$ ,  $j \leq u$  be integers and  $a_1, \dots, a_{u-j+1}, x_1, \dots, x_{u-j+1} \geq 0$ . Let  $x \geq \max\{|x_k| \vee 1; 1 \leq k \leq u - j + 1\}$  and let  $a \geq \max\{|a_k|; 1 \leq k \leq u - j + 1\}$ . Let  $B_{u,j}$  be a Bell polynomial. We have:*

$$|B_{u,j}(x_1 + a_1, \dots, x_{u-j+1} + a_{u-j+1}) - B_{u,j}(x_1, \dots, x_{u-j+1})| \leq Ca(x + a)^u$$

*Proof.* Let  $p \geq 1$  and  $l_1, \dots, l_p$  be integers. Then we have:

$$(x_1 + a_1)^{l_1} \dots (x_p + a_p)^{l_p} - x_1^{l_1} \dots x_p^{l_p} = \sum_{i=1}^p x_1^{l_1} \dots x_{i-1}^{l_{i-1}} (x_i + a_i)^{l_i} \dots (x_p + a_p)^{l_p} - x_1^{l_1} \dots x_i^{l_i} (x_{i+1} + a_{i+1})^{l_{i+1}} \dots (x_p + a_p)^{l_p}$$

$$(x_1 + a_1)^{l_1} \dots (x_p + a_p)^{l_p} - x_1^{l_1} \dots x_p^{l_p} = \sum_{i=1}^p x_1^{l_1} \dots x_{i-1}^{l_{i-1}} (x_{i+1} + a_{i+1})^{l_{i+1}} \dots (x_p + a_p)^{l_p} \left[ (x_i + a_i)^{l_i} - x_i^{l_i} \right]$$

(with the conventions  $x_1^{l_1} \dots x_0^{l_0} = 1$  and  $x_{p+1}^{l_{p+1}} \dots x_p^{l_p} = 1$ ).

Since  $(x_i + a_i)^{l_i} - x_i^{l_i} \leq l_i |a_i| (|x_i| + |a_i|)^{l_i - 1} \leq l_i a (|x_i| + a)^{l_i - 1}$ ,  $1 \leq l_i \leq u$  and  $x + a \geq 1$  (because  $x \geq 1$ ), we obtain:

$$|B_{u,j}(x_1 + a_1, \dots, x_{u-j+1} + a_{u-j+1}) - B_{u,j}(x_1, \dots, x_{u-j+1})| \leq a(u - j + 1)u B_{u,j}(x + a, \dots, x + a)$$

By the formula giving the Bell polynomials, we have:

$$B_{u,j}(x + a, \dots, x + a) \leq C(x + a)^u$$

□

To show lemma 4.5, For any  $0 \leq v \leq u$ , we write:

$$D^v(st)(x) - D^v(sR_\delta)(x) = \sum_{j=0}^v D^j s(t(x)) \left[ B_{v,j}(Dt(x), \dots, D^{v-j+1}t(x)) - B_{v,j}(DR_\delta(x), \dots, D^{v-j+1}R_\delta(x)) \right] +$$

$$\left[ D^j s(t(x)) - D^j s(R_\delta(x)) \right] B_{v,j} \left( DR_\delta(x), \dots, D^{v-j+1} R_\delta(x) \right)$$

We apply lemma 4.6 with  $a = \|t - R_\delta\|_u$  and  $x = \|R_\delta\|_u \geq 1$ . Since  $t \in D^u(\mathbb{T}^1)$ , then  $\|t\|_u \geq 1$ . We get:

$$\left| B_{v,j} \left( Dt(x), \dots, D^{v-j+1} t(x) \right) - B_{v,j} \left( DR_\delta(x), \dots, D^{v-j+1} R_\delta(x) \right) \right| \leq C \|t - R_\delta\|_u (1 + \|t - R_\delta\|_u)^u$$

$$\left| B_{v,j} \left( Dt(x), \dots, D^{v-j+1} t(x) \right) - B_{v,j} \left( DR_\delta(x), \dots, D^{v-j+1} R_\delta(x) \right) \right| \leq C \|t - R_\delta\|_u (2 + \|t\|_u)^u \leq C \|t - R_\delta\|_u \|t\|_u^u$$

□

### 4.3 Successive estimations

To prove lemma 4.3, we also need these successive estimations:

**Lemma 4.7.** *Let  $l \geq 3$  be an integer,  $f \in D^l(\mathbb{T}^1)$  of rotation number  $\alpha \in \mathbb{R}$ ,  $\eta > 0$ ,  $0 \leq u \leq l - 2 - \eta$ , and  $g \in D^l(\mathbb{T}^1)$  be such that  $fg = gf$ . Let  $(q_t)_{t \geq 0}$  be the sequence of denominators of the convergents of  $\alpha$ . Let  $\tilde{\alpha}$  be an irrational number of constant type,  $\lambda_0 \in \mathbb{R}$  the associated number and  $h$  the associated  $C^{l-1-\eta}$  diffeomorphism given by lemma 3.3. Let  $f' = h^{-1}fh$  and  $g' = h^{-1}gh$ . We have the estimates:*

$$A_{1,u} = \|h^{-1}\|_u \leq C \|h\|_u^C \frac{1}{(\min Dh)^C} \quad (8)$$

$$A_{2,u} = \|f'\|_u \leq CA_{1,u} \|f\|_u^C \|h\|_u^C \quad (9)$$

$$A_{3,u}(m) = \|f'^m\|_u \leq C^m A_{2,u}^{mC} \quad (10)$$

$$A_{4,u} = \|f' - R_\alpha\|_u \leq C \|h\|_{u+1}^C \|f\|_u^C \frac{1}{(\min Dh)^C} |\tilde{\alpha} - \alpha| \quad (11)$$

$$A_{5,u}(m) = \|f'^m - R_{m\alpha}\|_u \leq mCA_{4,u} A_{2,u}^C \max_{k \leq m-1} A_{3,u+1}(k) \quad (12)$$

$$A_{6,u} = \|g'\|_u \leq CA_{1,u} \|g\|_u^C \|h\|_u^C \quad (13)$$

and for any integer  $r \geq 0$ , we have:

$$A_{7,u} = \|g' - R_{g'(0)}\|_u \leq \frac{A_{6,u+1} + 1}{q_r} + \max_{m \leq 2q_r} \left( A_{6,u+1} A_{3,u}^C(m) A_{5,u}(m) + A_{6,u}^C A_{3,u+1}(m) A_{5,u}(m) \right) \quad (14)$$

$$A_{8,u} = \|g'h^{-1} - R_\alpha h^{-1}\|_u \leq CA_{6,u+1} A_{7,u} A_{1,u}^C \quad (15)$$

$$A_{9,u} = \|hg'h^{-1} - hR_{g'(0)}h^{-1}\|_u \leq C \|g\|_u^C A_{8,u} A_{1,u}^C \|h\|_{u+1} \quad (16)$$

The crucial estimate is (14), which is obtained by approaching modulo 1 each  $x \in \mathbb{R}$  by a  $m(x)\alpha$ , with  $m(x) \leq q_r$ . If  $q_r$  increases,  $x - m(x)\alpha$  is smaller modulo 1, but the bound on  $A_{3,u}(m(x))$  and  $A_{5,u}(m(x))$  increases. In the proof of theorem 4.1, we make a proper choice of  $r$  (and  $q_r$ ).

estimate (11) corresponds to estimate (5) of the proof of the result of quasi-reducibility.

The other estimations, namely, estimations (8),(9),(10), (12),(13), (15) and (16) are derived from applications of the Faa-di-Bruno formula: either corollary 3.6, lemma 4.4 or lemma 4.5.

*Proof of lemma 4.7.* For  $A_{1,u}$ , by estimate (3), we have:

$$\|h^{-1}\|_u \leq C \|h\|_u^C \frac{1}{(\min Dh)^C}$$

Hence estimate (8).

For  $A_{2,u}$ , by applying corollary 3.6 twice, we have,

$$\|f'\|_u \leq CA_{1,u} \|f\|_u^C \|h\|_u^C$$

Hence estimate (9).

For  $A_{3,u}$ , by applying corollary 3.6 again, we have, for any  $m$ ,

$$\|f'^{m+1}\|_u \leq C \|f'^m\|_u \|f'\|_u^C$$

and therefore, by iteration, we get:

$$\|f'^m\|_u \leq C^m \|f'\|_u^{mC}$$

Hence (10).

estimate (11) is a direct application of estimate (5).

For estimate (12), we observe that for any  $0 \leq v \leq u$ :

$$D^v f'^m - D^v R_{m\alpha} = D^v \sum_{k=0}^{m-1} f'^{m-k} R_{k\alpha} - f'^{m-k-1} R_{(k+1)\alpha}$$

$$D^v f'^m - D^v R_{m\alpha} = \sum_{k=0}^{m-1} D^v (f'^{m-k-1} f') R_{k\alpha} - D^v (f'^{m-k-1} R_\alpha) R_{k\alpha}$$

By applying lemma 4.5, and by noting that for any  $k$ ,  $\|f'^{m-k-1}\|_{u+1} \leq \max_{0 \leq k \leq m-1} \|f'^k\|_{u+1}$ , we get:

$$\|f'^m - R_{m\alpha}\|_u \leq mC \|f'\|_u^C \max_{0 \leq k \leq m-1} \|f'^k\|_{u+1} \|f' - R_\alpha\|_u$$

Hence (12).

For  $A_{6,u}$ , estimate (13) is the same as (9):

$$\|g'\|_u \leq C \|h^{-1}\|_u \|g\|_u^C \|h\|_u^C$$

Hence (13).

For  $A_{7,u}$ , let  $m \geq 0$  and  $u \geq v \geq 1$ . For any  $x$ ,  $D^v R_\alpha(x) = \int_0^1 D^v g'(y) dy$ . Therefore,

$$\begin{aligned} |D^v g'(x) - D^v R_\alpha(x)| &= \left| D^v g'(x) - \int_0^1 D^v g'(y) dy \right| = \\ & \left| \int_0^1 (D^v g'(x) - D^v g'(y)) dy \right| \leq \max_{x,y \in [0,1]} |D^v g'(x) - D^v g'(y)| \end{aligned}$$

On the other hand, we have:

$$\begin{aligned} D^v g'(x) - D^v g'(y) &= D^v g'(x) - D^v g'(y + m\alpha) + D^v g'(R_{m\alpha}(y)) - D^v(g' f^m(y)) + \\ & D^v(f^m g'(y)) - D^v g'(y) \end{aligned}$$

Moreover, we have:

$$|D^v g'(x) - D^v g'(y + m\alpha)| \leq |D^{u+1} g'|_0 |x - y - m\alpha|$$

By lemma 4.5, we also have:

$$|D^v g'(R_{m\alpha}(y)) - D^v(g' f^m(y))| \leq C \|g'\|_{u+1} \|f^m\|_u^C \|f^m - R_{m\alpha}\|_u$$

Finally, by lemma 4.4, we have:

$$|D^v(f^m g'(y)) - D^v(R_{m\alpha} g'(y))| \leq C \|f^m\|_{u+1} \|f^m - R_{m\alpha}\|_u \|g'\|_u^C$$

Since  $R_{m\alpha} g'(y) = g'(y) + m\alpha$ , and  $v \geq 1$ , then  $D^v(R_{m\alpha} g'(y)) = D^v(R_{m\alpha} g'(y))$ . Therefore, the same estimate holds for  $|D^v(f^m g'(y)) - D^v(g'(y))|$ .

By combining these estimations, we obtain:

$$\begin{aligned} |D^v g'(x) - D^v g'(y)| &\leq \|g'\|_{u+1} |x - y - m\alpha| + C \|g'\|_{u+1} \|f^m\|_u^C \|f^m - R_{m\alpha}\|_u + \\ & C \|f^m\|_{u+1} \|f^m - R_{m\alpha}\|_u \|g'\|_u^C \end{aligned}$$

Moreover, for any  $r \geq 0$ , any  $x, y \in \mathbb{R}$ , there is an integer  $m(x, y) \leq 2q_r$ , there are real numbers  $x', y'$  such that  $x' - x \in \mathbb{Z}$ ,  $y' - y \in \mathbb{Z}$  and such that  $|x' - y' - m(x, y)\alpha| \leq 1/q_r$ . Since  $v \geq 1$ , then  $|D^v g'(x) - D^v g'(y)| = |D^v g'(x') - D^v g'(y')|$ . We apply the former estimate with  $x'$  and  $y'$  and we get:

$$\max_{1 \leq v \leq u} \|D^v g' - D^v R_{g'(0)}\|_0 \leq \frac{A_{6,u+1} + 1}{q_r} + \max_{m \leq 2q_r} (A_{6,u+1} A_{3,u}^C(m) A_{5,u}(m) + A_{6,u}^C A_{3,u+1}(m) A_{5,u}(m))$$

If  $v = 0$ , we note that for any  $r \geq 0$ , any  $x \in \mathbb{R}$ , there is an integer  $m(x) \leq q_r$  and a real number  $x' \in \mathbb{R}$  such that  $x' - x \in \mathbb{Z}$ , and such that  $|x' - m(x)\alpha| \leq 1/q_r$ . Moreover, we have:  $g'(x) - R_{g'(0)}(x) = g'(x') - R_{g'(0)}(x')$ , and

$$g'(x') - R_{g'(0)}(x') = g'(x') - g'(m\alpha) + g'(m\alpha) - g' f^m(0) + f^m g'(0) - R_{m\alpha}(g'(0)) + R_{g'(0)}(m\alpha) - R_{g'(0)}(x')$$

Hence estimate (14).

For  $A_{8,u}$ , estimate (15) follows immediately from lemma 4.4.

For  $A_{9,u}$ , let  $x \in \mathbb{R}$ . Let  $0 \leq v \leq u$ . By the Faa-di-Bruno formula:

$$\begin{aligned} D^v (hg'h^{-1})(x) - D^v (hR_{g'(0)}h^{-1})(x) &= \\ & \sum_{j=0}^v D^j h(g'h^{-1}(x)) B_{v,j} (D(g'h^{-1})(x), \dots, D^{v-j+1}(g'h^{-1}(x))) - \\ & D^j h(g'h^{-1}(x)) B_{v,j} (D(R_{g'(0)}h^{-1})(x), \dots, D^{v-j+1}(R_{g'(0)}h^{-1}(x))) \\ &= \sum_{j=0}^v D^j h(g'h^{-1}(x)) \\ & \left[ B_{v,j} (D(g'h^{-1})(x), \dots, D^{v-j+1}(g'h^{-1}(x))) - B_{v,j} (D(R_{g'(0)}h^{-1})(x), \dots, D^{v-j+1}(R_{g'(0)}h^{-1}(x))) \right] - \\ & \left[ D^j h(R_{g'(0)}h^{-1}(x)) - D^j h(g'h^{-1}(x)) \right] B_{v,j} (D(R_{g'(0)}h^{-1})(x), \dots, D^{v-j+1}(R_{g'(0)}h^{-1}(x))) \end{aligned}$$

since  $\|h^{-1}\|_u \geq 1$ , then lemma 4.6 gives,

$$\left| B_{v,j} (D(g'h^{-1})(x), \dots, D^{v-j+1}(g'h^{-1}(x))) - B_{v,j} (D(R_{g'(0)}h^{-1})(x), \dots, D^{v-j+1}(R_{g'(0)}h^{-1}(x))) \right| \leq C \|g'h^{-1}\|_u^C \|g'h^{-1} - R_{g'(0)}h^{-1}\|_u$$

Since  $g'h^{-1} = h^{-1}g$  and  $\|h^{-1}g\|_u \leq C \|h^{-1}\|_u \|g\|_u^C$ , we get,

$$\left| D^v (hg'h^{-1})(x) - D^v (hR_{g'(0)}h^{-1})(x) \right| \leq C \|g\|_u^C \|h\|_u \|h^{-1}\|_u^C \|g'h^{-1} - R_{g'(0)}h^{-1}\|_u + C \|h\|_{u+1} \|g'h^{-1} - R_{g'(0)}h^{-1}\|_u \|h^{-1}\|_u^C$$

Hence estimate (16). This completes the proof of lemma 4.7.  $\square$

By combining these estimations, we obtain:

$$A_{9,u} \leq CA_{1,u+1}^C \|h\|_{u+1}^C \|g\|_{u+1}^C \left( \frac{1}{q_r} + \max_{m \leq 2q_r} (A_{3,u+1}^C(m) A_{5,u}(m)) \right)$$

$$A_{9,u} \leq C \|h\|_{u+1}^C \|f\|_u^C \|g\|_{u+1}^C \left( \frac{1}{q_r} + |\tilde{\alpha} - \alpha| \left( \frac{(C \|h\|_{u+1} \|f\|_{u+1})^{Cq_r}}{(\min Dh)^C} \right) \right)$$

Hence lemma 4.3. Notice the loss of one derivative for  $h$ .  $\square$



## References

- [1] V.I. Arnol'd. Small denominators I: Mappings of the circumference into itself. *American Mathematical Society Translations*, 46:213–284, 1965.
- [2] M. Benhenda. Estimates of the linearization of circle diffeomorphisms. <http://hal.archives-ouvertes.fr/hal-00628297>, October 2011.
- [3] A.D.D. Craik. Prehistory of Faà di Bruno's formula. *The American Mathematical Monthly*, 112(2):119–130, 2005.
- [4] K. Dajani and C. Kraaikamp. *Ergodic theory of numbers*, volume 29. The Mathematical Association of America, 2002.
- [5] B. Fayad and A. Katok. Constructions in elliptic dynamics. *Ergodic Theory and Dynamical Systems*, 24(5):1477–1520, 2004.
- [6] B. Fayad and K. Khanin. Smooth linearization of commuting circle diffeomorphisms. *Annals of Mathematics*, 170:961–980, 2009.
- [7] M.R. Herman. Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations. *Publications Mathématiques de L'IHÉS*, 49(1):5–233, 1979.
- [8] A. Katok and B. Hasselblatt. *Introduction to the modern theory of dynamical systems*, volume 54. Cambridge University Press, 1996.
- [9] Y. Katznelson. Sigma-finite invariant measures for smooth mappings of the circle. *Journal D'Analyse Mathématique*, 31(1):1–18, 1977.
- [10] J.C. Yoccoz. Petits diviseurs en dimension un. *Asterisque*, 231, 1995.