



Relatively compact Siegel disks with non-locally connected boundaries

Arnaud Chéritat

► To cite this version:

Arnaud Chéritat. Relatively compact Siegel disks with non-locally connected boundaries. *Mathematische Annalen*, 2011, 349 (3), pp.529-542. 10.1007/s00208-010-0527-1 . hal-00630976

HAL Id: hal-00630976

<https://hal.science/hal-00630976>

Submitted on 8 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Relatively compact Siegel disks with non-locally connected boundaries

Arnaud Chéritat¹²

Received: date / Revised version: date – © Springer-Verlag 2010

Abstract. We construct holomorphic maps f with a Siegel disk whose boundary is not locally connected (and is an indecomposable continuum), yet compactly contained in the domain of definition of the map. Our examples are injective and defined on a subset of \mathbb{C} .

1. Introduction

In [Han82], Handel constructed a C^∞ area preserving diffeomorphism of the plane that has a minimal set that is a pseudo circle (see section 3.5 for a definition of pseudo circles). In [Her86], Herman adapted the construction to produce a C^∞ diffeomorphism of the sphere that is conjugated to a rotation in the two complementary components of an invariant pseudo circle, and holomorphic in one of them. In the same spirit, in [PM97] Pérez Marco was able, using tube-log Riemann surfaces, to construct examples of injective holomorphic maps defined in a subset U of \mathbb{C} that have a Siegel disk compactly contained in U whose boundary is a C^∞ Jordan curve, which came as a surprise. Again the method is versatile and Kingshook Biswas used Pérez-Marco's construction to produce a set of interesting examples: [Bis05, Bis08]. Here we add an ingredient to this construction and get:

Theorem 1. *There exists a holomorphic map f defined in a simply connected open subset U of \mathbb{C} containing the origin, fixing 0 and having at 0 a Siegel disk Δ that is compactly contained in U and whose boundary is a pseudo circle.*

As Herman remarked, the construction is very flexible. The reader will find in the conclusion section more properties that these maps can be given. There, we also suggest possible other consequences of the method.

Remark 1. We ignore if the maps can be chosen entire (or entire meromorphic).

Centre National de la Recherche Scientifique.

Institut de Mathématiques de Toulouse, Université Paul Sabatier, 118 Route de Narbonne, 31062 Toulouse Cedex 9, France. e-mail: arnaud.cheritat@math.univ-toulouse.fr

1.1. The new ingredient

Compared to the previous constructions of Siegel disks and hedgehogs, the main new idea is to make use of Runge's theorem. Runge's theorem has been applied in a variety of situations to construct unexpected examples. We will make use of the following version:

Theorem (Runge, simply connected case). *For all holomorphic map f defined in a simply connected open set $U \subset \mathbb{C}$, there exists a sequence of polynomials that tends to f locally uniformly on U .¹*

It is easy to see that f can be approximated (in the sense above) by polynomials if and only if it can be approximated by entire maps. In our application, the fact that the approximating sequence consists in polynomials is not so important. We will only need entire maps.

1.2. Acknowledgements

The author would like to thank Kingshook Biswas with whom he had very useful discussions, Pérez-Marco for inventing his methods and explaining them to me, Yohann Genzmer for submitting me a stimulating math problem for which I found a complicated solution using iterated exponentials, Julien Duval for explaining me a simple solution to Genzmer's problem using Runge's theorem, thus bringing back this theorem to my memory and showing me its usefulness. The author would also like to thank the referee for useful comments.

2. Terminology

Saying that two holomorphic maps commute can mean different things. To be rigorous, we will say that “ f and g commute, domain included”, if the domain of definition of $f \circ g$ equals that of $g \circ f$ and if they take the same values on it. This is what $f \circ g = g \circ f$ is supposed to mean. Similarly we will say that “ $f \circ g = h \circ i$, domain included” if the domain of definition of $f \circ g$ equals that of $h \circ i$ and if they take the same values on it. We will say that “ f and g commute on U ” if U is included in the domain of $f \circ g$ and in the domain of $g \circ f$ and if both compositions take the same value on U . There is another natural notion that we will not use: “ f and g commute, wherever defined”, if $f \circ g(z) = g \circ f(z)$ for all z for which both hand sides are defined.

We will denote the domain of definition of a map f by $\text{dom}(f)$.

¹ a.k.a. “on every compact subset” of U

3. The construction

3.1. Working in the universal cover of \mathbb{C}^*

The map f will be the limit of a sequence of compositions $f_n = g_n \circ \cdots \circ g_1$ of maps g_i fixing the origin. However we prefer to work at the level of lifted coordinates: let

$$E(z) = \exp(2i\pi z).$$

The map E is a universal cover from \mathbb{C} to \mathbb{C}^* , with deck transformations $(\mathbb{Z}, +)$ and induces an isomorphism between the cylinder \mathbb{C}/\mathbb{Z} and \mathbb{C}^* . This map can be used to extend the Riemann surface \mathbb{C}/\mathbb{Z} at its upper end (we will not use the lower end in this article): denote $+i\infty$ the added point, corresponding to $0 \in \mathbb{C} \subset \mathbb{C}^*$.

Let T_v denote the translation by v : A holomorphic map F that commutes with T_1 , domain included, and such that

$$F(z) = z + t + o(1)$$

for some $t \in \mathbb{R}$ as $\text{Im}(z)$ tends² to $+\infty$, “projects” to a holomorphic map f with an erasable singularity at the origin, i.e. there exists f holomorphic defined on $\{0\} \cup E(\text{domain of } f)$, with $E \circ F = f \circ E$, $f(0) = 0$, $f'(0) = e^{2\pi it} \neq 0$, and $f(z) = 0 \implies z = 0$.

Conversely, a holomorphic map f defined in an open subset O of \mathbb{C} containing the origin, with $f(0) = 0$, $f'(0) \neq 0$, and $f(z) = 0 \implies z = 0$, will have a (non unique) lift $F : E^{-1}(O) \rightarrow \mathbb{C}$ that commutes with T_1 , domain included, with $E \circ F = f \circ E$ and $F(z) - z$ has a limit when $\text{Im}(z) \rightarrow +\infty$.

3.2. Presentation of the actors

Recall that T_v denotes the translation by v :

$$T_v(z) = z + v.$$

Let

$$\mathbb{H}_h = \{z \in \mathbb{C} \mid \text{Im}(z) > h\}, \quad \mathbb{H} = \mathbb{H}_0$$

that we will call *upper half planes*. Let $\mathbf{1}$ be the constant vector field in \mathbb{C} of expression $\mathbf{1}(z) = 1$.

Assume that maps \mathcal{R}_k are given for $k = 1, \dots, n$, satisfying the following conditions:

- \mathcal{R}_k is entire

² The filter understood by “ $\text{Im}(z) \longrightarrow +\infty$ ” is the one generated by all the half planes \mathbb{H}_h for $h \in \mathbb{R}$. We also imply that the domain of F contains such a half-plane.

- there exists $q_k \in \mathbb{N}^*$ such that $\mathcal{R}_k \circ T_1 = T_{q_k} \circ \mathcal{R}_k$
- $\mathcal{R}_k(z)$ is injective on some upper half-plane. Given the above condition, it is equivalent to “ $\mathcal{R}_k(z) = q_k z + c + o(1)$ as $\text{Im}(z) \rightarrow +\infty$ ”
- $\forall z \in \mathbb{C}, \mathcal{R}'_k(z) \neq 0$

We call these maps *renormalizations*. They will be inductively defined later in this article.

Let

- $X_n = \mathcal{R}_1^* \cdots \mathcal{R}_n^* \mathbf{1}$, i.e. the pull back of the vector field $\mathbf{1}$ by the composition $\mathcal{R}_n \circ \cdots \circ \mathcal{R}_1$
- G_n , the time 1 of the flow of the vector field X_n
- $F_n = G_n \circ \cdots \circ G_1$

Then X_n is a non-vanishing entire vector field. The maps G_n and F_n are holomorphic but not necessarily defined on all of \mathbb{C} .

These objects “project” to \mathbb{C}/\mathbb{Z} in the following sense:

- $T_1^* X_n = X_n$, i.e. X_n is T_1 -invariant
- G_n and F_n commute with T_1 , domain included
- recall that $\mathcal{R}_n \circ T_1 = T_{q_n} \circ \mathcal{R}_n$

In particular, they can be thought of objects living on \mathbb{C}/\mathbb{Z} .

Also, they are defined in a neighborhood³ of $+i\infty$, and we have the following expansions as $\text{Im}(z) \rightarrow +\infty$:

- $\mathcal{R}_n(z) = q_n z + c + o(1)$
- $X_n(z) = \frac{1}{q_1 \cdots q_n} + o(1)$
- $G_n(z) = z + \frac{1}{q_1 \cdots q_n} + o(1)$
- $F_n(z) = z + \sum_{k=1}^n \frac{1}{q_1 \cdots q_k} + o(1)$

Since each G_n , and thus F_n commute with T_1 , domains included, the map F_n projects by E to a map f_n , i.e. there exists a holomorphic map $f_n : \{0\} \cup E(\text{dom}(F_n)) \rightarrow \mathbb{C}$ such that $E \circ F_n = f_n \circ E$, domain included, and f_n fixes the origin.

Let Dom be the set of points that have a neighborhood on which all F_n are defined and converge uniformly. It is an open set. It might be empty. Let F be the map defined on Dom as the limit of the F_n . Then F is holomorphic and commutes with T_1 . The following lemma is elementary so we will not explain its proof: it is stated for reference.

³ i.e. on a set containing a half plane \mathbb{H}_h for some h that may depend on n

Lemma 1. *Assume that Dom contains an upper half plane. Define $dom = E(Dom) \cup \{0\}$. Then dom is an open subset of \mathbb{C} and f_n converges locally uniformly on dom to a holomorphic map f defined on dom and satisfying $f \circ E = E \circ F$, domain included.*

The set of domains containing an upper half plane and on which \mathcal{R}_n is a bijection to an upper half plane or to \mathbb{C} , is non-empty and has a greatest element V_n for inclusion. Let $U_0 = \mathbb{C}$ and for $n \geq 1$ let U_n be the set of points $z \in V_1$ such that for all $k < n$, $\mathcal{R}_k \circ \dots \circ \mathcal{R}_1(z) \in V_{k+1}$. Then $(U_n)_{n \in \mathbb{N}}$ forms a decreasing sequence for inclusion, of T_1 -invariant simply connected open subsets U_n of \mathbb{C} containing upper half planes, on which $\mathcal{R}_n \circ \dots \circ \mathcal{R}_1$ is an isomorphism to an upper half plane or to \mathbb{C} and on which $\mathcal{R}_k \circ \dots \circ \mathcal{R}_1$ is injective for all $k \leq n$. Let $\tilde{U}_n = E(U_n) \cup \{0\}$. It is a connected and simply connected open subset of \mathbb{C} . Note that each f_n is conjugated to a rotation in a neighborhood of the origin. Indeed, let $\theta_n = \sum_{k=1}^n 1/(q_1 \dots q_k)$. The map $z \mapsto \frac{1}{q_n} \mathcal{R}_n \circ \dots \circ \mathcal{R}_1(z)$ is a conjugacy from F_n on U_n to T_{θ_n} on an upper half plane or on \mathbb{C} , and commutes with T_1 on U_n , domain included. Therefore it projects by E to a conjugacy on \tilde{U}_n to the rotation of angle $2\pi\theta_n$ on a disk or on \mathbb{C} .

We will use the following lemma, applied to well chosen subsets \tilde{D}_n of \tilde{U}_n :

Lemma 2. *Assume f_n is a sequence of holomorphic maps defined on open subsets of \mathbb{C} containing the origin. Assume f_n fixes 0 and that $f'_n(0) = e^{i2\pi\theta_n}$ with $\theta_n \in \mathbb{R}$ and $\theta_n \rightarrow \theta \in \mathbb{R}$ as $n \rightarrow +\infty$. Assume \tilde{D}_n is a sequence of simply connected open subsets of the domain of f_n and containing 0, such that f_n is analytically conjugated on \tilde{D}_n to the rotation of angle $2\pi\theta_n$ on a disk or on \mathbb{C} . Assume that there exists a limit \tilde{D} to the sequence \tilde{D}_n in the sense of Caratheodory. Then f_n tends on \tilde{D} locally uniformly to a map f that is analytically conjugated on \tilde{D} to the rotation of angle θ on a disk or on \mathbb{C} .*

Proof. Let $R_\alpha(z) = e^{i2\pi\alpha}z$. The conjugacy ϕ_n on \tilde{D}_n can always be chosen so that $\phi'_n(0) = 1$. Then a theorem of Caratheodory says that ϕ_n converges locally uniformly on \tilde{D} to the unique conformal map ϕ from \tilde{D} to a disk or \mathbb{C} such that $\phi(0) = 0$ and $\phi'(0) = 1$. The reciprocal ϕ_n^{-1} is locally uniformly convergent to ϕ^{-1} . From $f_n = \phi_n^{-1} \circ R_{\theta_n} \circ \phi_n$ we deduce that f_n tends to $\phi^{-1} \circ R_\theta \circ \phi$ locally uniformly.

Depending on authors, the definition of Siegel disks may differ slightly. For us, it is the maximal domain of conjugacy to an irrational rotation. In the lemma above, if θ is irrational, then \tilde{D} is an invariant subset of the Siegel disk of f : could the Siegel disk of f be bigger? It necessarily requires the boundary of \tilde{D} to be an analytic Jordan curve⁴, compactly contained in dom . In fact it is equivalent. So if we get a boundary that is not a Jordan curve or not an analytic

⁴ this claim uses the fact that θ is irrational

one, then we know that \tilde{D} equals the Siegel disk of f . (This implies that \tilde{U}_n has the same Caratheodory limit as \tilde{D}_n .)

How to build examples with $\tilde{D} \subset \subset \text{dom}$? First we will ensure that Dom contains \mathbb{H}_{-h} for some $h > 0$. Second we will ensure that \tilde{D} is contained in \mathbb{H} .

3.3. Ensuring convergence on big domains

Here we give a slightly informal description, and a rigorous lemma that will be used in the more formal construction described in section 3.6.

It will be handy to use a notation for the set of maps satisfying the conditions we required on \mathcal{R}_n in the previous section. So let \mathcal{H}_1 denote the set of maps \mathcal{R} such that:

- \mathcal{R} is entire
- $\mathcal{R} \circ T_1 = T_1 \circ \mathcal{R}$
- $\mathcal{R}(z)$ is injective on some upper half-plane
- $\forall z \in \mathbb{C}, \mathcal{R}'(z) \neq 0$

So that $\mathcal{R}_k \in q_k \mathcal{H}_1$, i.e. \mathcal{R}_k is the product of a member of \mathcal{H}_1 with the scalar q_k .

For this, we will inductively define the maps \mathcal{R}_n using the following (elementary) lemma:

Lemma 3. *Fix $n \in \mathbb{N}$. Assume we are given $\mathcal{R}_k \in q_k \mathcal{H}_1$ for $k = 1, \dots, n-1$ and some $B_n \in \mathcal{H}_1$. Let $\mathcal{R}_n = q B_n$ for some $q \in \mathbb{N}^*$. The corresponding map f_n depends on q but f_{n-1} does not. Then, as $q \rightarrow +\infty$, f_n tends to f_{n-1} in the following sense: every compact subset K of $\text{dom}(f_{n-1})$ is eventually contained in $\text{dom}(f_n)$ and $f_n \rightarrow f_{n-1}$ uniformly on K . Note however, that the domain U_n defined in section 3.2 is independent of q .*

Proof. Indeed, $\mathcal{R}_n^* \mathbf{1} = (q B_n)^* \mathbf{1} = \frac{1}{q} B_n^* \mathbf{1}$. So X_n is the product of the scalar $\frac{1}{q}$ with an entire vector field that does not depend on q . The map G_n is the time $1/q$ of the flow of the latter vector field. The rest follows easily.

Fix some number $h > 0$. We will inductively define the sequence \mathcal{R}_n by first choosing B_1 , then q_1 , then B_2 , then q_2 , etc... We will be careful to choose q_k big enough so that the domain of definition of f_n contains the upper half plane $\mathbb{H}_{-h-\frac{1}{n}}$ and $|f_{n-1} - f_n| < 1/2^n$ on \mathbb{H}_{-h} . This is always possible by the lemma above. We can also choose q_n to ensure that the sum θ is convergent⁵ and has an irrational value: a sufficient condition for this is that q_n tends to $+\infty$. Then the half plane \mathbb{H}_{-h} is contained in the domain of definition Dom of F .

⁵ It already follows from the convergence of f_n , though.

It shall be stressed out that in the construction, the sequence (B_n) will not be fixed independently of the sequence (q_n) : the choice of B_{n+1} will depend on the choice of all the previous objects $B_1, q_1, B_2, q_2, \dots, B_n, q_n$.

3.4. Ensuring compactly contained bad boundaries

As in the previous section we give here a slightly informal description, and a rigorous lemma that will be used in the more formal construction described in section 3.6.

The following claim is where we use **Runge's theorem**:

Lemma 4. *Assume ϕ is a conformal bijection from \mathbb{H}/\mathbb{Z} to a domain $W \subset \mathbb{C}/\mathbb{Z}$ that extends to a conformal bijection fixing the upper end. Then there exists a sequence of holomorphic maps $\psi_n : \mathbb{C}/\mathbb{Z} \rightarrow \mathbb{C}/\mathbb{Z}$ such that:*

- ψ'_n does not vanish,
- $\psi_n(z) - z$ has a limit when $\text{Im}(z)$ tends to $+\infty$,
- as n tends to $+\infty$, ψ_n tends uniformly to ϕ^{-1} on every compact subset of W and in a neighborhood of the upper end of \mathbb{C}/\mathbb{Z} .

Proof. Conjugate the situation by E : let $\widetilde{W} = E(W) \cap \{0\}$ and $\Psi : \widetilde{W} \rightarrow \mathbb{D}$ be the conjugate of ϕ^{-1} , extended by setting $\Psi(0) = 0$, where \mathbb{D} denotes the unit disk. It is a conformal isomorphism. It can therefore be written as $\Psi(z) = ze^{v(z)}$ for some holomorphic map $v : \widetilde{W} \rightarrow \mathbb{C}$. As a conformal isomorphism its derivative does not vanish, hence $1 + zv'(z) \neq 0$. Thus there exists a map $u : \widetilde{W} \rightarrow \mathbb{C}$ such that $1 + zv'(z) = e^{u(z)}$. Since $e^{u(0)} = 1$, we can choose u such that $u(0) = 0$. So $u(z) = zw(z)$ for some holomorphic $w : \widetilde{W} \rightarrow \mathbb{C}$. Now apply Runge's theorem: there exists a sequence of entire maps w_n (even polynomials) that converge locally uniformly to w on \widetilde{W} . There exists an entire map v_n (unique) taking the same value as v at the origin and such that $1 + zv'_n = e^{zw_n}$. Indeed $(e^{zw_n} - 1)/z$ has a removable singularity at the origin, so it extends to an entire map, of which v_n is just the appropriate primitive. The maps $\psi_n(z) = z + v_n(e^z)$ satisfy the required conditions.

Now we describe informally how we can use this lemma to choose the functions B_n to get interesting examples. A more formal, but more specific, description is done in section 3.6.

Let $D_0 = \mathbb{H}/\mathbb{Z}$. Let W_1 be any open simply connected strict subset of $\mathbb{H}/\mathbb{Z} \cup \{+i\infty\}$ containing the upper end $+i\infty$. Let ϕ be any conformal map from $\mathbb{H}/\mathbb{Z} \cup \{+i\infty\}$ to W_1 that fixes the upper end. Let $\varepsilon > 0$ and let the Jordan curve J_1 be the image by ϕ of the Jordan curve of equation “ $\text{Im}(z) = \varepsilon$ ” in \mathbb{H}/\mathbb{Z} . If the boundary of W_1 is a convoluted Jordan curve (or something more complicated than a Jordan curve) and ε is small then J_1 will also be convoluted.

Apply lemma 4 to $W = W_1 - \{+i\infty\}$. By the uniform convergence of ψ_n to ϕ^{-1} on compact subsets of W and on a neighborhood of the upper end of \mathbb{C}/\mathbb{Z} , there is for n big enough a branch ϕ_n of ψ_n^{-1} defined for $\text{Im}(z) > \varepsilon/2$ and mapping the upper end to itself. For n big, the Jordan curve $J'_1 = \phi_n(\mathbb{R} + i\varepsilon)$, image by ϕ_n of the Jordan curve of equation “ $\text{Im}(z) = \varepsilon$ ” in \mathbb{H}/\mathbb{Z} , will be close to J_1 so it will also be convoluted.

So we will choose $B_1 = \psi_n(z) - i\varepsilon$ for some big value of n . Then we will chose q_1 as in section 3.4. Let D_1 be the domain bounded by J'_1 and containing the upper end. Among the integral lines of the vector field X_1 , there are the images of the horizontals by ϕ_n , which loop. This includes J'_1 and also a set of curves foliating the domain D_1 . The map $F_1 = G_1$, seen as acting on \mathbb{C}/\mathbb{Z} , is conjugate to a finite order⁶ rotation on D_1 . The renormalization \mathcal{R}_1 induces a conjugacy of the restriction of F_1 to D_1 to the translation T_1 on the half cylinder $\mathbb{H}/q_1\mathbb{Z}$.

The next step in the construction consists in choosing a new simply connected domain W_2 , contained in D_1 , containing the upper end, whose boundary is even more convoluted, yet invariant by F_1 , and very close to J'_1 . By the conjugacy \mathcal{R}_1 on D_1 it amounts to choosing a domain in $\mathbb{H}/q_1\mathbb{Z} \cup \{+i\infty\}$ that is invariant by T_1 , i.e. to choosing a simply connected domain W'_2 in $\mathbb{H}/\mathbb{Z} \cup \{+i\infty\}$ (by applying the cover $z + q_1\mathbb{Z} \in \mathbb{H}/q_1\mathbb{Z} \mapsto z + \mathbb{Z} \in \mathbb{H}/\mathbb{Z}$), and whose boundary is very close to \mathbb{R} .

Let $\phi : \mathbb{H}/\mathbb{Z} \cup \{+i\infty\} \rightarrow W'_2$ be a conformal map fixing the upper end. We choose a new value of ε , which gives a new Jordan curve $J_2 = D_1 \cap \mathcal{R}_1^{-1}(\phi(\mathbb{R} + i\varepsilon))$ close to the boundary of W_2 . We apply lemma 4 again and get a new sequence of entire maps ψ_n tending to ϕ^{-1} on W'_2 . We set $B_2 = \psi_n(z) - i\varepsilon$ for n big enough so that the curve $J'_2 = D_1 \cap \mathcal{R}_1^{-1}(\phi_n(\mathbb{R} + i\varepsilon))$ is close to J_2 . We choose q_2 as in section 3.4. The domain D_2 bounded by J'_2 is then invariant by G_1 and by the flow of the vector field X_2 . The maps G_1 and G_2 , will be thus conjugated by $\mathcal{R}_2 \circ \mathcal{R}_1$ on D_2 to the translations by respectively q_2 and 1 on $\mathbb{H}/q_1q_2\mathbb{Z}$.

And so on. . . We get a decreasing sequence of domains $D_n \subset \mathbb{H}/\mathbb{Z}$ on which $\mathcal{R}_n \circ \dots \circ \mathcal{R}_1$ is a conformal bijection to $\mathbb{H}/(q_1 \dots q_n)\mathbb{Z}$ and conjugating all the maps G_k , for $k \leq n$, to integer translations, and thus the map F_n too. Their boundaries form a sequence of Jordan curves extremely convoluted and close to each other. The construction has enough flexibility to allow for $D = \bigcap_n D_n$ to have a non locally connected boundary, as we will prove in the following sections (basically by the same method as Handel in [Han82]).

⁶ Recall that these maps are not entire. Some iterate will be the identity on the component of the domain for the iterate that contains D_1 .

3.5. Pseudo circles

Pseudo circles are amazing topological objects. The interested⁷ reader may look at [Bin51] (Example 2 p.48), [Fea70], [Rog70], [KG09]. It was chosen by Handel in [Han82] as a good example of extreme pathology that a minimal set can have even for an area preserving smooth diffeomorphism of the sphere. Herman did the same choice in [Her86] for his construction. So we will continue this tradition.

For our purpose, it will be enough to know the following:

- A pseudo circle is compact, connected, separates the plane into two components, but it is not locally connected thus it is not a Jordan curve.
- It is equal to the boundary of both components (in particular it has empty interior).
- Any set K obtained by the following procedure is a pseudo circle.

PROCEDURE: A *circular chain* in a topological space X is a sequence $e = (e_i)$ of open subsets of X , called *links*, indexed by $\mathbb{Z}/m\mathbb{Z}$ for some $m \geq 4$, such that $e_i \cap e_j \neq \emptyset$ if and only if i, j are adjacent or equal.⁸ For each $n \geq 1$, let Q_n be a circular chain in the plane⁹ such that:

- the diameters of the links of Q_n are finite for all n and their supremum for a given n tends to 0 as $n \rightarrow +\infty$
- Q_{n+1} is a refinement of Q_n : each link of the former is contained in a link of the latter
- Q_{n+1} is *crookedly embedded* in Q_n , which means the following: assume that $Q_n = (d_i)$ has m links, and that $Q_{n+1} = (e_i)$ has m' links. It will be convenient to let $i \in \mathbb{Z}$ and let d_i denote $d_{\bar{i}}$ where \bar{i} is the class of i modulo m . Similarly, for $i \in \mathbb{Z}$ let e_i denote $e_{\bar{i}}$ for the residue class modulo m' . To be *crookedly embedded* means that there exists a map $f : \mathbb{Z} \rightarrow \mathbb{Z}$ such that
 1. $\forall i \in \mathbb{Z}$, the closure of e_i is contained in $d_{f(i)}$
 2. $\forall i \in \mathbb{Z}$, $f(i+1) - f(i) \in \{-1, 0, 1\}$ (¹⁰)
 3. $\forall i \in \mathbb{Z}$, $f(i+m') = f(i) + m$ (¹¹)
 4. for all i, j with $f(i) + 2 < f(j) < f(i) + m$, there exists i' and j' with $f(i') = f(j) - 1$, $f(j') = f(i) + 1$ and either $i < i' < j' < j$ or $i > i' > j' > j$.

⁷ Let us mention the following facts: There exists a circularly chainable and hereditarily indecomposable planar continuum. For the definition of all these terms, the reader is referred to the introduction of [Rog93]. It was proved to be unique up to homeomorphism (but it is not homogeneous). Such an object is called a pseudo circle.

⁸ This is probably inspired from Čech cohomology: we want the *nerve* of the open cover to be a circle. So for chains of length 3 one would also require $e_{\bar{0}} \cap e_{\bar{1}} \cap e_{\bar{2}} = \emptyset$.

⁹ it does not matter whether the e_i are connected or not

¹⁰ By condition 1 it is already the case modulo m because Q_n and Q_{n+1} are circular chains.

¹¹ It translates the idea of Q_{n+1} having winding number 1 in Q_n .

Then define $K = \bigcap_{n \geq 1} Q_n$: it is a pseudo circle. END OF DESCRIPTION

Let us say that a Jordan curve, parameterized by $\iota(t)$, $t \in \mathbb{R}/\mathbb{Z}$, is *stably crooked* in a circular chain $Q_n = (e_i)$ with $i \in \mathbb{Z}/m\mathbb{Z}$, whenever:

- the curve is contained in the union of the links of the chain,
- it has winding number 1 in Q_n , oriented by ι ,⁽¹²⁾
- $\forall t_1, t_2 \in \mathbb{R}$ with $t_1 < t_2 < t_1 + 1$ and $\forall u, v \in \mathbb{Z}$, if $\iota(t_1) \in \bar{e}_u$ and $\iota(t_2) \in \bar{e}_v$ and if $u + 2 < v < u + m$ then there exists t'_1 and t'_2 with either $t_1 < t'_1 < t'_2 < t_2$ or $t_1 > t'_1 > t'_2 > t_2$ such that $\iota(t'_1) \in e_{v-1}$ and $\iota(t'_2) \in e_{u+1}$.

It is independent of the choice of parameterization of the Jordan curve. The subtlety of taking closures for \bar{e}_u and \bar{e}_v but not e_{u+1} and e_{v-1} is to ensure the following: *if a Jordan curve is stably crooked in Q_n then any nearby Jordan curve is*. By nearby we mean a Jordan curve which is parameterized by a map that is close to ι .

3.6. The core of the construction

Following Herman (with different conventions), for $m \geq 4$ consider the circular chain C_m of length m consisting of the rectangular links $e_i =]i - 1/4, i + 5/4[\times]0, 1[$ in the cylinder $\mathbb{C}/m\mathbb{Z}$. Then

- *There exists a Jordan curve in $\mathbb{C}/m\mathbb{Z}$ that is stably crooked in C_m and invariant by T_1 .* The proof is in [Han82] but let us mention that it is quite entertaining to work it out on one's own. Figure 1 gives examples of solutions for $m = 5, 6$ and 7 .
- *Fix such a Jordan curve J . Consider a conformal map ϕ from $\mathbb{H}/m\mathbb{Z} \cup \{+i\infty\}$ to the component of $\mathbb{C}/m\mathbb{Z} \cup \{+i\infty\} - J$ that contains the upper end, and that fixes the upper end. There exists such maps.¹³ *There exists ε such that the Jordan curve J_ε , image by ϕ of the curve of equation “ $\text{Im}(z) = \varepsilon$ ”, is stably crooked in C_m .* It follows from the definition of stably crooked and*

¹² The winding number of an oriented closed curve ι in a circular chain $Q = (d_j)$ of length $m \geq 4$ could be defined in terms of first Čech cohomology groups with coefficients in \mathbb{Z} by a diagram looking like this: $\mathbb{Z} \approx \check{H}^1(Q, \mathbb{Z}) \mapsto \check{H}^1(\bigcup_j d_j, \mathbb{Z}) \mapsto \check{H}^1(S^1, \mathbb{Z}) \approx \mathbb{Z}$, but this is sophisticated. It amounts to doing the following: cut the curve into a finite number of small consecutive pieces $e_{\bar{i}} = \iota([t_{\bar{i}}, t_{\bar{i}+1}])$ for some sequence $t_{\bar{i}}$ indexed by $\mathbb{Z}/p\mathbb{Z}$ for some $p > 0$. If they are small enough, they will all be contained in some $d_{\bar{j}}$. There is at most two possible values of \bar{j} . Choose anyone and call it $g(\bar{i})$. Since Q_n is a chain, $g(\bar{i} + 1)$ and $g(\bar{i})$ must differ of at most one in $\mathbb{Z}/m\mathbb{Z}$. There exists a lift $f : \mathbb{Z} \rightarrow \mathbb{Z}$ of g such that $f(i + 1)$ and $f(i)$ differ of at most one for all i . Then there exists $k \in \mathbb{Z}$ such that $f(i + p) = f(i) + km$ for all i . This k is the winding number. It is independent of all choices and of the oriented parameterization ι , and invariant by small perturbations of ι .

¹³ Apply the Riemann mapping theorem to the image of the situation by E .

from the following theorem of Caratheodory: ϕ extends continuously to the closure of $\mathbb{H}/m\mathbb{Z}$ and the extension is on $\mathbb{R}/m\mathbb{Z}$ a parameterization of the Jordan curve J .

- Let W be the preimage by the natural projection $\mathbb{C} \rightarrow \mathbb{C}/m\mathbb{Z}$ of the image of ϕ . Then W is connected and ϕ lifts to a conformal bijection, also denoted by ϕ , from \mathbb{H} to W . Since W is T_1 -invariant, ϕ commutes with T_1 , domain included. So ϕ induces an isomorphism from \mathbb{H}/\mathbb{Z} to W/\mathbb{Z} that we will also denote by ϕ .
- Now apply lemma 4 to ϕ and W/\mathbb{Z} . We get a sequence of holomorphic maps $\psi_n : \mathbb{C}/\mathbb{Z} \rightarrow \mathbb{C}/\mathbb{Z}$ such that:
 - ψ'_n does not vanish,
 - $\psi_n(z) - z$ has a limit when $\text{Im}(z)$ tends to $+\infty$,
 - as n tends to $+\infty$, ψ_n tends uniformly to ϕ^{-1} on every compact subset of W and in a neighborhood of the upper end of \mathbb{C}/\mathbb{Z} .
- By this uniform convergence, there is for n big enough a branch ϕ_n of ψ_n^{-1} defined for $\text{Im}(z) > \varepsilon/2$ and mapping the upper end to itself. *There exists $M \geq 4$ and $N \in \mathbb{N}$ such that for all $m' \geq M$ and $n \geq N$, the image of the chain $C_{m'}$ by $z \mapsto \phi_n(\frac{m}{m'}z + i\varepsilon)$ is a circular chain that is crookedly embedded in C_m .*

Proof. Denote $(e_{\bar{i}})$ this circular chain and $C_m = (d_{\bar{j}})$, with $\bar{i} \in \mathbb{Z}/m'\mathbb{Z}$ and $\bar{j} \in \mathbb{Z}/m\mathbb{Z}$. First there is M, N such that for $m' \geq M$ and $n \geq N$, for all link $e_{\bar{i}}$, the set of \bar{j} such that “ $e_{\bar{i}} \subset d_{\bar{j}}$ ” is not empty, for otherwise, taking subsequences, there would be a point of J_ε that is a limit of points not contained in any link, but links are open and J_ε is contained in their union, which leads to a contradiction. Since C_m is a chain, this set of \bar{j} is at most two consecutive integers modulo m . Let $g(\bar{i})$ be any of them (the “smallest” for instance). Because $(e_{\bar{i}})$ and $(d_{\bar{j}})$ are circular chains, we have necessarily $g(\bar{i} + 1) - g(\bar{i}) \in \{-1, 0, 1\}$. There is thus a unique lift of f from \mathbb{Z} to \mathbb{Z} that satisfies $f(i + 1) - f(i) \in \{-1, 0, 1\}$ and $f(0) \in \{0, \dots, m - 1\}$. This lift necessarily satisfies $f(i + m') = f(i) + km$ for some k independent of i . This k is necessarily equal to the winding number in C_m of the curve $\phi_n(x + i\varepsilon)$ that runs along the lower edges of the “rectangles” defining $(e_{\bar{i}})$, which is equal to the winding number of J_ε in C_m , that is 1, so we get condition (3). If condition (4) were not satisfied for all m', n big enough, then taking subsequences would contradict the fact that J_ε is stably crooked in C_m . \square

Let us sum up what we will use of the above analysis. Setting $\psi(z) = \psi_N(z) - i\varepsilon$ for the value of N mentioned above, we get:

Lemma 5. *For all $m \geq 4$ there exists $M \geq 4$ and a holomorphic map ψ such that*

- ψ is entire,
- ψ' does not vanish,




Fig. 1. Examples of T_1 -invariant stably crooked Jordan curves in circular chains (unrolled) of length 5, 6 and 7. To make the figure more readable, we replaced the rectangles of the chain C_m by smoother regions.

- $\psi \circ T_1 = T_1 \circ \psi$,
- $\psi(z) - z$ has a limit when $\text{Im}(z) \rightarrow +\infty$,
- ψ has an inverse branch ϕ defined for $\text{Im}(z) > -\varepsilon/2$ for some $\varepsilon > 0$, with $\phi(z) - z$ having a limit as $\text{Im}(z) \rightarrow +\infty$, and $\phi \circ T_1 = T_1 \circ \phi$,
- for all $m' \geq M$, the image of the circular chain $C_{m'}$ by $z \mapsto \phi(\frac{m}{m'}z)$ is a circular chain crookedly embedded in C_m .

3.7. Putting it all together

We can now apply this to build our example. Fix $h > 0$. Let $B_n \in \mathcal{H}_1$, $q_n \in \mathbb{N}^*$ be defined by induction as follows:

Let $B_1 = \text{id}$ on \mathbb{C} , and q_1 be any integer ≥ 5 .

Assume that $B_1, q_1, \dots, B_n, q_n$ have been fixed such that the following holds for all $k \leq n$:

1. As in section 3.2, let $\mathcal{R}_k = q_k B_k$, let V_k be biggest domain containing an upper half plane and on which \mathcal{R}_k is a bijection to an upper half plane, let $U_0 = \mathbb{C}$ and let U_n be the set of points $z \in V_1$ such that for all $k < n$, $\mathcal{R}_k \circ \dots \circ \mathcal{R}_1(z) \in V_{k+1}$. We already saw that $\mathcal{R}_k \circ \dots \circ \mathcal{R}_1$ are bijections from U_k to upper half planes. We require these half-planes to contain $\mathbb{H}_{-\varepsilon}$ for some $\varepsilon > 0$ which may depend on k .
2. Let Q_k be the circular chain defined as the preimage in \mathbb{C}/\mathbb{Z} by the restriction of $\mathcal{R}_k \circ \dots \circ \mathcal{R}_1$ to U_k/\mathbb{Z} of the chain $C_m \subset \mathbb{H}/m\mathbb{Z}$ with $m = q_1 \dots q_k$. If $k > 1$, we require Q_k to be crookedly embedded in Q_{k-1} .
3. The supremum of the diameters of the links of Q_k is $\leq 1/k$.
4. Let F_k be defined as in section 3.2. We require the domain of definition of F_k to contain $\mathbb{H}_{-h-1/k}$.
5. If $k > 1$, the supremum on \mathbb{H}_{-h} of $|F_k - F_{k-1}|$ is $\leq 1/2^k$.
6. $q_k \geq k$.

This set of conditions is satisfied for $n = 1$: for (1) we have $\mathcal{R}_1(z) = 5z$, $\mathbb{C} = V_1 = U_1 = \mathcal{R}_1(U_1)$; (2) is empty; for (3) the diameter of the links C_m is $\sqrt{13}/2$ thus the diameter of the links of Q_1 is $\sqrt{13}/2q_1$ and here $m = q_1 \geq 5$; (4) $F_1 = T_{1/q_1}$ and is defined on \mathbb{C} ; (5) is empty; (6) $q_1 \geq 5 \geq 1$.

If it is satisfied for some n , then let us explain why it is possible to choose (B_{n+1}, q_{n+1}) such that it is satisfied for $n + 1$:

- First, we define B_{n+1} . Let $m = q_1 \dots q_n$. Let ψ, ϕ, M be given by lemma 5. Let $B_{n+1}(z) = \psi(z)$ and for any $q_{n+1} \in \mathbb{N}^*$ set $m' = q_{n+1}m$. We obtain at once (1), and (2) follows as soon as q_{n+1} is big enough so that $m' \geq M$.
- That (3) holds when q_{n+1} is big enough follows merely from the continuity of ϕ .
- That (4) and (5) hold when q_{n+1} is big enough follows from section 3.3.
- Point (6) needs no comment.

Then by (5) the sequence F_n tends uniformly on \mathbb{H}_{-h} to some holomorphic map F , so $\mathbb{H}_{-h} \subset \text{Dom}$ in the notations of section 3.2. Let f the map associated to F in lemma 1. By (6) the number $\theta = \sum_n 1/(q_1 \dots q_n)$ is irrational. By (2) and (3) the set $K = \bigcap_n Q_n$ is a pseudo circle. Let D_n be the preimage of \mathbb{H} by the restriction of $\mathcal{R}_k \circ \dots \circ \mathcal{R}_1$ to U_k and let $\tilde{D}_n = E(D_n) \cup \{0\}$.

By (1), f_n is conjugated to a (rational¹⁴) rotation on \tilde{D}_n (see the discussion in section 3.2). The sequence \tilde{D}_n is decreasing for inclusion, and tends in the sense of Caratheodory to the connected component \tilde{D} of the complement of the pseudo circle $E(K)$ that contains the origin. The boundary of \tilde{D} is equal to $E(K)$ (it is a property that pseudo circles share with Jordan curves: they are equal to the boundary of both components of their complement). By lemma 2, \tilde{D} is contained in the Siegel disk of f . But since its boundary is not a Jordan curve (it is a pseudo circle) and the rotation number is irrational, \tilde{D} is the Siegel disk. It is contained in the unit disk \mathbb{D} , because $\tilde{D}_1 = \mathbb{H}$ and \tilde{D}_n is decreasing. So the Siegel disk of f is compactly contained in the domain of definition of f , because the latter contains $e^{2\pi h}\mathbb{D}$.

4. Conclusion

With the above proof, we realize that the map f we constructed is injective, as a limit of injective holomorphic maps. The domain of definition can be taken to contain $R\mathbb{D}$ for R as big as wanted (while the pseudo circle is contained in \mathbb{D}). By injectivity and properties of univalent maps, f will be very close to a rotation on, say, $2\mathbb{D}$ when R gets big. The pseudo circle can be chosen to be very close to the unit circle. Alternatively, with a slight modification of the construction it can be chosen to span between distance ε and $1 - \varepsilon'$.

In [PM97], it is mentioned that with Pérez Marco's examples, one can get a whole uncountable group of commuting maps sharing the same Siegel disk. Here the same holds: by taking the u_n small enough we can ensure that for any infinite subset J of \mathbb{N} , the infinite composition of the G_k over $k \in J$ will converge on big domains, and they will all leave the Siegel disk of f invariant, so they will have the same Siegel disk since their rotation number remains irrational.

We believe that the new flexibility allowed by Runge's theorem allows to prove the following claims. But this has to be carefully checked.

Claim. Prove that there exists an injective holomorphic map f defined in a simply connected open subset U of \mathbb{C} containing the origin, fixing 0 and having at 0 a hedgehog of positive Lebesgue measure compactly contained in U .

Claim. Prove that there exists an injective holomorphic map f defined in a doubly connected open subset U of \mathbb{C} and a Jordan curve J with positive Lebesgue measure contained in U that is invariant by f , and carries an invariant line field.

We still believe in the following conjecture:

¹⁴ Some iterate of f_n is thus the identity on the component of its domain that contains U_n . Recall that f_n is not entire: when f_n is iterated, its domain decreases and may disconnect.

Conjecture. *The boundary of the Siegel disks of all polynomials are Jordan curves.*

It seems very likely that the construction requires gigantic values of q_k so that we probably always get very Liouvillian rotation numbers (in particular non Brjuno). We can always arrange so that it is the case. So Herman's question in [Her86] concerning which rotation numbers allow for non Jordan curve boundaries is still quite topical.

5. Addendum: Anosov and Katok

After submitting the present article, the author learned by discussion with colleagues that Herman's construction is an instance of a general construction by Anosov and Katok [AK70a], [AK70b]. The following description of the Anosov and Katok construction is adapted from [FK04] (with the area preserving assumption removed and h_n replaced by its inverse): consider a (real) manifold M together with a smooth action of the group $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ by diffeomorphisms $S_\alpha : M \rightarrow M$ with $S_{\alpha+\beta} = S_\alpha S_\beta$. Typically, imagine $M =$ a sphere and S_α are rotations of angle $2\pi\alpha$ around a common axis. Then one tries to construct $f : M \rightarrow M$ obtained as a limit:

$$f = \lim_{n \rightarrow \infty} f_n, \text{ where } f_n = H_n^{-1} \circ S_{\alpha_{n+1}} \circ H_n$$

with $\alpha_n \in \mathbb{Q}$ and

$$H_n = h_n \circ \dots \circ h_1,$$

where every h_n is a diffeomorphism of M that satisfies

$$h_n \circ S_{\alpha_n} = S_{\alpha_n} \circ h_n.$$

The sequence f_n converges but usually not the sequence H_n . The h_n and α_n are chosen inductively in the following order:

$$\alpha_1, h_1, \alpha_2, h_2, \alpha_3, \dots$$

The choice of h_n is very free (within the condition $h_n \circ S_{\alpha_n} = S_{\alpha_n} \circ h_n$), it does not need to be close to the identity, and usually won't. To ensure convergence of the sequence f_n , it is enough to choose each time α_{n+1} very close to α_n , because then f_n is close to f_{n-1} .

This can be rewritten as follows:

$$f_n = g_n \circ \dots \circ g_1$$

with

$$g_n = H_n^{-1} \circ S_{\alpha_{n+1} - \alpha_n} \circ H_n.$$

Note that g_n is the time $\alpha_{n+1} - \alpha_n$ of the flow of a vector field $(H_n)^*X$, the pull back by $H_n = h_n \circ \cdots \circ h_1$ of the vector field X generating the action S .

For comparison, the construction in the present article also takes the form

$$f_n = g_n \circ \cdots \circ g_1$$

but g_n is the time $\alpha_{n+1} - \alpha_n$ of the flow of a vector field $(H_n)^*X$ where X generates the group of rotations on \mathbb{C} and) $H_n = h_n \circ \cdots \circ h_1$ where the h_n are entire holomorphic maps of \mathbb{C} with non-vanishing derivative but are *not* injective: they are only semi-conjugacies.

References

- [AK70a] D. V. Anosov and A. B. Katok. New examples in smooth ergodic theory. Ergodic diffeomorphisms. *Trudy Moskov. Mat. Obšč.*, 23:3–36, 1970.
- [AK70b] D. V. Anosov and A. B. Katok. New examples of ergodic diffeomorphisms of smooth manifolds. *Uspehi Mat. Nauk*, 25(4 (154)):173–174, 1970.
- [Bin51] R. H. Bing. Concerning hereditarily indecomposable continua. *Pacific J. Math.*, 1:43–51, 1951.
- [Bis05] Kingshook Biswas. Smooth combs inside hedgehogs. *Discrete Contin. Dyn. Syst.*, 12(5):853–880, 2005.
- [Bis08] Kingshook Biswas. Hedgehogs of Hausdorff dimension one. *Ergodic Theory Dynam. Systems*, 28(6):1713–1727, 2008.
- [Fea70] Lawrence Fearnley. The pseudo-circle is unique. *Trans. Amer. Math. Soc.*, 149:45–64, 1970.
- [FK04] Bassam Fayad and Anatole Katok. Constructions in elliptic dynamics. *Ergodic Theory Dynam. Systems*, 24(5):1477–1520, 2004.
- [Han82] Michael Handel. A pathological area preserving C^∞ diffeomorphism of the plane. *Proc. Amer. Math. Soc.*, 86(1):163–168, 1982.
- [Her86] Michael-R. Herman. Construction of some curious diffeomorphisms of the Riemann sphere. *J. London Math. Soc. (2)*, 34(2):375–384, 1986.
- [KG09] Krystyna Kuperberg and Kevin Gammon. A short proof of nonhomogeneity of the pseudo-circle. *Proc. Amer. Math. Soc.*, 137(3):1149–1152, 2009.
- [PM97] Ricardo Pérez Marco. Siegel disks with smooth boundary. *Submitted*, 1997.
- [Rog70] James Ted Rogers, Jr. Pseudo-circles and universal circularly chainable continua. *Illinois J. Math.*, 14:222–237, 1970.
- [Rog93] James T. Rogers, Jr. Intrinsic rotations of simply connected regions and their boundaries. *Complex Variables Theory Appl.*, 23(1-2):17–23, 1993.